

## THE EXT CLASS OF AN APPROXIMATELY INNER AUTOMORPHISM

AKITAKA KISHIMOTO AND ALEX KUMJIAN

ABSTRACT. Let  $A$  be a simple unital  $AT$  algebra of real rank zero. It is shown below that the range of the natural map from the approximately inner automorphism group to  $KK(A, A)$  coincides with the kernel of the map  $KK(A, A) \rightarrow \bigoplus_{i=0}^1 \text{Hom}(K_i(A), K_i(A))$ .

### 1. INTRODUCTION AND PRELIMINARIES

1.1. An automorphism  $\alpha$  of a unital  $C^*$ -algebra  $A$  is said to be an approximately inner automorphism if there is a sequence of unitaries  $u_n \in A$  such that  $\alpha(a) = \lim \text{Ad } u_n(a)$  for all  $a \in A$ . It follows that the induced map on  $K_*(A)$  is the identity map; there is, however, an invariant of  $K$ -theoretical interest which can occur. Nontrivial extensions may arise in the six-term periodic sequence for the  $K$ -theory of the mapping torus. We show below that every extension does arise if  $A$  is a simple unital  $AT$  algebra of real rank zero. As an immediate corollary we obtain a stronger form of Elliott's classification theorem for such algebras: an invertible  $KK$  element that preserves positivity and the class of the unit lifts to an isomorphism.

1.2. Recall that a unital  $C^*$ -algebra  $A$  is said to be a unital  $AT$  algebra if it is expressible as the inductive limit of finite direct sums of algebras of the form  $M_n \otimes C(\mathbf{T})$  with unital embeddings. Let  $A$  be a unital  $AT$  algebra and let  $\alpha$  be an approximately inner automorphism of  $A$ . The mapping torus of  $\alpha$  is the  $C^*$ -algebra

$$M_\alpha = \{x \in C[0, 1] \otimes A : \alpha(x(0)) = x(1)\}.$$

It will be convenient to identify  $SA$ , the suspension of  $A$ , with the kernel of the map of evaluation at zero,  $e_0 : M_\alpha \rightarrow A$ . From the short exact sequence:

$$0 \rightarrow SA \rightarrow M_\alpha \xrightarrow{e_0} A \rightarrow 0$$

one obtains the usual six-term periodic exact sequence:

$$\begin{array}{ccccc} K_0(SA) & \rightarrow & K_0(M_\alpha) & \rightarrow & K_0(A) \\ & & \uparrow & & \downarrow \\ K_1(A) & \leftarrow & K_1(M_\alpha) & \leftarrow & K_1(SA) \end{array}$$

In identifying  $K_i(SA)$  with  $K_{i+1}(A)$  the index maps become  $\text{id} - \alpha_*$  (cf. [B, 10.4.1]); further, as  $\alpha$  is an approximately inner automorphism one has  $\alpha_* = \text{id}$  and the index

---

Received by the editors August 5, 1996.

1991 *Mathematics Subject Classification*. Primary 46L40; Secondary 46L80, 46L35.

maps are zero. Consequently, the six-term periodic exact sequence reduces to a pair of short exact sequences:

$$0 \rightarrow K_{i+1}(A) \rightarrow K_i(M_\alpha) \rightarrow K_i(A) \rightarrow 0$$

for  $i = 0, 1$ . Let  $\eta_i(\alpha)$  denote the class of this sequence in  $\text{Ext}(K_i(A), K_{i+1}(A))$ . Since  $A$  is an inductive limit of type I  $C^*$ -algebras,  $A$  is in the bootstrap class, and thus the universal coefficient theorem of Rosenberg and Schochet (see [RS, 1.17]) applies: that is, the following is exact:

$$\begin{aligned} 0 \rightarrow \bigoplus_{i=0}^1 \text{Ext}(K_i(A), K_{i+1}(A)) &\xrightarrow{\delta} KK(A, A) \\ \xrightarrow{\gamma} \bigoplus_{i=0}^1 \text{Hom}(K_i(A), K_i(A)) &\rightarrow 0. \end{aligned}$$

Rørdam observed (see [Rø, p. 435]) that  $\delta(\eta_0(\alpha), \eta_1(\alpha)) = 1 - [\alpha]$ , where  $[\alpha]$  denotes the class of  $\alpha$  in  $KK(A, A)$ . Since the product of any two elements in  $\ker \gamma$  is 0 ([RS, 7.10]), one has  $\eta_i(\alpha\beta) = \eta_i(\alpha) + \eta_i(\beta)$  for approximately inner automorphisms  $\alpha, \beta$ .

1.3. Since  $A$  is a unital  $AT$  algebra, both  $K_0(A)$  and  $K_1(A)$  are torsion free and so the extensions above are pure (i.e. locally trivial). Given a pair of countable abelian groups  $P, Q$ ; if  $Q = \varinjlim Q_n$  (with connecting maps  $f_n : Q_n \rightarrow Q_{n+1}$ ) one has the following short exact sequence for  $\text{Ext}(Q, P)$  (cf. [Ro]):

$$0 \rightarrow \varprojlim^1 \text{Hom}(Q_n, P) \rightarrow \text{Ext}(Q, P) \rightarrow \varprojlim \text{Ext}(Q_n, P) \rightarrow 0.$$

If  $Q$  is torsion free, the  $Q_n$  may be chosen to be isomorphic to  $\mathbf{Z}^{k_n}$ ; in this case  $\text{Ext}(Q_n, P) = 0$  and one has the isomorphism

$$\varprojlim^1 \text{Hom}(Q_n, P) \cong \text{Ext}(Q, P).$$

In showing that every Ext class arises from the  $KK$ -class of an approximately inner automorphism, it will be useful to have an explicit form for this isomorphism. Note that  $\varprojlim^1 \text{Hom}(Q_n, P)$  may be identified with the cokernel of the map,  $\prod_n \text{Hom}(Q_n, \tilde{P}) \rightarrow \prod_n \text{Hom}(Q_n, P)$ , where  $(g_n) \mapsto (g_n - g_{n+1}f_n)$ . Given  $(g_n)$ , we construct an extension

$$0 \rightarrow P \rightarrow E \rightarrow Q \rightarrow 0$$

as follows: set  $E_n = P \oplus Q_n$  and define  $h_n : E_n \rightarrow E_{n+1}$  by  $h_n(p, q) = (p + g_n(q), f_n(q))$ ; then  $E = \varinjlim E_n$  gives the desired extension. If  $P = \varinjlim P_n$  then, by passing to a suitable subsequence (and relabeling), one has  $E \cong \varinjlim P_n \oplus Q_n$ .

We will have occasion to consider inductive systems of short exact sequences; such an inductive system will consist of a sequence of short exact sequences:

$$0 \rightarrow P_n \rightarrow E_n \rightarrow Q_n \rightarrow 0$$

together with maps between them such that the following diagram commutes:

$$\begin{array}{ccccccccc} 0 & \rightarrow & P_n & \rightarrow & E_n & \rightarrow & Q_n & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & P_{n+1} & \rightarrow & E_{n+1} & \rightarrow & Q_{n+1} & \rightarrow & 0 \end{array}$$

Note that the limit of the inductive system is again a short exact sequence:

$$0 \rightarrow P \rightarrow E \rightarrow Q \rightarrow 0.$$

1.4. In §2 we compute the class of the two extensions arising from the  $K$ -theory of the mapping torus for certain approximately inner automorphisms of a unital  $\mathbf{AT}$  algebra (see Theorem 2.4). Here we do not assume that the  $C^*$ -algebra  $A$  is simple or has real rank zero. The unitality is not essential either if we define an automorphism to be approximately inner when it is so in the algebra  $A^+$  obtained by adjoining a unit to  $A$ , since  $A^+$  is again an  $\mathbf{AT}$  algebra (see [E, §1]). The maps that determine the extensions keep track of the  $K_1$ -class of  $u_{n+1}u_n^*$  for the various partial embeddings (this gives the maps  $K_0(A_n) \rightarrow K_1(A)$ ) as well as the Bott number for approximately commuting unitaries (see [Ex]): it may be assumed that  $u_{n+1}$  approximately commutes with the canonical central unitaries of  $A_n$  (this gives the maps  $K_1(A_n) \rightarrow K_0(A)$ ).

**Definition.** Given two unitary  $n \times n$  matrices  $U$  and  $V$  with  $\|VUV^*U^* - 1\| < 2$ , there is a selfadjoint matrix  $H$  with  $\|H\| < 1/2$  such that  $VUV^*U^* = e^{2\pi iH}$ . Then define the Bott number of the pair by  $B(U, V) = \text{Tr}(H)$ .

Since  $\det(VUV^*U^*) = 1$ , it follows that  $\text{Tr}(H) \in \mathbf{Z}$ . Note that  $B(U, V) = \omega(U, V)$ , where  $\omega(U, V)$  is the winding number of the loop (see [Ex, Lemma 3.1] and [EL]):

$$t \mapsto \det((1 - t)UV + tVU) = \det(UV) \det(1 - t + tVUV^*U^*).$$

Thus, the Bott number is invariant under homotopy of pairs of unitaries for which the norm of the commutator is less than 2. Moreover, if  $\|V_iUV_i^*U^* - 1\| < 1$  for  $i = 1, 2$ , then  $B(U, V_1V_2) = B(U, V_1) + B(U, V_2)$ .

1.5. We proceed to the main result, Theorem 3.1, in §3. Let  $A$  be a simple unital  $\mathbf{AT}$  algebra of real rank zero; we show for  $i = 0, 1$  that, given

$$\eta \in \text{Ext}(K_i(A), K_{i+1}(A))$$

there is an approximately inner automorphism  $\alpha$  for which  $\eta_i(\alpha) = \eta$  and  $\eta_{i+1}(\alpha) = 0$ . The proof is divided into the two cases  $i = 0, 1$  and proceeds through a sequence of lemmas; one must for example show that an extension is expressible as an inductive limit of groups of the form  $\mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n}$  in a way that makes all relevant diagrams commute.

1.6. With  $A$  as above,  $K_0(A)$  is a simple dimension group and  $K_1(A)$  is a (countable) torsion free group.  $K_i(A)$  is thus expressible as the inductive limit of a system

$$\mathbf{Z}^{k_1} \xrightarrow{\chi_1^i} \mathbf{Z}^{k_2} \xrightarrow{\chi_2^i} \mathbf{Z}^{k_3} \rightarrow \dots$$

with  $\chi_n^0(j, k) > 0$  for all  $j, k$  and  $n$ . We may assume that image of  $\mathbf{Z}^{k_1}$  contains the class of the unit  $[1_A] \in K_0(A)$ . We show below that, under mild assumptions on the  $\chi_n^i$ ,  $A$  is expressible as an inductive limit of certain maps  $\varphi_n : A_n \rightarrow A_{n+1}$ , where  $K_i(A_n) = \mathbf{Z}^{k_n}$  and  $(\varphi_n)_* = (\chi_n^0, \chi_n^1)$  and the partial embeddings of  $\varphi_n$  are given by the standard maps described below. These maps are closely related to those arising in the path model constructed by Deaconu in [D].

Let  $k_0, k_1$  be integers with  $k_0 \gg 0$ . Define  $\varphi_{k_0, k_1} : C(\mathbf{T}) \rightarrow M_{k_0} \otimes C(\mathbf{T})$  by

$$\varphi_{k_0, k_1}(u)(z) = \begin{pmatrix} 0 & 0 & \dots & z^{k_1} \\ 1 & 0 & \dots & 0 \\ & \ddots & \ddots & \\ 0 & \dots & 1 & 0 \end{pmatrix}$$

for  $k_1 \neq 0$ , and define  $\varphi_{k_0,0}$  by

$$\varphi_{k_0,0}(u)(z) = \begin{pmatrix} \varphi_{\lfloor k_0/2 \rfloor,1} & 0 \\ 0 & \varphi_{\lceil k_0/2 \rceil,-1} \end{pmatrix}.$$

Note that when  $K_i(C(\mathbf{T}))$  is identified with  $\mathbf{Z}$ ,  $(\varphi_{k_0,k_1})_*$  is given by multiplication by  $k_i$ . We will refer to  $\varphi_{k_0,k_1}$  as a standard embedding of type  $(k_0, k_1)$ . Note that under the assumption  $k_0 \gg 0$  the image of the unitary generator  $v$  of  $C(\mathbf{T})$  has small spectral variation in  $M_{k_0} \otimes C(\mathbf{T})$ ; indeed, if  $k_1 \neq 0$  the spectrum of  $\varphi_{k_0,k_1}(v)(z)$  consists of the  $k_0^{th}$  roots of  $z^{k_1}$ , where  $z \in \mathbf{T}$ . To avoid cumbersome notation we let  $\varphi_{k_0,k_1}$  also denote the map  $M_n \otimes C(\mathbf{T}) \rightarrow M_n \otimes M_{k_0} \otimes C(\mathbf{T})$  (where  $x \otimes a \mapsto x \otimes \varphi_{k_0,k_1}(a)$ ).

A unitary  $u \in M_n \otimes C(\mathbf{T})$  is said to be normal if there are integers  $p, q$  with  $p \neq 0, q > 0$  and  $q|n$  such that the eigenvalues of  $u_t$  (for  $t \in \mathbf{T} = \mathbf{R}/\mathbf{Z}$ ) are  $e^{2\pi i(pt+r)/q}$  for  $r = 0, 1, \dots, q - 1$ , and each eigenvalue occurs with multiplicity  $n/q$ . Observe that the spectral variation of  $u$ ,

$$\sup_{s,t \in \mathbf{T}} \text{dist}(\sigma(u_s), \sigma(u_t)),$$

is  $\pi/q$ , where  $\sigma(u)$  denotes the spectrum of the unitary  $u$  counted with multiplicity and distance is given by the arc length metric. Note that  $\varphi_{q,p}(v) \in M_q \otimes C(\mathbf{T})$  (where  $v$  is the unitary generator of  $C(\mathbf{T})$ ) is normal with parameters  $p, q$ .

**Fact.** Suppose  $v \in M_n \otimes C(\mathbf{T})$  is a normal unitary with parameters  $p, q$ , and

$$\varphi_{k_0,k_1} : M_n \otimes C(\mathbf{T}) \rightarrow M_n \otimes M_{k_0} \otimes C(\mathbf{T})$$

is a standard embedding with  $k_1 \neq 0$ . Then  $\varphi_{k_0,k_1}(v)$  is normal in  $M_n \otimes M_{k_0} \otimes C(\mathbf{T})$  with parameters  $pk_1/l, qk_0/l$ , where  $l = \text{gcd}(p, k_0)$ . Hence if  $k_0 \gg |p|$ , then  $\varphi_{k_0,k_1}(v)$  has small spectral variation.

We proceed to the construction of  $A$  using standard embeddings. By passing to a subsequence if necessary we may assume that  $M(\chi_{n+1}^0) \geq \max(2L(\chi_n^1), 4)$ , where

$$M(\chi) = \min_{i,j} \chi_{i,j} \text{ and } L(\chi) = \max_{i,j} |\chi_{i,j}|$$

for a matrix  $\chi$ . Choose a preimage of the unit  $[1_A] \in K_0(A)$  in  $\mathbf{Z}^{k_1}$  and denote its  $i^{th}$  coordinate by  $[1, i]$ ; set  $[n+1, i] = \sum_j \chi_n^0(i, j)[n, j]$ ,  $A_n = \bigoplus_{k=1}^{k_n} M_{[n,k]} \otimes C(\mathbf{T})$  and define  $\varphi_n : A_n \rightarrow A_{n+1}$  to be the block diagonal sum of standard embeddings of type  $(\chi_n^0(i, j), \chi_n^1(i, j))$  (that is, the partial embedding from the  $j^{th}$  summand of  $A_n$  to the  $i^{th}$  summand of  $A_{n+1}$  is of the above type). Since the partial embedding from a central summand in  $A_n$  to a central summand in  $A_m$  with  $m > n$  is a sum of composites of standard embeddings, a central unitary is mapped to a sum of normal unitaries which by the above assumption must have uniformly small spectral variation; it decreases by a factor of two at least when embedded into the next level (we require  $M(\chi_{n+1}^0) \geq 4$  to deal with the case  $\chi_n^1(i, j) = 0$ ). Then  $\lim A_n$  is a simple  $A\mathbf{T}$  algebra of real rank zero (small spectral variation guarantees real rank zero — see [BBEK, 1.3]). Hence, by Elliott’s classification theorem for simple real rank zero  $A\mathbf{T}$  algebras (cf. [E, 7.1]),  $A \cong \lim A_n$ .

1.7. In Corollary 3.13 we establish a sharper version of Elliott’s theorem: given two unital simple  $A\mathbf{T}$  algebras of real rank zero,  $A$  and  $B$ , then an invertible element in  $KK(A, B)$  which preserves positivity and the class of the unit lifts to an isomorphism.

1.8. The attempt to understand an invariant for approximately inner automorphisms of simple unital  $AT$  algebras of real rank zero introduced by Elliott and Rørdam (see [ER, 4.5]) stimulated our interest in the question addressed in the present work; their analysis of the invariant for the case of the Bunce-Deddens algebra (see [ER, 4.12i]) was particularly helpful. We thank George Elliott for several valuable conversations; we also wish to thank the staff of the Fields Institute, where much of this work was done, for their hospitality.

1.9. Some notational conventions: as usual we let  $\mathbf{N}$ ,  $\mathbf{Z}$ ,  $\mathbf{R}$  and  $\mathbf{T}$  denote the natural numbers, the integers, the reals, and the complex numbers of unit modulus. For  $m \in \mathbf{N}$ , let  $\{e_i\}$  denote the canonical basis in  $\mathbf{Z}^m$ ; a homomorphism  $\chi : \mathbf{Z}^n \rightarrow \mathbf{Z}^m$  will be confused with the  $m \times n$  matrix  $\chi_{i,j}$  (or  $\chi(i,j)$ ) for which  $\chi(e_j) = \sum_i \chi_{i,j} e_i$ .

2. THE EXTENSION CLASS OF AN APPROXIMATELY INNER AUTOMORPHISM

2.1. Let  $A$  be an  $AT$  algebra given as the limit of an inductive system of circle algebras  $\{A_n, \varphi_{mn}\}$ , where  $A_n = \bigoplus_{k=1}^{k_n} A_{n,k}$ ,  $A_{n,k} = M_{[n,k]} \otimes C(\mathbf{T})$  and  $\varphi_{mn} : A_n \rightarrow A_m$  is a unital homomorphism for  $m > n$ ; to simplify notation write  $\varphi_n$  in place of  $\varphi_{n+1,n}$ . Also, let  $\bar{\varphi}_n$  denote the canonical homomorphism from  $A_n$  to  $A$ . Let  $v_{n,k}$  be the unitary in  $A_n$  which restricts to the canonical central unitary in the  $k^{th}$  summand and the identity in the other summands. Let  $\{e_{ij}^{n,k}\}$  denote the canonical family of matrix units for the  $k^{th}$  summand and let  $p_{n,k}$  denote the projection onto the  $k^{th}$  summand (so  $p_{n,k} = \sum_i e_{ii}^{n,k}$ ). Set  $R_n = \{e_{ij}^{n,k} : 1 \leq i, j \leq [n,k], 1 \leq k \leq k_n\} \cup \{v_{n,k} : 1 \leq k \leq k_n\}$  and  $S_n = R_n \cup \varphi_{n-1}(S_{n-1})$ , where  $S_0$  is the empty set.

2.2. We consider a family of approximately inner automorphisms and determine the resulting Ext classes in the following theorem. Let  $\alpha_n = \text{Ad } u_n$  be an inner automorphism of  $A_n$ , where  $u_n$  is a unitary in  $A_n$ ;  $\alpha_n$  and  $\delta_n > 0$  are chosen inductively as follows: If  $\varphi$  and  $\varphi'$  are homomorphisms from  $A_n$  to  $A_{n+1}$  such that  $\|\varphi(x) - \varphi'(x)\| < \delta_n$  for  $x \in R_n$ , then  $\|\varphi(x) - \varphi'(x)\| < 2^{-n}$  for  $x \in S_n$ . Choose  $\alpha_{n+1}$  such that

$$(1) \quad \begin{aligned} \|\alpha_{n+1}^l \circ \varphi_n(e_{ij}^{n,k}) - \varphi_n \circ \alpha_n^l(e_{ij}^{n,k})\| &< \delta_n/32[n,k]k_n, \\ \|\alpha_{n+1}^l \circ \varphi_n(v_{n,k}) - \varphi_n \circ \alpha_n^l(v_{n,k})\| &< \delta_n/32 \end{aligned}$$

for  $l = \pm 1$ . Since  $R_n$  generates  $A_n$ , the limit

$$\lim_{m \rightarrow \infty} \bar{\varphi}_m \circ \alpha_m \circ \varphi_{mn}(x) = \lim_{m \rightarrow \infty} \text{Ad } \bar{\varphi}_m(u_m)(\bar{\varphi}_n(x))$$

exists for all  $x \in R_n$ . One defines a homomorphism  $\alpha : \bigcup_{n=1}^\infty \bar{\varphi}_n(A_n) \rightarrow A$  by

$$\alpha(\bar{\varphi}_n(x)) = \lim_{m \rightarrow \infty} \bar{\varphi}_m \circ \alpha_m \circ \varphi_{mn}(x)$$

for  $x \in A_n$ ; note that  $\alpha$  extends to a unital endomorphism. Since  $\beta \circ \bar{\varphi}_n(x) = \lim \bar{\varphi}_m \circ \alpha_m^{-1} \circ \bar{\varphi}_{mn}(x)$  for  $x \in A_n$  defines a unital endomorphism  $\beta$  of  $A$  and  $\alpha \circ \beta \circ \bar{\varphi}_n(x) = \bar{\varphi}_n(x)$  for  $x \in S_n$ , it follows that  $\alpha$  is an automorphism of  $A$ .

2.3. We view elements of  $A_n$  as matrix valued functions on  $k_n$  copies of  $\mathbf{T}$  (the size of the matrix may vary). Since

$$\|\alpha_{n+1} \circ \varphi_n(v_{n,j}) - \varphi_n(v_{n,j})\| < 2^{-n},$$

the Bott number  $B(\varphi_n(v_{n,j})(\iota_i), u_{n+1}(\iota_i))$  (see 1.4) is well-defined (where  $\iota_i$  is the basepoint of the  $i^{\text{th}}$  copy of  $\mathbf{T}$  in the spectrum of  $A_{n+1}$ ). For the purposes of computing the Bott number we may suppose that  $u_{n+1}(\iota_i)$  commutes with  $\varphi_n(p_{n,j})(\iota_i)$  (by perturbing it slightly) and so regard both  $\varphi_n(v_{n,j})(\iota_i)$  and  $u_{n+1}(\iota_i)$  as unitaries in  $\varphi_n(p_{n,j})(\iota_i)M_{[n+1,i]}\varphi_n(p_{n,j})(\iota_i)$ . Since the image of  $A_{n,j}$  in

$$\varphi_n(p_{n,j})(\iota_i)M_{[n+1,i]}\varphi_n(p_{n,j})(\iota_i)$$

almost commutes with  $\varphi_n(v_{n,j})(\iota_i)$  and  $u_{n+1}\varphi_n(u_n^*)(\iota_i)$ , it follows that

$$B(\varphi_n(v_{n,j})(\iota_i), u_{n+1}(\iota_i)) = B(\varphi_n(v_{n,j})(\iota_i), u_{n+1}(\iota_i)\varphi_n(u_n^*)(\iota_i))$$

is a multiple of  $[n, j]$ . For  $i = 1, \dots, k_{n+1}$  and  $j = 1, \dots, k_n$  set

$$\psi_n^0(i, j) = \begin{cases} B(\varphi_n(v_{n,j})(\iota_i), u_{n+1}(\iota_i))/[n, j] & \text{if } \varphi_n(p_{n,j})p_{n+1,i} \neq 0, \\ 0 & \text{otherwise} \end{cases}$$

(note that this is an integer) and

$$\psi_n^1(i, j) = \begin{cases} [\varphi_n(p_{n,j})u_{n+1}p_{n+1,i}\varphi_n(u_n^*p_{n,j})]_1/[n, j] & \text{if } \varphi_n(p_{n,j})p_{n+1,i} \neq 0, \\ 0 & \text{otherwise} \end{cases}$$

where  $[\cdot]_1$  denotes the class of an invertible element in

$$K_1(\varphi_n(p_{n,j})A_{n+1,i}\varphi_n(p_{n,j})) \cong \mathbf{Z}.$$

Since  $u_{n+1}p_{n+1,i}\varphi_n(u_n^*p_{n,j})$  almost commutes with the image of  $A_{n,j} = M_{[n,j]}$ ,  $\psi_n^1(i, j)$  is again an integer (for  $n$  sufficiently large).

$K_i(A)$  is given as the inductive limit of the sequence

$$\mathbf{Z}^{k_1} \xrightarrow{\chi_1^i} \mathbf{Z}^{k_2} \xrightarrow{\chi_2^i} \mathbf{Z}^{k_3} \rightarrow \dots,$$

where  $\chi_n^i = (\varphi_n)_*$ , viewed as a  $k_{n+1} \times k_n$  matrix with integer entries (note that  $\chi_n^0(i, j) \geq 0$ ).

**2.4. Theorem.** *Let  $A, A_n, \alpha, \alpha_n, \psi_n^i, \chi_n^i$  be as above. Then for  $i = 0, 1$  the extension*

$$0 \rightarrow K_{i+1}(A) \rightarrow K_i(M_\alpha) \rightarrow K_i(A) \rightarrow 0$$

is obtained as the inductive limit of the sequence

$$\begin{array}{ccccccccccc} \vdots & & \vdots & & \vdots & & \vdots & & \vdots & & \vdots \\ 0 & \rightarrow & \mathbf{Z}^{k_n} & \xrightarrow{\iota} & \mathbf{Z}^{k_n} & \oplus & \mathbf{Z}^{k_n} & \xrightarrow{\rho} & \mathbf{Z}^{k_n} & \rightarrow & 0 \\ & & \chi_n^{i+1} \downarrow & & \chi_n^{i+1} \downarrow & \swarrow \psi_n^{i+1} & \downarrow \chi_n^i & & \downarrow \chi_n^i & & \\ 0 & \rightarrow & \mathbf{Z}^{k_{n+1}} & \xrightarrow{\iota} & \mathbf{Z}^{k_{n+1}} & \oplus & \mathbf{Z}^{k_{n+1}} & \xrightarrow{\rho} & \mathbf{Z}^{k_{n+1}} & \rightarrow & 0 \\ & & \downarrow & & \downarrow & \swarrow & \downarrow & & \downarrow & & \\ \vdots & & \vdots & & \vdots & & \vdots & & \vdots & & \vdots \end{array}$$

where  $\iota$  is the embedding onto the first summand  $\mathbf{Z}^{k_n}$  and  $\rho$  is the projection onto the second summand  $\mathbf{Z}^{k_n}$  for each  $n$ ; note that in each case  $K_0$  is regarded as an ordered abelian group (while  $K_1$  is regarded as an abelian group).

2.5. We proceed now to the proof of the theorem. For the inner automorphism  $\alpha_n$  of  $A_n$  the six-term periodic sequence for the mapping torus  $M_{\alpha_n}$  reduces to two trivial short exact sequences:

$$0 \rightarrow K_{i+1}(A_n) \rightarrow K_i(M_{\alpha_n}) \rightarrow K_i(A_n) \rightarrow 0$$

for  $i = 0, 1$ ; note that  $K_i(A_n)$  is naturally isomorphic to  $\mathbf{Z}^{k_n}$  for  $i = 0, 1$ . The identification of  $K_0(M_{\alpha_n})$  (resp.  $K_1(M_{\alpha_n})$ ) with  $\mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n}$  as ordered abelian groups (resp. abelian groups) proceeds as follows: Let  $v : [0, 1] \rightarrow U(M_2 \otimes A_n)$  be a continuous path of unitaries with

$$v_0 = \begin{pmatrix} u_n & 0 \\ 0 & u_n^* \end{pmatrix}, \quad v_1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

For  $x \oplus y \in M_{\alpha_n} \oplus M_{\alpha_n^{-1}}$  set

$$(2) \quad (x \oplus y) \sim(t) = v_t \begin{pmatrix} x(t) & 0 \\ 0 & y(t) \end{pmatrix} v_t^*.$$

One checks that

$$(x \oplus y) \sim(0) = (x \oplus y) \sim(1);$$

thus  $(x \oplus y) \sim$  may be regarded as an element of  $C(\mathbf{T}) \otimes M_2 \otimes A_n$ . This defines an embedding of  $M_{\alpha_n} \oplus M_{\alpha_n^{-1}}$  into  $C(\mathbf{T}) \otimes M_2 \otimes A_n$ . Since the image of the map  $M_{\alpha_n} \rightarrow C(\mathbf{T}) \otimes M_2 \otimes A_n$  (by  $x \mapsto (x \oplus 0) \sim$ ) is a full hereditary subalgebra, we obtain an isomorphism,

$$K_*(M_{\alpha_n}) \simeq K_*(C(\mathbf{T}) \otimes M_2 \otimes A_n).$$

To identify  $K_i(C(\mathbf{T}) \otimes M_2 \otimes A_n)$  with  $\mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n}$  it suffices to identify  $K_i(C(\mathbf{T}) \otimes M_2 \otimes A_{n,k})$  with  $\mathbf{Z} \oplus \mathbf{Z}$  in the usual way — note that

$$C(\mathbf{T}) \otimes M_2 \otimes A_{n,k} = C(\mathbf{T}) \otimes M_2 \otimes M_{[n,k]} \otimes C(\mathbf{T}) \simeq M_{2[n,k]} \otimes C(\mathbf{T}^2).$$

The class of a unitary  $u \in M_{2[n,k]} \otimes C(\mathbf{T}^2)$  is identified with  $([u(\cdot, 1)]_1, [u(1, \cdot)]_1)$  (where the first variable is identified with the parameter in the mapping torus construction) and the class of a projection,  $p \in M_{2[n,k]} \otimes C(\mathbf{T}^2)$ , is identified with  $(\text{ch}(p), \dim(p))$ , where  $\text{ch}(p)$  is the first Chern class of  $p$  and  $\dim(p)$  is the dimension of  $p$ . Under this identification a nonzero element  $(m, n)$  is positive if and only if  $n > 0$ .

**2.6. Lemma.** *With identifications as above the extension*

$$0 \rightarrow K_{i+1}(A_n) \rightarrow K_i(M_{\alpha_n}) \rightarrow K_i(A_n) \rightarrow 0,$$

for  $i = 0, 1$  is isomorphic to

$$0 \rightarrow \mathbf{Z}^{k_n} \xrightarrow{\iota} \mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n} \xrightarrow{\rho} \mathbf{Z}^{k_n} \rightarrow 0$$

where  $\iota(a) = (a, 0)$  and  $\rho(a, b) = b$  for  $a, b \in \mathbf{Z}^{k_n}$ .

*Proof.* Let  $u$  be a unitary in  $A_n$  and let  $w : [0, 1] \rightarrow U(M_2 \otimes A_n)$  be a continuous path of unitaries with

$$w_0 = \begin{pmatrix} u & 0 \\ 0 & u^* \end{pmatrix}, \quad w_1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Set  $e_t = w_t(1 \oplus 0)w_t^*$ ; then  $e$  is a projection in  $M_2 \otimes (SA)^\sim$  and the map  $K_1(A_n) \rightarrow K_0(M_{\alpha_n})$ , is given by  $[u] \mapsto [e] - [1]$  (see [B, 8.2.2]). By the above identification the second component of  $[e] - [1]$  zero. Since the second component of  $K_0(M_{\alpha_n})$

corresponds to dimension, the map  $K_0(M_{\alpha_n}) \rightarrow K_0(A_n)$  is just the projection onto the second component.

Let  $p$  be a projection in  $A_n$ . Set  $u_t = (e^{2\pi it} - 1)p + 1$ . Then  $u$  is a unitary in  $(SA)^\sim \subset M_{\alpha_n}$ , and  $[p]$  maps to  $[u]$  under the map  $K_0(A_n) \rightarrow K_1(M_{\alpha_n})$ . Set  $\tilde{u}(t) = v_t(u_t \oplus 1)v_t^*$ , where  $\{v_t\}$  is as above (see equation (2)); we now compute  $[\tilde{u}]_1$  (in  $K_1(C(\mathbf{T}) \otimes M_2 \otimes A_n)$ ). Since  $\tilde{u}(1) = 1$ , the second component of  $[\tilde{u}]$  must be zero. The winding number of

$$t \mapsto \det \tilde{u}(t)(\iota_i) = \det((e^{2\pi it} - 1)p(\iota_i) + 1)$$

with  $\tilde{u}(t)$  evaluated at the point  $\iota_i$  in the spectrum of  $A_n$  is just  $\dim p(\iota_i) = \dim(p \cdot p_{n,i})$ . Thus the first component of  $[\tilde{u}]$  is equal to  $[p]$ . Let  $u$  be a unitary in  $M_{\alpha_n}$ , and set

$$\tilde{u}(t) = v_t(u_t \oplus 1)v_t^*.$$

Then, since  $\tilde{u}(1) = u_1 \oplus 1$ ,  $[u] = (*, [u_1])$  maps to  $[u_1]$  under the map  $K_1(M_{\alpha_n}) \rightarrow K_1(A_n)$ . □

2.7. We now define a homomorphism from  $M_{\alpha_n}$  to  $M_{\alpha_{n+1}}$ . This is done by using the fact that  $\varphi_n \circ \alpha_n$  and  $\alpha_{n+1} \circ \varphi_n$  are homotopic (as homomorphisms from  $A_n$  to  $A_{n+1}$ ). The induced map,  $K_i(M_{\alpha_n}) \rightarrow K_i(M_{\alpha_{n+1}})$  (for  $i = 0, 1$ ), is computed in the following lemma.

Recall that we have chosen  $\{\alpha_n\}$  so that

$$\begin{aligned} \|\alpha_{n+1} \circ \varphi_n(e_{ij}^{n,k}) - \varphi_n \circ \alpha_n(e_{ij}^{n,k})\| &< \delta_n/32[n, k]k_n, \\ \|\alpha_{n+1} \circ \varphi_n(v_{n,k}) - \varphi_n \circ \alpha_n(v_{n,k})\| &< \delta_n/32. \end{aligned}$$

We assume that  $\delta_n > 0$  is sufficiently small in the following. Set

$$x = \sum_{i,k} \alpha_{n+1} \circ \varphi_n(e_{ii}^{n,k})\varphi_n \circ \alpha_n(e_{ii}^{n,k});$$

then  $\|x - 1\| < \delta_n/32$ , and the unitary  $v_1 = x|x|^{-1}$  in  $A_{n+1}$  satisfies

$$\text{Ad } v_1 \circ \varphi_n \circ \alpha_n(e_{ii}^{n,k}) = \alpha_{n+1} \circ \varphi_n(e_{ii}^{n,k})$$

and  $\|v_1 - 1\| < \delta_n/16$  (for large  $n$ ). It follows that for  $i \neq j$

$$\|\text{Ad } v_1 \circ \varphi_n \circ \alpha_n(e_{ij}^{n,k}) - \alpha_{n+1} \circ \varphi_n(e_{ij}^{n,k})\| < 5\delta_n/32[n, k]k_n < 5\delta_n/32.$$

Now set

$$v_2 = \sum_{i,k} \text{Ad } v_1 \circ \varphi_n \circ \alpha_n(e_{i1}^{n,k})\alpha_{n+1} \circ \varphi_n(e_{1i}^{n,k});$$

then  $v_2$  is a unitary such that  $\|v_2 - 1\| \leq 5\delta_n/32$  and  $\text{Ad } v_2 v_1 \circ \varphi_n \circ \alpha_n(e_{ij}^{n,k}) = \alpha_{n+1} \circ \varphi_n(e_{ij}^{n,k})$ . Since  $\|v_2 v_1 - 1\| < 7\delta_n/32$ , there is a self-adjoint  $h \in A_{n+1}$  such that  $2\pi\|h\| \leq \delta_n/4$  and  $v_2 v_1 = e^{2\pi ih}$ . Since

$$\|\text{Ad } e^{2\pi ih} \circ \varphi_n \circ \alpha_n(v_{n,k}) - \alpha_{n+1} \circ \varphi_n(v_{n,k})\| < 15\delta_n/32,$$

there is a self-adjoint  $h_k \in A_{n+1}$  such that  $2\pi\|h_k\| < 16\delta_n/32$  and  $e^{2\pi ih_k} = \alpha_{n+1} \circ \varphi_n(v_{n,k})\text{Ad } e^{2\pi ih} \circ \varphi_n \circ \alpha_n(v_{n,k}^*)$ . For  $t \in [0, 1]$  we define a homomorphism  $\nu_t : A_n \rightarrow A_{n+1}$  as follows: for  $0 \leq t \leq 1/2$ , set

$$\nu_t = \text{Ad } e^{4\pi it h} \circ \varphi_n \circ \alpha_n = \text{Ad } (e^{4\pi it h} \varphi_n(u_n)) \circ \varphi_n,$$



and for  $1/2 < t \leq 1$ , set

$$\begin{aligned} \nu_t(e_{ij}^{n,k}) &= \alpha_{n+1} \circ \varphi_n(e_{ij}^{n,k}), \\ \nu_t(v_{n,k}) &= e^{2\pi i(2t-1)h_k} \text{Ad } e^{2\pi ih} \circ \varphi_n \circ \alpha_n(v_{n,k}). \end{aligned}$$

Note that  $\nu_t$  is well defined,  $t \mapsto \nu_t(x)$  is continuous for all  $x \in A_n$ ,  $\nu_0 = \varphi_n \circ \alpha_n$  and  $\nu_1 = \alpha_{n+1} \circ \varphi_n$ . Moreover,  $\|\nu_t(x) - \varphi_n \circ \alpha_n(x)\| \leq 2^{-n}$  for all  $x \in S_n$  (since  $\|\nu_t(x) - \varphi_n \circ \alpha_n(x)\| < \delta_n$  for  $x \in R_n$ ). Define  $\Phi_n : M_{\alpha_n} \rightarrow M_{\alpha_{n+1}}$  by

$$\Phi_n(x)(t) = \begin{cases} \varphi_n(x(\frac{t}{1-2^{-n}})), & 0 \leq t \leq 1 - 2^{-n}, \\ \nu_{2^n(t-1+2^{-n})}(x(0)), & 1 - 2^{-n} < t \leq 1. \end{cases}$$

**2.8. Lemma.** *Let  $\Phi_n$  be as above. Then, for  $i = 0, 1$ ,  $(\Phi_n)_* : K_i(M_{\alpha_n}) \rightarrow K_i(M_{\alpha_{n+1}})$  is given by*

$$\begin{array}{ccc} \mathbf{Z}^{k_n} & \oplus & \mathbf{Z}^{k_n} \\ \chi_n^{i+1} \downarrow & \swarrow \psi_n^{i+1} & \downarrow \chi_n^i \\ \mathbf{Z}^{k_{n+1}} & \oplus & \mathbf{Z}^{k_{n+1}} \end{array}$$

*Proof.* We deal with the case  $i = 0$  first. Let  $u \in A_n$  be a unitary and  $\{w_t\}$  a continuous path of unitaries in  $M_2 \otimes A_n$  from  $u \oplus u^*$  to 1. Set  $e_t = w_t(1 \oplus 0)w_t^*$ . Since  $e_0 = 1 \oplus 0$ ,  $\Phi_n(e)$  is equivalent to the projection defined by  $t \mapsto \varphi_n(e_t)$ , which is defined by  $\varphi_n(u)$  in the same way as  $e$  is defined by  $u$ . Since  $\Phi_n(1) = 1$  and  $[u]$  is mapped to  $[e] - [1]$ , this shows that  $\chi_n^1$  gives the map from the first summand of  $K_0(M_{\alpha_n})$  to the first summand of  $K_0(M_{\alpha_{n+1}})$  (a similar argument shows that  $\chi_n^0$  is the analogous map for  $i = 1$ ).

Let  $p_{n,k}$  be the central projection in  $A_n$  as before. Regarding  $p_{n,k}$  as a projection in  $M_{\alpha_n}$ , we have

$$[p_{n,k}] = ([u_n p_{n,k}], [n, k]) \in \mathbf{Z} \oplus \mathbf{Z} \subset \mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n},$$

where  $\mathbf{Z} \oplus \mathbf{Z}$  is mapped to the  $k^{\text{th}}$  summand of  $(\mathbf{Z} \oplus \mathbf{Z})^{k_n} = \mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n}$ . Let  $\{v_t\}$  be a continuous path of unitaries in  $M_2 \otimes A_{n+1}$  from  $u_{n+1} \oplus u_{n+1}^*$  to 1. Since  $u_{n+1} \oplus u_{n+1}^*$  almost commutes with  $\varphi_n(p_{n,k}) \oplus \varphi_n(p_{n,k})$ , we can assume that  $v_t$  almost commutes with  $\varphi_n(p_{n,k}) \oplus \varphi_n(p_{n,k})$  for all  $t \in [0, 1]$ . It follows that the class of  $\Phi_n(p_{n,k})$  in  $\mathbf{Z}^{k_{n+1}} \oplus \mathbf{Z}^{k_{n+1}}$  is the equivalence class of  $t \mapsto v_t(\varphi_n(p_{n,k}) \oplus 0)v_t^*$  composed with a short path from  $\varphi_n(p_{n,k}) \oplus 0$  to  $\alpha_{n+1} \circ \varphi_n(p_{n,k}) \oplus 0$ . Since this corresponds to the invertible element  $\varphi_n(p_{n,k})u_{n+1}\varphi_n(p_{n,k}) + 1 - \varphi_n(p_{n,k})$ , the first component of  $[\Phi_n(p_{n,k})]$  is

$$\begin{aligned} [\varphi_n(p_{n,k})u_{n+1}\varphi_n(p_{n,k})]_1 &= [\varphi_n(p_{n,k})u_{n+1}\varphi_n(u_n^*)\varphi_n(p_{n,k})]_1 + [\varphi_n(u_n)\varphi_n(p_{n,k})]_1 \\ &= (\psi_n^1(i, k)[n, k] + \chi_n^1(i, k)[u_n p_{n,k}])_i, \end{aligned}$$

where  $[\cdot]_1$  denotes the  $K_1$ -class of an invertible element. The second component of  $[\Phi_n(p_{n,k})]$  is the  $K_0$ -class of  $\varphi_n(p_{n,k})$ , i.e.  $(\chi_n^0(i, k)[n, k])$ . This takes care of the case  $i = 0$ .

Regarding  $v_{n,k}$  as an element of  $M_{\alpha_n}$ , it follows that  $[\Phi_n(v_{n,k})]$  is the class of

$$U : t \mapsto v_t(\Phi_n(v_{n,k})(t) \oplus 1)v_t^*$$

in  $K_1(C(\mathbf{T}) \otimes M_2 \otimes A_{n+1})$ . Evaluate  $U$  at  $\iota_i$  (the base point of the  $i^{\text{th}}$  copy of  $\mathbf{T}$  in the spectrum of  $A_{n+1}$ ) and let  $w_i$  be the winding number of

$$\begin{aligned} t &\mapsto \det(v_t(\iota_i)(\Phi_n(v_{n,k})(t, \iota_i) \oplus 1)v_t(\iota_i)^*) \\ &= \det(\Phi_n(v_{n,k})(t, \iota_i) \oplus 1). \end{aligned}$$

By the definition of  $\Phi_n(v_{n,k})$ ,  $w_i$  is equal to  $\text{Tr}(h_k(\iota_i))$ , where  $h_k$  is defined by

$$e^{2\pi i h_k} = \alpha_{n+1} \circ \varphi_n(v_{n,k}) \text{Ad } e^{2\pi i h} \circ \varphi_n \circ \alpha_n(v_{n,k}^*)$$

with  $h_k$  small. Since

$$e^{-2\pi i h} e^{2\pi i h_k} e^{2\pi i h} = e^{-2\pi i h} u_{n+1} \cdot \varphi_n(v_{n,k}) \cdot (e^{-2\pi i h} u_{n+1})^* \cdot \varphi_n(v_{n,k})^*,$$

it follows that  $w_i$  is equal to

$$B(\varphi_n(v_{n,k})(\iota_i), e^{-2\pi i h(\iota_i)} u_{n+1}(\iota_i)) = B(\varphi_n(v_{n,k})(\iota_i), u_{n+1}(\iota_i)) = \psi_n^0(i, k)[n, k].$$

Note that  $[U(0)] = [\varphi_n(v_{n,k})]$ ; thus,  $[\Phi_n(v_{n,k})] = (\psi_n^0(i, k)[n, k], \chi_n^1(i, k)[n, k])_i$ . Since  $[v_{n,k}] = (0, [n, k]) \in \mathbf{Z} \oplus \mathbf{Z} \subset \mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n}$ , the proof in the case  $i = 1$  is complete.  $\square$

**2.9. Lemma.** *For any  $x \in M_{\alpha_n}$ , the limit*

$$\lim_{m \rightarrow \infty} \overline{\varphi}_{n+m} \circ \Phi_{n+m-1} \circ \cdots \circ \Phi_n(x)$$

*exists in  $C[0, 1] \otimes A$  and defines a homomorphism of  $M_{\alpha_n}$  to  $M_\alpha$ .*

*Proof.* Given  $x \in M_{\alpha_n}$ , then for all  $t \in [0, 1]$ ,  $\overline{\varphi}_{n+m} \circ \Phi_{n+m-1} \circ \cdots \circ \Phi_n(x)(t)$  converges, say to  $\tilde{x}(t)$ ; one has  $\tilde{x}(1) = \alpha(\tilde{x}(0))$  and  $\tilde{x}(0) = x(0)$ . It also follows that convergence is uniform on  $[0, t]$  for all  $t < 1$ . For  $x \in S_n$  the family of functions,  $\{\overline{\varphi}_{n+m} \circ \Phi_{n+m-1} \circ \cdots \circ \Phi_n(x)(t) : m > 0\}$  is uniformly continuous in a neighbourhood of  $t = 1$ . Hence convergence is uniform for all  $x \in M_{\alpha_n}$ .  $\square$

2.10. We conclude the proof of the theorem. By the previous lemma we have a homomorphism  $\mu_n : M_{\alpha_n} \rightarrow M_\alpha$  such that  $\mu_n = \mu_{n+1} \circ \Phi_n$ . Let  $E$  denote the C\*-inductive limit,  $\lim_n M_{\alpha_n}$ ; there is a homomorphism  $\mu : E \rightarrow M_\alpha$  such that  $\mu_n = \mu \circ \overline{\Phi}_n$ , where  $\overline{\Phi}_n : M_{\alpha_n} \rightarrow E$  is the canonical map. We show that  $\mu$  is an isomorphism.

If  $\ker \mu \neq 0$ , then  $\ker \mu \cap \text{im } \overline{\Phi}_n \neq 0$  for some  $n$ . Let  $x \in M_{\alpha_n}$  be such that  $\mu \circ \overline{\Phi}_n(x) = \mu_n(x) = 0$ ; then, since  $\mu_n(x)(t) = \overline{\varphi}_n(x(t/s_n))$  for  $0 \leq t \leq s_n$ , where  $s_n = \prod_{k=1}^\infty (1 - 2^{-k}) > 1 - 2^{-n+1}$ , it follows that  $\overline{\varphi}_n(x(t)) = 0$ , and thus  $\overline{\Phi}_n(x)(t) = 0$  for  $0 \leq t \leq 1 - 2^{-n+1}$ . Since  $\overline{\Phi}_n(x) = \overline{\Phi}_m \circ \Phi_{mn}(x)$  for  $m > n$ , it follows that  $\overline{\Phi}_n(x) = 0$ ; thus,  $\mu$  is injective.

Let  $x \in M_\alpha$  with  $x(0) \in \overline{\varphi}_k(S_k)$  for some  $k$  and  $\epsilon > 0$ . There exist an  $n$  and  $y \in C[0, 1] \otimes A_n$  such that  $\|x(t) - \overline{\varphi}_n(y(t))\| < \epsilon$ . Since

$$\begin{aligned} & \| \overline{\varphi}_m \circ \alpha_m \circ \varphi_{mn}(y(0)) - \overline{\varphi}_n(y(1)) \| \\ & \leq \| \overline{\varphi}_m \circ \alpha_m \circ \varphi_{mn}(y(0)) - \alpha \circ \overline{\varphi}_m \circ \varphi_{mn}(y(0)) \| \\ & \quad + \| \alpha \circ \overline{\varphi}_n(y(0)) - \alpha(x(0)) \| + \| x(1) - \overline{\varphi}_n(y(1)) \| \\ & \leq \epsilon_m + 2\epsilon, \end{aligned}$$

where  $\epsilon_m \rightarrow 0$  as  $m \rightarrow \infty$ , we may take a small modification of  $y$  around  $t = 1$  and obtain, for sufficiently large  $m > n$ ,  $y \in C[0, 1] \otimes A_m$  such  $y(0) \in S_n$ ,  $\overline{\varphi}_m(y(0)) = x(0)$ ,  $y \in M_{\alpha_m}$ , and  $\|x(t) - \overline{\varphi}_m(y(t))\| < 3\epsilon$ . If  $m$  is sufficiently large we have that  $\|\overline{\varphi}_m(y(t)) - \mu_m(y)(t)\| < 7\epsilon$ , since  $\mu_m(y)$  is obtained by extending  $t \mapsto \overline{\varphi}_m(y(t))$  beyond  $t = 1$  up to  $1/s_m$  with  $s_m$  defined above and rescaling it; the extended part is within a sphere of radius  $2^{-m+1}$  centered at  $\overline{\varphi}_m(y(1))$ . Thus we have that  $\|x - \mu_m(y)\| < 11\epsilon$ . This implies that  $\mu$  is surjective.

3. REALIZATION OF AN EXTENSION CLASS  
BY AN APPROXIMATELY INNER AUTOMORPHISM

3.1. The proof of the following theorem will be divided into two cases; the case  $i = 1$  will be dealt with in sections 3.8 and 3.9 and the case  $i = 0$  will be handled in 3.11.

**Theorem.** *Let  $A$  be a simple unital AT algebra of real rank zero, and let  $i = 0$  or 1. Then, given an extension*

$$0 \rightarrow K_{i+1}(A) \rightarrow E \rightarrow K_i(A) \rightarrow 0,$$

there is an approximately inner automorphism  $\alpha$  of  $A$  such that

$$0 \rightarrow K_{i+1}(A) \rightarrow K_i(M_\alpha) \rightarrow K_i(A) \rightarrow 0$$

is equivalent to the above extension and

$$0 \rightarrow K_i(A) \rightarrow K_{i+1}(M_\alpha) \rightarrow K_{i+1}(A) \rightarrow 0$$

is a trivial extension. That is, for every element  $\eta \in \text{Ext}(K_i(A), K_{i+1}(A))$  there is an approximately inner automorphism  $\alpha$  of  $A$  such that  $\eta_i(\alpha) = \eta$  and  $\eta_{i+1}(\alpha) = 0$ .

**3.2. Lemma.** *Given  $\chi \in M_{mn}(\mathbf{Z})$  and a subgroup  $W$  of  $\mathbf{Z}^m$  with  $\text{im } \chi \cap W = \{0\}$ , there is a constant  $c \geq 0$  such that for any  $\psi \in M_{mn}(\mathbf{Z})$  with  $\ker \psi \supset \ker \chi$  there is a  $\gamma \in M_n(\mathbf{Z})$  such that*

$$\ker \gamma \supset W \quad \text{and} \quad |(\psi - \gamma\chi)(i, j)| \leq c$$

for  $i = 1, \dots, m, j = 1, \dots, n$ .

*Proof.* Let  $r = \text{rank } \chi$  and let  $b_1, \dots, b_r$  be generators for  $\mathbf{Z}^n$  such that  $\chi b_1, \dots, \chi b_r$  generate  $\chi\mathbf{Z}^n$  and  $\chi b_i = 0$  for  $i = r+1, \dots, n$ . We denote by  $U$  the invertible matrix in  $M_n(\mathbf{Z})$  defined by  $(b_1 \cdots b_n)$ .

Let  $\tilde{W}$  be a maximal subgroup of  $\mathbf{Z}^m$  such that

$$\tilde{W} \cap \text{im } \chi = \{0\}, \quad \tilde{W} \supset W.$$

Then  $\tilde{W}$  is isomorphic to  $\mathbf{Z}^{m-r}$ . Let  $a_i = \chi b_i$  for  $i = 1, \dots, r$ , and let  $a_{r+1}, \dots, a_m$  be generators for  $\tilde{W}$ . Define a matrix  $V \in M_m(\mathbf{Z})$  by  $V = (a_1 \cdots a_m)$ . Note that the determinant  $\det V$  is non-zero and that  $|\det V| > 1$  if  $\text{im } \chi + \tilde{W} \neq \mathbf{Z}^m$ . Then  $\chi$  can be expressed by

$$\chi = V\tilde{E}_r U^{-1},$$

where  $\tilde{E}_r$  is the canonical rank  $r$  matrix in  $M_{mn}(\mathbf{Z})$  with 1 on the first  $r$  diagonal elements and 0 elsewhere. Since  $\ker \psi \supset \ker \chi$ ,  $\psi$  can be expressed as

$$\begin{aligned} \psi &= (\psi(b_1) \cdots \psi(b_r) 0 \cdots 0)\tilde{E}_r U^{-1} \\ &= (\psi(b_1) \cdots \psi(b_r) 0 \cdots 0)V^{-1}\chi. \end{aligned}$$

Note that for any  $\eta_1, \dots, \eta_r$  in  $\mathbf{Z}^m$  the matrix

$$\gamma = (\eta_1 \cdots \eta_r 0 \cdots 0)(\det V)V^{-1}$$

is such that  $\gamma \in M_m(\mathbf{Z})$  and  $\ker \gamma \supset \tilde{W} \supset W$ . We choose  $\eta_i \in \mathbf{Z}^m$  so that

$$\|(\det V)^{-1}\psi(b_i) - \eta_i\|_\infty \leq 1/2,$$

where  $\|x\|_\infty = \max |x_j|$  for  $x \in \mathbf{R}^m$ . Then defining  $\gamma$  as above, we obtain that

$$\psi - \gamma\chi = \{(\det V)^{-1}(\psi(b_1) \cdots \psi(b_r) 0 \cdots 0) - (\eta_1 \cdots \eta_r 0 \cdots 0)\}(\det V)V^{-1}\chi$$

and

$$|(\psi - \gamma\chi)(i, j)| \leq c,$$

where

$$c = \max_j 1/2 \sum_{k,l} |\det V| \cdot |V^{-1}(k, l)\chi(l, j)|,$$

which does not depend on  $\psi$ . □

3.3. Let  $G_1$  be a countable torsion-free abelian group. Then there exist a sequence  $l_n$  of positive integers and an inductive system

$$\mathbf{Z}^{l_1} \xrightarrow{\chi_1^1} \mathbf{Z}^{l_2} \xrightarrow{\chi_2^1} \mathbf{Z}^{l_3} \rightarrow \dots,$$

for which the inductive limit is isomorphic to  $G_1$ , such that

$$\text{rank } \chi_{n+1}^1 \chi_n^1 = \text{rank } \chi_n^1$$

for  $n = 1, 2, 3, \dots$ . Since  $\text{rank } \chi_{n+1,m}^1 = \text{rank } \chi_{nm}^1$  for  $n > m$ , we have that  $\ker \chi_n^1 \cap \text{im } \chi_{nm}^1 = \{0\}$  for  $n > m$ . Thus we obtain a constant  $c_{nm} = c$  for the pair  $\chi_{nm}^1$  and  $\ker \chi_n^1$  by the previous lemma. We set

$$c_n = \max\{c_{km} : n \geq k > m\},$$

which forms an increasing sequence. We always assume that  $c_n \geq 1$ .

Later, for a sequence  $k_n$  of positive integers with  $k_n \geq l_n$ , we define another inductive system

$$\mathbf{Z}^{k_1} \xrightarrow{\tilde{\chi}_1^1} \mathbf{Z}^{k_2} \xrightarrow{\tilde{\chi}_2^1} \mathbf{Z}^{k_3} \rightarrow \dots,$$

by extending  $\chi_n^1$  with 0's, i.e., defining  $\tilde{\chi}_n^1$  by

$$\mathbf{Z}^{k_n} \xrightarrow{\rho} \mathbf{Z}^{l_n} \xrightarrow{\chi_n^1} \mathbf{Z}^{l_{n+1}} \xrightarrow{\iota} \mathbf{Z}^{k_{n+1}},$$

where  $\rho$  (resp.  $\iota$ ) is an obvious projection (resp. embedding). The inductive limit is the same as before. Since  $\ker \tilde{\chi}_{n+1}^1 = \iota(\ker \chi_{n+1}^1) \oplus \mathbf{Z}^{k_{n+1}-l_{n+1}}$  and  $\text{im } \tilde{\chi}_n^1 \subset \mathbf{Z}^{l_n} \oplus 0$ , we retain the same constant  $c_n$  as before for this new inductive system.

Let  $G_0$  be a simple dimension group other than  $\mathbf{Z}$ . Then there exist an increasing sequence  $m_n$  of positive integers and an inductive system

$$\mathbf{Z}^{m_1} \xrightarrow{\chi_1^0} \mathbf{Z}^{m_2} \xrightarrow{\chi_2^0} \mathbf{Z}^{m_3} \rightarrow \dots,$$

for which the inductive limit is isomorphic to  $G_0$  (as ordered abelian groups), such that

$$m_n \geq l_n, \quad M(\chi_n^0) \geq nc_n,$$

where  $M(\chi) = \min_{i,j} \chi(i, j)$  for a matrix  $\chi$  with non-negative entries. The extra conditions placed on the inductive system can be handled easily.

**3.4. Lemma.** *Given an extension*

$$0 \rightarrow K_0(A) \xrightarrow{\iota} E \xrightarrow{q} K_1(A) \rightarrow 0$$

as in the theorem, there exist an increasing sequence  $k_n$  of positive integers and inductive systems

$$\mathbf{Z}^{k_1} \xrightarrow{\chi_1^i} \mathbf{Z}^{k_2} \xrightarrow{\chi_2^i} \mathbf{Z}^{k_3} \rightarrow \dots$$

for  $i = 0, 1$  and

$$\mathbf{Z}^{k_1} \oplus \mathbf{Z}^{k_1} \xrightarrow{\psi_1} \mathbf{Z}^{k_2} \oplus \mathbf{Z}^{k_2} \xrightarrow{\psi_2} \mathbf{Z}^{k_3} \oplus \mathbf{Z}^{k_3} \rightarrow \dots$$

such that  $M(\chi_n^0) \geq nc_n$  with  $c_n$  defined for the sequence  $\chi_n^1$  as in 3.3,  $\psi_n$  is of the form  $\psi_n(a, b) = (\chi_n^0(a) + \psi_n^0(b), \chi_n^1(b))$ ,  $\ker \psi_n^0 \supset \ker \chi_n^1$ ,  $\text{rank } \chi_{n+1}^1 \chi_n^1 = \text{rank } \chi_n^1$ , and the given short exact sequence is isomorphic to the inductive limit of the sequence

$$\begin{array}{ccccccccc} 0 & \rightarrow & \mathbf{Z}^{k_1} & \xrightarrow{\iota} & \mathbf{Z}^{k_1} \oplus \mathbf{Z}^{k_1} & \xrightarrow{\rho} & \mathbf{Z}^{k_1} & \rightarrow & 0 \\ & & \chi_n^0 \downarrow & & \psi_1 \downarrow & & \downarrow \chi_n^1 & & \\ 0 & \rightarrow & \mathbf{Z}^{k_2} & \xrightarrow{\iota} & \mathbf{Z}^{k_2} \oplus \mathbf{Z}^{k_2} & \xrightarrow{\rho} & \mathbf{Z}^{k_2} & \rightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ \vdots & & \vdots & & \vdots & & \vdots & & \vdots \end{array}$$

We may also assume that  $M(\chi_{n+1}^0) \geq \max(2L(\chi_n^1), 4)$  for all  $n$ .

*Proof.* For  $G_i = K_i(A)$  we choose inductive systems as in 3.3,  $\mathbf{Z}^{m_1} \rightarrow \mathbf{Z}^{m_2} \rightarrow \dots$  for  $i = 0$  and  $\mathbf{Z}^{l_1} \rightarrow \mathbf{Z}^{l_2} \rightarrow \dots$  for  $i = 1$ , with all the properties specified there. We shall choose a subsequence  $k(n)$  of positive integers and homomorphisms  $\zeta_n : \mathbf{Z}^{m_{k(n)}} \oplus \mathbf{Z}^{l_n} \rightarrow E$ ,  $\psi_n : \mathbf{Z}^{m_{k(n)}} \oplus \mathbf{Z}^{l_n} \rightarrow \mathbf{Z}^{m_{k(n+1)}} \oplus \mathbf{Z}^{l_{n+1}}$  for each  $n$  such that the diagrams

$$\begin{array}{ccccccccc} 0 & \rightarrow & \mathbf{Z}^{m_{k(n)}} & \xrightarrow{\iota} & \mathbf{Z}^{m_{k(n)}} \oplus \mathbf{Z}^{l_n} & \xrightarrow{\rho} & \mathbf{Z}^{l_n} & \rightarrow & 0 \\ & & \bar{\chi}_{k(n)}^0 \downarrow & & \zeta_n \downarrow & & \downarrow \bar{\chi}_n^1 & & \\ 0 & \rightarrow & K_0(A) & \longrightarrow & E & \longrightarrow & K_1(A) & \rightarrow & 0 \end{array}$$

and

$$\begin{array}{ccc} \mathbf{Z}^{m_{k(n)}} \oplus \mathbf{Z}^{l_n} & \xrightarrow{\psi_n} & \mathbf{Z}^{m_{k(n+1)}} \oplus \mathbf{Z}^{l_{n+1}} \\ \zeta_n \downarrow & & \downarrow \zeta_{n+1} \\ E & = & E \end{array}$$

are commutative,  $M(\chi_{k(n+1)k(n)}^0) \geq nc_n$  (and  $M(\chi_{k(n+1)k(n)}^0) \geq \max(2L(\chi_{n-1}^1), 4)$ , for  $n > 1$ ) and

$$\ker \psi_n^0 \supset \ker \chi_n^1, \quad \ker \zeta_n \supset \ker(\bar{\chi}_{k(n)}^0 \oplus \bar{\chi}_n^1),$$

where  $\psi_n(a, b) = (\chi_{k(n+1)k(n)}^0(a) + \psi_n^0(b), \chi_n^1(b))$ . Once this is done, we set  $k_n = m_{k(n)} (\geq l_n)$  and replace the inductive system  $\mathbf{Z}^{l_1} \rightarrow \mathbf{Z}^{l_2} \rightarrow \dots$  for  $K_1(A)$  by

$$\mathbf{Z}^{k_1} \xrightarrow{\chi_1^1} \mathbf{Z}^{k_2} \rightarrow \dots$$

as specified in 3.3. We also redefine  $\zeta_n, \psi_n$  in the obvious way. The required properties are retained under this change, and thus the proof will be complete.

We prove the above assertion by induction. Let  $k(1) = 1$ . Since  $\text{im } \bar{\chi}_1^1$  is torsion-free, it is isomorphic to  $\mathbf{Z}^r$  for some  $r \leq l_1$ . Hence we can find a homomorphism  $\phi$  of  $\text{im } \bar{\chi}_1^1$  into  $E$  such that  $q \circ \phi = \text{id}$  and define  $\zeta_1$  by

$$\zeta_1(a, b) = \iota \circ \bar{\chi}_1^0(a) + \phi \circ \bar{\chi}_1^1(b).$$

Then the required properties for  $\zeta_1$  are immediate. Suppose that we have constructed  $k(1), \dots, k(n), \zeta_1, \dots, \zeta_n$ , and  $\psi_1, \dots, \psi_{n-1}$  with the required properties. We then find a homomorphism  $\phi : \text{im } \bar{\chi}_{n+1}^1 \rightarrow E$  such that  $q \circ \phi = \text{id}$ . Choose a basis  $b_1, \dots, b_{l_n} \in \mathbf{Z}^{l_n}$  so that  $\bar{\chi}_n^1(b_1), \dots, \bar{\chi}_n^1(b_r)$  generates  $\text{im } \bar{\chi}_n^1 \cong \mathbf{Z}^r$  and  $\bar{\chi}_n^1(a_i) = 0$  for  $r < i \leq l_n$ . Then it follows that

$$\zeta_n(0, b_i) - \phi \circ \bar{\chi}_n^1(b_i) \in \iota(K_0(A)).$$

for all  $1 \leq i \leq l_n$  (note that  $\zeta_n(0, b_i) = 0 = \phi \circ \bar{\chi}_n^1(b_i)$  for  $i > r$ ). Since  $\bigcup_{n=1}^\infty \text{im } \bar{\chi}_n^0 = K_0(A)$ , there is a  $k(n+1) > k(n)$  such that the left hand side is contained in

$\iota(\text{im } \bar{\chi}_{k(n+1)}^0)$  for all  $i$  — we also require that  $M(\chi_{k(n+1)k(n)}^0) \geq \max(2L(\chi_{n-1}^1), 4)$  for  $n > 1$ .

We define  $\zeta_{n+1} : \mathbf{Z}^{m_{k(n+1)}} \oplus \mathbf{Z}^{l_{n+1}} \rightarrow E$  by

$$\zeta_{n+1}(a, b) = \iota \circ \bar{\chi}_{k(n+1)}^0(a) + \phi \circ \bar{\chi}_{n+1}^1(b);$$

it follows by construction that  $\ker \zeta_{n+1} \supset \ker(\bar{\chi}_{k(n+1)}^0 \oplus \bar{\chi}_{n+1}^1)$ . Then there exist  $p_i \in \mathbf{Z}^{m_{k(n+1)}}$  such that

$$\zeta_n(0, b_i) - \zeta_{n+1}(0, \chi_n^1(b_i)) = \zeta_n(0, b_i) - \phi \circ \bar{\chi}_n^1(b_i) = \iota \circ \bar{\chi}_{k(n+1)}^0(p_i).$$

We assume that  $p_i = 0$  for  $i > r$  and define  $\psi_{n+1}^0 : \mathbf{Z}^{l_n} \rightarrow \mathbf{Z}^{m_{k(n+1)}}$  by

$$\psi_{n+1}^0(b_i) = p_i.$$

Since the  $b_i$ 's form a basis,  $\psi_{n+1}^0$  is well-defined; moreover,  $\ker \psi_{n+1}^0 \supset \ker \bar{\chi}_{n+1}^1 = \ker \chi_{n+1}^1$ , where the last equality follows from the assumption  $\text{rank } \chi_{n+1}^1 \chi_n^1 = \text{rank } \chi_n^1$ . Since  $k(n+1) \geq n+1$  and hence  $c_{k(n+1)} \geq c_{n+1}$  (from the definition of  $c_n$ ), it follows that

$$M(\chi_{k(n+1)}^0) \geq k(n+1)c_{k(n+1)} \geq (n+1)c_{n+1},$$

where the first inequality follows from the choice made at the beginning of the proof. Since the commutativity of the diagrams follows immediately, this completes the induction.  $\square$

3.5. By passing to a subsequence one may further assume that  $M(\chi_n^0) > L_n c_n$  for arbitrarily large  $L_n$ . First note that for  $n > m$ ,

$$\psi_{nm}(a, b) = \psi_{n-1} \circ \dots \circ \psi_m(a, b) = (\chi_{nm}^0(a) + \psi_{nm}^0(b), \chi_{nm}^1(b)),$$

where  $\psi_{nm}^0 : \mathbf{Z}^{k_m} \rightarrow \mathbf{Z}^{k_{n+1}}$  is defined by

$$\psi_{nm}^0(b) = \sum_{k=m}^{n-1} \chi_{n,k+1}^0 \circ \psi_k \circ \chi_{km}^1(b).$$

Thus it follows that  $\ker \psi_{nm}^0 \supset \ker \bar{\chi}_m^1 = \ker \chi_m^1$ . In this way we show that by passing to a subsequence  $n_j$  in the situation of 3.4 all the algebraic conditions are retained and the estimate on  $M(\chi_n^0)$  is improved as follows:

$$M(\chi_{n_{j+1}n_j}^0) \geq M(\chi_{n_{j+1}-1}^0) \geq (n_{j+1} - 1)c_{n_{j+1}-1},$$

where  $(c_{n_{j+1}-1})$  can be regarded as  $(c_j)$  for the sequence  $(\chi_{n_{j+1}n_j}^1)$  and  $n_{j+1} - 1$  can be made arbitrarily large.

**3.6. Lemma.** *Suppose that the extension*

$$0 \rightarrow K_0(A) \xrightarrow{\iota} E \xrightarrow{q} K_1(A) \rightarrow 0$$

is obtained as the inductive limit of

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbf{Z}^{k_1} & \xrightarrow{\iota} & \mathbf{Z}^{k_1} \oplus \mathbf{Z}^{k_1} & \xrightarrow{\rho} & \mathbf{Z}^{k_1} \rightarrow 0 \\ & & \chi_1^0 \downarrow & & \psi_1 \downarrow & & \downarrow \chi_1^1 \\ 0 & \rightarrow & \mathbf{Z}^{k_2} & \xrightarrow{\iota} & \mathbf{Z}^{k_2} \oplus \mathbf{Z}^{k_2} & \xrightarrow{\rho} & \mathbf{Z}^{k_2} \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \vdots & & \vdots & & \vdots & & \vdots \end{array}$$

where  $\psi_n$  is of the form  $\psi_n(a, b) = (\chi_n^0(a) + \psi_n^0(b), \chi_n^1(b))$ . Let  $\gamma_n \in M_{k_n}(\mathbf{Z})$  be given for all  $n \geq 1$  and define a homomorphism  $\varphi_n : \mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n} \rightarrow \mathbf{Z}^{k_{n+1}} \oplus \mathbf{Z}^{k_{n+1}}$  in the same way as  $\psi_n$  with  $\psi_n^0$  replaced by

$$\varphi_n^0 = \psi_n^0 + \chi_n^0 \gamma_n - \gamma_{n+1} \chi_n^1.$$

Then the extension obtained as the inductive limit of the above system with  $\varphi_n$  in place of  $\psi_n$  is isomorphic to the original extension.

*Proof.* Define a homomorphism  $\nu_n : \mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n} \rightarrow \mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n}$  by

$$\nu_n(a, b) = (a + \gamma_n(b), b).$$

Then  $\nu_{n+1} \circ \varphi_n = \psi_n \circ \nu_n$  by computation. Thus  $(\nu_n)$  induces an isomorphism of the inductive limit  $E'$  for  $(\varphi_n)$  onto  $E$  as required. One can show this by direct computation, or it also follows from the following commutative diagram by knowing that both rows are exact:

$$\begin{array}{ccccccccc} 0 & \rightarrow & K_0(A) & \rightarrow & E' & \rightarrow & K_1(A) & \rightarrow & 0 \\ & & \parallel & & \downarrow & & \parallel & & \\ 0 & \rightarrow & K_0(A) & \rightarrow & E & \rightarrow & K_1(A) & \rightarrow & 0 \end{array}$$

□

**3.7. Lemma.** *Let  $\varphi$  be the standard  $n$ -times around embedding of  $C(\mathbf{T})$  into  $M_n \otimes C(\mathbf{T})$  and let  $u$  be the canonical unitary of  $C(\mathbf{T})$ . Then for any  $k \in \{0, 1, \dots, n-1\}$  there is a unitary  $v \in M_n \otimes C(\mathbf{T})$  with  $[v] = 0$  such that  $v\varphi(u)v^* = e^{2\pi ik/n}\varphi(u)$ .*

*Proof.* Recall that  $\varphi$  has the form

$$\varphi(u)(z) = \begin{pmatrix} 0 & 0 & \cdots & z \\ 1 & 0 & \cdots & 0 \\ & \ddots & \ddots & \\ 0 & \cdots & 1 & 0 \end{pmatrix},$$

and define

$$v = \begin{pmatrix} \omega^0 & 0 & \cdots & 0 \\ 0 & \omega^1 & \cdots & 0 \\ & & \ddots & \\ 0 & 0 & \cdots & \omega^{n-1} \end{pmatrix},$$

where  $\omega = e^{2\pi i/n}$ . Then  $\text{Ad } v(\varphi(u)) = \omega\varphi(u)$ . □

**3.8. Proof of the theorem for the case  $i = 1$ .** By Lemma 3.4 we may express the short exact sequence.

$$0 \rightarrow K_0(A) \xrightarrow{\iota} E \xrightarrow{q} K_1(A) \rightarrow 0$$

as the limit of an inductive system of short exact sequences:

$$\begin{array}{ccccccccc} 0 & \rightarrow & \mathbf{Z}^{k_1} & \xrightarrow{\iota} & \mathbf{Z}^{k_1} \oplus \mathbf{Z}^{k_1} & \xrightarrow{\rho} & \mathbf{Z}^{k_1} & \rightarrow & 0 \\ & & \chi_1^0 \downarrow & & \chi_1^0 \downarrow \swarrow \psi_1^0 \downarrow \chi_1^1 & & \downarrow \chi_1^1 & & \\ 0 & \rightarrow & \mathbf{Z}^{k_2} & \xrightarrow{\iota} & \mathbf{Z}^{k_2} \oplus \mathbf{Z}^{k_2} & \xrightarrow{\rho} & \mathbf{Z}^{k_2} & \rightarrow & 0 \\ & & \downarrow & & \downarrow \swarrow \downarrow & & \downarrow & & \\ \vdots & & \vdots & & \vdots \quad \vdots \quad \vdots & & \vdots & & \vdots \end{array}$$

where  $M(\chi_n^0) > L_n c_n$  for all  $n$  and some constant  $L_n \geq n$ ,  $\ker \psi_n^0 \supset \ker \chi_n^1$ ,  $\text{rank } \chi_{n+1}^1 \chi_n^1 = \text{rank } \chi_n^1$  and  $M(\chi_{n+1}^0) \geq \max(2L(\chi_n^1), 4)$  for all  $n$ . Note by 3.5 that these properties hold with arbitrarily large  $L_n$  when passing to a subsequence.

We choose a subsequence of the above inductive system, replace  $\psi_n^0$  so that  $|\psi_n^0(i, j)| \leq c_n$  holds (Lemma 3.2) without changing the inductive limit (Lemma 3.6), and construct circle algebras  $A_n = \bigoplus_{k=1}^{k_n} M_{[n,k]} \otimes C(\mathbf{T})$  for each level of the subsequence, embeddings  $\varphi_n : A_n \rightarrow A_{n+1}$  by using standard embeddings (see 1.6), and unitaries  $u_n \in A_n$  such that  $\varphi_n$  induces  $\chi_n^i : K_i(A_n) \rightarrow K_i(A_{n+1})$ ,  $\text{Ad } u_{n+1} \circ \varphi_n \approx \varphi_n \circ \text{Ad } u_n$  as described in the first theorem (which depends on how small  $|\psi_n^0(i, j)|/M(\chi_n^0) \leq L_n^{-1}$  is),  $[u_{n+1}, \varphi_n(p_{n,j})] = 0$ ,

$$[u_{n+1} p_{n+1,i} \varphi_n(u_n^* p_{n,j})]_1 = 0$$

in  $K_1(\varphi_n(p_{n,j})A_{n+1,i}\varphi_n(p_{n,j}))$  and

$$B(\varphi_n(v_{n,j})(\iota_i), u_{n+1}(\iota_i)) = \psi_n^0(i, j)[n, j].$$

Let  $A_\infty$  denote the inductive limit of  $A_1 \rightarrow A_2 \rightarrow A_3 \rightarrow \dots$ ; then, by Elliott's classification theorem (cf. [E, 7.1]),  $A_\infty$  is isomorphic to  $A$  (by 1.6 the condition  $M(\chi_{n+1}^0) \geq \max(2L(\chi_n^1), 4)$  for all  $n$  guarantees that  $A_\infty$  has real rank zero), and the limit  $\alpha = \lim \text{Ad } \bar{\varphi}_n(u_n)$  defines the desired automorphism  $\alpha$  of  $A_\infty$ . Note that  $\eta_0(\alpha) = 0$  by the above properties of  $[u_n]$ .

Let  $u_1$  be an arbitrary unitary in  $A_1$  with  $[u_1] = 0$ . The construction now proceeds inductively; suppose that  $A_1, \dots, A_n, \varphi_1, \dots, \varphi_{n-1}, \delta_1, \dots, \delta_{n-1}$ , and  $u_1, \dots, u_n$  have been chosen so that condition (1) on  $\alpha_{m+1} \circ \varphi_m \approx \varphi_m \circ \alpha_m$  holds with  $\alpha_m = \text{Ad } u_m$ . Recall that  $S_{m+1} = R_{m+1} \cup \varphi_m(S_m)$  and  $R_m$  are the generators for  $A_m$ . □

3.9. We note the following lemma, without giving its proof.

**Lemma.** *Let  $F$  be a finite subset of  $A_n$ . For any  $\epsilon > 0$  there is a  $\delta > 0$  such that for any two homomorphisms  $\varphi, \varphi'$  from  $A_n$  into another  $C^*$ -algebra  $B$ , if  $\|\varphi(x) - \varphi'(x)\| < \delta$  for all  $x \in R_n$ , then  $\|\varphi(x) - \varphi'(x)\| < \epsilon$  for  $x \in F$ .*

By applying the above lemma for  $F = S_n$  and  $\epsilon = 2^{-n}$  we obtain  $\delta = \delta_n > 0$ . Then, by passing to a subsequence (see 3.5) and modifying  $\psi_n^0, \psi_{n+1}^0$  by using 3.2 and 3.6, we may assume that

$$2\pi(L(\psi_n^0) + 1)/M(\chi_n^0) < \delta_n/32,$$

where  $L(\psi) = \max_{i,j} |\psi(i, j)|$ . Note that the modified  $\psi_{n+1}^0$  satisfies the condition  $\ker \psi_{n+1}^0 \supset \ker \chi_{n+1}^1$ . Define an embedding  $\varphi_n : A_n \rightarrow A_{n+1}$  so that the partial embedding from  $A_{n,j}$  to  $A_{n+1,i}$  is a standard embedding of type  $(\chi_n^0(i, j), \chi_n^1(i, j))$  (see 1.6). Using 3.7 choose a unitary

$$v_{n+1} \in \varphi_n\left(\bigoplus_{j=1}^{k_n} M_{[1,j]} \otimes 1\right)' \cap \bigoplus_{i=1}^{k_{n+1}} M_{[n+1,i]} \otimes 1 \subset A_{n+1}$$

such that

$$\text{Ad } v_{n+1} \circ \varphi_n(v_{n,j})p_{n+1,i} = e^{2\pi i \psi_n^0(i,j)/\chi_n^0(i,j)} \varphi_n(v_{n,j})p_{n+1,i}$$

in the case  $\chi_n^1(i, j) \neq 0$ , and

$$\text{Ad } v_{n+1} \circ \varphi_n(v_{n,j})p_{n+1,i} \cong e^{2\pi i a_1/b_1} \varphi_{b_1,1}(v_{n,j}) \oplus e^{2\pi i a_2/b_2} \varphi_{b_2,-1}(v_{n,j})$$



in the case  $\chi_n^1(i, j) = 0$ , where  $\varphi_n(v_{nj}) \cong \varphi_{b_1,1}(v_{nj}) \oplus \varphi_{b_2,-1}(v_{nj})$ ,  $a_1 = \lfloor \psi_n^0(i, j)/2 \rfloor$ ,  $b_1 = \lfloor \chi_n^0(i, j)/2 \rfloor$ ,  $a_1 + a_2 = \psi_n^0(i, j)$ , and  $b_1 + b_2 = \chi_n^0(i, j)$  (see 1.6). In either case it follows that  $[v_{n+1}\varphi_n(p_{n,j})]_1 = 0$  in  $K_1(\varphi_n(p_{n,j})A_{n+1}\varphi_n(p_{n,j}))$ , and

$$\begin{aligned} B(\varphi_n(v_{nj})(l_i), u_{n+1}(l_i)) &= \psi_n^0(i, j)[n, j], \\ \|\text{Ad } v_{n+1} \circ \varphi_n(v_{n,j}) - \varphi_n(v_{n,j})\| &< \delta_n/32, \\ \text{Ad } v_{n+1} \circ \varphi_n(e_{ij}^{n,k}) &= \varphi_n(e_{ij}^{n,k}). \end{aligned}$$

Setting  $u_{n+1} = v_{n+1}\varphi_n(u_n)$  completes the induction.

**3.10. Lemma.** *Given an extension as in the theorem,*

$$0 \rightarrow K_1(A) \xrightarrow{\iota} E \xrightarrow{q} K_0(A) \rightarrow 0,$$

there exists an increasing sequence  $k_n$  of positive integers and inductive systems

$$\mathbf{Z}^{k_1} \xrightarrow{\chi_1^i} \mathbf{Z}^{k_2} \xrightarrow{\chi_2^i} \mathbf{Z}^{k_3} \rightarrow \dots$$

for  $i = 0, 1$  and

$$\mathbf{Z}^{k_1} \oplus \mathbf{Z}^{k_1} \xrightarrow{\psi_1} \mathbf{Z}^{k_2} \oplus \mathbf{Z}^{k_2} \xrightarrow{\psi_2} \mathbf{Z}^{k_3} \oplus \mathbf{Z}^{k_3} \rightarrow \dots$$

where  $\chi_n^0$  is positive, and  $\psi_n$  is of the form  $\psi_n(a, b) = (\chi_n^1(a) + \psi_n^1(b), \chi_n^0(b))$  for each  $n$ , such that the given short exact sequence is isomorphic to the inductive limit of the sequence

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbf{Z}^{k_1} & \xrightarrow{\iota} & \mathbf{Z}^{k_1} \oplus \mathbf{Z}^{k_1} & \xrightarrow{\rho} & \mathbf{Z}^{k_1} \rightarrow 0 \\ & & \chi_1^1 \downarrow & & \psi_1 \downarrow & & \downarrow \chi_1^0 \\ 0 & \rightarrow & \mathbf{Z}^{k_2} & \xrightarrow{\iota} & \mathbf{Z}^{k_2} \oplus \mathbf{Z}^{k_2} & \xrightarrow{\rho} & \mathbf{Z}^{k_2} \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ \vdots & & \vdots & & \vdots & & \vdots \end{array}$$

Before going into the proof, note that  $E$  has a natural order,  $a \in E$  is positive if  $q(a)$  is a non-zero positive element of  $K_0(A)$  or  $a = 0$ , and that  $E$  is a dimension group with respect to this order. Define an order on  $\mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n}$  by  $(a, b) \geq 0$  if  $b \geq 0$ ,  $b \neq 0$  or  $(a, b) = 0$ ; observe that  $\psi_n$  preserves order and that the inductive limit is isomorphic to  $E$  as an ordered abelian group. This is how  $K_0(M_\alpha)$  is obtained as an inductive limit in 2.4.

*Proof.* There is an increasing sequence  $\{l_n\}$  and inductive systems

$$\mathbf{Z}^{l_1} \xrightarrow{\xi_1^i} \mathbf{Z}^{l_2} \xrightarrow{\xi_2^i} \mathbf{Z}^{l_3} \rightarrow \dots$$

with limit  $K_i(A)$  for  $i = 0, 1$  (as an ordered group for  $i = 0$ ).

First, we find a homomorphism  $\zeta_1$  such that the diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbf{Z}^{l_1} & \xrightarrow{\iota} & \mathbf{Z}^{l_1} \oplus \mathbf{Z}^{l_1} & \xrightarrow{\rho} & \mathbf{Z}^{l_1} \rightarrow 0 \\ & & \bar{\xi}_1^1 \downarrow & & \zeta_1 \downarrow & & \downarrow \bar{\xi}_1^0 \\ 0 & \rightarrow & K_1(A) & \longrightarrow & E & \longrightarrow & K_0(A) \rightarrow 0 \end{array}$$

is commutative. Set  $k_1 = l_1, m(1) = 2$  and  $\theta_1^i = \bar{\xi}_1^i$ ; suppose that we have an increasing sequence  $\{k_1, \dots, k_n\}$  of positive integers and commutative diagrams:

$$(3) \quad \begin{array}{ccccccc} \mathbf{Z}^{l_{m(1)}} & \longrightarrow & \mathbf{Z}^{l_{m(2)}} & \longrightarrow & \dots & \longrightarrow & \mathbf{Z}^{l_{m(n)}} \longrightarrow \dots \\ \nearrow \gamma_1^i & \delta_1^i \downarrow & \nearrow \gamma_2^i & \delta_2^i \downarrow & \dots & & \nearrow \gamma_n^i \\ \mathbf{Z}^{k_1} & \xrightarrow{\chi_1^i} & \mathbf{Z}^{k_2} & \xrightarrow{\chi_2^i} & \mathbf{Z}^{k_3} & \longrightarrow \dots & \longrightarrow \mathbf{Z}^{k_n} \end{array}$$

and

$$\begin{array}{ccccccc} \mathbf{Z}^{k_1} \oplus \mathbf{Z}^{k_1} & \xrightarrow{\psi_1} & \mathbf{Z}^{k_2} \oplus \mathbf{Z}^{k_2} & \xrightarrow{\psi_2} & \dots & \longrightarrow & \mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n} \\ \downarrow E & = & \downarrow E & = & \dots & = & \downarrow E \end{array}$$

and

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbf{Z}^{k_j} & \xrightarrow{\iota} & \mathbf{Z}^{k_j} \oplus \mathbf{Z}^{k_j} & \xrightarrow{\rho} & \mathbf{Z}^{k_j} \longrightarrow 0 \\ & & \theta_j^1 \downarrow & & \zeta_j \downarrow & & \downarrow \theta_j^0 \\ 0 & \longrightarrow & K_0(A) & \longrightarrow & E & \longrightarrow & K_1(A) \longrightarrow 0 \end{array}$$

for  $j = 1, 2, \dots, n$ , where  $\psi_j$  is of the form  $\psi_j(a, b) = (\chi_j^1(a) + \psi_j^1(b), \chi_j^0(b))$ , and  $\theta_j^i = \bar{\xi}_{m(j)}^i \circ \gamma_j^i$ .

Choose  $a_i \in E$  such that  $q(a_i) = \bar{\xi}_{m(n)}^{-0}(e_i)$ . By the commutativity of the diagram

$$\begin{array}{ccccc} \mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n} & \xrightarrow{\rho} & \mathbf{Z}^{k_n} & \xrightarrow{\gamma_n^0} & \mathbf{Z}^{l_{m(n)}} \\ \zeta_n \downarrow & & \theta_n^0 \downarrow & & \downarrow \bar{\xi}_{m(n)}^0 \\ E & \xrightarrow{q} & K_0(A) & = & K_0(A) \end{array}$$

we have for  $i = 1, \dots, k_n$

$$\zeta_n(0, e_i) - \sum_j \gamma_n^0(j, i) a_j \in \iota(K_1(A)).$$

Since  $K_1(A) = \bigcup_n \text{im } \bar{\xi}_n^{-1}$ , these elements are contained in  $\iota(\text{im } \bar{\xi}_{m(n+1)}^{-1})$  for some  $m(n+1) > m(n)$ . Let  $k_{n+1} = l_{m(n)} l_{m(n+1)}$ , and index the coordinates in  $\mathbf{Z}^{k_{n+1}}$  by  $(s, t)$ ,  $s = 1, \dots, l_{m(n+1)}$ ,  $t = 1, \dots, l_{m(n)}$ . Define  $\delta_n^i : \mathbf{Z}^{l_{m(n)}} \rightarrow \mathbf{Z}^{k_{n+1}}$  by

$$\delta_n^1((s, t), t') = \begin{cases} \xi_{m(n+1)m(n)}^1(s, t) & \text{if } t = t', \\ 0 & \text{otherwise,} \end{cases}$$

and

$$\delta_n^0((s, t), t') = \begin{cases} 1 & \text{if } t = t', \\ 0 & \text{otherwise.} \end{cases}$$

Define  $\gamma_{n+1}^i : \mathbf{Z}^{k_{n+1}} \rightarrow \mathbf{Z}^{l_{m(n+1)}}$  by

$$\gamma_{n+1}^1(s', (s, t)) = \begin{cases} 1 & \text{if } s = s', \\ 0 & \text{otherwise,} \end{cases}$$

and

$$\gamma_{n+1}^0(s', (s, t)) = \begin{cases} \xi_{m(n+1)m(n)}^0(s, t) & \text{if } s = s', \\ 0 & \text{otherwise.} \end{cases}$$

These are defined so that  $\gamma_{n+1}^i \circ \delta_n^i = \xi_{m(n+1)m(n)}$ , the quotient  $\mathbf{Z}^{k_{n+1}} / \text{im } \delta_n^0$  has no torsion, and  $\gamma_{n+1}^1$  is surjective. Diagram (3) may now be extended in the following

way:

$$(4) \quad \begin{array}{ccccccc} & & & \longrightarrow & \mathbf{Z}^{l_{m(n)}} & \longrightarrow & \mathbf{Z}^{l_{m(n+1)}} & \longrightarrow & \dots \\ & & & & \nearrow \gamma_n^i & \delta_n^i \downarrow & \nearrow \gamma_{n+1}^i & & \\ \dots & \longrightarrow & \mathbf{Z}^{k_n} & \xrightarrow{\chi_n^i} & \mathbf{Z}^{k_{n+1}} & & & & \end{array}$$

where  $\chi_n^i = \delta_n^i \gamma_n^i$ .

Since  $\delta_n^0(e_t) = \sum_s e_{(s,t)}$  and thus  $\bar{\xi}_{m(n)}^0(e_t) = \theta_{n+1}^0(\sum_s e_{(s,t)})$  (where  $\theta_{n+1}^i = \bar{\xi}_{m(n+1)}^i \circ \gamma_{n+1}^i$ ), there exist  $a_{(s,t)} \in E$  such that  $a_t = \sum_s a_{(s,t)}$  and  $q(a_{(s,t)}) = \theta_{n+1}^0(e_{(s,t)})$ . Since

$$\bar{\xi}_{m(n+1)}^1(\mathbf{Z}^{l_{m(n+1)}}) = \theta_{n+1}^1(\mathbf{Z}^{k_{n+1}})$$

and

$$\chi_n^0((s,t), i) = \delta_n^0((s,t), t) \gamma_n^0(t, i) = \gamma_n^0(t, i),$$

we obtain

$$\zeta_n(0, e_i) - \sum_t \gamma_n^0(t, i) \sum_s a_{(s,t)} = \zeta_n(0, e_i) - \sum_{s,t} \chi_n^0((s,t), i) a_{(s,t)} \in \theta_{n+1}^1(\mathbf{Z}^{k_{n+1}}).$$

Thus there exist  $(\psi_n^1((s,t), i))$  such that the left hand side equals

$$\sum_{s,t} \psi_n^1((s,t), i) \theta_{n+1}^1(e_{(s,t)}).$$

Define  $\psi_n : \mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n} \rightarrow \mathbf{Z}^{k_{n+1}} \oplus \mathbf{Z}^{k_{n+1}}$  by

$$\psi_n(a, b) = (\chi_n^1(a) + \psi_n^1(b), \chi_n^0(b))$$

and define  $\zeta_{n+1} : \mathbf{Z}^{k_{n+1}} \oplus \mathbf{Z}^{k_{n+1}} \rightarrow E$  by

$$\zeta_{n+1}(e_{(s,t)}, e_{(s',t')}) = \iota \circ \theta_{n+1}^1(e_{(s,t)}) + a_{(s',t')}.$$

Then one checks that

$$\begin{array}{ccc} \mathbf{Z}^{k_n} \oplus \mathbf{Z}^{k_n} & \xrightarrow{\psi_n} & \mathbf{Z}^{k_{n+1}} \oplus \mathbf{Z}^{k_{n+1}} \\ \zeta_n \downarrow & & \zeta_{n+1} \downarrow \\ E & = & E \end{array}$$

and

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbf{Z}^{k_{n+1}} & \xrightarrow{\iota} & \mathbf{Z}^{k_{n+1}} \oplus \mathbf{Z}^{k_{n+1}} & \xrightarrow{\rho} & \mathbf{Z}^{k_{n+1}} & \rightarrow & 0 \\ & & \theta_{n+1}^1 \downarrow & & \zeta_{n+1} \downarrow & & \downarrow \theta_{n+1}^0 & & \\ 0 & \rightarrow & K_1(A) & \longrightarrow & E & \longrightarrow & K_0(A) & \rightarrow & 0 \end{array}$$

are commutative.

For example, since

$$\begin{aligned} \psi_n(e_i, 0) &= \left( \sum_{s,t} \chi_n^1((s,t), i) e_{(s,t)}, 0 \right), \\ \chi_n^1((s,t), i) &= \xi_{m(n+1)m(n)}^1(s,t) \gamma_n^1(t, i), \\ \theta_{n+1}^1(e_{(s,t)}) &= \bar{\xi}_{m(n+1)}^1(e_s), \end{aligned}$$

it follows that

$$\begin{aligned} \zeta_{n+1} \circ \psi_n(e_i, 0) &= \sum_{s,t} \chi_n^1((s,t), i) \iota \circ \theta_{n+1}^1(e_{(s,t)}) = \sum_t \gamma_n^1(t, i) \iota \circ \bar{\xi}_{m(n)}^1(e_t) \\ &= \iota \circ \bar{\xi}_{m(n)}^1 \circ \gamma_n^1(e_i) = \iota \circ \theta_n^1(e_i) = \zeta_n(e_i, 0). \quad \square \end{aligned}$$

3.11. *Proof of the theorem for the case  $i = 0$ .* By Lemma 3.10 we may assume that the extension

$$0 \rightarrow K_1(A) \rightarrow E \rightarrow K_0(A) \rightarrow 0$$

is obtained as the inductive limit of the sequence

$$\begin{array}{ccccccc} 0 & \rightarrow & \mathbf{Z}^{k_1} & \xrightarrow{\iota} & \mathbf{Z}^{k_1} \oplus \mathbf{Z}^{k_1} & \xrightarrow{\rho} & \mathbf{Z}^{k_1} \rightarrow 0 \\ & & \chi_1^1 \downarrow & & \chi_1^1 \downarrow \swarrow \psi_1^1 \downarrow \chi_1^0 & & \downarrow \chi_1^0 \\ 0 & \rightarrow & \mathbf{Z}^{k_2} & \xrightarrow{\iota} & \mathbf{Z}^{k_2} \oplus \mathbf{Z}^{k_2} & \xrightarrow{\rho} & \mathbf{Z}^{k_2} \rightarrow 0 \\ & & \downarrow & & \downarrow \swarrow \downarrow & & \downarrow \\ \vdots & & \vdots & & \vdots \quad \vdots \quad \vdots & & \vdots \quad \vdots \end{array}$$

By passing to a subsequence we may assume that  $M(\chi_{n+1}^0) \geq \max(2L(\chi_n^1), 4)$ ; this will ensure that the inductive limit algebra to be constructed has real rank zero. As in 1.6, we construct an inductive system for  $A$  with

$$A_n = \bigoplus_{k=1}^{k_n} M_{[n,k]} \otimes C(\mathbf{T})$$

and  $\varphi_n : A_n \rightarrow A_{n+1}$  so that  $\varphi_n$  induces  $\chi_n^i : K_i(A_n) \rightarrow K_i(A_{n+1})$  and the partial embeddings are standard embeddings of type  $(\chi_n^0(i, j), \chi_n^1(i, j))$ .

We define  $\alpha$  using a sequence of unitaries  $u_n \in A_n$  chosen recursively such that

$$[\varphi_n(p_{n,j})u_{n+1}p_{n+1,i} \varphi_n(u_n^*p_{n,j})]_1 = [n, j]\psi_n^1(i, j),$$

where  $[\cdot]_1$  denotes the class of a unitary in

$$K_1(\varphi_n(p_{n,j})A_{n+1,i}\varphi_n(p_{n,j})) \cong \mathbf{Z}$$

(see 2.3). Set  $q_{ij}^{n+1} = \varphi_n(p_{n,j})p_{n+1,i}$  and put

$$\begin{aligned} A_{ij}^{n+1} &= q_{ij}^{n+1}A_{n+1}q_{ij}^{n+1}, \\ B_{ij}^{n+1} &= q_{ij}^{n+1}\varphi_n(A_n)' \cap A_{n+1}q_{ij}^{n+1}; \end{aligned}$$

one checks that, under the identification  $K_1(A_{ij}^{n+1}) \cong \mathbf{Z}$ , the range of the map  $K_1(B_{ij}^{n+1}) \rightarrow K_1(A_{ij}^{n+1})$  (induced by the embedding  $B_{ij}^{n+1} \subset A_{ij}^{n+1}$ ) is  $[n, j]\mathbf{Z}$ . Set  $u_1 = 1$ ; given  $u_n$ , construct  $u_{n+1}$  as follows. Choose a unitary  $w_{ij}^{n+1} \in B_{ij}^{n+1} \subset A_{ij}^{n+1}$  such that  $[w_{ij}^{n+1}]_1 = [n, j]\psi_n^1(i, j)$ . Set  $u_{n+1} = (\bigoplus w_{ij}^{n+1})u_n$ ; since  $\bigoplus w_{ij}^{n+1} \in A_{n+1} \cap \varphi_n(A_n)'$ , one has  $\alpha_{n+1} \circ \varphi_n = \varphi_n \circ \alpha_n$  (where  $\alpha_n = \text{Ad } u_n$ ). Then  $\alpha = \lim \text{Ad } \bar{\varphi}_n(u_n)$  has the desired properties.  $\square$

**3.12. Corollary.** *Let  $A$  be a simple unital AT algebra of real rank zero and  $g \in KK(A, A)$  such that  $\gamma(g) = 0$ . Then there is an approximately inner automorphism  $\alpha$  of  $A$  such that  $g = 1 - [\alpha]$ .*

*Proof.* This follows immediately from the preceding theorem and the universal coefficient theorem (see [RS, 1.17]).  $\square$

**3.13. Corollary.** *Let  $A$  and  $B$  be simple unital AT algebras of real rank zero and  $g$  be an invertible element in  $KK(A, B)$  such that  $\gamma(g)$  preserves positivity and the class of the unit. Then there is an isomorphism  $\varphi : A \rightarrow B$  such that  $g = [\varphi]$ .*

*Proof.* Since  $g$  is invertible,  $\gamma(g)_i \in \text{Hom}(K_i(A), K_i(B))$  is an isomorphism for  $i = 0, 1$ ; further,  $\gamma_0([1_A]) = [1_B]$  and  $\gamma_0(K_0(A)^+) \subset K_0(B)^+$ . Hence, by Elliott's classification theorem (cf. [E, 7.1]) there is an isomorphism,  $\psi : A \rightarrow B$ , such that  $\gamma(g) = \gamma([\psi])$ . By the preceding corollary it follows that there is an approximately inner automorphism  $\alpha$  such that  $g = [\psi \circ \alpha]$ ; set  $\varphi = \psi \circ \alpha$ .  $\square$

3.14. *Remark.* Let  $\text{Inn}(A)$  denote the group of inner automorphisms of a  $C^*$ -algebra  $A$  and let  $\overline{\text{Inn}}(A)$  denote its closure, the group of approximately inner automorphisms of  $A$ ; further, let  $\overline{\text{Inn}}_0(A)$  denote the subgroup of approximately inner automorphisms which can be approximated by automorphisms of the form  $\text{Ad } u$  with  $u$  in the connected component of the unitary group. In [ER, 4.5] Elliott and Rørdam show that for  $A$  a simple unital  $AT$  algebra of real rank zero one has,

$$(5) \quad \overline{\text{Inn}}(A)/\overline{\text{Inn}}_0(A) \cong \varprojlim K_1(A)/n_j K_1(A)$$

where  $n_j$  ranges over the directed set of divisors of  $[1]$  in  $K_0(A)$  (we assume that  $n_j | n_{j+1}$ ) and the isomorphism commutes with the canonical homomorphism from  $K_1(A)$  to both groups. They ask whether  $\overline{\text{Inn}}_0(A)$  must be arc-connected. By construction the  $\alpha$  we use in the proof of Theorem 3.1 for the case  $i = 1$  (see 3.8) is in  $\overline{\text{Inn}}_0(A)$ . Thus, to answer the question in the negative it suffices to find a simple unital  $AT$  algebra of real rank zero for which  $\text{Ext}(K_1(A), K_0(A)) \neq 0$ ; there is then an  $\alpha \in \overline{\text{Inn}}_0(A)$  for which  $\eta_1(\alpha) \neq 0$  (note that  $\eta_1$  is a homotopy invariant). It is also possible that  $\eta_0(\alpha) \neq 0$  for such  $\alpha$ , as shown below.

An outer form of the Elliott-Rørdam invariant for  $\alpha \in \overline{\text{Inn}}(A)$  may be obtained from  $\eta_0(\alpha)$  as follows: taking the quotient of both sides of the above isomorphism (5) by the image of  $K_1(A)$ , one obtains

$$\begin{aligned} (\overline{\text{Inn}}(A)/\text{Inn}(A))/(\overline{\text{Inn}}_0(A)/\overline{\text{Inn}}_0(A) \cap \text{Inn}(A)) &\cong (\varprojlim K_1(A)/n_j K_1(A))/K_1(A) \\ &\cong \varprojlim^1 n_j K_1(A). \end{aligned}$$

With  $n_j$  as above, let  $D$  denote the inductive limit  $\lim D_j$ , where  $D_j = \mathbf{Z}$  and the map  $D_j \rightarrow D_{j+1}$  is given by multiplication by  $n_{j+1}/n_j$ ; we identify  $D$  with the divisible hull of  $\mathbf{Z}[1_A]$  in  $K_0(A)$  via the map  $\iota : D \rightarrow K_0(A)$ . It follows that

$$\varprojlim^1 n_j K_1(A) \cong \varprojlim^1 \text{Hom}(D_j, K_1(A)) \cong \text{Ext}(D, K_1(A)).$$

With identifications as above the Elliott-Rørdam invariant for  $\alpha$  is  $\iota^*(\eta_0(\alpha))$ .

## REFERENCES

- [B] B. Blackadar, *K-theory for operator algebras*, MSRI Publications **5**, Springer-Verlag, New York, 1986. MR **88g**:46082
- [BBEK] B. Blackadar, O. Bratteli, G. A. Elliott and A. Kumjian, *Reduction of real rank in inductive limits of  $C^*$ -algebras*, Math. Ann. **292** (1992) 111–126. MR **93a**:46112
- [D] V. Deaconu, *A path model for circle algebras*, J. Operator Theory **34** (1995) 57–89. MR **96m**:46113
- [E] G. A. Elliott, *On the classification of  $C^*$ -algebras of real rank zero*, J. Reine Angew. Math. **443** (1993) 179–219. MR **94i**:46074
- [ER] G. A. Elliott and M. Rørdam, *The automorphism group of the irrational rotation  $C^*$ -algebra*, Commun. Math. Phys. **155** (1993) 3–26. MR **94j**:46059
- [Ex] R. Exel, *The soft torus and applications to almost commuting matrices*, Pacific J. Math. **160** (1993) 207–217. MR **94f**:46091
- [EL] R. Exel and T. Loring, *Invariants of almost commuting matrices*, J. Funct. Anal. **95** (1991) 364–376. MR **92a**:46083

- [Ro] J. E. Roos, *Sur les foncteurs dérivés de  $\varprojlim$ . Applications*, C. R. Acad. Sci. Paris **252** (1961) 3702–3704. MR **24**:A1938
- [RS] J. Rosenberg and C. Schochet, *The Künneth theorem and the universal coefficient theorem for Kasparov's generalized K-functor*, Duke Math. J. **55** (1987) 431–474. MR **88i**:46091
- [Rø] M. Rørdam, *Classification of certain infinite simple  $C^*$ -algebras*, J. Funct. Anal. **131** (1995) 415–458. MR **96e**:46080a

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO 060 JAPAN  
*E-mail address*: `kishi@math.sci.hokudai.ac.jp`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEVADA, RENO, NEVADA 89557  
*E-mail address*: `alex@unv.edu`