

ON THE STRUCTURE OF WEIGHT MODULES

IVAN DIMITROV, OLIVIER MATHIEU, AND IVAN PENKOV

ABSTRACT. Given any simple Lie superalgebra \mathfrak{g} , we investigate the structure of an arbitrary simple weight \mathfrak{g} -module. We introduce two invariants of simple weight modules: the shadow and the small Weyl group. Generalizing results of Fernando and Futorny we show that any simple module is obtained by parabolic induction from a cuspidal module of a Levi subsuperalgebra. Then we classify the cuspidal Levi subsuperalgebras of all simple classical Lie superalgebras and of the Lie superalgebra $\mathbf{W}(n)$. Most of them are simply Levi subalgebras of \mathfrak{g}_0 , in which case the classification of all finite cuspidal representations has recently been carried out by one of us (Mathieu). Our results reduce the classification of the finite simple weight modules over all classical simple Lie superalgebras to classifying the finite cuspidal modules over certain Lie superalgebras which we list explicitly.

INTRODUCTION

In order to be able to explain the topic of this paper and to state the results, we need to start with a few definitions. The remaining definitions needed are given in section 1. Some of the results apply to arbitrary finite dimensional Lie superalgebras and will be stated in this way.

Consider a finite dimensional Lie superalgebra $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ defined over an algebraically closed field of characteristic zero. Let $\mathfrak{h} = \mathfrak{h}_0 \oplus \mathfrak{h}_1$ be a Cartan subsuperalgebra and Q be the root lattice. A \mathfrak{g} -module M is called a *generalized weight module* if $M = \bigoplus_{\lambda} M^{(\lambda)}$, where for $\lambda \in \mathfrak{h}_0^*$, $M^{(\lambda)}$ denotes the maximal subspace of M on which $h - \lambda(h)$ acts locally nilpotently for any $h \in \mathfrak{h}_0$.

By a *triangular decomposition* T of \mathfrak{g} we mean a decomposition $\mathfrak{g} = \mathfrak{g}_T^+ \oplus \mathfrak{g}_T^0 \oplus \mathfrak{g}_T^-$ such that there exists a linear map $l : Q \rightarrow \mathbf{Z}$ for which $\mathfrak{g}_T^+ = \bigoplus_{l(\alpha) > 0} \mathfrak{g}^{(\alpha)}$, $\mathfrak{g}_T^0 = \bigoplus_{l(\alpha) = 0} \mathfrak{g}^{(\alpha)}$ and $\mathfrak{g}_T^- = \bigoplus_{l(\alpha) < 0} \mathfrak{g}^{(\alpha)}$. This requirement is weaker than the usual one, see for instance [PS]. Correspondingly, we will call the subsuperalgebra \mathfrak{g}_T^0 a *Levi subsuperalgebra of \mathfrak{g}* , or the *Levi component of T* , which will sometimes be in contradiction with the accepted terminology in the theory of Lie algebras.

Any simple generalized weight \mathfrak{g}_T^0 -module Ω can be viewed as a $\mathfrak{g}_T^0 \oplus \mathfrak{g}_T^+$ -module with a trivial action of \mathfrak{g}_T^+ . Let $M_T(\Omega) = \text{ind}(\mathfrak{g}_T^0 \oplus \mathfrak{g}_T^+, \mathfrak{g}; \Omega)$ be the induced module. It is easy to prove that $M_T(\Omega)$ has a unique simple quotient, denoted by $L_T(\Omega)$. When the decomposition T is proper (i.e. when $\mathfrak{g}_T^0 \neq \mathfrak{g}$), we call the \mathfrak{g} -module $L_T(\Omega)$ *parabolically induced*. A *cuspidal \mathfrak{g} -module* is by definition a simple generalized weight module which is not parabolically induced.

Received by the editors October 8, 1997.
2000 *Mathematics Subject Classification*. Primary 17B10.

Let M be any simple generalized weight module. We associate to M an invariant, called its shadow. This is a triple $(\mathfrak{g}_M^+, \mathfrak{g}_M^I, \mathfrak{g}_M^-)$ consisting of three Lie subsuperalgebras defined in section 3. We call a triangular decomposition T of \mathfrak{g} M -adapted if its Levi component is \mathfrak{g}_M^I and $\mathfrak{g}_M^\pm \subset \mathfrak{g}_T^\pm$. The first step of the classification of simple generalized weight modules is the following basic statement (see Theorem 3.6 below):

(T1) *For any simple generalized weight \mathfrak{g} -module M , there is at least one M -adapted triangular decomposition T and a cuspidal \mathfrak{g}_T^0 -module Ω such that $M \simeq L_T(\Omega)$.*

In general, there are many pairs (T, Ω) such that $M \simeq L_T(\Omega)$. To get a more precise statement, one can assume that \mathfrak{g}_0 is reductive and define certain canonical sets \mathcal{C} of good triangular decompositions (see section 6). For the simple classical Lie superalgebras there are explicit descriptions of such canonical sets. To any simple weight module M we attach then a subgroup W_M of the Weyl group W of \mathfrak{g}_0 , called its small Weyl group, see section 5. The following stronger version of (T1) provides an explicit reduction of the classification of simple weight modules to the classification of cuspidal modules.

(T2) *For any simple generalized weight \mathfrak{g} -module M , there is $T \in \mathcal{C}$ and a cuspidal \mathfrak{g}_T^0 -module Ω such that $M \simeq L_T(\Omega)$. Moreover the pair (T, Ω) is unique up to W_M -conjugacy.*

A generalized weight module is called *finite* if the dimensions of all its weight spaces are finite. The classification of the finite generalized weight modules over all simple Lie algebras is carried out in [M]. We complete the present paper by making the first steps towards classifying all finite generalized weight modules over the simple Lie superalgebras. More precisely, we define a Levi subsuperalgebra to be *cuspidal* if it admits a finite cuspidal module, and then we classify the cuspidal Levi subsuperalgebras of all classical simple Lie superalgebras. It turns out that most of them are simply Levi subalgebras of \mathfrak{g}_0 , and hence the corresponding cuspidal modules are already classified in [M]. In this way, we reduce the classification problem for the simple finite weight modules over all classical Lie superalgebras to the problem of finding all cuspidal modules over certain Lie superalgebras which are listed in section 8.

Here is a quick guide through the contents of the paper. In section 1 we fix the notations and section 2 is a brief introduction into the general facts about generalized weight modules. In section 3 we prove (T1). Section 4 contains a cohomological characterization of finite cuspidal modules. From section 5 and on we restrict ourselves to the superalgebras with reductive even part, and in section 5 we introduce the small Weyl group. In section 6 we prove (T2). In the final section 7 we present a complete list of cuspidal Levi subalgebras of all classical simple Lie superalgebras and of $W(n)$.

Remark. When \mathfrak{g} is a reductive Lie algebra and M is finite, the statement (T1) is due to S. Fernando, [Fe]. When \mathfrak{g} is a reductive Lie algebra without factors of type E_8 , it is due to V. Futorny, [Fu], and A. Cylke, V. Futorny and S. Ovsienko, [CFO]. Our general approach provides a simpler proof.

ACKNOWLEDGMENT

The support and hospitality of the International Erwin Schrödinger Institute (Vienna) in the Spring of 1996 (where this work was started) is gratefully acknowl-

edged. We thank G. Benkart, V. Futorny, F. Malikov and V. Serganova for very helpful comments on the subject of this paper. I.D. has been supported partially by NSF Grants DMS 9500755 and INT 9511943, O.M. has been supported by UA1 of the CNRS, and I.P. has been supported partially by NSF Grant DMS 9500755.

1. NOTATIONS AND CONVENTIONS

1.1. Lie superalgebras. All Lie superalgebras will be defined over an algebraically closed field K of characteristic 0 and will be assumed finite dimensional. All underlying vector spaces of Lie superalgebras and their representations are automatically assumed to be \mathbf{Z}_2 -graded, and the even and odd part of a \mathbf{Z}_2 -graded vector space V will be denoted by V_0 and V_1 . If $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ is a Lie superalgebra, a Lie subsuperalgebra $\mathfrak{k} \subset \mathfrak{g}$ will be called a *Lie subalgebra* iff $\mathfrak{k}_1 = 0$.

1.2. Cartan subsuperalgebras. A *Cartan subsuperalgebra* $\mathfrak{h}_0 \oplus \mathfrak{h}_1$ of a Lie superalgebra \mathfrak{g} is by definition a self normalizing nilpotent Lie subsuperalgebra of \mathfrak{g} . According to Scheunert, [Sch] (see also [PS]), $\mathfrak{h} = \mathfrak{h}_0 \oplus \mathfrak{h}_1$ is a Cartan subsuperalgebra iff \mathfrak{h}_0 is a Cartan subalgebra of \mathfrak{g}_0 and \mathfrak{h}_1 is the generalized weight space of weight 0 of the \mathfrak{h}_0 -module \mathfrak{g}_1 . It should be noted that the definition of a generalized weight module (see the Introduction) involves only the action of \mathfrak{h}_0 .

1.3. Roots $\Delta, \Delta_0, \Delta_1$. Let \mathfrak{g} be a Lie superalgebra and let \mathfrak{h} be a Cartan subsuperalgebra. For $i \in \mathbf{Z}_2$, we denote by Δ_i the set of all $\alpha \in \mathfrak{h}_0^*$ such that $\alpha \neq 0$ and $\mathfrak{g}_i^{(\alpha)} \neq 0$. Set $\Delta := \Delta_0 \cup \Delta_1$. The elements of Δ are the *roots* of \mathfrak{g} . A root α is *even* (respectively *odd*) if $\alpha \in \Delta_0$ (respectively $\alpha \in \Delta_1$). A root can be simultaneously even and odd. The *root lattice* Q is the subgroup of \mathfrak{h}^* generated by Δ .

1.4. Triangular decompositions. The definition of a triangular decomposition T of \mathfrak{g} , $\mathfrak{g} = \mathfrak{g}_T^+ \oplus \mathfrak{g}_T^0 \oplus \mathfrak{g}_T^-$, is given in the Introduction. Let $\Delta^T = \Delta_0^T \cup \Delta_1^T$ be the set of roots of \mathfrak{g}_T^0 . We will say that T is a *good triangular decomposition* iff

- (i) the monoid generated by Δ_0^T is a group (denoted by Q_0^T), and
- (ii) for any $\beta \in \Delta_1^T$, there is $m > 0$ so that $m\beta \in Q_0^T$.

In such a case, we will say that \mathfrak{g}_T^0 is a *good Levi subsuperalgebra* of \mathfrak{g} .

1.5. Dense and torsion-free modules. The *support* of a generalized weight module M is the set of all $\lambda \in \mathfrak{h}_0^*$ such that $M^{(\lambda)} \neq 0$, and it is denoted by $\text{supp } M$. We denote by $\text{inj } M$ the set of all roots $\alpha \in \Delta_0$ for which there is some $x \in \mathfrak{g}_0^{(\alpha)}$ such that x acts injectively on M .

We define M to be *torsion-free* if the monoid generated by $\text{inj } M$ is a subgroup of finite index in Q (Fernando's original definition, [Fe], requires that $\text{inj } M = \Delta$). We say that M is *dense* if $\text{supp } M$ is a finite union of Q' -cosets, for some subgroup Q' of finite index in Q (Futorny's original definition, [Fu], requires that $Q = Q'$). For simple generalized weight modules, we will show in section 3 that the notions of density and torsion-freeness are both equivalent to cuspidality. With the original definitions of density and torsion-freeness, the latter is true only for finite modules over reductive Lie algebras.

2. GENERALITIES ABOUT GENERALIZED WEIGHT MODULES

Let \mathfrak{g} be a Lie superalgebra with a fixed Cartan subsuperalgebra \mathfrak{h} and let U be the enveloping algebra of \mathfrak{g} . We denote by $\mathcal{GW}(\mathfrak{g})$ the category of all generalized weight \mathfrak{g} -modules and by $\mathcal{GW}_f(\mathfrak{g})$ the subcategory of finite generalized weight modules. Throughout the rest of the paper, M will denote a fixed generalized weight \mathfrak{g} -module, i.e. an object of $\mathcal{GW}(\mathfrak{g})$. M will not be assumed simple unless the contrary is stated explicitly. We have $U^{(\lambda)} \cdot M^{(\mu)} \subset M^{(\lambda+\mu)}$, for any $\lambda, \mu \in \mathfrak{h}_0^*$. In particular $U^{(0)}$ is a subalgebra of U and each $U^{(\lambda)}$ is a $U^{(0)}$ -bimodule.

Lemma 2.1. (i) *Let $\lambda \in Q$. Then $U^{(\lambda)}$ is finitely generated as a left as well as a right $U^{(0)}$ -module.*

(ii) *Let $M \in \mathcal{GW}(\mathfrak{g})$ be simple, let $\lambda \in \text{supp } M$ and let $\beta \in Q$. Then $M^{(\lambda)}$ is a simple $U^{(0)}$ -module and $U^{(\beta)} \cdot M^{(\lambda)} = M^{(\lambda+\beta)}$.*

Proof. (i) Set $H := \text{Hom}_{\mathbf{Z}}(Q, K^*)$. The torus H acts on U and on its associated graded algebra S . The generalized weight space decomposition under \mathfrak{h} is identical with its generalized weight space decomposition under H . Hence by Hilbert’s invariant theorem, $S^{(0)}$ is finitely generated and each $S^{(\lambda)}$ is a finitely generated $S^{(0)}$ -module. This implies that $U^{(\lambda)}$ is finitely generated as a left as well as a right $U^{(0)}$ -module.

(ii) Let $m \in M^{(\lambda)}$, $m \neq 0$. We have $U \cdot m = M$. Hence $U^{(\beta)} \cdot m = M^{(\beta+\lambda)}$, for every $\beta \in Q$. Therefore $U^{(\beta)} \cdot M^{(\lambda)} = M^{(\beta+\lambda)}$, and $M^{(\lambda)}$ is simple as a $U^{(0)}$ -module. □

A *weight module* is a module such that $M = \bigoplus_{\lambda} M^{\lambda}$, where M^{λ} is the common kernel of $h - \lambda(h)$ for all $h \in \mathfrak{h}_0$.

Proposition 2.2. *Let M be simple.*

- (i) *supp M is contained in a single Q -coset.*
- (ii) *If \mathfrak{g} is semi-simple as an \mathfrak{h}_0 -module, then M is a weight module.*
- (iii) *If $M^{(\lambda)}$ is finite dimensional for some $\lambda \in \text{supp } M$, then M is finite.*

Proof. Assertion (i) follows from Lemma 2.1, (ii). If \mathfrak{h}_0 acts semi-simply on \mathfrak{g} , $\bigoplus_{\lambda} M^{\lambda}$ is obviously a \mathfrak{g} -submodule. Thus M is a weight module and Assertion (ii) is proved. Let $\beta \in Q$. By Lemma 2.1,(i) there is a finite set $\{u_1, \dots, u_k\}$ generating the right $U^{(0)}$ -module $U^{(\beta)}$. Hence we have

$$M^{(\beta+\lambda)} = U^{(\beta)} \cdot M^{(\lambda)} = \sum_{1 \leq i \leq k} (u_i \cdot U^{(0)} \cdot M^{(\lambda)}) = \sum_{1 \leq i \leq k} (u_i \cdot M^{(\lambda)}).$$

Therefore $\dim M^{(\beta+\lambda)} \leq k \dim M^{(\lambda)} < \infty$, which proves Assertion (iii). □

Let T be a triangular decomposition of \mathfrak{g} and let Q^T be the root lattice of \mathfrak{g}_T^0 . For any object Ω of $\mathcal{GW}(\mathfrak{g}_T^0)$, set $M_T(\Omega) = \text{ind}(\mathfrak{g}_T^0 \mathfrak{g}_T^+, \mathfrak{g}; \Omega)$, Ω being viewed as a $\mathfrak{g}_T^0 \oplus \mathfrak{g}_T^+$ -module with a trivial action of \mathfrak{g}_T^+ .

Lemma 2.3. *Assume that the support of Ω is included in a single Q^T -coset.*

- (i) *The generalized weight \mathfrak{g} -module $M_T(\Omega)$ is finite iff Ω is finite.*
- (ii) *There exists a unique maximal \mathfrak{g} -submodule $Z_T(\Omega)$ in $M_T(\Omega)$ such that its intersection with Ω (considered as a $\mathfrak{g}_T^0 \oplus \mathfrak{g}_T^+$ -submodule in $M_T(\Omega)$) equals zero.*

Proof. Let $l : Q \rightarrow \mathbf{Z}$ be a linear map defining T and let U^- be the enveloping algebra of \mathfrak{g}_T^- . We have $U^- = \bigoplus_{n \geq 0} U^n$, where $U^n = \bigoplus_{l(\beta) = -n} (U^-)^{\beta}$. Set $M' =$

$\bigoplus_{n>0} U^n \otimes \Omega$. By Poincaré-Birkhoff-Witt's theorem, we have $M_T(\Omega) \simeq \Omega \oplus M'$ and the supports of Ω and M' are disjoint. Therefore any \mathfrak{g} -submodule having trivial intersection with Ω is a subspace of M' , and $Z_T(\Omega)$ is the sum of all those modules. Assertion (ii) follows. Assertion (i) follows from the fact that each U^n is finite dimensional. \square

For any object Ω of $\mathcal{GW}(\mathfrak{g}_T^0)$ whose support is included in a single Q^T -coset, we set $L_T(\Omega) := M_T(\Omega)/Z_T(\Omega)$. The correspondence $\Omega \rightsquigarrow L_T(\Omega)$ is a kind of Harish-Chandra induction functor. The corresponding Harish-Chandra restriction is the functor $M \rightsquigarrow H^0(\mathfrak{g}_T^+, M)$.

Corollary 2.4. *Let Ω be an object of $\mathcal{GW}(\mathfrak{g}_T^0)$ whose support is contained in a single Q^T -coset.*

- (i) $H^0(\mathfrak{g}_T^+, L_T(\Omega)) = \Omega$.
- (ii) $H_0(\mathfrak{g}_T^-, L_T(\Omega)) = \Omega$.
- (iii) Ω is simple iff $L_T(\Omega)$ is simple.
- (iv) Ω is finite iff $L_T(\Omega)$ is finite.

Proof. Let v be a generalized weight vector in $H^0(\mathfrak{g}_T^+, L_T(\Omega))$. If $v \notin \Omega$, then the support of the \mathfrak{g} -submodule of $L_T(\Omega)$ generated by v is disjoint from the support of Ω , which is a contradiction. This proves Assertion (i). The proof of Assertion (ii) is similar. Any non-zero submodule of $L_T(\Omega)$ intersects Ω non-trivially, and Ω generates $L_T(\Omega)$. Hence $L_T(\Omega)$ is simple whenever Ω is simple. For any \mathfrak{g}_T^0 -submodule $\Omega' \subset \Omega$, we have $L_T(\Omega') \subset L_T(\Omega)$, which proves also that Ω is simple whenever $L_T(\Omega)$ is simple. Assertion (iii) is proved. Assertion (iv) follows from Lemma 2.3,(ii) and from the fact that $\Omega \subset L_T(\Omega)$. \square

3. THE SHADOW AND A GENERALIZATION OF FERNANDO-FUTORNY'S THEOREM

Lemma 3.1. *Let M be simple and let $x \in \mathfrak{g}_0^\alpha$ for some $\alpha \in \Delta_0$. Then $x : M \rightarrow M$ is either injective or locally nilpotent.*

Proof. The set M' of all vectors $m \in M$ such that $x^N \cdot m = 0$ for some $N \geq 1$ is a \mathfrak{g} -submodule. If $M' = 0$, then x acts injectively. Otherwise $M' = M$ and x acts locally nilpotently. \square

We define a *cone* as a finitely generated submonoid of Q . For any cone C , its *saturation* \overline{C} is the cone of all $\lambda \in Q$ such that $m\lambda \in C$ for some positive integer m . Any cone contains a maximal subgroup $G(C)$ ($G(C)$ is nothing but $C \cap -C$) and we have $G(\overline{C}) = \overline{G(C)}$. For any M , we denote by C_M^1 the cone generated by $\text{inj } M$, and by C_M^2 the set of all $\beta \in Q$ such that $\beta + \text{supp } M \subset \text{supp } M$. It is obvious that $C_M^1 \subset C_M^2$.

Proposition 3.2. *Let M be simple. There is a finite set Θ such that $\text{supp } M = C_M^1 + \Theta$. In particular $\overline{C_M^1} = \overline{C_M^2}$.*

Proof. For any $i \in \mathbf{Z}_2$ and any $\alpha \in \Delta_i$, choose a basis $B_i^\alpha = \{x_{i,1}^\alpha, x_{i,2}^\alpha, \dots\}$ of \mathfrak{g}_i^α , with the requirement that $x_{i,1}^\alpha$ acts injectively on M whenever α belongs to $\text{inj } M$. Let $U(\alpha)$ be the span of all monomials $(x_{i,1}^\alpha)^{m_1} \cdot (x_{i,2}^\alpha)^{m_2} \cdot \dots$ where $i \in \mathbf{Z}_2$ is such that $\Delta_i \neq \emptyset$, and m_1, m_2, \dots are non-negative integers not exceeding 1 in case $i = 1$.

Set $j = \# \text{inj } M$, $a = \#\Delta_0$, $b = \#\Delta_1$ ($\#$ standing for cardinality) and write $\Delta_0 \cup \Delta_1 = \{\alpha_1, \alpha_2, \dots, \alpha_{a+b}\}$, where the first j roots are the roots in $\text{inj } M$ and

the first a roots are the roots in Δ_0 . Choose furthermore $\lambda \in \text{supp } M$, put $N := U(\alpha_{j+1}) \cdot U(\alpha_{j+2}) \cdot \dots \cdot U(\alpha_{a+b}) \cdot M^{(\lambda)}$ and let Θ be the support of the finite dimensional \mathfrak{h}_0 -module N . By Poincare-Birkhoff-Witt's theorem, we have $M = U(\alpha_1) \cdot U(\alpha_2) \cdot \dots \cdot U(\alpha_j) \cdot N$, and therefore $\text{supp } M = C_M^1 + \Theta$. This implies that the saturations of C_M^1 and C_M^2 are the same. \square

In what follows we will denote the saturation of the cones C_M^1 and C_M^2 by C_M and will call it the *cone of M* . Using the cone C_M (M being assumed simple) one can decompose Δ into four disjoint sets $\Delta_M^I, \Delta_M^F, \Delta_M^+$ and Δ_M^- , defined as follows:

$$\begin{aligned} \Delta_M^I &:= \{\alpha \in \Delta \mid \alpha \in C_M, -\alpha \in C_M\}, \\ \Delta_M^F &:= \{\alpha \in \Delta \mid \alpha \notin C_M, -\alpha \notin C_M\}, \\ \Delta_M^+ &:= \{\alpha \in \Delta \mid \alpha \notin C_M, -\alpha \in C_M\}, \\ \Delta_M^- &:= \{\alpha \in \Delta \mid \alpha \in C_M, -\alpha \notin C_M\}. \end{aligned}$$

The decomposition $\Delta = \Delta_M^I \sqcup \Delta_M^F \sqcup \Delta_M^+ \sqcup \Delta_M^-$ is by definition the M -decomposition of Δ . Recalling that if $\alpha \in \Delta$ and $\lambda \in \text{supp } M$, the α -string through λ is the set $\{x \in \mathbf{Q} \mid \lambda + x\alpha \in \text{supp } M\}$, one verifies

- Lemma 3.3.** (i) $\alpha \in \Delta_M^F$ iff the α -string through any $\lambda \in \text{supp } M$ is bounded;
 (ii) $\alpha \in \Delta_M^I$ iff the α -string through any $\lambda \in \text{supp } M$ is unbounded in both directions;
 (iii) $\alpha \in \Delta_M^+$ iff the α -string through any $\lambda \in \text{supp } M$ is bounded from above only;
 (iv) $\alpha \in \Delta_M^-$ iff the α -string through any $\lambda \in \text{supp } M$ is bounded from below only.

Corollary 3.4. If M is simple, the α -string through λ is bounded (respectively bounded from above only, bounded from below only, or unbounded in both directions) for every $\lambda \in \text{supp } M$ whenever it is bounded (respectively bounded from above only, bounded from below only, or unbounded in both directions) for some $\mu \in \text{supp } M$.

It is an immediate consequence from the definition of the M -decomposition that the following three subspaces of \mathfrak{g}

$$(*) \quad \mathfrak{g}_M^+ := \bigoplus_{\alpha \in \Delta_M^+} \mathfrak{g}^{(\alpha)}, \quad \mathfrak{g}_M^I := \mathfrak{h} \oplus \left(\bigoplus_{\alpha \in \Delta_M^I} \mathfrak{g}^{(\alpha)} \right), \quad \mathfrak{g}_M^- := \bigoplus_{\alpha \in \Delta_M^-} \mathfrak{g}^{(\alpha)}$$

are Lie subsuperalgebras of \mathfrak{g} . We define the *shadow of M onto \mathfrak{g}* (or simply the *shadow of M*) as the triple $(*)$. It is clear that the shadow of M reconstructs the M -decomposition of Δ and vice versa.

We are ready now to prove a generalization of Theorem 4.18 in [Fe] as well as of the main result of [CFO]. We start with

Lemma 3.5. Let $C \subset Q$ be a saturated cone and let $D \subset Q$ be a finite subset. There is a linear form $l : Q \rightarrow \mathbf{Z}$ such that $\ker l \cap (C \cup D) = G(C)$.

Proof. Exercise which uses the convexity of C . \square

Recalling (see the Introduction) that a triangular decomposition T of \mathfrak{g} is M -adapted if $\mathfrak{g}_M^I = \mathfrak{g}_T^0$ and $\mathfrak{g}_M^\pm \subset \mathfrak{g}_T^\pm$, we have

Theorem 3.6. Let M be a simple generalized weight \mathfrak{g} -module.

- (i) \mathfrak{g}_M^I is a good Levi subsuperalgebra of \mathfrak{g} .
 (ii) There is at least one and at most finitely many M -adapted triangular decompositions T .

(iii) For any adapted triangular decomposition T , there exists a unique cuspidal \mathfrak{g}_M^I -module Ω such that $M \simeq L_T(\Omega)$.

Proof. Set $\text{inv } M = \text{inj } M \cap \Delta_M^I$. Assertion (i) follows from the fact that the monoid generated by $\text{inv } M$ is the group $G(C_M^1)$, which is of finite index in the root lattice Q^T of \mathfrak{g}_M^I .

The existence of an M -adapted triangular decomposition follows from Lemma 3.3. There are only finitely many triangular decompositions, so only finitely many are M -adapted. Assertion (ii) is proved.

It remains to prove Assertion (iii). Let $l : Q \rightarrow \mathbf{Z}$ be a linear map defining the M -adapted triangular decomposition T and let X be the Q -coset containing $\text{supp } M$. There is a map $d : X \rightarrow \mathbf{Z}$ such that $d(\lambda + \beta) = d(\lambda) + l(\beta)$ for any $\lambda \in X, \beta \in Q$. By the definition of l we have $l(\beta) \leq 0$ for any $\beta \in C_M^1$. As $\text{supp } M = C_M^1 + \Theta$ for some finite set Θ , the function d is bounded from above on $\text{supp } M$. Set $\Omega = \bigoplus_{\beta \in Q^T} M^{(\lambda^0 + \beta)}$, where $\lambda^0 \in \text{supp } M$ is such that $d(\lambda^0)$ is maximal. The fact that $d(\lambda^0)$ is a maximum of d implies that $\mathfrak{g}_T^+ \cdot \Omega = 0$. Thus Ω is a $\mathfrak{g}_T^0 \oplus \mathfrak{g}_T^+$ -module with trivial action of \mathfrak{g}_T^+ . The inclusion $\Omega \subset M$ extends to a morphism of \mathfrak{g} -modules $\phi : M_T(\Omega) \rightarrow M$. As M is simple, ϕ is onto and $\phi(L_T(\Omega)) = 0$. Hence we have $M \simeq L_T(\Omega)$. By Corollary 2.4,(iii), Ω is simple. Finally, the support of Ω is invariant by $G(C_M^1)$ which is a subgroup of finite index in Q^I , hence Ω is cuspidal. The uniqueness of Ω follows from Corollary 2.4,(ii). \square

Corollary 3.7. For a simple generalized weight \mathfrak{g} -module M , the following assertions are equivalent:

- (i) M is cuspidal;
- (ii) $C_M = Q$;
- (iii) M is dense.

4. A COHOMOLOGICAL CHARACTERIZATION OF FINITE CUSPIDAL MODULES

The main result of this section, Proposition 4.3, is analogous to cohomological characterizations of cuspidal representations of p -adic groups (vanishing of the Jacquet functors) and of finite Chevalley groups (vanishing of Harish-Chandra restrictions). It enhances Theorem 3.6. Proposition 4.3 will not be used in the rest of the paper.

Throughout section 4, M is assumed to be finite. An *admissible* \mathfrak{g} -module (or \mathfrak{g}_0 -module) is by definition a finite generalized weight \mathfrak{g} -module (or \mathfrak{g}_0 -module) such that the dimensions of its generalized weight spaces are uniformly bounded.

Lemma 4.1. Let M be cuspidal.

- (i) Let $\alpha \in \text{inj } M$ and let $x \in \mathfrak{g}_0^{(\alpha)}$ act injectively on M . Then there is $y \in U(\mathfrak{g}_0)^{(-\alpha)}$ such that $x \cdot y$ acts bijectively on M .
- (ii) M is admissible.

Proof. By Theorem 3.6,(ii), C_M^1 is a group. Hence for some $\beta_1, \dots, \beta_k \in C_M^1, \alpha + \beta_1 + \dots + \beta_k = 0$. Choose elements $x_i \in \mathfrak{g}_0^{(\beta_i)}$ which act injectively on M , and set $y = x_1 \cdot \dots \cdot x_k$. Then $x \cdot y$ acts injectively on M and maps each generalized weight space to itself. Hence $x \cdot y$ acts bijectively on M . Therefore x itself acts bijectively and the function $\lambda \mapsto \dim M^{(\lambda)}$ is C_M^1 invariant. By Proposition 3.2, the support of M is a finite union of C_M^1 -cosets, hence M is admissible. \square

Lemma 4.2. *Let \mathfrak{u} be a nilpotent Lie superalgebra and let N be a \mathfrak{u} -module. Assume that \mathfrak{u}_0 contains an element x which acts bijectively on N . Then*

$$H^*(\mathfrak{u}, N) = H_*(\mathfrak{u}, N) = 0.$$

Proof. Set $\mathfrak{v} = \mathfrak{u}/(Kx)$ and let $\Lambda^*\mathfrak{v}$ be the exterior superalgebra of \mathfrak{v} . There is a spectral sequence, [HS], computing $H_*(\mathfrak{u}, N)$ and such that

$$E_{p,q}^1 = H_p(Kx, C^q\mathfrak{v} \otimes N).$$

As x acts bijectively on $C^q\mathfrak{v} \otimes M$, we have $E_{p,q}^1 = 0$. Hence $H_*(\mathfrak{u}, N) = 0$. The argument for the vanishing of $H^*(\mathfrak{g}, N)$ is similar. □

Proposition 4.3. *The following are equivalent:*

- (i) M is cuspidal;
- (ii) $H_0(\mathfrak{g}_T^+, M) = 0$ for every proper triangular decomposition T ;
- (iii) $H^0(\mathfrak{g}_T^+, M) = 0$ for every proper triangular decomposition T ;
- (iv) $H_*(\mathfrak{g}_T^+, M) = H^*(\mathfrak{g}_T^+, M) = 0$ for every proper triangular decomposition T .

Proof. Let T be a triangular decomposition. By Corollary 2.4, we have

$$H^0(\mathfrak{g}_T^+, L_T(\Omega)) = \Omega$$

for any simple object Ω of $\mathcal{GW}(\mathfrak{g}_T^0)$. Hence Assertion (iii) implies Assertion (i). Similarly, Assertion (ii) implies Assertion (i). Obviously Assertion (iv) implies Assertion (ii) and (iii).

Assume that Assertion (i) holds, and let $l : Q \rightarrow \mathbf{Z}$ be a linear function defining the triangular decomposition T . As the monoid generated by $\text{inj } M$ is a finite index subgroup of Q , we have $l(\alpha) > 0$ for some $\alpha \in \text{inj } M$. By Lemma 4.1, \mathfrak{g}_T^+ contains a generalized root vector acting bijectively on M . Therefore Lemma 4.2 implies Assertion (iv). □

5. THE SMALL WEYL GROUP

To be able to further strengthen the claim of Theorem 3.6 we will assume throughout sections 5 and 6 that \mathfrak{g}_0 is reductive. Let M be simple. By Proposition 2.2, (ii), M is automatically a weight module. By definition, the *small Weyl group of M* is the subgroup W_M of the Weyl group W of \mathfrak{g}_0 generated by the reflections s_α , where α runs over $\Delta_M^F \cap \Delta_0$. By Q_M^F and Q_M^I we denote respectively the saturations of the subgroups of Q generated by $\Delta_M^F \cap \Delta_0$ and Δ_M^I .

Lemma 5.1. (i) *The sets $\text{supp } M$ and C_M are W_M -invariant.*

- (ii) *Any $\beta \in Q_M^I$ is W_M -invariant.*
- (iii) $Q_M^F \cap C_M = 0$.

Proof. For $\alpha \in \Delta_M^F$, there is a subalgebra \mathfrak{s} in \mathfrak{g}_0 isomorphic to $\mathfrak{sl}(2)$ with roots $\pm\alpha$. Since M is locally finite as an \mathfrak{s} -module, Assertion (i) follows.

To prove Assertion (ii), fix $\alpha \in \Delta_M^F$ and $\beta \in Q_M^I$. As $\beta - s_\alpha(\beta)$ is a multiple of α and belongs to Q_M^I , we have $\beta - s_\alpha(\beta) = 0$. Hence β is W_M -invariant.

For $\alpha \in \Delta_M^F$, we have $\alpha + s_\alpha(\alpha) = 0$. Therefore $\sum_{w \in W_M} w(\beta) = 0$ for any $\beta \in Q_M^F$. Hence $Q_M^F \cap C_M$ is a subgroup of C_M , and $Q_M^F \cap C_M \subset Q_M^I$. But since Q_M^F contains no W_M -invariant vectors, we have $Q_M^F \cap Q_M^I = 0$ and therefore also $Q_M^F \cap C_M = 0$. □

Set $(\mathfrak{g}_0)_M^F := \mathfrak{h} \oplus (\bigoplus_{\alpha \in Q_M^F} \mathfrak{g}_0^\alpha)$, $(\mathfrak{g}_0)_M^+ := \mathfrak{g}_M^+ \cap \mathfrak{g}_0$, $(\mathfrak{g}_0)_M^I := \mathfrak{g}_M^I \cap \mathfrak{g}_0$ and $\mathfrak{p}(M) := ((\mathfrak{g}_0)_M^F + (\mathfrak{g}_0)_M^I) \oplus \mathfrak{g}_M^+$. For any Lie algebra \mathfrak{a} , put $\mathfrak{a}' := [\mathfrak{a}, \mathfrak{a}]$.

Corollary 5.2. (i) $(\mathfrak{g}_0)_M^F$ is a reductive Lie subalgebra of \mathfrak{g} , its root system is $\Delta_M^F \cap \Delta_0$ and its Weyl group is W_M .

(ii) The Lie algebras $((\mathfrak{g}_0)_M^F)'$ and $((\mathfrak{g}_0)_M^I)'$ commute.

(iii) $\mathfrak{p}(M)$ is a parabolic subalgebra of \mathfrak{g}_0 .

Proof. Assertion (i) is a direct consequence of the definition of $(\mathfrak{g}_0)_M^F$, and Assertions (ii) and (iii) follow from the fact that $Q_M^F \cap C_M = 0$, as proved in Lemma 5.1. □

For any triangular decomposition T , set $\mathfrak{p}(T) := (\mathfrak{g}_T^0 \oplus \mathfrak{g}_T^+) \cap \mathfrak{g}_0$ and note that $\mathfrak{p}(T)$ is a parabolic subalgebra of \mathfrak{g}_0 .

Proposition 5.3. Let T, T' be triangular decompositions, and let Ω and Ω' be respectively a cuspidal \mathfrak{g}_0^T -module and a cuspidal $\mathfrak{g}_0^{T'}$ -module so that $M = L_T(\Omega) \simeq L_{T'}(\Omega')$. Then the parabolic subalgebras $\mathfrak{p}(T)$ and $\mathfrak{p}(T')$ are conjugated under W_M .

Proof. We have $(\mathfrak{g}_0)_M^I \oplus (\mathfrak{g}_0)_M^+ \subset \mathfrak{p}(T) \subset \mathfrak{p}(M)$. It follows from Corollary 4.2 that the parabolic subalgebra $\mathfrak{p}(T)$ is determined by its intersection with $(\mathfrak{g}_0)_M^I$ which is a Borel subalgebra of $(\mathfrak{g}_0)_M^I$. As any two such intersections are conjugated by W_M , the Proposition is proved. □

6. REDUCTION OF THE CLASSIFICATION PROBLEM

Denote by \mathcal{G} the set of all good triangular decompositions T of \mathfrak{g} (with \mathfrak{g}_T^0 containing the fixed Cartan subsuperalgebra \mathfrak{h}), and let \mathcal{P} be the set of all parabolic subalgebras in \mathfrak{g}_0 which contain \mathfrak{h}_0 . We define a *canonical set of good triangular decompositions* as a W -invariant subset $\mathcal{C} \subset \mathcal{G}$ such that the map $T \mapsto \mathfrak{p}(T)$ is a bijection between \mathcal{C} and \mathcal{P} . For any classical simple Lie superalgebra one can define explicitly at least one canonical set of good triangular decompositions. This is an easy consequence of the classification of all good Levi subsuperalgebras given in the next section. Here are two examples. For instance, if $\mathfrak{g} = \mathfrak{psq}(n)$, (see section 7; in [K] $\mathfrak{psq}(n)$ is denoted by $Q(n-1)$), then $\Delta_0 = \Delta_1$ and therefore \mathcal{G} itself is a canonical set of good triangular decompositions. If $\mathfrak{g} = \mathfrak{sl}(m|n)$ (see [K]), there is a natural \mathbf{Z} -grading of \mathfrak{g} as $\mathfrak{g}^{-1} \oplus \mathfrak{g}^0 \oplus \mathfrak{g}^1$ (where $\mathfrak{g}^0 = \mathfrak{g}_0$), and for $m \neq n$ the set of all triangular decompositions T such that $\mathfrak{g}_T^+ \supset \mathfrak{g}_1$ is a canonical set of good triangular decompositions.

Assume now that \mathcal{C} is any fixed canonical set of triangular decompositions. Here is a stronger version of Theorem 3.6.

Theorem 6.1. Let M be simple.

(i) There is a triangular decomposition $T \in \mathcal{C}$ and a cuspidal \mathfrak{g}_T^0 -module Ω such that $M \simeq L_T(\Omega)$.

(ii) For any $T' \in \mathcal{C}$ and any cuspidal \mathfrak{g}_T^0 -module Ω' such that $M \simeq L_{T'}(\Omega')$, there is $w \in W_M$ for which $T' = T^w$, $\Omega' = \Omega^w$.

Proof. Theorem 6.1 follows from Theorem 3.6 and Proposition 5.3. □

Theorem 6.1 reduces the general classification problem to the problem of classifying cuspidal representations. The results of [M] together with Proposition 6.3 below provide a classification of finite cuspidal representations of \mathfrak{g} up to a finite indeterminacy.

The following lemma is proved in [M] (Lemma 3.3):

Lemma 6.2. *Any admissible \mathfrak{g}_0 -module has finite length.*

Setting $\Omega^e := \text{ind}(\mathfrak{g}_0, \mathfrak{g}; \Omega)$ for any \mathfrak{g}_0 -module Ω , we have now

Proposition 6.3. (i) *For any finite cuspidal \mathfrak{g}_0 -module Ω , Ω^e contains at least one and only finitely many non-isomorphic cuspidal submodules.*

(ii) *For any finite cuspidal \mathfrak{g} -module M , there is at least one and only finitely many non-isomorphic cuspidal \mathfrak{g}_0 -modules Ω such that $M \subset \Omega^e$.*

Proof. Note first that if Ω is a finite cuspidal \mathfrak{g}_0 -module, for any $\alpha \in \Delta_0$ there is an element $e_\alpha \in \mathfrak{g}_0^\alpha$ which acts bijectively on both Ω and Ω^e . Furthermore Ω^e is an admissible \mathfrak{g}_0 -module and is therefore of finite length by Lemma 6.2. Thus Ω^e may only have finitely many non-isomorphic irreducible \mathfrak{g} -submodules. Each of them is necessarily cuspidal since by Lemma 4.1 e_α acts bijectively on it. Assertion (i) is proved. The proof of Assertion (ii) is similar. □

7. A CLASSIFICATION OF CUSPIDAL LEVI SUBSUPERALGEBRAS

In this section, we will use Kac’s classification, [K], to classify all cuspidal Levi subsuperalgebras (see the Introduction) of all simple classical Lie superalgebras (including the exceptional ones) and of $\mathbf{W}(n)$. For the remaining series $\mathbf{S}(n)$, $\tilde{\mathbf{S}}(n)$, $\mathbf{H}(n)$ the problem is still open.

The classification is based on the following three remarks:

(i) By Theorem 3.6, any cuspidal Levi subsuperalgebra is good. Furthermore, the map $\mathfrak{s} \mapsto \mathfrak{s}_0$ is a bijection between the set of good Levi subsuperalgebras of \mathfrak{g} and the set of good Levi subalgebras of \mathfrak{g}_0 . The inverse map is described as follows. Let \mathfrak{s}_0 be a good Levi subalgebra of \mathfrak{g}_0 and let Q' be the saturation in Q of its root lattice. Set $\mathfrak{s}_1 = \bigoplus_{\beta \in Q'} \mathfrak{s}_1^{(\beta)}$. Then $\mathfrak{s}_0 \oplus \mathfrak{s}_1$ is the unique good Levi subsuperalgebra whose even part is \mathfrak{s}_0 .

(ii) Let \mathfrak{s} be a Levi subsuperalgebra such that \mathfrak{s}_0 is reductive. A result of Fernando (Theorem 5.2 in [Fe]) together with Proposition 6.3 implies that \mathfrak{s} is cuspidal iff all simple factors of \mathfrak{s}_0 are of type A or C .

(iii) If \mathfrak{g} is a classical simple Lie superalgebra or $\mathfrak{g} = \mathbf{W}(n)$, and \mathfrak{s} is a good Levi subsuperalgebra of \mathfrak{g} , then \mathfrak{s}_0 is reductive.

In the rest of this section, we present the list of all good Levi subsuperalgebras and the list of all cuspidal Levi subsuperalgebras. All simple Lie superalgebras which we do not define here are studied in detail in [K]. In addition, we will use the following notations:

- a *partition* \mathbf{n} of a positive integer n is a k -tuple of positive integers (n_1, \dots, n_k) with $\sum_i n_i = n$; the partition is *proper* iff $k \neq 1$;

- for a partition \mathbf{n} we set

$$\mathfrak{sl}(\mathbf{n}) := \{(g_1 + \dots + g_k) \in \mathfrak{gl}(\mathbf{n}) := \mathfrak{gl}(n_1) \oplus \dots \oplus \mathfrak{gl}(n_k) \mid \sum_i \text{tr}(g_i) = 0\},$$

$$\mathfrak{sq}(\mathbf{n}) := \{(g_1 + \dots + g_k) \in \mathfrak{q}(\mathbf{n}) := \mathfrak{q}(n_1) \oplus \dots \oplus \mathfrak{q}(n_k) \mid \sum_i \text{otr}(g_i) = 0\},$$

$$\mathfrak{psq}(\mathbf{n}) := \mathfrak{sq}(\mathbf{n})/K \cdot \text{id},$$

where $\mathfrak{q}(l)$ is the Lie superalgebra of block matrices with $l \times l$ -blocks

$$\alpha = \begin{pmatrix} A & B \\ B & A \end{pmatrix}$$

with arbitrary $l \times l$ -matrices A and B , and $\text{otr } \alpha := \text{tr } B$.

7.1. Let $\mathfrak{g} = \mathfrak{sl}(n|m)$, with $n > m$. Then $\mathfrak{g}_0 \simeq \mathfrak{sl}(n, m)$ and $\mathfrak{g}_1 \simeq U \otimes V^* \oplus U^* \otimes V$, where U and V are respectively the natural representations of $\mathfrak{gl}(n)$ and $\mathfrak{gl}(m)$. Any good Levi subsuperalgebra is a Lie subalgebra of \mathfrak{g}_0 , it is necessarily cuspidal, and moreover it is isomorphic to $\mathfrak{sl}(\mathbf{n} \cup \mathbf{m})$ where \mathbf{n} , \mathbf{m} are partitions of n and m respectively.

7.2. Let $\mathfrak{g} = \mathfrak{psl}(n|n)$. Then $\mathfrak{g}_0 \simeq \mathfrak{sl}(n) \oplus \mathfrak{sl}(n)$ and $\mathfrak{g}_1 \simeq U \otimes V^* \oplus U^* \otimes V$, where U and V are respectively the natural representations of the two factors $\mathfrak{sl}(n)$. Any good Levi subsuperalgebra is cuspidal, and moreover it is isomorphic either to \mathfrak{g} or to $\mathfrak{sl}(\mathbf{n}) \oplus \mathfrak{sl}(\mathbf{n}')$, \mathbf{n} and \mathbf{n}' being partitions respectively of n such that at least one of them is proper.

7.3. Let $\mathfrak{g} = \mathfrak{osp}(n|2m)$. Then $\mathfrak{g}_0 \simeq \mathfrak{o}(n) \oplus \mathfrak{sp}(2m)$ and $\mathfrak{g}_1 \simeq U \otimes V$, where U and V are respectively the natural representations of $\mathfrak{o}(n)$ and $\mathfrak{sp}(2m)$. Any good Levi subsuperalgebra is of the type $\mathfrak{osp}(n'|2m') \oplus \mathfrak{sl}(\mathbf{n}) \oplus \mathfrak{sl}(\mathbf{m})$, where $n - n'$ is even, \mathbf{n} and \mathbf{m} are respectively partitions of $n - n'$ and $m - m'$. Such a subsuperalgebra is cuspidal iff $n' \leq 6$. The latter follows from the fact that $\mathfrak{o}(k)$ is of type A or C for $k \leq 6$.

7.4. Let $\mathfrak{g} = \mathfrak{sp}(n)$, see [P] (in [K] $\mathfrak{sp}(n)$ is denoted by $P(n - 1)$). Then $\mathfrak{g}_0 \simeq \mathfrak{sl}(n)$ and $\mathfrak{g}_1 \simeq S^2V \oplus \wedge^2V^*$, where V is the natural representation of $\mathfrak{sl}(n)$. Any good Levi subsuperalgebra is cuspidal. If n is odd, all such subsuperalgebras are isomorphic to \mathfrak{g} or to $\mathfrak{sl}(\mathbf{n})$, where \mathbf{n} is a proper partition of n . If n is even, all such subsuperalgebras are isomorphic to \mathfrak{g} , to $\mathfrak{sl}(\mathbf{n})$ for a proper partition \mathbf{n} of n other than $(n/2, n/2)$, or to $\mathfrak{sl}(n/2|n/2)$.

7.5. Let $\mathfrak{g} = \mathfrak{psq}(n)$. Then $\mathfrak{g}_0 \simeq \mathfrak{sl}(n)$ and \mathfrak{g}_1 is isomorphic to the adjoint representation. Any good Levi subsuperalgebra cuspidal and is isomorphic to $\mathfrak{psq}(\mathbf{n})$, where \mathbf{n} is a partition of n .

7.6. Let $\mathfrak{g} = \mathbf{D}(2, 1, \alpha)$. Then $\mathfrak{g}_0 \simeq \mathfrak{sl}(2) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(2)$ and $\mathfrak{g}_1 \simeq V^1 \otimes V^2 \otimes V^3$, where V^1, V^2 and V^3 are respectively the natural representations of the three factors. Any good Levi subsuperalgebra is cuspidal and is isomorphic to \mathfrak{g} or to any proper Levi subalgebra of \mathfrak{g}_0 .

7.7. Let $\mathfrak{g} = \mathbf{G}(3)$. Then $\mathfrak{g}_0 \simeq \mathfrak{sl}(2) \oplus \mathbf{G}_2$ and $\mathfrak{g}_1 \simeq V \otimes U$, V being the natural representation of $\mathfrak{sl}(2)$ and U being the 7-dimensional \mathbf{G}_2 -module. Any good Levi subsuperalgebra is isomorphic to one of the following:

- (i) \mathfrak{g} ;
- (ii) $\mathfrak{gl}(1) \oplus \mathbf{G}_2$;
- (iii) a Levi subalgebra of \mathfrak{g}_0 which does not contain a short root of \mathbf{G}_2 ;
- (iv) a Levi subalgebra of \mathfrak{g}_0 which is isomorphic to $\mathfrak{sl}(2, 1, 1)$ and contains a long root of \mathbf{G}_2 ;
- (v) $\mathfrak{o}(4) \ltimes H_4$, where H_4 is the Heisenberg superalgebra of dimension $1|4$.

The good cuspidal Levi subsuperalgebras are the ones of types (iii), (iv) and (v).

7.8. Let $\mathfrak{g} = \mathbf{F}(4)$. Then $\mathfrak{g}_0 \simeq \mathfrak{sl}(2) \oplus \mathfrak{o}(7)$ and $\mathfrak{g}_1 \simeq V \otimes U$, V being the natural representation of $\mathfrak{sl}(2)$ and U being the 8-dimensional spin representation of $\mathfrak{o}(7)$. Any good Levi subsuperalgebra is isomorphic to one of the following:

- (i) \mathfrak{g} ;
- (ii) $\mathfrak{gl}(1) \oplus \mathfrak{o}(7)$;
- (iii) $\mathfrak{osp}(4|2)$;

(iv) a Levi subalgebra of \mathfrak{g}_0 which does not contain $\mathfrak{sl}(2) \oplus \mathfrak{sl}(2) \oplus \mathfrak{sl}(2)$.

The good cuspidal Levi subsuperalgebras are the ones of types (iii) and (iv).

7.9. Let $\mathfrak{g} = \mathbf{W}(n)$. Then $\mathfrak{gl}(n)$ is embedded canonically into \mathfrak{g}_0 . Any good Levi subsuperalgebra is simply a Levi subalgebra of $\mathfrak{gl}(n)$ and is necessarily cuspidal. Moreover, such a subalgebra is isomorphic to $\mathfrak{gl}(\mathbf{n})$ for an arbitrary partition \mathbf{n} of n .

8. CONCLUSION

The present paper, together with [M] provides the following classification results for simple finite weight \mathfrak{g} -modules.

8.1. A complete classification of simple finite weight modules over any reductive Lie algebra \mathfrak{g} . A simple finite module of \mathfrak{g} is isomorphic to a tensor product of simple finite weight modules over the simple components of \mathfrak{g} , and for a simple Lie algebra the classification is carried out explicitly in [M].

Moreover, if \mathfrak{g} is any finite dimensional Lie superalgebra with reductive even part, Proposition 6.3 reduces the classification of simple finite weight modules to a finite indeterminacy.

8.2. A complete classification of simple finite weight modules over $\mathfrak{sl}(m|n)$ with $m \neq n$ and over $\mathbf{W}(n)$. Indeed, if $\mathfrak{g} = \mathfrak{sl}(m|n)$ or $\mathfrak{g} = \mathbf{W}(n)$, there is a natural \mathbf{Z} -grading of \mathfrak{g} as $\mathfrak{g}^{-1} \oplus \mathfrak{g}^0 \oplus \mathfrak{g}^1 \oplus \dots$, where $\mathfrak{g}^0 \simeq \mathfrak{sl}(m, n)$ for $\mathfrak{g} = \mathfrak{sl}(m|n)$, and $\mathfrak{g}^0 \simeq \mathfrak{gl}(n)$ for $\mathfrak{g} = \mathbf{W}(n)$. Moreover, when $m \neq n$, the map $M \mapsto H^0(\mathfrak{g}^{-1}, M)$ is a bijection between all simple weight \mathfrak{g} -modules (modulo the parity change) and all simple weight \mathfrak{g}_0 -modules. Thus all simple finite weight \mathfrak{g} -modules are classified.

8.3. A complete classification of non-cuspidal simple finite weight modules over $\mathfrak{psl}(n|n)$, $\mathfrak{sp}(2k+1)$ and $\mathbf{D}(2, 1, \alpha)$. It follows from the results in sections 6 and 7. More generally, if \mathfrak{g} is any Lie superalgebra whose even part \mathfrak{g}_0 is reductive, we obtain a complete classification of all finite simple weight modules M with $\mathfrak{g}_M^I \subset \mathfrak{g}_0$.

In order to finish the classification of finite simple weight modules over all simple classical Lie superalgebras and over $\mathbf{W}(n)$, it is necessary to classify the finite cuspidal modules over $\mathfrak{psl}(n|n)$, $\mathfrak{osp}(n|2m)$ (with $n \leq 6$), $\mathfrak{psq}(n)$, $\mathbf{D}(2, 1, \alpha)$, $\mathbf{G}(3)$, and $\mathfrak{o}(4) \ltimes H_4$. For the remaining three Cartan-type series of simple Lie superalgebras $\mathbf{S}(n)$, $\tilde{\mathbf{S}}(n)$, $\mathbf{H}(n)$ one should first classify their cuspidal Levi subsuperalgebras and then try to complete the classification.

REFERENCES

- [CFO] A. Cylke, V. Futorny and S. Ovsienko, On the support of irreducible non-dense modules for finite-dimensional Lie algebras, *Canad. J. Math.* (to appear).
- [Fe] S. Fernando, Lie algebra modules with finite dimensional weight spaces I, *TAMS* 322 (1990), 757-781. MR **91c**:17006
- [Fu] V. Futorny, The weight representations of semisimple finite dimensional Lie algebras, Ph.D. Thesis, Kiev University, 1987.
- [HS] G. Hochschild, J.-P. Serre, Cohomology of Lie algebras, *Ann. Math.* 57 (1953), 591-603. MR **14**:943c
- [K] V. Kac, Lie superalgebras, *Adv. Math.* 26 (1977), 8-96. MR **58**:5803
- [M] O. Mathieu, Classification of irreducible weight modules, *Ann. Inst. Fourier* (to appear).
- [P] I. Penkov, Characters of strongly generic irreducible Lie superalgebra representations, *Internat. J. Math.* 9 (1998), 331-366. MR **99d**:17011

- [PS] I. Penkov, V. Serganova, Generic irreducible representations of finite-dimensional Lie superalgebras, *Internatl. J. Math.* 5 (1994), 389-419. MR **95c**:17015
- [Sch] M. Scheunert, Invariant supersymmetric multilinear forms and the Casimir elements of P -type Lie superalgebras, *J. Math. Phys.* 28 (1987), 1180-1191. MR **89b**:17008

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT RIVERSIDE, RIVERSIDE, CALIFORNIA 92521

Current address: Department of Mathematics, University of California, Los Angeles, California 90095-1555

E-mail address: dimitrov@math.ucla.edu

UNIVERSITÉ LOUIS PASTEUR, IRMA, 7 RUE RENÉ DESCARTES, 67000 STRASBOURG, FRANCE

E-mail address: mathieu@math.u-strasbg.fr

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT RIVERSIDE, RIVERSIDE, CALIFORNIA 92521

E-mail address: penkov@math.ucr.edu