

DEGREE OF STRATA OF SINGULAR CUBIC SURFACES

RAFAEL HERNÁNDEZ AND MARÍA J. VÁZQUEZ-GALLO

ABSTRACT. We determine the degree of some strata of singular cubic surfaces in the projective space \mathbf{P}^3 . These strata are subvarieties of the \mathbf{P}^{19} parametrizing all cubic surfaces in \mathbf{P}^3 . It is known what their dimension is and that they are irreducible. In 1986, D. F. Coray and I. Vainsencher computed the degree of the 4 strata consisting on cubic surfaces with a double line. To work out the case of isolated singularities we relate the problem with (stationary) multiple-point theory.

1. INTRODUCTION

The purpose of this work is to determine the degree of certain strata of singular cubic surfaces in \mathbf{P}_C^3 . In the nineteenth century L. Schläfli [26] and A. Cayley [6] classified the cubic surfaces in \mathbf{P}^3 according to the type of their singularities. We consider this classification with the point of view given by J. W. Bruce and C. T. C. Wall [5] 100 years later, based on the modern theory of singularities [2], [3].

Let us suppose that the cubic surfaces are irreducible and have only isolated singularities. Fixed a singular point P of a cubic surface C , there are 4 types of surfaces, according to the rank of a quadratic form associated to each one of them: surfaces with a conic node, a binode, a uninode and a triple point at P .

Inside each one of these classes, the different singularities presented by a surface C are determined by the possible coincidences of the 6 lines passing through the fixed singular point P and contained in C .

Besides of these, there are 4 strata of irreducible surfaces with non isolated singularities (that are, in fact, ruled surfaces with a double line) and 9 of reducible cubic surfaces.

The strata resulting from this classification is the closure of quasi-projective subvarieties of the \mathbf{P}^{19} parametrizing all cubic surfaces in \mathbf{P}^3 . Regarding geometric properties of strata, it is known what their dimension is [5] and that the strata are irreducible [4]. Our aim here is to determine their degree.

First of all, the degree of strata parametrizing reducible surfaces is easy to compute [30]. In the case of the 4 big strata given by the rank of a quadratic form, the degree can be obtained at least in two ways [30], [28]. On the other hand, in 1986, D. F. Coray and I. Vainsencher calculated the degree of the 4 strata of ruled cubic surfaces having a double line [8]. Their work motivates ours.

Received by the editors March 15, 1999.

2000 *Mathematics Subject Classification*. Primary 14N05, 14C17; Secondary 14C05, 14C15.

Key words and phrases. Enumerative geometry, strata of singular cubic surfaces, stationary multiple-point theory.

In this paper, we determine the degree of 7 of the 11 strata of cubic surfaces with a conic node, using a method that must work, essentially in the same way, with some strata of cubics with a binode or a uninode.

Our approach to the problem is to relate it with stationary multiple-point theory, whose main goal is the enumeration of k -tuples of points of a scheme Z with the same image under a morphism of schemes $f : Z \rightarrow S$ and such that the points have collapsed in accordance with a given partition of k .

We construct first some suitable parameter spaces and then we consider the lines whose coincidences determine the strata in our case, as stationary multiple points of a morphism between two of these parameter spaces.

One of the first examples of multiple-point formulas related with our problem is that of De Jonquières (1866), allowing us to consider coincidences of points in the zero scheme of sections of a line bundle on a given curve. One century later, I. Vainsencher [29] (1981) proved a generalized De Jonquières' formula, valid for a general line bundle. Some years after, S. J. Colley [7] obtained more formulas for stationary multiple points using the iteration method by S. L. Kleiman [16], [17], [18], who had already given some multiple-point formulas for the non-stationary cases.

We cannot apply Vainsencher's results in order to compute the strata's degree because we are dealing with coincidences of points in the plane. On the other hand, in Colley's work there are no formulas for the cases we need (stationary 6-tuple points).

We use then the more general multiple-point formula by Z. Ran [24] (1984), based on the residual intersection formulas of W. Fulton and D. Laksov [10], that allows us to treat coincidences of points in the zero scheme, let's say X , of sections of a vector bundle with rank equal to $\text{codim}(X)$. This turns out to be our case with a rank 2 vector bundle while in [24] all the examples worked out deal with line bundles.

Then, in §5 we prove results justifying the computation of the degree of strata of cubic surfaces with a conic node, as a product of cycles in the intersection ring of a suitable space of 6-tuples.

The cases not covered here are the strata given by a collapse with less than 3 different lines. Those correspond in our construction to 6-tuples of points that are not curvilinear (i.e. that don't live in a smooth curve) and, in general, the parameter space Z_k used to enumerate them only parametrizes ordered and curvilinear k -tuples of a scheme Z over another one S [17], [24].

Regarding to the effective computation of these degrees, the method employed here leads to calculate in $A^*(Z_k)$ (the intersection ring of Z_k) the degree of a zero-dimensional cycle, obtained by taking up cycles in $A^*(Z_l)$, with $l < k$, through the inductive construction of Z_k .

Given a morphism from Z to S , if $A^*(S)$ is known and $A^*(Z)$ is a finitely generated $A^*(S)$ -module, we are able to give an inductive method to compute the degree of zero-dimensional cycles of $A^*(Z_k)$ in the previous conditions (see Proposition 6.1).

We note that, whenever Z is a projective bundle over S , we can explicitly describe the intersection ring of the space Z_k (see §6). In our case $A^*(Z_6)$ is a polynomial ring over Z with 23 generators modulo 38 relations and effective computations in this ring, using Gröbner basis' method, is much more costly than our inductive method.

We implement this method in a computer program written in Maple [20]. This program could be useful each time that one uses the parameter spaces Z_k in order to study enumerative questions depending on what is happening on the fibers of Z over S .

This paper is organized as follows: §2 is devoted to the classification of singular cubic surfaces in \mathbf{P}^3 . In §3, we construct some parameter spaces appropriated for the problem. We deal in §4 with the spaces of k -tuples and the secant bundles of Ran's work [24]. In §5 we state and prove our main result giving the degree of some strata of cubic surfaces with a conic node. Finally, in §6 we work with the intersection ring of spaces of k -tuples and the effective computation of strata's degree.

This work is part of the second author's Ph.D. Thesis. We would like to thank E. Arrondo and I. Vainsencher for suggestions and comments.

Notation 1.1. We shall tacitly assume that all schemes are defined over an algebraically closed field of characteristic 0. Our notation follows the book of W. Fulton [9], in particular we will write $P(E)$ for the projectivization of a vector bundle E over a scheme Z , defined as $P(E) = \text{Proj}(\text{Sym } E^*)$. If Z is a non-singular scheme, the intersection ring we shall use is the Chow ring of classes of rational equivalence of cycles, graduated by codimension and denoted by $A^*(Z)$. We will denote by Gr the grassmannian of lines of \mathbf{P}^3 .

2. CLASSIFICATION OF SINGULAR CUBIC SURFACES

The classification of singular cubic surfaces in \mathbf{P}^3 given by J. W. Bruce and C. T. C. Wall [5] corresponds to another one in accordance with the incidence relations of the lines contained in a cubic surface, studied by several authors [6], [25]. A cubic surface is non-singular if and only if it contains 27 different lines. When it is singular, some of the lines collapse. The surface given by a general point of each stratum presents a certain type of coincidence between these lines.

We recall here the main points of Bruce's and Wall's work. If P is a singular point of a cubic surface C and $(x_0 : x_1 : x_2 : x_3)$ are homogeneous coordinates in \mathbf{P}^3 , one can assume, doing a change of coordinates if necessary, that P is the point $(0 : 0 : 0 : 1)$ and then, the equation of the surface takes the form

$$(2.1) \quad x_3 f_2(x_0, x_1, x_2) + f_3(x_0, x_1, x_2) = 0$$

where f_i is homogeneous of degree i . The classification depends, first on the rank of the quadratic form f_2 , and second, on the coincidences between the collection of 6 lines common to the two cones given by $f_2 = 0$ and $f_3 = 0$. These are lines through P , contained in C and they are equivalent to the intersection points of the two plane curves, let's say V_2 and V_3 , given by $f_2(x_0, x_1, x_2) = 0$ and $f_3(x_0, x_1, x_2) = 0$ respectively.

Let us suppose that the surface is irreducible and has only isolated singularities. According to the rank of f_2 there are 4 types of singularity at the fixed point P : conic node, binode, uninode and triple point.

When the rank of f_2 is maximum, the singular point is a *conic node*. There are 11 strata in accordance with the partition of 6 given by the coincidences of the intersection points (lines) of V_2 and V_3 , namely

	1^6	21^4	31^3	2^21^2	41^2	321	2^3
Sing.	A_1	$2A_1$	A_1A_2	$3A_1$	A_1A_3	$2A_1A_2$	$4A_1$
Param.	3	2	1	1	0	0	0
	51	42	33	6			
Sing.	A_1A_4	$2A_1A_3$	A_12A_2	A_1A_5			
Param.	0	0	0	0			

In this table, the partition called, for example, 2^21^2 corresponds to a collection of 6 points with 2 of them being double points. At a singular point of type A_k , the normal form of the surface's equation is $x_0^{k+1} + \sum_{j=1}^3 x_j^2 = 0$. The third row indicates the number of parameters on which each type of surface depends, up to projective transformations.

A *binode* corresponds to $\text{rank}(f_2) = 2$, in this case, P is a singularity of type A_k with $k \geq 2$. There are 13 possible configurations of common points to the cubic and the line pair and 6 of them correspond to new strata.

	$1^3.1^3$	$1^3.21$	$1^3.3$	21.21	21.3	3.3	$1^2.1^2$
Sing.	A_2	A_2A_1	$2A_2$	A_22A_1	$2A_2A_1$	$3A_2$	A_3
Param.	2	1	1	0	0	0	1
	1 ² .2	2.2	1 ² .1	2.1	1 ² .0	2.0	
Sing.	A_3A_1	A_32A_1	A_4	A_4A_1	A_5	A_5A_1	
Param.	0	0	0	0	0	0	

When $\text{rank}(f_2) = 1$, P is a *uninode* and one gets three more strata called D_4 , D_5 and E_6 corresponding to the cases in which the points in $V_2 \cap V_3$ are three distinct points, two and one, respectively.

Finally, in the case of $f_2 = 0$, P is a *triple point*. It must happen that $f_3 = 0$ defines a non-singular cubic in order to have an isolated singularity at P . The surface is then a cone over this plane cubic.

Until now the singular point P is fixed on the cubic surface. When P moves in \mathbf{P}^3 , one gets strata in $\mathbf{P}^{19} \times \mathbf{P}^3$ of cubic surfaces with a distinguished singular point. Its projection onto the first factor gives the strata of singular cubic surfaces, living in \mathbf{P}^{19} , that we are interested in.

3. PARAMETER SPACES

We first construct a subscheme of $\mathbf{P}^{19} \times \mathbf{P}^3$, that we will call S , parametrizing cubic surfaces in \mathbf{P}^3 with a distinguished singular point.

In this way, strata of cubic surfaces with a singular point can be seen as subschemes of S and the degree of corresponding strata of singular cubic surfaces in \mathbf{P}^{19} , obtained by projection onto the first factor, could be computed, using the projection formula (see 3.2.(c) of [9]), as a product in the Chow ring $A^*(S)$.

3.1. Cubics with a distinguished singular point. The construction of the parameter space S for cubic surfaces with a distinguished singular point is analogous to others appearing in the literature for plane curves (see, for instance, [22] and [23]).

To fix notation, let V be a vector space of dim 4 over \mathbf{C} and $\mathbf{P}^3 = P(V)$ the projective space of lines of V . We set

$$(3.1) \quad 0 \longrightarrow L \longrightarrow V|_{\mathbf{P}^3} \longrightarrow Q \longrightarrow 0$$

for the tautological exact sequence with $\text{rank}(L) = 1$ and $\text{rank}(Q) = 3$ and we denote by $h = c_1(L^*)$ the first Chern class of $L^* = \mathcal{O}_{\mathbf{P}^3}(1)$ (h is the hyperplane class of \mathbf{P}^3) and by S^3V^* the third symmetric power of V^* .

Definition 3.1. Let F be the subbundle of $S^3V^*|_{\mathbf{P}^3}$ whose fiber over a point of \mathbf{P}^3 is the linear subspace of S^3V^* of cubic forms that have multiplicity at least 2 at such point. Denote by S the projective bundle associated to F .

With this definition $S = P(F) \subset P(S^3V^*|_{\mathbf{P}^3}) \cong \mathbf{P}^{19} \times \mathbf{P}^3$ is the incidence subvariety whose points consist in couples (f, P) , where P is at least a double point for the cubic surface given by $f = 0$. Since $\text{rank}(F) = 16$, then $\dim(S) = 18$. In order to determine the intersection ring of $S = P(F)$, we start with a preliminary lemma.

Lemma 3.2. *With the previous notation, the sequence of vector bundles over \mathbf{P}^3*

$$(3.2) \quad 0 \rightarrow S^2Q^* \otimes Q^* \xrightarrow{u} S^2Q^* \otimes V^*|_{\mathbf{P}^3} \oplus S^3Q^* \xrightarrow{v} F \rightarrow 0$$

where $u(\alpha \otimes \beta) = (\alpha \otimes \beta, -\alpha\beta)$ and $v(\alpha \otimes \omega, \beta) = \alpha\omega + \beta$, is exact. Therefore $c(F) = 1 - 8h + 40h^2 - 160h^3$.

Proof. The first part is similar to the analogous result for plane curves (see [22]). The second one is a straightforward computation applying Whitney's formula (see 3.2.(e) of [9]) to this resolution and to the tautological exact sequence over \mathbf{P}^3 above (3.1). \square

Proposition 3.3. *Let π be the projection from S to \mathbf{P}^3 . The intersection ring of $S = P(F)$ is*

$$A^*(P(F)) \cong \frac{\mathbf{Z}[b, \mu]}{(b^4, \mu^{16} - 8b\mu^{15} + 40b^2\mu^{14} - 160b^3\mu^{13})}$$

where $b = \pi^*h$ and $\mu = c_1(\mathcal{O}_{P(F)}(1))$. Moreover, the following relations hold:

$$b^3\mu^{15} = 1, b^2\mu^{16} = 8, b\mu^{17} = 24 \text{ and } \mu^{18} = 32.$$

Proof. The description of $A^*(S)$ as a polynomial ring is a direct consequence of the result giving the Chow ring of a projective bundle $P(E)$ in terms of the total Chern class of E and the intersection ring of the base space (see 3.3.(b) and Ex. 8.3.4. of [9]). In order to verify the relations in $A^*(S)$ we take down the products in this ring to $A^*(\mathbf{P}^3)$, using the projection formula and standard results about Chern classes (see Ex. 3.2.1. of [9]). \square

3.2. More parameter spaces. We build now a space, say X , that fixes a singular point P for each cubic surface C , and parametrizes the lines contained in C and passing through P . The coincidences between these lines determine the stratum where the surface lives.

To get X , we first construct another space Z in $\mathbf{P}^{19} \times \mathbf{P}^3 \times Gr$. This Z turns out to be a projective bundle over S and we will define X as a subscheme of Z . If we write f for the restriction to X of the projection from Z to S , it is natural to consider the stationary multiple points of f to solve our problem concerning the strata of singular cubic surfaces, because the condition defining the strata of cubics is the collapse produced between the points in the fiber of X over S .

Remark 3.4. One has to take care of three pairs of strata such that both strata on each pair have the same codimension in the \mathbf{P}^{19} of cubic surfaces and they correspond to the same type of coincidence between points of X over S . These pairs are $(4A_1, D_4)$, $(2A_1 A_3, D_5)$ and $(A_1 A_5, E_6)$. In this paper, we don't compute the degree of the strata of surfaces having singularities of the types appearing on the two last pairs but, in order to compute the degree of surfaces with four conic nodes ($4A_1$) one must subtract the contribution given by surfaces with a uninode (D_4) (see Theorem 5.1).

3.2.1. Construction of Z . We consider the incidence of points and lines of \mathbf{P}^3 , $I \subset \mathbf{P}^3 \times Gr$, consisting of pairs (P, l) , $P \in L$, where l is the point of Gr corresponding to the line $L \subset \mathbf{P}^3$.

Definition 3.5. If p_1 (resp. p_2) is the projection from I to \mathbf{P}^3 (resp. Gr), Z is defined as the scheme $I \times_{\mathbf{P}^3} S$ (see B.2.3. of [9]).

$$\begin{array}{ccccc} Z & \xrightarrow{\pi} & I & \xrightarrow{p_2} & Gr \\ \overline{p_1} \downarrow & & \downarrow p_1 & & \\ S & \xrightarrow{\pi} & \mathbf{P}^3 & & \end{array}$$

This Z represents collections (f_C, P, l) with $f_C = 0$ the equation of the cubic surface C , P a double point of C and L a line through P . By definition, Z is a projective bundle over S : the projection p_1 identifies the incidence I with $P(T_{\mathbf{P}^3})$, ($T_{\mathbf{P}^3}$ is the tangent bundle of \mathbf{P}^3), so we have also (see B.5.5. of [9])

$$I \cong P(T_{\mathbf{P}^3}(-2)) \cong P(Q(-1))$$

and therefore $Z \cong P(\pi^*Q(-1))$ and $\dim(Z) = 20$.

It is also easy to see that $p_2^*\mathcal{O}_{Gr}(1) \cong \mathcal{O}_{P(Q(-1))}(1)$, $p_1^*\mathcal{O}_{\mathbf{P}^3}(1) \cong \mathcal{O}_{P(U)}(1)$ and $p_2 : I \rightarrow Gr$ can be identified with the projective bundle $P(U)$ over Gr , being U the tautological subbundle of rank 2 fitting in the tautological exact sequence over Gr :

$$(3.3) \quad 0 \longrightarrow U \longrightarrow V|_{Gr} \longrightarrow Q_{Gr} \longrightarrow 0.$$

These facts will be useful on later computations (see 3.2.2).

3.2.2. Construction of X . If C is a cubic surface in \mathbf{P}^3 given as the zeroes of a section f_C of $\mathcal{O}_{\mathbf{P}^3}(3)$, the Fano scheme of lines contained in C , $F(C)$, can be seen as the scheme of zeroes of the section induced by f_C in $p_{2*}(p_1^*\mathcal{O}_{\mathbf{P}^3}(3)) = S^3U^*$. The line bundle $p_1^*\mathcal{O}_{\mathbf{P}^3}(3)$ over I corresponds to the divisor $(p_1^*f_C)_0$ with fiber at a general $l \in Gr$ the three intersection points of the cubic surface C with the line L of \mathbf{P}^3 .

This is a particular case of the Fano scheme of pairs (V, l) where V is a hypersurface in \mathbf{P}^n of given degree d and l parametrizes a line L contained in V , defined in [1]. There it is proved that, if H is the projective space of all hypersurfaces of degree d , G is the grassmannian of lines of \mathbf{P}^n and U is the tautological subbundle over G , then the Fano scheme in $H \times G$ is given by the scheme of zeroes of the section of $S^dU^* \otimes \mathcal{O}_H(1)$, induced by the section of $\mathcal{O}_{\mathbf{P}^n}(d) \otimes \mathcal{O}_H(1)$ defining the universal divisor of $H \times \mathbf{P}^n$ (see Thm. 3.3. of [1]).

To obtain a space $X \subset Z$ consisting on the points (f_C, P, l) of Z such that $L \subset C$, we adapt this result to our case for which lines must be contained in the surface C

and must pass through the distinguished singular point $P \in C$ (see Proposition 3.9 below).

$$\begin{array}{ccccccc} X & \hookrightarrow & Z & \hookrightarrow & \mathbf{P}^{19} \times \mathbf{P}^3 \times Gr \\ & \searrow & \downarrow & & & & \\ & & S & \hookrightarrow & \mathbf{P}^{19} \times \mathbf{P}^3. & & \end{array}$$

First, we define a space T over Z which plays the role of the incidence I as projective bundle over Gr in the case of a concrete cubic surface: we want the fiber of T over a point (f_C, P, l) parametrized by Z to represent the \mathbf{P}^1 of points of the line L . We then consider the diagram

$$(3.4) \quad \begin{array}{ccccccc} T & \xrightarrow{\overline{\pi}} & Y & \xrightarrow{pr_2} & I' & \xrightarrow{p'_1} & \mathbf{P}^3 \\ \overline{pr_1} \downarrow & & \downarrow pr_1 & & \downarrow p'_2 & & \\ Z & \xrightarrow{\overline{\pi}} & I & \xrightarrow{p_2} & Gr & & \\ \overline{p_1} \downarrow & & \downarrow p_1 & & & & \\ S & \xrightarrow{\pi} & \mathbf{P}^3 & & & & \end{array}$$

where p'_2 is the projection of $P(U) \rightarrow Gr$, $Y := I \times_{Gr} I' \cong P(p_2^* U) \subset \mathbf{P}^3 \times Gr \times \mathbf{P}^3$ and $T := Y \times_I Z \cong P(\overline{\pi}^* p_2^* U) \subset \mathbf{P}^{19} \times \mathbf{P}^3 \times Gr \times \mathbf{P}^3$.

In this way, Y parametrizes collections (P, l, P') such that $P \in L$ and $P' \in L$ and T collections (f_C, P, l, P') where P is a double point of the surface C and $P, P' \in L$. Now, we build an invertible sheaf N over T which is the analogue of $p_1^* \mathcal{O}_{\mathbf{P}^3}(3)$ in the case of $F(C)$.

Lemma 3.6. *Let N be the line bundle $\overline{pr_1}_* \overline{p_1}^* \mathcal{O}_S(1) \otimes \mathcal{O}_T(3)$ over T . Then N has an associated section s_N such that $D := (s_N)_0$ parametrizes collections (f_C, P, l, P') with $P' \in C$, besides P being a double point of C and $P, P' \in L$.*

Proof. Let π_1 and π_2 be the projections from $S \times \mathbf{P}^3$ to its factors. If ϕ is the morphism from T to $S \times \mathbf{P}^3$ given by the diagram (3.4) then, by construction, N is the pullback by ϕ of $\pi_1^* \mathcal{O}_S(1) \otimes \pi_2^* \mathcal{O}_{\mathbf{P}^3}(3)$. The section s_N on the statement above is ϕ^* of the section of $\pi_1^* \mathcal{O}_S(1) \otimes \pi_2^* \mathcal{O}_{\mathbf{P}^3}(3)$ vanishing on the universal surface in $S \times \mathbf{P}^3$. \square

Remark 3.7. For each point $z \in Z$, the fiber of $(s_N)_0 \subset T$ consists of two points: P the double point of C and, let's say R , the remaining point in the intersection of a general line L through this P , and the surface C .

Let D_P be the divisor of T given by the point P when z moves in Z . Let D_R be the residual divisor of D_P on D (see 9.2.1. of [9]), given by the point R when z moves in Z . We will get X as the scheme of zeroes of a section of a rank 2-vector bundle over Z , defined as $\overline{pr_1}_*$ of the line bundle over T corresponding to the divisor D_R .

Definition 3.8. Let E be the sheaf $\overline{pr_1}_*(\mathcal{O}_T(D_R))$, where $\overline{pr_1}$ is the projection from T to Z and D_R is defined in the previous remark.

Proposition 3.9. *With the previous notation:*

1. *The pairs consisting of a flat family of cubic surfaces of \mathbf{P}^3 with a distinguished singular point and a flat family of lines contained in the surface and passing through its singular point, both parametrized by the same scheme W are the W -points of a closed subscheme X of Z .*

2. The space X is the scheme of zeroes of a regular section, say s_E , of the locally free sheaf E of rank 2 over Z .

Proof. The proof is analogous to that of Thm. 3.3 (i), (iii) of [1] except for the following concerning 2:

By definition, $D = 2D_P + D_R$, so $M := \mathcal{O}_T(D_R) = N \otimes \mathcal{O}_T(-2D_P)$ and the section s_N of N induces another one of M , let's say s_M , vanishing on D_R . The sought section is equal to $s_E = \overline{pr_1}_* s_M$.

E is a rank 2-vector bundle because by Grauert's theorem (see III.12.9 of [12]), $E = R^0\overline{pr_1}_*(\mathcal{O}_T(D_R))$ is locally free if $\dim H^0(T_z, \mathcal{O}_T(D_R)|_{T_z})$ is constant $\forall z \in Z$ and in our case, $h^0(T_z, \mathcal{O}_T(D_R)|_{T_z}) = h^0(\mathbf{P}^1, \mathcal{O}_{\mathbf{P}^1}(1)) = 2, \forall z \in Z$. \square

We write j for the inclusion $X \subset Z$, p for the projection from Z to S (called before $\overline{p_1}$) and f for the composition $p \circ j$.

The situation we get is

$$\begin{array}{ccccc} & & E & & \\ & & \downarrow \uparrow s_E & & \\ X = (s_E)_0 & \xrightarrow{j} & Z & \hookrightarrow & \mathbf{P}^{19} \times \mathbf{P}^3 \times Gr \\ & \searrow f & \downarrow p & & \\ & & S & \hookrightarrow & \mathbf{P}^{19} \times \mathbf{P}^3 \end{array}$$

with the space X parametrizing, for each cubic surface with a given double point, the lines passing through that point and contained in the surface. In this way, the strata of singular cubic surfaces with a conic node of given type (see §2) depend on the coincidences between points on the fiber of $f : X \rightarrow S$, keeping in mind Remark 3.4. We will show in §5 that their degree can be determined using a suitable cycle of stationary multiple points of f and computing in the intersection ring of the space of 6-tuples Z_6 (see §4).

In §5 we will need to know $c(E)$, the total Chern class of the vector bundle E and also that X is smooth.

Lemma 3.10. *With the previous notation, if $\xi = c_1(\mathcal{O}_Z(1))$ is the hyperplane class of Z and we denote by b, μ the pullback by $p : Z \rightarrow S$ of the same classes in $A^*(S)$, then $E = \overline{\pi}^* p_2^* U^* \otimes \overline{p_1}_* \mathcal{O}_S(1) \otimes \overline{\pi}^* \mathcal{O}_{P(U)}(-2) \otimes \overline{\pi}^* \mathcal{O}_{P(Q(-1))}(2)$ and therefore $c(E) = 1 + (5\xi - 4b + 2\mu) + (6\xi^2 + 5\mu\xi - 9b\xi + \mu^2 - 4\mu b + 3b^2)$.*

Proof. Since $E := \overline{pr_1}_*(\mathcal{O}_T(D_R))$ and $\mathcal{O}_T(D_R) = \mathcal{O}_T(D) \otimes \mathcal{O}_T(-2D_P)$, we first determine $[D_P] \in A^1(T)$. If ξ_T is the hyperplane class of $T = P(\overline{\pi}^* p_2^* U)$ over Z , then

$$[D_P] = a \cdot \xi_T + \overline{pr_1}_*[D']$$

with $D' \in A^1(Z)$ (see Ex. 8.3.4. of [9]). Now, by construction, $[D_P] \cdot [T_z] = 1$ and this implies $a = 1$ (for $\xi_T \cdot [T_z] = 1$ and the fibers of T over Z don't intersect each other).

To get D' , we compute $\overline{pr_1}_*([D_P] \cdot \xi_T)$ in two ways:

1. $\overline{pr_1}_*([D_P] \cdot \xi_T) = \overline{pr_1}_*(\xi_T^2) + [D'] \cdot \overline{pr_1}_*(\xi_T) = \overline{\pi}^* p_2^* c_1(Q_{Gr}) + [D']$, where the first equality is the projection formula and the second is Ex. 3.2.1 of [9], being Q_{Gr} the tautological quotient of rank 2 over the grassmannian.

2. Let s_P be the section of T over Z such that $Im(s_P) = D_P$. Then

$$\overline{pr_1}_*([D_P] \cdot \xi_T) = c_1(s_P^* \mathcal{O}_T(1)).$$

From 1 and 2 above, $[D'] = c_1(s_P^* \mathcal{O}_T(1)) - \bar{\pi}^* c_1(\mathcal{O}_{P(Q(-1))}(1)) \in A^1(T)$, recalling that $c_1(Q_{Gr}) = c_1(\mathcal{O}_{Gr}(1))$ and $p_2^* \mathcal{O}_{Gr}(1) = \mathcal{O}_{P(Q(-1))}(1)$.

To compute $c_1(s_P^* \mathcal{O}_T(1))$, we use the upper part of diagram (3.4):

$$\begin{array}{ccccccc} T & \xrightarrow{\bar{\pi}} & Y & \xrightarrow{pr_2} & I' & \xrightarrow{p'_1} & \mathbf{P}^3 \\ \overline{pr_1} \downarrow \uparrow s_P & & t_P \downarrow \uparrow pr_1 & & \downarrow p'_2 & & \\ Z & \xrightarrow{\bar{\pi}} & I & \xrightarrow{p_2} & Gr & & \end{array}$$

where we define t_P as the section of $Y = I \times_{Gr} I'$ whose image is the diagonal of Y . Looking at the diagram we see that $s_P^* \mathcal{O}_T(1) = \bar{\pi}^* t_P^* \mathcal{O}_Y(1)$ (as $T = P(\bar{\pi}^* p_2^* U)$). Now, by construction, $t_P^* c_1(\mathcal{O}_Y(1)) = c_1(\mathcal{O}_I(1))$ with $I = P(U)$ over Gr .

Putting all these things together $[D'] = \bar{\pi}^*(c_1(\mathcal{O}_I(1)) - c_1(\mathcal{O}_{P(Q(-1))}(1)))$ and:

- $\mathcal{O}_T(D_P) = \mathcal{O}_T(1) \otimes \overline{pr_1}^* \bar{\pi}^* \mathcal{O}_{P(U)}(1) \otimes \overline{pr_1}^* \bar{\pi}^* \mathcal{O}_{P(Q(-1))}(-1)$,
- $\mathcal{O}_T(D_R) = \mathcal{O}_T(1) \otimes \overline{pr_1}^* (\overline{p_1}^* \mathcal{O}_S(1) \otimes \bar{\pi}^* \mathcal{O}_{P(U)}(-2) \otimes \bar{\pi}^* \mathcal{O}_{P(Q(-1))}(2))$.

Finally, by the projection formula:

$$E = \overline{pr_1}_* (\mathcal{O}_T(D_R)) = \bar{\pi}^* p_2^* U^* \otimes \overline{p_1}^* \mathcal{O}_S(1) \otimes \bar{\pi}^* \mathcal{O}_{P(U)}(-2) \otimes \bar{\pi}^* \mathcal{O}_{P(Q(-1))}(2).$$

From this expression, the computation of $c(E)$ is straightforward. \square

We see now that X is smooth (even though X is not smooth over S).

Proposition 3.11. *With the previous notation, X is smooth.*

Proof. We are going to verify that, if $P \in \mathbf{P}^3$, then the fiber X_P can be identified with the total space of a projective bundle over \mathbf{P}^2 and therefore X is smooth.

Since $X = (s_E)_0$ with s_E a section of the vector bundle E over Z , then $X_P \subset Z_P$ can be identified with the scheme of zeroes of the section of $E|_{Z_P}$ induced by s_E . Now $Z_P \cong \mathbf{P}^{15} \times \mathbf{P}^2$ and, if τ_1 y τ_2 are the projections from $\mathbf{P}^{15} \times \mathbf{P}^2$ to its factors, then

$$E|_{Z_P} = \tau_1^* \mathcal{O}_{\mathbf{P}^{15}}(1) \otimes \tau_2^* (\mathcal{O}_{\mathbf{P}^2}(2) \oplus \mathcal{O}_{\mathbf{P}^2}(3)).$$

To see this we use the explicit description of E given in the previous lemma, noting that:

- The fiber I_P consists on the plane in Gr of lines through the point P .
- $p_2^* U^*|_{I_P} \cong \mathcal{O}_{\mathbf{P}^2} \oplus \mathcal{O}_{\mathbf{P}^2}(1)$. This can be checked recalling that

$$U^* \cong p_{2*}(p_1^* \mathcal{O}_{\mathbf{P}^3}(1)),$$

and so, every section ρ of U^* is induced by a section η of $\mathcal{O}_{\mathbf{P}^3}(1)$ in such a way that ρ vanishes on the plane Π_Λ of Gr parametrizing the lines contained in the plane Λ which is the zero locus of η . In this way, the restriction of the Koszul exact sequence (see B.3.4. of [9]):

$$0 \rightarrow \mathcal{O}_{Gr} \rightarrow U^* \rightarrow \mathcal{I}_{\Pi_\Lambda}(1) \rightarrow 0,$$

to I_P is

$$0 \rightarrow \mathcal{O}_{\mathbf{P}^2} \rightarrow U^*|_{\mathbf{P}^2} \rightarrow \mathcal{O}_{\mathbf{P}^2}(1) \rightarrow 0,$$

for the intersection in Gr of a plane of lines through a general point of \mathbf{P}^3 and a plane of lines contained in a general plane of \mathbf{P}^3 is empty (see Ex. 14.7.2. of [9]).

- $\overline{p_1}^* \mathcal{O}_S(1)|_{Z_P} \cong \mathcal{O}_{\mathbf{P}^{15}}(1)$ (recalling that $S_P \cong \mathbf{P}^{15}$).

- The restriction of $\mathcal{O}_{P(U)}(-1)$ to I_P is trivial since $I \cong P(U) \xrightarrow{p_2} Gr$ and

$$\mathcal{O}_{P(U)}(1) = p_1^* \mathcal{O}_{\mathbf{P}^3}(1).$$

- $\mathcal{O}_{P(Q(-1))}(2)|_{I_P} = \mathcal{O}_{\mathbf{P}^2}(2)$, for $I \cong P(Q(-1)) \xrightarrow{p_1} \mathbf{P}^3$.

Now, if we call $R := \mathcal{O}_{\mathbf{P}^2}(2) \oplus \mathcal{O}_{\mathbf{P}^2}(3)$, then $P(H^0(R)) \cong \mathbf{P}^{15}$ and the sequence of vector bundles over \mathbf{P}^2 :

$$0 \rightarrow K \rightarrow H^0(R) \otimes \mathcal{O}_{\mathbf{P}^2} \xrightarrow{ev} R \rightarrow 0,$$

given by the evaluation of sections of R , is exact. The key observation is that the tautological section Γ of $\tau_1^* \mathcal{O}_{\mathbf{P}^{15}}(1) \otimes \tau_2^* R = E|_{Z_P}$ is precisely the restriction to the fiber Z_P of s_E .

The situation is

$$\begin{array}{ccc} & \tau_1^* \mathcal{O}_{\mathbf{P}^{15}}(1) \otimes \tau_2^* R & \\ & \downarrow \Gamma & \\ (\Gamma)_0 & \longrightarrow & P(H^0(R)) \times \mathbf{P}^2 \\ & \searrow & \downarrow \tau_2 \\ & & P(H^0(R)) \end{array}$$

and so $X_P \cong (\Gamma)_0$ which is isomorphic to the projective bundle $P(K)$ over \mathbf{P}^2 . Therefore X is smooth. \square

4. SPACES FOR k -TUPLES AND SECANT BUNDLES

In §5, we shall use the spaces for k -tuples Z_k and the secant bundles E_k , for Z and E defined in §3, to compute the degree of some strata of cubic surfaces with a conic node. We describe them here taking as reference the work by Z. Ran [24].

Spaces for k -tuples. The space Z_k is built inductively as a suitable blow-up of an ordered k -product of Z .

- Steps 0, 1. One denotes $Z_0 := S$, $Z_1 := Z$, $p_0^1 : Z_1 \rightarrow Z_0$ is the morphism p and one defines $\iota_j : Z_j \rightarrow Z_j$ as the identity i for $j = 0, 1$.
- Step 2. Considering the fiber product defined by the cartesian diagram (see B.2.3. of [9]):

$$\begin{array}{ccc} Z_1 \times_{Z_0} Z_1 & \xrightarrow{pr_2} & Z_1 \\ \downarrow pr_1 & & \downarrow p \\ Z_1 & \xrightarrow{p} & Z_0, \end{array}$$

one defines a diagonal $\Delta_1 = (i \times \iota_1)(Z_1) \hookrightarrow Z_1 \times_{Z_0} Z_1$; b_2 , the blow-up of $Z_1 \times_{Z_0} Z_1$ along Δ_1 with exceptional divisor $D_{1,2}^2$,

$$\begin{array}{ccc} D_{1,2}^2 & \hookrightarrow & Z_2 \\ \downarrow & & \downarrow b_2 \\ \Delta_1 & \hookrightarrow & Z_1 \times_{Z_0} Z_1 \end{array}$$

and new projections from Z_2 to Z_1 ; $p_{1,1}^2 := pr_1 \circ b_2$ y $p_{2,2}^2 := pr_2 \circ b_2$.

Since p is smooth, $\Delta_1 \hookrightarrow Z_1 \times_{Z_0} Z_1$ is a regular inclusion (see B.7.3. of [9]) and the new projections are also smooth.

Out of the exceptional divisor, Z_2 parametrizes couples (z_1, z_2) of different points of Z having the same image in S , i.e., strict double points of f . In $D_{1,2}^2$ live the couples of points of Z having the same image by p that are infinitely near $(b_2(D_{1,2}^2))$ consists on pairs (z, z) where z is a ramification point of p .

One notes that $(\iota_1 \circ pr_2, \iota_1 \circ pr_1)$ is an involution of $Z_1 \times_{Z_0} Z_1$ reversing the order in (z_1, z_2) and fixing the diagonal Δ_1 , so it lifts to an involution of Z_2 that will be denoted by ι_2 .

- Induction hypothesis. One supposes that a space Z_k of k -tuples and associated objects $(D_{1,k}^k, p_{1,k-1}^k, p_{2,k}^k, \iota_k)$ are built as before.
- Step $k+1$. One considers the fiber product defined by

$$(4.1) \quad \begin{array}{ccc} Z_k \times_{Z_{k-1}} Z_k & \xrightarrow{pr_2} & Z_k \\ \downarrow pr_1 & & \downarrow g = \iota_{k-1} \circ p_{1,k-1}^k \\ Z_k & \xrightarrow{g \circ \iota_k = p_{2,k}^k} & Z_{k-1} \end{array}$$

where $p_{1,k-1}^k := pr_1 \circ b_k$, $p_{2,k}^k := pr_2 \circ b_k$ and $b_k : Z_k \rightarrow Z_{k-1} \times_{Z_{k-2}} Z_{k-1}$ is the previous blow-up.

Next, one defines $\Delta_k = (i \times \iota_k)(Z_k)$, where ι_k is $(\iota_{k-1} \circ pr_2, \iota_{k-1} \circ pr_1)$ lifted to Z_k (ι_k is a regular inclusion by B.7.3. of [9]) and the blow-up along Δ_k with exceptional divisor $D_{1,k+1}^{k+1}, b_{k+1} : Z_{k+1} \rightarrow Z_k \times_{Z_{k-1}} Z_k$.

In this way, the points of Z_k correspond to ordered k -tuples of points of Z having the same image by p in S . By B.7.3. of [9], it holds

$$N_{\Delta_k} Z_k \times_{Z_{k-1}} Z_k \cong \iota_k^* T_{\iota_{k-1} \circ p_{1,k-1}^k} \cong T_{p_{2,k}^k}.$$

This will be useful in §6, in order to make computations in the intersection ring $A^*(Z_k)$.

Remark 4.1. Note that in [24] g is taken as $p_{1,k-1}^k$ and so $\Delta_k = (i \times \iota_k)(Z_k)$ is not contained in $Z_k \times_{Z_{k-1}} Z_k$.

Regarding the enumeration of stationary multiple points of the morphism p , given a partition $\sigma = (k_1, k_2, \dots, k_r)$ of k with $\sum_{i=1}^r k_i = k$, Ran defines a subscheme of Z_k called V_σ , parametrizing k -tuples of Z/S with the first k_1 points being infinitely near to each other (i.n.), the next k_2 points being also i.n., etc. For example, the subscheme of Z_4 parametrizing pairs of points of type $(2, 2)$ is $V_{(2,2)} = (p_{1,2}^3 \circ p_{1,3}^4)^* D_{1,2}^2 \cap (p_{2,4}^4)^* D_{1,3}^3$ for points in Z_4 correspond to collections $(z_1, z_2, z_2, z_3, z_3, z_2, z_2, z_4)$ of Z/S .

Anyway, for each partition σ of k , one can define a subscheme of Z_k parametrizing k -tuples of type σ as a suitable intersection of (pullbacks by projections of) exceptional divisors reflecting the coincidences given by σ .

Remark 4.2. The space Z_k had already been defined by S. L. Kleiman (see p. 390 of [16] and [17]). For the construction of Z_{k+1} , Kleiman considers the fiber product $Z_k \times_{Z_{k-1}} Z_k$ containing the usual diagonal $(i \times i)(Z_k)$.

For the justification of Z_k as parameter spaces, see [18] and [24] where a relation between Z_k and the relative Hilbert scheme $Hilb^k(Z/S)$ is established, showing that Z_k is a parameter space for ordered and curvilinear k -tuples of points of Z with the same image in S (a collection of points is curvilinear if the points live in a smooth curve). Consequently, the (stationary) multiple points formulas given by Colley, Kleiman and Ran cannot enumerate the non-curvilinear points of a morphism.

Secant bundles. The notion of secant bundle, introduced by R. L. E. Schwarzenberger [27] in 1964, generalizes the concept of tangent bundle of a projective variety Y in the following sense: the secant bundle of Y is defined over a product scheme of Y so that its restriction to the diagonal in this product is the tangent bundle of

Y . The Chern classes of secant bundles are related with multiple-point cycles (see [21], [19] and [29]). By definition, a secant bundle is always associated to a line bundle over a scheme.

In [24], given a vector bundle E of arbitrary rank over a smooth scheme Z , a *secant bundle* E_k over Z_k is defined, allowing us to compute the class $[X_k]$ in $A^*(Z_k)$, when X is the zero subscheme of a section of E . The E_k are defined inductively and they fit into an exact sequence of vector bundles over Z_k (analogous to the one given by Schwarzenberger for its secant bundles) from which one can compute its Chern classes.

Regarding the definition of E_k : for $k = 1$, $E_1 := E$; if $k = 2$, E_2 is defined by the exact sequence over Z_2 :

$$0 \rightarrow E_2 \xrightarrow{(\lambda_2^1, \lambda_2^2)} (p_{1,1}^2)^*E \oplus (p_{2,2}^2)^*E \xrightarrow{\rho_2} (p_{1,1}^2)^*E|_{D_{1,2}^2} \rightarrow 0,$$

with $\rho_2(a, b) = res(a) - res(b)$, and res the restriction to the exceptional divisor $D_{1,2}^2$. E_2 is a vector bundle over Z_2 by Nakayama's lemma and from its definition it is easy to see that

$$0 \rightarrow (p_{2,2}^2)^*E \otimes L_{-D_{1,2}^2} \rightarrow E_2 \rightarrow (p_{1,1}^2)^*E \rightarrow 0$$

is exact, where in general L_D is the line bundle associated to the divisor D .

When $k \geq 3$, by induction, one supposes that, $\forall j \leq k$, vector bundles E_j and applications

$$\left\{ \begin{array}{l} \lambda_j^1 : E_j \rightarrow (\iota_{j-1} \circ p_{1,j-1}^j)^*E_{j-1} \\ \lambda_j^2 : E_j \rightarrow (p_{2,j}^j)^*E_{j-1} \end{array} \right\},$$

are defined. Then

Definition 4.3. Let $J_{k+1} := (\iota_k \circ p_{1,k}^{k+1})^*E_k \oplus (p_{2,k+1}^{k+1})^*E_k$ and let K_{k+1} be the vector bundle given by the exact sequence over Z_{k+1} :

$$0 \rightarrow K_{k+1} \xrightarrow{(\kappa^1, \kappa^2)} J_{k+1} \xrightarrow{\rho_{k+1}} (p_{2,k+1}^{k+1})^*E_k|_{D_{1,k+1}^{k+1}} \rightarrow 0,$$

with $\rho_{k+1}(a, b) := res(a) - res(b)$. Let E_{k+1} be defined by

$$0 \rightarrow E_{k+1} \xrightarrow{(\lambda_{k+1}^1, \lambda_{k+1}^2)} K_{k+1} \xrightarrow{\nu_{k+1}} (p_{2,k}^{k+1})^*E_{k-1} \otimes L_{-D_{1,k+1}^{k+1}} \rightarrow 0,$$

with

$$\begin{aligned} \nu_{k+1} &= ((p_{1,k}^{k+1})^*(\lambda_k^2) - (p_{2,k+1}^{k+1})^*(\lambda_k^1)), \\ p_{2,k}^{k+1} &:= \iota_{k-1} \circ p_{1,k-1}^k \circ p_{2,k+1}^{k+1} : Z_{k+1} \rightarrow Z_{k-1}. \end{aligned}$$

A standard computation using commutative diagrams shows that with this definition the sequence of vector bundles over Z_{k+1} :

$$(4.2) \quad 0 \rightarrow (p_{k+1,k+1}^{k+1})^*E \otimes L_{-D^{k+1,k+1}} \rightarrow E_{k+1} \rightarrow (p_{1,k}^{k+1})^*E_k \rightarrow 0$$

where $p_{k+1,k+1}^{k+1} := p_{2,2}^2 \circ \dots \circ p_{2,k}^k \circ p_{2,k+1}^{k+1}$ and

$$D^{k+1,k+1} := D_{1,k+1}^{k+1} + (p_{2,k+1}^{k+1})^*D_{1,k}^k + \dots + (p_{2,3}^3 \circ p_{2,4}^4 \circ p_{2,5}^5 \circ \dots \circ p_{2,k+1}^{k+1})^*D_{1,2}^2$$

is exact (for details see [30]).

Remark 4.4. Note that our definition of E_k for $k \geq 3$ is different from the one given in [24] for which, an easy computation, using Whitney's formula (3.2.(e) of [9]), shows that the total Chern class of E_{k+1} is not the one given by the exact sequence (4.2) with which, in some examples, one recovers known results (see [30]).

5. DEGREE OF SOME STRATA OF CUBIC SURFACES WITH A CONIC NODE

Let $\sigma = (k_1, k_2, \dots, k_r)$ be a partition of 6 with $k_j \geq k_l$, if $j \leq l$, C_σ the stratum in \mathbf{P}^{19} of cubic surfaces with a conic node of type σ , C_U the one of surfaces with a uninode and $\alpha_\sigma := \dim(C_\sigma)$.

In principle, we would get $\deg(C_\sigma)$ as $\#(C_\sigma \cap \Lambda)$ being $\Lambda \subset \mathbf{P}^{19}$ a general linear space of codimension α_σ .

The stratum C_σ depends on coincidences between points on the fibers of the map $f : X \rightarrow S$, unless in the case of a coincidence of type $\sigma = (2, 2, 2)$ that corresponds to the union of $C_{(2,2,2)}$ and C_U (see Remark 3.4). Since $X \subset Z$ is not smooth over S , we consider the space Z_6 of 6-tuples of Z/S treated in §4. Even though f is not smooth, the inclusion $j : X \hookrightarrow Z$ induces, by construction, inclusions $j_k : X_k \hookrightarrow Z_k$.

Let ϕ_1 denote the projection from Z_6 to Z assigning to each 6-tuple of Z_6 its first element and τ the projection from S to \mathbf{P}^{19} . Let ϕ be the composite projection $\tau \circ p \circ \phi_1 : Z_6 \rightarrow \mathbf{P}^{19}$ and V_σ the subscheme of Z_6 corresponding to 6-tuples with a coincidence of type σ (see §4). A diagram of the situation is:

$$\begin{array}{ccc} X_6 & \xrightarrow{j_6} & Z_6 \\ \downarrow & & \downarrow \phi_1 \\ X & \xrightarrow{j} & Z \\ & \searrow f & \downarrow p \\ & & S \\ & & \downarrow \tau \\ & & \mathbf{P}^{19} \end{array}$$

The result giving the degree of some strata of cubic surfaces with a conic node is the following

Theorem 5.1. *Let $\sigma = (k_1, \dots, k_r)$ be a partition of 6 with $r \geq 3$ and let Λ be a linear space of \mathbf{P}^{19} of dimension $7 - r$. If $\Pi = \phi^{-1}(\Lambda)$, then the degree of the stratum C_σ of cubic surfaces with a conic node of type σ is*

$$\deg(C_\sigma) = \frac{1}{n(\sigma)} \int_{Z_6} [X_6] \cdot [V_\sigma] \cdot [\Pi],$$

except for the case $\sigma = (2, 2, 2)$ where

$$\deg(C_\sigma) = \frac{1}{n(\sigma)} \int_{Z_6} [X_6] \cdot [V_\sigma] \cdot [\Pi] - \frac{1}{4} \deg(C_U).$$

In both formulas, $n(\sigma)$ is an integer determined by the partition σ .

Proof. We divide the proof into 4 steps.

1. We first translate the computation of $\deg(C_\sigma)$ to $A^*(S)$ showing that, if $S_\sigma \subset S$ is the closure of the locus of cubic surfaces with a distinguished singular point such that the associated collection of lines through this point have

collapsed in accordance with the partition σ (i.e. $\tau(S_\sigma) = C_\sigma$), then if $\sigma \neq (2, 2, 2)$,

$$m \cdot \deg(C_\sigma) = \int_S [S_\sigma \cap \Lambda_S],$$

where m is the number of conic nodes of a cubic surface in the stratum C_σ and $\Lambda_S = \tau^{-1}(\Lambda)$ (see Lemma 5.2).

When $\sigma = (2, 2, 2)$, $\tau(S_\sigma) = C_\sigma \cup C_U$ and therefore

$$4 \cdot \deg(C_\sigma) + \deg(C_U) = \int_S [S_\sigma \cap \Lambda_S].$$

2. We define a subscheme $W \subset Z_6$ by $W = X_6 \cap V_\sigma \cap \Pi$ (in this way $(p \circ \phi_1)(W) = S_\sigma \cap \Lambda_S$) and we prove that $[W] \in A^*(Z_6)$ enumerates the ordered 6-tuples of $\phi_1(W)$ over S if and only if $\dim(\Lambda_S) \leq 3$, which is equivalent to $r \geq 3$ above (see Lemma 5.3). For the rest of the proof we then suppose that $\dim(\Lambda_S) \leq 3$.
3. We show in Lemma 5.4 that the cycle $[W] \in A^*(Z_6)$ pushes forward to $\tilde{n}(\sigma) \cdot [S_\sigma \cap \Lambda_S]$ in $A^*(S)$, and we compute $\tilde{n}(\sigma)$ according to the partition σ .
4. We verify in Lemma 5.5 that $[W] = [X_6 \cap V_\sigma \cap \Pi] = [X_6] \cdot [V_\sigma] \cdot [\Pi]$. In this way, the theorem is proved with $n(\sigma) = m \cdot \tilde{n}(\sigma)$ depending only on the partition σ . \square

Lemma 5.2. *With the previous notation, let m be the number of conic nodes presented by a surface belonging to the stratum C_σ , then if $\sigma \neq (2, 2, 2)$,*

$$m \cdot \deg(C_\sigma) = \int_S [S_\sigma \cap \Lambda_S]$$

and if $\sigma = (2, 2, 2)$,

$$m \cdot \deg(C_\sigma) + \deg(C_U) = \int_S [S_\sigma \cap \Lambda_S].$$

Proof. Suppose that $\sigma \neq (2, 2, 2)$: if the type of singularities presented by a cubic surface corresponding to a point of S_σ is $m A_1 A_q$ (see §2) then $\deg(\tau|_{S_\sigma}) = m$ (for the projection τ forgets the double point) and consequently $\tau_*([S_\sigma]) = m \cdot [C_\sigma]$.

By construction, $\mu := c_1(\mathcal{O}_S(1)) = \tau^* H$, where H is the hyperplane class of \mathbf{P}^{19} . Applying the projection formula, we then have

$$m \cdot \deg(C_\sigma) = m \int_{\mathbf{P}^{19}} [C_\sigma] \cdot H^{\alpha_\sigma} = \int_S [S_\sigma] \cdot \mu^{\alpha_\sigma} = \int_S [S_\sigma \cap \Lambda_S].$$

The last equality is $[S_\sigma \cap \Lambda_S] = [S_\sigma] \cdot [\Lambda_S]$. This is true because the intersection $S_\sigma \cap \Lambda_S$ consists of points and it is transversal at each one of them. (If this last claim were false, it would be also false for $\tau(S_\sigma \cap \Lambda_S) = C_\sigma \cap \Lambda$ and we know that a general linear space in a projective space has a non-empty intersection with any subvariety of complementary dimension and one can choose it so the intersection is transversal by Bertini's theorem (see II.8.18 of [12])).

If $\sigma = (2, 2, 2)$, a surface corresponding to a point of S_σ can have 4 conic nodes or else a uninode, so this time $\tau_*([S_\sigma]) = 4 \cdot [C_\sigma] + [C_U]$ and the rest is as before. \square

With the notation given in §4, if $p : Z \rightarrow S$, a cycle $[A] \in Z_k$ enumerates all the k -tuples of $\phi_1(A) \subset Z$ with the same image in S if and only if $p|_{\phi_1(A)}$ is curvilinear. We recall that a map $p : Z \rightarrow S$ is called curvilinear when $\dim(\Omega_p^1(z)) \leq 1, \forall z \in Z$

(with Ω_p^1 the relative differentials (see B.2.7 of [9])) and that, if $\dim(Z) = n$, $\dim(S) = m$ and

$$\overline{S_i}(p) := \{z \in Z : \text{rango}(dp) \leq j\} = \{z \in Z : \dim(\Omega_p^1(z)) \geq n - j\},$$

where $j = \min(n, m) - i$, is the locus of Thom-Boardman singularities of p (see [16]), then if $n \leq m$, p curvilinear $\Leftrightarrow \overline{S_2}(p) = \emptyset$, and if $n \geq m$, p curvilinear implies that $\overline{S_2}(p) = \emptyset$.

Lemma 5.3. *The cycle $[W] = [X_6 \cap V_\sigma \cap \Pi]$ enumerates all the (ordered) 6-tuples of $\phi_1(W)$ with the same image in S if and only if $\dim(\Lambda_S) \leq 3$.*

Proof. Taking suitable scheme-theoretic inverse images of X through the inductive construction of X_6 and then going down to X by appropriated projections, one can define a subscheme $X_\sigma \subset X$ parametrizing the points $x \in X$ such that $\exists x_1, x_2, \dots, x_5 \in X$ with $f(x) = f(x_1) = \dots = f(x_5)$, and the first k_1 points (included x) are i.n., the next k_2 points are i.n., and the last k_r points are i.n., where i.n. means infinitely near points (see 2.3. of [7]).

Recalling the previous definitions, we have

$$\phi_1(W) = j(X_\sigma \cap f^{-1}\Lambda_S).$$

Since j is a regular inclusion and $f = p \circ j$, then $p|_{\phi_1(W)}$ will be curvilinear when $f|_{X_\sigma \cap f^{-1}\Lambda_S}$ is curvilinear. Now, f curvilinear $\Leftrightarrow \overline{S_2}(f) = \emptyset$, for $\dim(X) = \dim(S)$. We then study the Thom-Boardman singularities' locus of f ,

$$\overline{S_2}(f) = \left\{x \in X : \dim(\Omega_f^1(x)) \geq 2\right\} = \left\{x \in X : \text{rk}(df_x) \leq \dim(X) - 2\right\},$$

with $df_x : T_x X \rightarrow T_{f(x)} S$, the map induced by f on the tangent spaces.

By definition of $\overline{S_2}(f)$, we have $\text{codim}(\overline{S_2}(f), X) \leq 2 \cdot 2 = 4$ (see (III, 5) of [16]). Checking that, in fact, for our f this codimension is 4, the proof is finished because in that case, when $\dim(\Lambda_S) \leq 3$, $f^{-1}(\Lambda_S)$ can be chosen so that $f^{-1}(\Lambda_S) \cap \overline{S_2}(f) = \emptyset$ and it is easy to see that $\dim(\Lambda_S) \leq 3 \Leftrightarrow r \geq 3$.

Let us verify that $\text{codim}(\overline{S_2}(f), X) = 4$. We recall from (2.1) that, fixed $P \in \mathbf{P}^3$, a cubic surface of \mathbf{P}^3 with P singular, has equation $x_3 f_2 + f_3 = 0$, up to change of coordinates. The plane where the conic and cubic curves associated to f_2 and f_3 , respectively, live is fixed and all possible surfaces correspond to all possible pairs of conic and cubic.

If $R = \mathcal{O}_{\mathbf{P}^2}(2) \oplus \mathcal{O}_{\mathbf{P}^2}(3)$ (see Proposition 3.11), then one gets $X_P \hookrightarrow P(H^0(R)) \times \mathbf{P}^2 \cong \mathbf{P}^{15} \times \mathbf{P}^2$ intersecting every pair of conic and cubic curves and, consequently, the part of $T_x X$ living in $T_w \mathbf{P}^2$, which eventually will contribute to the locus $\overline{S_2}(f)$ is the intersection of the Zariski tangent spaces of the two curves at the point w . In that way, $x \in \overline{S_2}(f)$ when this intersection has dimension 2 and so $\text{codim}(\overline{S_2}(f), X)$ is the number of conditions for a pair of plane curves of degree 2 and 3 having an intersection of their tangent spaces at a common point of dimension 2. It is easy to see that this number is 4 (because that happens if and only if the two curves are singular at the common point). \square

Lemma 5.4. *With the previous notation:*

$$\int_{Z_6} [W] = \int_S \tilde{n}(\sigma) \cdot [S_\sigma \cap \Lambda_S],$$

where $\tilde{n}(\sigma)$ is an integer determined by the partition σ according to the following table:

σ	$\tilde{n}(\sigma)$
(1, 1, 1, 1, 1, 1)	$6! = 5! \cdot 6$
(2, 1, 1, 1, 1)	$24 = 4! \cdot 1$
(3, 1, 1, 1)	$6 = 3! \cdot 1$
(2, 2, 1, 1)	$4 = 2 \cdot 2$
(4, 1, 1)	$2 = 2 \cdot 1$
(3, 2, 1)	1
(2, 2, 2)	$6 = 2 \cdot 3$

Proof. By construction, $p|_{\phi_1(W)} = S_\sigma \cap \Lambda_S$, so

$$\int_{Z_6} [W] = \int_S \deg((p \circ \phi_1)|_W) [S_\sigma \cap \Lambda_S],$$

and the factor $\tilde{n}(\sigma)$ is $\deg((p \circ \phi_1)|_W) = \deg(\phi_1|_W) \cdot \deg(p|_{\phi_1(W)})$.

When $p : Z \rightarrow S$ is curvilinear, there is a natural closed inclusion $Z_k \subset Z^{\times k} = Z \times_S \cdots \times_S Z$, which is an isomorphism outside the diagonals. Each permutation between the factors of $Z^{\times k}$ induces an automorphism of Z_k (see 4.7.2. of [18]). In our case, when $\sigma = (k_1, \dots, k_r)$ and $r \geq 3$, then $p|_{\phi_1(W)}$ is curvilinear; W is included in the intersection

$$\Delta_\sigma = \Delta_{1, \dots, k_1} \cap \Delta_{k_1+1, \dots, k_1+k_2} \cap \cdots \cap \Delta_{k_1+\dots+k_{r-1}+1, \dots, 6} \subset Z^{\times 6}$$

(see 3.3 of [7]) and, this time, the automorphisms of W are the permutations interchanging diagonals of the same size.

Since the image by ϕ_1 of a 6-tuple in W is its first element, then:

- $\deg(\phi_1|_W)$ is the number of automorphisms of W fixing the first coordinate (see 3.3 of [7]),

- if the first element on the partition $\sigma = (k_1, k_2, \dots, k_r)$ is repeated l times, then $\deg(p|_{\phi_1(W)}) = l$. \square

Lemma 5.5. *With the previous notation,*

$$\int_{Z_6} [X_6 \cap V_\sigma \cap \Pi] = \int_{Z_6} [X_6] \cdot [V_\sigma] \cdot [\Pi].$$

Proof. The cycle $[X_6 \cap V_\sigma \cap \Pi]$ is zero-dimensional (we have seen that

$$\int_{Z_6} [X_6 \cap V_\sigma \cap \Pi] = \int_S \tilde{n}(\sigma) \cdot [S_\sigma \cap \Lambda_S]$$

and that $[S_\sigma \cap \Lambda_S]$ is zero-dimensional). We recall from 8.1.11 of [9] that if U and V are two non-singular varieties, it is enough that the intersection is transversal at generic points in order to have $[U \cap V] = [U] \cdot [V]$. In our case, this result applies because:

- $\Pi = \phi^{-1}(\Lambda)$ is the inverse image by the smooth morphism ϕ of a linear space $\Lambda \subset \mathbf{P}^{19}$ and therefore Π intersects every component of $X_6 \cap V_\sigma$.

- The full intersection is transversal at each point, because their image by ϕ in \mathbf{P}^{19} is transversal.

- $X_6 \cap V_\sigma$ is also transversal since, if it were not, $X_6 \cap V_\sigma \cap \Pi$ and its image in \mathbf{P}^{19} will be non-reduced and this is not the case. \square

Now, the proof of Theorem 5.1 is complete and the problem of getting the degree of a stratum C_σ consists in determining $[X_6]$ and computing $[X_6] \cdot [V_\sigma] \cdot [\Pi]$ in the ring $A^*(Z_6)$. We are going to get $[X_6]$ using a result by Z. Ran [24] based on the formula of residual intersection of W. Fulton and D. Laksov [10].

Proposition 5.6. *With the previous notation,*

$$[X_6] = \prod_{j=1}^6 \left((p_{j,j}^j)^*([X]) - [D^{j,6}] \left\{ \frac{(p_{j,j}^j)^* c(E)}{1 + [D^{j,6}]} \right\}_1 \right)$$

where $p_{j,j}^j = p_{2,2}^2 \circ p_{2,3}^3 \circ \cdots \circ p_{2,j}^j$, $D^{j,6} = D_{1,j}^6 + D_{2,j}^6 + \cdots + D_{j-1,j}^6$ and $D_{i,j}^6 = (p_{i,j}^6)^* D_{1,j-i+1}^{j-i+1}$ with $p_{i,j}^6$ a composition of projections from Z_6 to Z_{j-i+1} onto the first factor if $i = 1$ and onto the second factor if $i > 1$.

Proof. To get $[X_6]$ we can apply Theorem (4.2) from [24] because:

- since $X = (s_E)_0$ where s_E is a section of the vector bundle E over Z , then $c(N_X Z) = j^* c(E)$.

- the morphisms $X_t \rightarrow X_{t-1}$ induced by $f : X \rightarrow S$ are local complete intersection (l.c.i.) of the same codimension, for $t = 1, \dots, k$, so the residual intersection formula of [10] applies. To see this, we first note that f is l.c.i. of codimension 0 because $X \hookrightarrow Z$ is a closed regular inclusion of codimension 2 and Z is smooth over S with relative dimension 2. Now, in general, if X is Cohen-Macaulay and f is l.c.i. of codimension n , then a necessary and sufficient condition for every morphism $X_t \rightarrow X_{t-1}$ with $t = 1, \dots, k$, to be l.c.i. of codimension n is that all of these morphisms are of the same codimension as f (see [17]). As we saw in §3, X is smooth and X is also the zero scheme of a regular section s_E of the vector bundle E over Z . In this case, by the inductive constructions of §4, each X_t is again smooth and it is also the zero locus of a regular section of the secant bundle E_t over Z_t , so $\text{codim}(X_t, Z_t) = \text{rank}(E_t)$ (for Z_t is Cohen-Macaulay (in fact, smooth)) and, finally $X_t \rightarrow X_{t-1}$ is l.c.i. of codimension 0 for all t . The formula of residual intersection ((4.2) of [24]) says that

$$[X_k] = \prod_{j=1}^k \left((p_{j,j}^j)^*([X]) - [D^{j,k}] \left\{ \frac{(p_{j,j}^j)^* c(E)}{1 + [D^{j,k}]} \right\}_{j-1} \right),$$

where $\{\cdot\}_r$ denotes the homogeneous component of $\dim r$ and $c(E)$ and $[X] = c_2(E)$ are known (see Lemma 3.10). \square

Lemma 5.7. *If V_σ is the intersection of (the pullback of) the exceptional divisors E_1, \dots, E_j , then $[V_\sigma] = [E_1] \cdots [E_j]$.*

Proof. Using 8.1.11 of [9], as in Lemma 5.5, it is enough to note that the intersection of exceptional divisors $V_\sigma = \bigcap_{l=1}^j E_l$ is not only transversal but it is smooth. \square

6. EFFECTIVE COMPUTATION OF DEGREE AND INTERSECTION RING OF SPACES FOR k -TUPLES

In order to determine $\deg(C_\sigma)$ we have to compute $[X_6] \cdot [V_\sigma] \cdot [\Pi]$ in $A^*(Z_6)$.

In general, if Z is a projective bundle over S and $A^*(S)$ is known, we can determine completely $A^*(Z_k)$ for any $k = 1, 2, \dots$ (see Proposition 6.2) and, in our case, it turns out to be a polynomial ring on 23 variables, modulo 38 relations, so effective computations in this ring, using Gröbner bases' methods are very costly.

In spite of this fact, in order to effectively compute the product $[X_6] \cdot [V_\sigma] \cdot [\Pi]$ in $A^*(Z_6)$ it is not necessary to know the whole ring, as we will see in a moment.

In this section the notation is the same as in §4, writing $ec_{k,1} := [D_{1,k}^k]$ for the class of the exceptional divisor in Z_k . We will apply several times a result by S. Keel [15] about the intersection ring of a blown-up variety, saying that, if $i : X \hookrightarrow Y$ is a regular inclusion of codimension d , $i^* : A^*(Y) \rightarrow A^*(X)$ is surjective and $\tilde{Y} \rightarrow Y$ is the blow up of Y along X with exceptional divisor \tilde{X} , then

$$(6.1) \quad A^*(\tilde{Y}) \cong \frac{A^*(Y)[T]}{(P(T), T \cdot \ker(i^*))}$$

where $P(T) \in A^*(Y)[T]$ is any polynomial with constant term equal to $[X]$ and whose restriction to $A^*(X)$ is the Chern polynomial of the normal bundle $N_X Y$.

Proposition 6.1. *Let $p : Z \rightarrow S$ be a smooth morphism and suppose that $A^*(Z)$ is a finitely generated $A^*(S)$ -module. For $k = 1, 2, \dots$ let $a \in A^*(Z_k)$ be a zero-dimensional cycle consisting in a product of pullbacks of cycles from $A^*(Z_j)$ with $j \leq k$ and some power of the class of the exceptional divisor in Z_k . Then, the degree of the cycle a can be computed as the degree of a zero-dimensional cycle in $A^*(Z)$.*

Proof. We prove this by induction in k . The cases $k = 1, 2$ are easy to work out, so let's suppose that we can determine the degree of a zero-dimensional cycle in $A^*(Z_k)$ going down to $A^*(Z_{k-1})$ and let's prove that we can do the same with one in $A^*(Z_{k+1})$. By Keel's result (6.1) described before, $a = ec_{k+1,1}^\alpha \cdot b_{k+1}^* c$ with $c \in A^*(Z_k \times_{Z_{k-1}} Z_k)$. We distinguish two cases:

1. $\alpha = 0$. In this case, as $b_{k+1*} b_{k+1}^* c = c$ (see 6.7.(b) of [9]), we have

$$\int_{Z_{k+1}} a = \int_{Z_{k+1}} b_{k+1}^* c = \int_{Z_k \times_{Z_{k-1}} Z_k} c = \int_{Z_k} pr_{1*} c.$$

To get $pr_{1*} c$, we recall the diagram

$$\begin{array}{ccccccc} E_{k+1} & \xrightarrow{j_{k+1}} & Z_{k+1} & & & & \\ g_{k+1} \downarrow & & \downarrow b_{k+1} & & & & \\ \Delta_k & \xrightarrow{l_k} & Z_k \times_{Z_{k-1}} Z_k & \xrightarrow{pr_2} & Z_k & & \\ & & \downarrow pr_1 & & \downarrow \iota_{k-1} \circ p_{1,k-1}^k & & \\ & & Z_k & \xrightarrow{p_{2,k}^k} & Z_{k-1} & & \end{array}$$

Since, by hypothesis $c \in A^*(Z_k \times_{Z_{k-1}} Z_k)$ can be written as $c = pr_1^* d \cdot pr_2^* e$ with $d, e \in A^*(Z_k)$ then, denoting by $q_1 := \iota_{k-1} \circ p_{1,k-1}^k$ and $q_2 := p_{2,k}^k$ and using the projection formula, we have

$$\int_{Z_k} pr_{1*} c = \int_{Z_k} d \cdot pr_{1*} pr_2^* e = \int_{Z_k} d \cdot q_2^* q_{1*} e,$$

where the last equality is (1.7) of [9].

To know the product $d \cdot q_2^* q_{1*} e \in A^*(Z_k)$ we need to determine $q_{1*} e$. Now, by Keel's result (6.1) again, $e \in A^*(Z_k)$ must be $e = ec_{k,1}^\gamma \cdot b_k^* u$ for some γ , with $u \in A^*(Z_{k-1} \times_{Z_{k-2}} Z_{k-1})$. Then:

- (a) If $\gamma > 0$, $q_{1*} e = \iota_{k-1*} \circ pr_{1*} \circ b_{k*}(ec_{k,1}^\gamma \cdot b_k^* u) = \iota_{k-1*}(pr_{1*}(u \cdot b_{k*}(ec_{k,1}^\gamma)))$ by projection formula. Writing $U := pr_{1*}(u \cdot b_{k*}(ec_{k,1}^\gamma))$, we have that $U = (-1)^{\gamma-1} m_{\gamma-2} \cdot l_{k-1}^* u \in A^*(Z_{k-1}) \cong A^*(\Delta_{k-1})$, being $m_{\gamma-2}$ the term of degree $\gamma - 2$ en $c(N_{\Delta_{k-1}})^{-1}$ (see 4.2.2. of [9]).

It is easy to get \overline{U} such that $U = \iota_{k-1}^* \overline{U}$, and then: $q_{1*}e = \overline{U}$ so

$$\int_{Z_{k+1}} a = \int_{Z_k \times_{Z_{k-1}} Z_k} c = \int_{Z_k} d \cdot q_2^* \overline{U},$$

which is known by induction.

- (b) If $\gamma = 0$, $q_{1*}e = \iota_{k-1*} \circ pr_{1*} \circ b_{k*}b_k^*u = \iota_{k-1*} \circ pr_{1*}u$. Since $u \in A^*(Z_{k-1} \times_{Z_{k-2}} Z_{k-1})$, we have again, by hypothesis, that $u = pr_1^*f \cdot pr_2^*h$ with $f, h \in A^*(Z_{k-1})$ and, consequently $pr_{1*}u = f \cdot pr_{1*}pr_2^*h$. Calling now $V := pr_{1*}u$, and working as in (a), we find

$$\int_{Z_{k+1}} a = \int_{Z_k \times_{Z_{k-1}} Z_k} c = \int_{Z_k} d \cdot q_2^* \overline{V},$$

which is known again by induction.

2. If $a = ec_{k+1,1}^\alpha \cdot b_{k+1}^*c$ with $\alpha > 0$, then the situation is similar to (a) when we computed U , so using 4.2.2. of [9], we conclude

$$\int_{Z_{k+1}} a = \int_{Z_k} pr_{1*} \circ b_{k+1*}a = \int_{Z_k} (-1)^{\alpha-1} m_{\alpha-2} \cdot l_k^*c$$

(where again $m_{\alpha-2}$ is the term of degree $\alpha - 2$ in $c(N_{\Delta_k})^{-1}$), and this is known by induction. \square

In order to effectively apply Proposition 6.1, we need to determine $c(N_{\Delta_k})$ for every k . From §4, we know that N_{Δ_k} is isomorphic to the relative tangent bundle $T_k := T_{p_{2,k}^k}$. The total Chern class of T_k can be computed inductively, starting with $c(T_1)$ the class of a relative tangent bundle (given by B.5.8. of [9]). Using standard results about Chern classes (see 15.4.(iv), 15.4.2. of [9]), one gets

$$c(T_k) = (1 + ec_{k,1}) \sum_{i=0}^2 (1 - ec_{k,1})^i (p_{1,k-1}^k)^* c_{2-i}(T_{k-1}).$$

Implementing the inductive method given in Proposition 6.1 in Maple [20] we finally get the degree of the strata of singular cubic surfaces with a conic node mentioned in Theorem 5.1:

σ	1^6	21^4	31^3	$2^2 1^2$	41^2	321	2^3
Singularity	A_1	$2A_1$	$A_1 A_2$	$3A_1$	$A_1 A_3$	$2A_1 A_2$	$4A_1$
Parameters	3	2	1	1	0	0	0
$deg(C_\sigma)$	32	280	1200	800	2200	3420	305

Note that for $\sigma = (2, 2, 2)$, we have subtracted the contribution of surfaces with a uninode which is $\frac{1}{4}deg(C_U)$, according to Theorem 5.1. The degree of the stratum C_U , 260, can be computed separately at least in two ways. First, as a particular case of some formulas for the degree of varieties parametrizing quadrics with vertex of specified dimension, given by I. Vainsencher [28] (1979). Secondly, as we do in [30], using a result by J. Harris and L. W. Tu [11] (1984) about the cohomology class of the degeneracy locus of a symmetric bundle map.

For completeness, we describe here the intersection ring of the space Z_k of k -tuples of Z with the same image in S (defined in §4), when $Z = P(E) \xrightarrow{\pi} S$ is the projectivization of a vector bundle E over a smooth scheme S .

Proposition 6.2. *Let Z be a projective bundle over a smooth scheme S and let Z_k be the space of k -tuples of Z over S defined in §4. Then*

$$A^*(Z_k) \cong \frac{A^*(S)[g_1, \dots, g_m]}{(r_1, \dots, r_n)},$$

where $m = \sum_{j=1}^k j$, $n = \sum_{j=1}^k (2j - 1)$; k of the generators g_j are (pullback of) the hyperplane class of Z ; the rest are (pullback of) exceptional divisors' classes and the relations r_j are determined by 3.3.(b), Ex. 8.3.4. of [9] and Keel's result (6.1) above.

Proof. We see this by induction on k , using repeatedly 3.3.(b), Ex. 8.3.4. of [9], to determine the ring of a projective bundle and Keel's work (6.1) from [15], for the ring of a blown-up scheme. One starts with the projective bundle Z and its Chow ring $A^*(Z)$ written in terms of $A^*(S)$. For $k = 2$, by construction, the fiber product $Z \times_S Z$ is another projective bundle, one finds that $A^*(Z \times_S Z) \cong A^*(Z) \otimes_{A^*(S)} A^*(Z)$, and $Z_2 = Bl_\Delta(Z \times_S Z)$ is a blown-up projective bundle. When $k = 3$ checking that $Z_2 \times_Z Z_2$ is a projective bundle over Z_2 (by 3.4. of [13]) one has that Z_3 is a projective bundle blown-up two times.

At the step $(k - 1)$, one supposes that Z_{k-1} is a projective bundle over Z_{k-2} blown-up $(k - 1)$ times and therefore its Chow ring can be written as it is said in the proposition. Now, again by definition, the fiber product $Z_{k-1} \times_{Z_{k-2}} Z_{k-1}$ is another projective bundle over Z_{k-1} blown-up $(k - 1)$ times and, finally, for Z_k one adds another blow-up getting $A^*(Z_k)$ as the ring described in the proposition (for details see [30]). \square

REFERENCES

1. A. B. Altman and S. L. Kleiman, *Foundations of the theory of Fano schemes*, Compos. Math. **34** (1977), 3-47. MR 58:27967
2. V. I. Arnold, *Normal forms of functions near degenerate critical points, the Weyl groups A_k , D_k and E_k and Lagrangian singularities*, Func. Anal. Appl. **6** (1972), 254-272. MR 50:8595
3. V. I. Arnold, *Normal forms of functions in the neighborhood of degenerate critical points*, Russ. Math. Surv. **29** (2) (1974), 11-49. MR 58:24324
4. J. W. Bruce, *A stratification of the space of cubic surfaces*, Math. Proc. Cambridge **87** (1980), 427-441. MR 81c:14022
5. J. W. Bruce and C. T. C. Wall, *On the classification of cubic surfaces*, J. Lond. Math. Soc. (2) **19** (1979), 245-256. MR 80f:14021
6. A. Cayley, *A memoir on cubic surfaces*, Phil. Trans. Roy. Soc. **159** (1869), 231-236.
7. S. J. Colley, *Enumerating stationary multiple-points*, Adv. Math. **66** (1987), 149-170. MR 89g:14042
8. D. F. Coray and I. Vainsencher, *Enumerative formulae for ruled cubic surfaces and rational quintic curves*, Comment. Math. Helv. **61** (1986), 510-518. MR 87m:14061
9. W. Fulton, *Intersection Theory*, Springer-Verlag, (1984). MR 85k:14004
10. W. Fulton and D. Laksov, *Residual intersections and the double point formula*, Real and complex singularities, ed. P. Holm, Sijthoff & Nordhoff, (1977), 171-177. MR 58:27971
11. J. Harris and L. W. Tu, *On symmetric and skew-symmetric determinantal varieties*, Topology, Vol. **23**, No.1 (1984), 71-84. MR 85c:14032
12. R. Hartshorne, *Algebraic Geometry*, Springer-Verlag, (1977). MR 57:3116
13. S. Iitaka, *Algebraic Geometry*, Springer Verlag, (1982). MR 84j:14001
14. S. Katz and S. A. Stromme, SCHUBERT: a MAPLE package for intersection theory. Available by anonymous ftp from ftp.math.okstate.edu or linus.mi.uib.no, cd pub/schubert.
15. S. Keel, *Intersection theory of moduli space of stable N -pointed curves of genus zero*, Trans. Amer. Math. Soc. **330** (1992), 545-574. MR 92f:14003
16. S. L. Kleiman, *The enumerative theory of singularities*, Real and complex singularities, ed. P. Holm, Sijthoff & Nordhoff, (1977), 297-396. MR 58:27960

17. S. L. Kleiman, *Multiple-point formulas. I: Iteration*, Acta Math. **147** (1981), 13-49. MR **83j**:14006
18. S. L. Kleiman, *Multiple-point formulas. II: the Hilbert scheme*, Enumerative Geometry, (Proc. Conf., Sitges, 1987), ed. S. Xambó Descamps, LNM **1436**, Springer-Verlag (1989), 101-138. MR **92a**:14062
19. D. Laksov, *Secant bundles and Todd's formula for the double points of maps into \mathbf{P}^n* , Proc. Lond. Math. Soc. **37** (1978), 120-142. MR **80c**:14007
20. MAPLE V, Copyright (c), 1981-1990, University of Waterloo.
21. A. Mattuck, *Secant bundles on symmetric products*, Am. J. Math. **87** (1965), 779-797. MR **33**:7345
22. J. M. Miret and S. Xambó Descamps, *On the geometry of nodal plane cubics: the condition p* , Enumerative Geometry: Zeuthen Symposium, ed. S. L. Kleiman y A. Thorup, Contemp. Math. **123**, A.M.S. (1991), 169-187. MR **93e**:14064
23. J. M. Miret and S. Xambó Descamps, *Rational equivalence on some families of plane curves*, Ann. I. Fourier, Grenoble, **44**, 2 (1994), 323-345. MR **95g**:14006
24. Z. Ran, *Curvilinear enumerative geometry*, Acta Math. **155** (1985), 81-101. MR **86m**:14040
25. L. G. Roberts, *Lines in singular cubics*, Queen's Papers in Pure and Appl. Math.
26. L. Schläfli, *On the distribution of surfaces of the third order into species*, Phil. Trans. Roy. Soc. **153** (1864), 193-247.
27. R. L. E. Schwarzenberger, *The secant bundle of a projective variety*, Proc. London. Math. Soc. **14** (1964), 369-384. MR **28**:3042
28. I. Vainsencher, *The degrees of certain strata of the dual variety*, Compos. Math. **38** (1979), 241-252. MR **81g**:14024
29. I. Vainsencher, *Counting divisors with prescribed singularities*, Tran. Amer. Math. Soc., Vol. **267**, No. 2, (1981), 399-422. MR **83b**:14021
30. María J. Vázquez-Gallo, *Grado de variedades de superficies singulares*, Ph.D. thesis, University Autonoma of Madrid (1997).

DEPARTAMENTO DE MATEMATICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD AUTONOMA DE MADRID, MADRID, 28049, SPAIN

E-mail address: rafael.hernandez@uam.es

DEPARTAMENTO DE MATEMATICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD AUTONOMA DE MADRID, MADRID, 28049, SPAIN

E-mail address: mjesus.vazquez@uam.es