

TOPOLOGICAL HORSESHOES

JUDY KENNEDY AND JAMES A. YORKE

ABSTRACT. When does a continuous map have chaotic dynamics in a set Q ? More specifically, when does it factor over a shift on M symbols? This paper is an attempt to clarify some of the issues when there is no hyperbolicity assumed. We find that the key is to define a “crossing number” for that set Q . If that number is M and $M > 1$, then Q contains a compact invariant set which factors over a shift on M symbols.

1. INTRODUCTION

The most basic and striking results in chaotic dynamics are perhaps the results about Smale horseshoes [S]. The theory concerns a set Q (usually diffeomorphic to a rectangle) in a two-dimensional manifold M and a diffeomorphism $f: Q \rightarrow M$. Using only hypotheses on the first iterate of f on Q , one obtains conclusions about all iterates of f . In particular, Smale concludes that there is a compact invariant set Q_I in Q which is homeomorphic to a shift on M symbols, where M is often taken to be 2. In this paper we will call M the “crossing number” and our definition of this number for more general maps and sets Q is central to our analysis of the chaos present in our dynamical systems.

Results such as Smale’s are useful for numerical studies of systems with positive entropy because there are unavoidable errors in the computation of $f(Q)$ that can be made small by careful computation, but the errors in the computation of $f^n(Q)$ increase exponentially fast as n increases, due to the positive entropy. Hence hypotheses on $f(Q)$ are easier to verify directly than hypotheses on $f^n(Q)$.

However, Smale’s assumption of hyperbolicity of f on Q is in practice difficult to verify numerically. When the hyperbolicity assumption is discarded, the dynamics remain just as rich as before in that there is a compact invariant set Q_I in Q that factors over a shift on M symbols, although the map need not be one-to-one. K. Burns and H. Weiss [BW] call such maps “geometric horseshoes,” and the term “topological horseshoes” is also often used. (In their paper aimed at discussing situations where stable and unstable manifolds cross but do not cross necessarily transversally, they concluded that there is a geometric horseshoe for some iterate of the map.) K. Mischaikov et al. [CKM, CM, MM] and A. Szymczak [S] investigate such situations using simplicial decompositions that can be analyzed rigorously numerically, showing for example that there is a subshift of finite type in the Lorenz

Received by the editors March 16, 1998 and, in revised form, December 21, 1998.

1991 *Mathematics Subject Classification*. Primary 58F12, 54F20; Secondary 54F50, 58F20.

Key words and phrases. Topological horseshoe, geometric horseshoe, chaos, shift dynamics, connection, preconnection, crossing number, expander, symbol set.

This research was supported by the National Science Foundation, Division of Mathematical Sciences.

system. Example 3(c) in Section 3 of this paper is one in which shift dynamics is predicted by our results but to which the results of Szymczak and Mischaikov et al do not apply (because no isolating neighborhood exists).

Our goal here is to formulate the idea of the crossing number M and to show that with this formulation and a few other hypotheses, it follows that there exists an invariant set Q_I in Q that factors over an M -shift. (The M -shift is one-sided if f is only assumed to be continuous, or two-sided if f is a homeomorphism.) In particular, we make no assumption that there exists a Markov partition on Q , nor do we assume that the continuous map f is differentiable at any point.

If f is continuous, we say $\mathcal{S} := \{S_i\}_{i=1}^M$ is a *set of symbol sets* if for each sequence $(j_i)_{i=0}^\infty$ of integers in $\{1, 2, \dots, M\}$, there is a trajectory (x_i) (where $x_{i+1} = f(x_i)$ for all i) such that x_i is in S_{j_i} . If f is a homeomorphism, we say $\mathcal{S} = \{S_i\}_{i=1}^M$ is a *set of symbol sets* if for each sequence $(j_i)_{i=-\infty}^\infty$ of integers in $\{1, 2, \dots, M\}$, there is a trajectory (x_i) such that x_i is in S_{j_i} . If the S_i are disjoint, it follows that the invariant set S contained in $\bigcup S_i$ can be mapped continuously onto a shift on M symbols. Each S_i in \mathcal{S} is, naturally enough, a *symbol set*.

Establishing the existence and dynamics of Q_I can be broken into two problems: first the construction of the symbol sets S_i , and second showing that they yield the required dynamics. The set of symbol sets S_i we construct are mutually disjoint. The construction is carried out in two lemmas. The second step, where they yield the desired dynamics, is carried out via a lemma that we call the ‘‘Chaos Lemma’’. It was formulated with Sahin Koak to be able to handle a variety of problems which will be investigated in a separate paper [KKY]. We only sketch the proof in this paper. We also give a number of examples showing how our results can be applied. In the last section, we show that our results can be applied in a still wider variety of situations, by proving two lemmas that demonstrate that the horseshoe hypotheses Ω (listed below) hold in the presence of a weaker set of hypotheses.

Throughout this paper (except in the Chaos Lemma, and the last section, where we relax our hypotheses further) we make the following five assumptions:

Ω_X : X is a separable metric space.

Ω_Q : $Q \subset X$ is locally connected and compact.

Ω_f : The map $f: Q \rightarrow X$ is continuous.

Ω_E The set $end_0 \subset Q$ and $end_1 \subset Q$ are disjoint and compact, and each component of Q intersects both end_0 and end_1 .

Ω_M : Q has crossing number (defined below) $M \geq 2$.

We refer to the five hypotheses $\Omega_X, \Omega_Q, \Omega_f, \Omega_E$, and Ω_M collectively as the *horseshoe hypotheses* Ω . A *connection* Γ is a compact connected subset of Q that intersects both end_0 and end_1 . (We use the term ‘connection’ because one might say Γ connects end_0 to end_1 .) Note that by hypothesis Ω_E , each component of Q is a connection. An example of a connection is a path in Q intersecting both end_0 and end_1 . A *preconnection* γ is a compact connected subset of Q for which $f(\gamma)$ is a connection. We define the crossing number M to be the largest number such that every connection contains at least M mutually disjoint preconnections. We say a set S is *invariant* if $f(S) = S$.

Our main result is the following theorem, which follows from the Chaos Lemma and Lemma 12.

Theorem 1. *Assume the horseshoe hypotheses Ω . Then there is a closed invariant set $Q_I \subset Q$ for which $f|_{Q_I}$ is semiconjugate to a one-sided M -shift. (If f is a homeomorphism, then $f|_{Q_I}$ is also semiconjugate to a two-sided M -shift.)*

By assumption Ω_M , each connection contains at least M mutually disjoint preconnections. We prove the theorem by constructing M compact mutually disjoint sets S_1, S_2, \dots, S_M such that each connection Γ contains a preconnection in $S_i \cap \Gamma$ for each $i = 1, \dots, M$. This construction is quite difficult despite the fact that in most examples it is quite easy to find S_1, \dots, S_M . Once the sets S_1, \dots, S_M are shown to exist, we can apply the following lemma by letting E be the set of all connections. As stated below however, the lemma is more general than needed in that the sets in E need not be connected.

Lemma 2. The Chaos Lemma. *Assume Ω_X, Ω_Q , and Ω_f . Let M be an integer greater than or equal to 2. Let E be a nonempty set of nonempty compact sets such that for each $\gamma \in E$ there exists a compact $\gamma_i \subset \gamma \cap S_i$ for which $f(\gamma_i) \in E$. Let Q_I be the largest invariant set in $\bigcup S_i$. Then $f|_{Q_I}$ is semiconjugate to a one-sided shift on M symbols. If $f|_{Q_I}$ is one-to-one on Q_I , the restricted map $f|_{Q_I}$ is semiconjugate to a two-sided shift on M symbols.*

2. NOTATION, DEFINITIONS, BACKGROUND

We use the term ‘space’ to mean ‘separable metric space’. The word ‘component’ means ‘connected component’. In the results that follow, we consider the topology on the closed subset Q of the space X only. Thus, ‘open’ means ‘open relative to Q ’ and ‘closed’ means ‘closed relative to Q ’. The sets in which we are interested are all contained in Q , and this means the restriction to Q avoids our having to add repeatedly ‘ $\cap Q$ ’ to the sets discussed. If A is a subset of Q , then we use the notation A° , \overline{A} , and ∂A to denote the interior, closure, and boundary of A in Q , respectively. A *continuum* is a compact, connected metric space. In this paper we allow a continuum to consist of just one point or to be the empty set. A subset of a continuum which is itself a continuum is a *subcontinuum*. If A and B are nonempty closed subsets of a space X , and K is a continuum in X which intersects both A and B , then K is a *continuum irreducible from A to B* if no proper subcontinuum of K intersects both A and B . If A and B are nonempty closed subsets of a space X , and K is a continuum in X which intersects both A and B , then K contains a continuum K' irreducible from A to B . A *chain* is a finite collection $C = \{C_0, C_1, \dots, C_n\}$ of sets listed so that $C_i \cap C_j \neq \emptyset$ if and only if $|i - j| \leq 1$. The members C_i of the chain C are called *links*. A *tiling chain* is a chain $C = \{C_0, C_1, \dots, C_n\}$ each member of which is a closed set with nonempty interior and which has the property that for each $i \neq j$, $C_i \cap C_j = \partial C_i \cap \partial C_j$. For $0 \leq i < n$, we say that $\partial C_i \cap C_{i+1}$ is the *right boundary* of the link C_i , and for $0 \leq i \leq n$, we say that $\partial C_i \cap C_{i-1}$ is the *left boundary* of the link C_i .

Note that in the discussion of the horseshoe hypotheses in the previous section, the sets end_0 and end_1 need not be connected subsets of Q —they must only be disjoint closed subsets of Q . We denote the set $end_0 \cup end_1$ with the notation *ends*, and we also call the sets end_0 and end_1 the “ends” of Q .

3. EXAMPLES

We begin with the most obvious and arguably simplest example, the Smale horseshoe with crossing number 2. This system serves as a nice example that




FIGURE 1. **Smale horseshoe with crossing number $M = 2$.** The set Q is the rectangle pictured, with end_0 being the left boundary of Q and end_1 being the right boundary of Q . The shaded region is the image $f(Q)$.

satisfies the horseshoe hypotheses Ω , and it is not hard to see how the ends and the set Q should be chosen, nor what the natures of the connections and preconnections are. We then move to systems on the unit interval, to illustrate the necessity of the properties we require a map f and associated set Q have to satisfy the horseshoe hypotheses Ω on Q . The last examples are less obvious applications of our results.

Example 3. Smale horseshoe with crossing number $M = 2$. The set Q and its image are illustrated in Figure 1. The horseshoe map is named f . Figure 2 shows Q and two different connections, both of which are arcs, with their preconnections. Note that Q itself is a connection, and that the vertical bands $(Q \cap f^{-1}(Q))$ are preconnections in Q (although they are not unique).

Example 4. Numerically obtained horseshoe. The topological horseshoe pictured in Figure 3 was obtained as a part of an investigation of a model of fluid flow past a sequence of cylinders. (See [SKGY, KSYG] for detailed discussions (mathematical and physical) of the model, and what can be proved rigorously when assumptions based on the numerical information are used.) The space for the Poincaré map of the flow given by the model is the plane. As is typical of such investigations, once the appropriate quadrangle Q_0 is found (that can be, and was, tedious), verifying horseshoe type behavior of the Poincaré map is usually straightforward. The map is an area preserving diffeomorphism on the plane (a property it inherits from the model), but verifying that the map is hyperbolic in Q_0 is quite difficult.

Example 5. Maps from the unit interval to itself. In these examples the entire space is the set Q . Let $X = [0, 1] = Q$, $end_0 = \{0\}$, and $end_1 = \{1\}$. See Figure 4 for the graphs of these maps. Each map possesses exactly one connection, $[0, 1]$. The first map, f , does not satisfy Ω because $[0, 1]$ does not contain two disjoint preconnections, although it does contain two preconnections. The second map, g , satisfies Ω on $Q = [0, 1]$ with crossing number $M = 2$. Note that $g^{-1}(0)$ is the union of a subinterval I_1 containing 0 and subinterval I_3 containing 1, while




FIGURE 2. **Connections and their preconnections for the Smale horseshoe.** Any continuum that extends from end_0 to end_1 in Q is a connection. Two possible connections are pictured above. Both are arcs. Connection 1 has the minimum number of preconnections possible, namely 2, while connection 2 has 4, due to the wiggly path it takes in travelling from end_0 to end_1 . The preconnections in this case are arcs in the grey-shaded regions which connect the two sides of a grey band.

$g^{-1}(1)$ is a subinterval I_2 in $(0, 1)$. This map possesses two disjoint preconnections, the obvious choice for those being (1) the smallest subinterval of $[0, 1]$ that contains a point of I_1 and a point of I_2 , and (2) the smallest subinterval of $[0, 1]$ that contains a point of I_2 and a point of I_3 . In this example, both end_0 and end_1 are contained in the interior of $g^{-1}(end_0)$. The third map, h , also satisfies Ω with $M = 2$. There are points $p < q$ in $(0, 1)$ such that $h^{-1}(0)$ consists of two points, 0 and q , while $h^{-1}(1)$ consists of two points, p and 1. This map possesses two disjoint preconnections, the only choice for those being (1) the smallest subinterval of $[0, 1]$ that contains 0 and p , and (2) the smallest subinterval of $[0, 1]$ that contains q and 1. The shift dynamics for Example 3(c) are implied by our results, but since there is no isolating neighborhood here (every open set containing the invariant Cantor set also contains periodic orbits not in the Cantor set), the shift dynamics do not follow from Szymczak’s results [S].

Example 6. Fewer conditions to satisfy. One reason locating the quadrangle Q_0 in Example 2 was so tedious was that it was necessary to find one that satisfied the following properties:

1. The quadrangle Q_0 must have sides labelled s_1, s_2, s_3, s_4 and vertices labelled v_1, v_2, v_3, v_4 such that $s_i \cap s_{i+1} = \{v_i\}$ for $i = 1, 2, 3$ and $s_4 \cap s_1 = \{v_4\}$.
2. Two opposite sides, say s_1 and s_3 , map outside Q_0 under F , i.e., $(F(s_1) \cap F(s_3)) \cap Q_0 = \emptyset$.
3. The image of the remaining two sides must not intersect those sides, i.e., $(F(s_2) \cap F(s_4)) \cap (s_2 \cup s_4) = \emptyset$.
4. The set $Q \cap F(Q)$ must contain at least two connected components, each of which intersects both s_1 and s_3 .




FIGURE 3. **Horseshoe from a fluid-flow model.** The figure shows a topological horseshoe obtained from the Poincaré map of a 2-dimensional area preserving fluid flow model. The crosses are the images of the vertices of the quadrilateral Q_0 under the action of the Poincaré map F . The map F is an area preserving diffeomorphism on \mathbf{R}^2 .

Figure 5 illustrates a map f on \mathbf{R}^2 and a rectangle Q in \mathbf{R}^2 such that f satisfies Ω on Q with crossing number $M = 2$, but properties 2 and 3 above are not satisfied (and it is not necessary to choose Q so very carefully). The map f might well be a homeomorphism or even a diffeomorphism, but it most probably is not hyperbolic in Q . Even so, if f is a continuous map, applying our theorem tells us that there is a “Cantor-like” invariant set Q_I in Q that factors over a two-sided two shift. However, without hyperbolicity, preimages of points in the two shift Cantor set need not be points—they need only be closed sets.

Example 7. It does not look like a horseshoe. The map f indicated on the circular region Q in Figure 6 satisfies Ω on Q with $M = 2$. Note that the image of any continuum in Q that intersects both $end(0)$ and $end(1)$, in other words, any connection, must have an image that intersects both $f(end(0))$ and $f(end(1))$, and this means that the connection itself must contain at least two disjoint preconnections.

Example 8. A more complicated example with $M = 14$. We can think of $f: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ as a composition of maps on Q (f is not specified outside Q —it does not matter as long as it is continuous).

1. First apply the map $f_1: Q \rightarrow S^2$ (where S^2 is the unit sphere in \mathbf{R}^3), with f_1 mapping the region between the boundary ∂Q and $end(1)$ (the outer annular region) to the point $(0, 0, 1)$, f_1 mapping the region inside $end(0)$ (the inner annular region) to the point $(0, 0, -1)$, and f_1 being one-to-one on the rest of Q . We can do this so that the set $f_1(end(0))$ is the set of all points (x, y, z) in S^2 with $z \geq 0.999$, and $f_1(end(1))$ is the set of all points (x, y, z) in S^2 with $z \leq -0.999$.




FIGURE 4. **Interval maps.** The graphs of three maps from $[0, 1]$ to itself are pictured above. In these examples, $Q = [0, 1] = X$, and Q is the only connection.

2. Let $B = \{(x, y) : x^2 + y^2 \leq 1\}$, and let \mathbf{v} denote a vector in 3-space with direction chosen so that no line in the same direction as \mathbf{v} intersects $f_1(\text{end}(0)) \cup f_1(\text{end}(1))$ more than once. Then apply the map $f_2 : S^2 \rightarrow B$ by identifying points in S^2 that are on the same line L in the same direction as \mathbf{v} . With an appropriate choice of \mathbf{v} , this can be done so that $f_2(f_1(\text{end}(0)) \cup f_1(\text{end}(1))) \subset B^\circ = \{(x, y) : x^2 + y^2 < 1\}$. Note that no point p in B has more than two points in its preimage $f_2^{-1}(p)$.
3. Next apply a homeomorphism f_3 to contract B in, say, the y -direction, and stretch it in the x -direction, and contract the sets $f_2(f_1(\text{end}(0)))$, and $f_2(f_1(\text{end}(1)))$ so that they have small diameter, obtaining a long thin tape.
4. Finally, apply the map f_4 , which places the long thin band $f_3 \circ f_2 \circ f_1(Q)$ back in the plane, by looping it 8 times on Q while crossing over $\text{end}(0)$ and $\text{end}(1)$, as shown.

Let $f = f_4 \circ f_3 \circ f_2 \circ f_1$. Since any connection has an image that stretches from $f(\text{end}(0))$ to $f(\text{end}(1))$, it must contain at least 14 mutually disjoint preconnections. Note that one of the loops does not completely cross the inner annular region $\text{end}(0)$, and that affects the crossing number $M = 14$. See Figure 7.

Example 9. Expanding balls. Suppose that $X = B_n = \{x \in \mathbf{R}^n : d(x, \mathbf{0}) \leq 1\}$ (where $\mathbf{0}$ denotes the origin and $n \geq 2$). Let $\text{balls} = \{I, II, III, IV, V\}$ denote a mutually disjoint collection of smaller closed balls in $(X)^\circ = \{x \in \mathbf{R}^n : d(x, \mathbf{0}) < 1\}$, as pictured in Figure 8. The map f takes each of the smaller balls homeomorphically onto X , and takes the boundary $\partial X = \{x \in \mathbf{R}^n : d(x, \mathbf{0}) = 1\}$ into itself. For simplicity, we also assume that if $\langle b_{i_j} \rangle_{j=0}^\infty$ is a sequence whose members are the small balls in $\{I, II, III, IV, V\}$, then $\bigcap_{j=0}^\infty f^{-j}(b_{i_j})$ is a single point. Now although there is an invariant Cantor set in X on which the dynamics are conjugate to those of the two-sided 5-shift, f does not satisfy $\Omega(X)$ for any choice of $\text{end}(0)$ or $\text{end}(1)$. With an appropriate choice of Q , $\text{end}(0)$ and $\text{end}(1)$, however, f satisfies Ω . We carefully choose an arc Q which contains the invariant Cantor set: Start by choosing 6 disjoint arcs P_J , $I \leq J \leq VI$, in $\overline{X \setminus (\bigcup \text{balls})}$ as follows. Choose P_I to be an arc which has a one point intersection with ∂X and a one point intersection with I ; for $I < J < VI$, choose P_J to be an arc which has a one point intersection with $J - I$ and a one point intersection with J ; and choose P_{VI} to be an arc which has a one point intersection with V and a one point intersection with ∂X . Let $P = \bigcup_{j=I}^{VI} P_j$




FIGURE 5. **No side or end conditions.** If Q is the rectangular region, the map f illustrated above satisfies $\Omega(Q)$ with crossing number $M = 2$. The sets $end(0)$ and $end(1)$ are strips along the left and right boundaries, respectively. Note that the connected component of $Q \cap f(Q)$ does not “count” since no subinterval of the top boundary of $Q \cap f^{-1}(Q)$ that is contained in maps to the top boundary of the top connected component.

and let $P' = \bigcup_{j=I}^V P_j$. Next define $Q_0 = \bigcup_{j=I}^{VI} P_j$. Then let $Q_{1,b} = f^{-1}(Q_0) \cap b$ for $b \in balls$. This can be done so that $Q_1 = \bigcup_{b \in balls} Q_{1,b}$ does not intersect Q_0 . To make sure that the end result is connected, for $b \in balls$ choose 2 disjoint arcs $A_{b,1}$ and $A_{b,2}$ in ∂b such that $A_{b,1} \cap P_b$ consists of one point, $A_{b,2} \cap P_{b+I}$ consists of one point, $A_{b,1} \cap (f^{-1}(P_I) \cap b)$ consists of one point, and $A_{b,2} \cap (f^{-1}(P_{VI}) \cap b)$ consists of one point. If $A = \bigcup \{A_{b,i} : b \in balls, i = 1, 2\}$, $L_0 = A \cup Q_0 \cup Q_1$ consists of 6 disjoint arcs. For $n > 1$, let $L_n = f^{-n}(L_0)$ and let $\tilde{Q} = \overline{\bigcup_{n \geq 0} L_n}$. By construction, \tilde{Q} is an arc. Finally we shorten \tilde{Q} to make the set Q be an irreducible between some point in the “interior” of Q : define Q to be $\tilde{Q} \setminus P_{VI}$. Suppose $end_0 = \partial X \cap Q$, and $end_1 = \partial V \cap f^{-1}(P_{VI})$. Thus, end_0 is a single point on the boundary ∂X , end_1 is a single point on the boundary ∂V , and f satisfies the horseshoe hypotheses Ω on Q relative to ends end_0 and end_1 with crossing number 5.

Example 10. The invariant set may contain no periodic points. Suppose that $f: \mathbf{R}^2 \rightarrow \mathbf{R}^2$ is the canonical Smale horseshoe map with crossing number 2 (as in the first example), and $r: S^1 \rightarrow S^1$ is a rotation by an irrational number. If Q denotes the rectangular region in Example 1, with end $end(0)$ and $end(1)$ chosen to be the vertical sides of Q , then the map $f \times r: \mathbf{R}^2 \times S^1 \rightarrow \mathbf{R}^2 \times S^1$ satisfies $\Omega(Q \times S^1)$ relative to ends $end(0) \times S^1$ and $end(1) \times S^1$ with crossing number 2. However, there are no periodic points in $Q \times S^1$ under the action of $f \times r$.




FIGURE 6. **Stretched, squeezed and folded disk.** In this example, the ends are in the interior of Q . The map f is not a homeomorphism, because the image “crosses itself,” i.e., there are points x in $f(Q) \cap Q$ such that $f^{-1}(x)$ contains at least two points.

4. THE RESULTS

Recall that in the results that follow, we consider the topology on Q only. The sets in which we are interested are all contained in Q ; this restriction to Q avoids our having to add repeatedly “ $\cap Q$ ” to the sets discussed.

Since Q is locally connected we may choose the distance function d on Q so that for $x \in Q$, $D_\varepsilon(x) = \{y \in Q : d(x, y) < \varepsilon\}$ is connected as long as $D_\varepsilon(x)$ is contained in the component of Q that contains x . (See [Ku] for a proof.)

Lemma 11. *Suppose that $f: X \rightarrow X$ is continuous, f satisfies the horseshoe hypotheses Ω on the set Q with crossing number M relative to ends end_0 and end_1 . Then there is $\varepsilon > 0$ with the following properties:*

1. *If Com is a component of Q , either $E_0 = f^{-1}(end_0) \setminus D_\varepsilon(ends)$ separates Com between end_0 and end_1 ; or $E_1 = f^{-1}(end_1) \setminus D_\varepsilon(ends)$ separates Com between end_0 and end_1 .*
2. *The set $E = f^{-1}(ends) \setminus D_\varepsilon(ends)$ separates Q between end_0 and end_1 .*
3. *If $D = \{D_0, D_1, \dots, D_n\}$ is a tiling chain such that*
 - (a) *each link D_i of D is the closure of a nonempty open set;*
 - (b) *$\bigcup D \subset Q$ is a continuum;*
 - (c) *$D_0 \cap end_0 \neq \emptyset$;*
 - (d) *$D_n \cap end_1 \neq \emptyset$;*
 - (e) *for $0 < i < n$, $D_i \cap D_\varepsilon(ends) = \emptyset$;*
 - (f) *for $0 \leq i \leq n$, $(D_i^\circ \setminus D_\varepsilon(ends)) \cap f^{-1}(ends) = \emptyset$; and*
 - (g) *either*
 - i. *for $0 \leq i < n$, the right boundary of the link D_i is contained in $f^{-1}(end_0)$, or the right boundary of the link D_i is contained in $f^{-1}(end_1)$, and*

- ii. for $0 < i \leq n$, the left boundary of the link D_i is contained in $f^{-1}(end_0)$, or the left boundary of the link D_i is contained in $f^{-1}(end_1)$,

then D has at least $2M - 1$ links, and there is a mutually disjoint subcollection D' of D of which each member contains a preconnection and which contains at least M members.

Proof. Note that no component of $f^{-1}(end_0)$ contains both a point of end_0 and a point of end_1 . There is some $\varepsilon > 0$ such that $\overline{D_{2\varepsilon}(end_0)} \cap \overline{D_{2\varepsilon}(end_1)} = \emptyset$, $\overline{D_{2\varepsilon}(f^{-1}(end_0))} \cap \overline{D_{2\varepsilon}(f^{-1}(end_1))} = \emptyset$, and such that if $x \in Q$, then $\overline{D_\varepsilon(x)}$ contains no preconnection. Suppose that Com is a component of Q .

To prove the first property, that either $E_0 = f^{-1}(end_0) \setminus D_\varepsilon(ends)$ separates Com between end_0 and end_1 , or $E_1 = f^{-1}(end_1) \setminus D_\varepsilon(ends)$ separates Com between end_0 and end_1 , suppose that $E_0 = f^{-1}(end_0) \setminus D_\varepsilon(ends)$ does not separate Com between end_0 and end_1 , and consider $E_1 = f^{-1}(end_1) \setminus D_\varepsilon(ends)$. If $E_1 = f^{-1}(end_1) \setminus D_\varepsilon(ends)$ does not separate Com between end_0 and end_1 either, then there is an arc A irreducible from end_0 to end_1 in Com such that $A \cap \overline{D_\varepsilon(end_0)} \subset \overline{D_\varepsilon(x_0)}$ for some $x_0 \in end_0$, $A \cap \overline{D_\varepsilon(end_1)} \subset \overline{D_\varepsilon(x_1)}$ for some $x_1 \in end_1$, $A \setminus (\overline{D_\varepsilon(end_0)} \cup \overline{D_\varepsilon(end_1)}) = A'$ is an arc, and $A' \subset Com \setminus E_1$. But then $A \cap \overline{D_{+\varepsilon}(end_0)}$ contains no preconnection, $A \cap \overline{D_\varepsilon(end_1)}$ contains no preconnection, and $A \setminus (\overline{D_\varepsilon(end_0)} \cup \overline{D_\varepsilon(end_1)})$ contains no preconnection.

Since A is a connection, it must contain at least M mutually disjoint preconnections. The only way this can happen is for $A \cap \overline{D_\varepsilon(end_0)}$ to intersect $f^{-1}(end_1)$, $A \cap \overline{D_\varepsilon(end_1)}$ to intersect $f^{-1}(end_0)$, and A' to intersect $f^{-1}(end_0)$. Further, this means that $A \cap \overline{D_\varepsilon(end_0)}$ does not intersect $f^{-1}(end_0)$ (because $\overline{D_{2\varepsilon}(f^{-1}(end_0))} \cap \overline{D_{2\varepsilon}(f^{-1}(end_1))} = \emptyset$), $A \cap \overline{D_\varepsilon(end_1)}$ does not intersect $f^{-1}(end_0)$, and $M = 2$. Since E_0 does not separate Com between end_0 and end_1 , there is an arc \hat{A} with endpoints the endpoints of A' , which is contained in $Com \setminus E_0$. Then $(A \cap (\overline{D_\varepsilon(end_0)} \cup \overline{D_\varepsilon(end_1)})) \cap \hat{A}$ is an arc and a connection, but it fails to intersect $f^{-1}(end_0)$, so that it contains no preconnection. This is a contradiction. Thus, either E_0 separates Com between end_0 and end_1 , or E_1 does, and the first property is proved. The second property follows immediately from the first.

To prove the third property, suppose that $D = \{D_0, D_1, \dots, D_n\}$ is a tiling chain satisfying the conditions listed in the statement. Since $\bigcup D$ is a connection, we can choose a continuum $K = \bigcup_{i=0}^n K_i$ irreducible from end_0 to end_1 with the following properties:

1. The set K_0 is a continuum in D_0 irreducible from end_0 to D_1 .
2. The set K_n is a continuum in D_n irreducible from D_{n-1} to end_1 .
3. For $0 < i < n$, K_i is a continuum in D_i irreducible from D_{i-1} to D_{i+1} .
4. For $0 \leq i < n$, $K_i \cap K_{i+1} \neq \emptyset$.

Then K must contain M mutually disjoint irreducible preconnections $\{P_{\alpha_0}, P_{\alpha_1}, P_{\alpha_2}, \dots, P_{\alpha_M}\}$. Because of the properties of the chain D and of the continuum K , we can assume that the indexing set $\{\alpha_0, \alpha_1, \alpha_2, \dots, \alpha_M\}$ is a subsequence of $\{0, 1, \dots, n\}$, and that $P_{\alpha_i} \subset D_{\alpha_i}$. Further, it must be the case that for $0 < \alpha_i < n$, $P_{\alpha_i} = K_{\alpha_i}$; that if $\alpha_1 = 0$, then $P_{\alpha_M} \subset K_{\alpha_M}$, $P_{\alpha_M} \cap D_\varepsilon(end_1) \neq \emptyset$, and $P_{\alpha_M} \cap D_{n-1} \neq \emptyset$. Since $P_{\alpha_i} \cap P_{\alpha_{i+1}} = \emptyset$, $\alpha_{i+1} - \alpha_i \geq 2$. Thus, D must have a minimum of $2M - 1$ links, and it must contain a subcollection D' of M mutually disjoint links each of which contain a preconnection. □




FIGURE 7. **Stretch, squeeze and loop 8 times.** The elliptical region is the set Q , with $end(0)$ and $end(1)$ the two annular regions in the interior of Q , as indicated. The dark shaded winding region is the image of Q under f , with the images of the ends the small disk-like regions indicated.

Lemma 12. *Suppose that $f: X \rightarrow X$ is continuous, and f satisfies the horseshoe hypotheses Ω on Q relative to ends end_0 and end_1 and has crossing number M . Then there is a collection $S = \{S_1, S_2, \dots, S_M\}$ of M mutually disjoint closed neighborhoods of Q having the property that $f(S_i) \cap S_j \neq \emptyset$ for each pair $(i, j) \in \{1, 2, \dots, M\}^2$. Thus S is a set of symbol sets for Q .*

Proof. Suppose Com is a component of Q . Without loss of generality, suppose that $Q_1 \subset f^{-1}(end_1)$ separates $Com \setminus E_1$ between end_0 and end_1 (using the ε, E_1, E_0 of the last lemma). The boundary of each component of $Com \setminus E_1$ is contained in $f^{-1}(end_1)$, but it may not be the case that

$$\overline{\bigcup \{C : C \text{ is a component of } Com \setminus E_1\}} = Com,$$

because it is possible that $E_1^\circ \neq \emptyset$. Then consider the collection \mathcal{S}_{Com} consisting of all members of $\mathcal{S}_{1Com} \cup \mathcal{S}_{2Com} \cup \mathcal{S}_{3Com}$, where $\mathcal{S}_{1Com} = \{\overline{C} : C \text{ is a component of } Com \setminus E_1, \text{ and } f^{-1}(end_0) \text{ separates } C\}$, $\mathcal{S}_{3Com} = \{\overline{C} : C \text{ is a component of } Com \setminus E_1, \text{ and } f^{-1}(end_0) \cap C \neq \emptyset, \text{ and } f^{-1}(end_0) \text{ does not separate } C\}$, and $\mathcal{S}_{2Com} = \{\overline{C} : C \text{ is a component of } Com \setminus (\bigcup (\mathcal{S}_{1Com} \cup \mathcal{S}_{2Com}))\}$. Because Com is locally connected, and $\partial \overline{C}$, for $\overline{C} \in \mathcal{S}_{1Com}$, is continued in $f^{-1}(end_1)$, each member of $\mathcal{S}_{1Com} \cup \mathcal{S}_{2Com}$ must contain an open set of diameter ε . Thus, $\mathcal{S}_{1Com} \cup \mathcal{S}_{2Com}$ is finite.




FIGURE 8. **Expanding balls.** Suppose that X is homeomorphic to a closed n -ball, and $I, II, III, IV,$ and V are mutually disjoint closed n -balls contained in the interior of X . The map $f: X \rightarrow X$ maps each interior ball homeomorphically onto X , and points in the complement of the n -balls to the boundary ∂X .




FIGURE 9. **Choosing Q .** The figure shows how Q can be chosen in order to satisfy the horseshoe hypotheses and pick up the 5-shift dynamics. Here, as long as each component of the invariant set is a single point Q can be a carefully chosen arc.

Construct inductively the tiling chain $T'_{Com} = \{T'_{Com,0}, T'_{Com,1}, \dots, T'_{Com,N}\}$ as follows: Let $T'_{Com,0} = \bigcup\{\bar{C} : C \in \mathcal{S}_{Com} \text{ and } C \cap \text{end}_0 \neq \emptyset\}$. Let $T'_{Com,1} = \bigcup\{\bar{C} : C \in \mathcal{S}_{Com}, \bar{C} \cap T'_{Com,0} \neq \emptyset \text{ and } C \cap (T'_{Com,0})^\circ = \emptyset\}$. Let

$$T'_{Com,2} = \bigcup\{\bar{C} : C \in \mathcal{S}_{Com}, \bar{C} \cap (T'_{Com,0} \cup T'_{Com,1}) \neq \emptyset,$$

and

$$C \cap (T'_{Com,0} \cup T'_{Com,1})^\circ = \emptyset\}.$$

Let $T'_{Com,3} = \bigcup\{C : C \in \mathcal{S}_{Com}, \overline{C} \cap (T'_{Com,0} \cup T'_{Com,1} \cup T'_{Com,2}) \neq \emptyset, \text{ and } C \cap (T'_{Com,0} \cup T'_{Com,1} \cup T'_{Com,2})^\circ = \emptyset\}$. Since $\mathcal{S}_{1Com} \cup \mathcal{S}_{2Com}$ is finite, and members of \mathcal{S}_{3Com} must intersect some member of $\mathcal{S}_{1Com} \cup \mathcal{S}_{2Com}$, but do not intersect any other member of \mathcal{S}_{3Com} except in points contained in $\mathcal{S}_{1Com} \cup \mathcal{S}_{2Com}$, the construction must be a finite process. Continue this finite process until each member of \mathcal{S}_{Com} is in some member of T'_{Com} , thereby obtaining, for some positive integer N , the desired chain $T'_{Com} = \{T'_{Com,0}, T'_{Com,1}, \dots, T'_{Com,N}\}$. Note that $\bigcup T'_{Com} = Com$, and that $\partial T'_{Com,l} \subset f^{-1}(end_1)$ for each l .

We can define a partial ordering \prec on the members of \mathcal{S}_{Com} by defining $C \prec C'$ in \mathcal{S}_{Com} if there exists a tiling chain $D = \{D_0, D_1, \dots, D_{N'}\} \subset \mathcal{S}_{Com}$ such that for some $i < j$, $C = D_i \subset T'_{Com,i}$ and $C' = D_j \subset T'_{Com,j}$. Note that if $C \in \mathcal{S}_{Com}$, there is some member C^b of \mathcal{S}_{Com} such that $C^b \prec C$ and $C^b \cap end_0 \neq \emptyset$. For $1 \leq i \leq N$, and $C \in \mathcal{S}_{Com}$, we define the *left boundary* $\partial_l C$ of C to be $C \cap T'_{Com,i-1}$. For $0 \leq i \leq N - 1$, and $C \in \mathcal{S}_{Com}$, we define the *right boundary* $\partial_r C$ of C to be $C \cap T'_{Com,i+1}$.

As for E_1 and $f^{-1}(end_1)$, it is possible that $f^{-1}(end_0)$ has nonempty interior in Q . Next we “split” each member D of \mathcal{S}_{1Com} into continua and replace D with those sets as follows: For each $D \in \mathcal{S}_{1Com}$ such that $D \cap (end_0 \cup end_1) = \emptyset$, let $\mathcal{D}_{lb} = \{\overline{C} : C \text{ is a component of } D \setminus f^{-1}(end_0) \text{ and } D \cap \partial D_l \neq \emptyset\}$. Because Q is locally connected, the components D of \mathcal{D}_{lb} that intersect both $f^{-1}(end_0)$ and $f^{-1}(end_1)$ form a finite collection. There may be infinitely many components of \mathcal{D}_{lb} that intersect only $f^{-1}(end_1)$, but these must be ‘small’ and ‘do not matter’ (relative to this construction). Let $\mathcal{D}_{rb} = \{\overline{C} : C \text{ is a component of } D \setminus (\bigcup \mathcal{D}_{lb})\}$. Again, \mathcal{D}_{rb} may be infinite, but has only finitely many components that ‘matter’, namely those that intersect both $f^{-1}(end_0)$ and $f^{-1}(end_1)$. For each $D \in \mathcal{S}_{1Com}$ such that $D \cap end_0 \neq \emptyset$, let $\mathcal{D}_{rb} = \{\overline{C} : C \text{ is a component of } D \setminus f^{-1}(end_0) \text{ and } D \cap \partial D_r \neq \emptyset\}$ and let $\mathcal{D}_{lb} = \{\overline{C} : C \text{ is a component of } D \setminus (\bigcup \mathcal{D}_{rb})\}$. For each $D \in \mathcal{S}_{1Com}$ such that $D \cap end_1 \neq \emptyset$, let $\mathcal{D}_{lb} = \{\overline{C} : C \text{ is a component of } D \setminus f^{-1}(end_0) \text{ and } D \cap \partial D_l \neq \emptyset\}$ and let $\mathcal{D}_{rb} = \{\overline{C} : C \text{ is a component of } D \setminus (\bigcup \mathcal{D}_{lb})\}$.

Note that for each $D \in \mathcal{S}_{1Com}$, $D = (\bigcup \mathcal{D}_{lb}) \cup (\bigcup \mathcal{D}_{rb})$. Let $\mathcal{C}_{Com} = \mathcal{S}_{2Com} \cup \mathcal{S}_{3Com} \cup \{\mathcal{D}_{lb} : D \in \mathcal{S}_{Com}\} \cup \{\mathcal{D}_{rb} : D \in \mathcal{S}_{Com}\}$.

We can “extend” the partial ordering \prec on \mathcal{S}_{Com} to a partial ordering \triangleleft on the members of \mathcal{C}_{Com} in a natural way, and we can also define the left boundary for members of \mathcal{C}_{Com} that do not contain a point of end_0 , as well as the right boundary for members of \mathcal{C}_{Com} which are followed by other members of \mathcal{C}_{Com} (relative to the partial ordering \triangleleft). Let $\mathcal{D}_{Com} = \{\underline{D} = \{D_0, D_1, \dots, D_{N'}\} : \underline{D} \text{ is a tiling chain, each link } D_i \text{ in } \underline{D} \text{ is a member of } \mathcal{C}_{Com}\}$, and $\mathcal{D}'_{Com} = \{\underline{D} = \{D_0, D_1, \dots, D_{N'}\} : \underline{D} \in \mathcal{D}'_{Com}, D_0 \text{ contains a point of } end_0, D_{N'} \text{ contains a point of } end_1, \text{ and no link } D_i, i < N', \text{ contains a point of } end_1\}$.

Then \mathcal{C}_{Com} has the following properties:

1. Each nonempty left boundary of a member of \mathcal{C}_{Com} that does not intersect end_0 is contained in either $f^{-1}(end_0)$ or it is contained in $f^{-1}(end_1)$.
2. Each nonempty right boundary of a member of \mathcal{C}_{Com} is contained in either $f^{-1}(end_0)$ or it is contained in $f^{-1}(end_1)$.
3. If $D \in \mathcal{D}_{Com}$, then no two consecutive links of D have both the (nonempty) left and (nonempty) right boundaries contained in $f^{-1}(end_0)$.

If $\underline{D} = \{D_0, D_1, \dots, D_{N'}\}$ is a tiling chain composed of members of \mathcal{C}_{Com} and D_0 intersects end_0 , $D_{N'}$ intersects end_1 , but no link D_i , $0 < i < N'$, intersects $ends$, then we call \underline{D} an *irreducible end_0 - end_1 \mathcal{C}_{Com} tiling chain*. If $C \in \mathcal{C}_{Com}$ such that

1. $C \cap (end_0 \cup end_1) = \emptyset$, and each continuum in C which intersects the nonempty left boundary of C and the nonempty right boundary of C contains a preconnection,
2. $C \cap end_1 \neq \emptyset$, and each continuum in C which intersects end_0 and the nonempty right boundary of C contains a preconnection, or
3. $C \cap end_1 \neq \emptyset$, and each continuum in C which intersects the nonempty left boundary of C and end_1 contains a preconnection,

then we call C a *preconnection link* in \mathcal{C}_{Com} . Note that if C is a preconnection link, then each continuum in C irreducible from $f^{-1}(end_0)$ to $f^{-1}(end_1)$ is a subset of $Q \cap f^{-1}(Q)$. To each irreducible end_0 - end_1 \mathcal{C}_{Com} tiling chain $\underline{D} = \{D_0, D_1, \dots, D_{N'}\}$, we associate a connection $K_{\underline{D}}$ constructed as follows:

1. Let $L_0 = end_0$ and for $0 < i \leq N'$, let $L_i = \partial D_{i-1} \cap \partial D_i$. For $0 \leq i < N'$, let $R_i = \partial D_{i+1} \cap \partial D_i$, and let $R_{N'} = end_1$.
2. The continuum $K_{\underline{D}}$ is a union of continua contained in the links D_i of \underline{D} , that is, $K_{\underline{D}} = \bigcup_{i=0}^{N'} K_{\underline{D},i}$, where each $K_{\underline{D},i}$ is a continuum contained in D_i which intersects both L_i and R_i .
3. If for some i , D_i is a preconnection link, and so is D_j , where $D_j = D_{i+1}$ or $D_j = D_{i-1}$, and there is a continuum K in $D_i \cup D_j$ which does not contain two disjoint preconnections, but which (1) intersects both L_i and R_j if $j = i+1$, or (2) intersects both L_j and R_i if $j = i-1$, then we say that D_i is an *adjoining preconnection link*, and that D_i *adjoins* D_j . Suppose then that D_i and D_{i+1} are adjoining preconnection links. Then there is an irreducible continuum \tilde{K} in $D_i \cup D_{i+1}$ which intersects both L_i and R_{i+1} , but does not contain two disjoint preconnections. There is an irreducible subcontinuum Λ_i of \tilde{K} in D_i which intersects both L_i and R_{i+1} . Then $\Lambda_i \cap \Lambda_{i+1} \neq \emptyset$, for otherwise \tilde{K} does contain two disjoint preconnections. Since \tilde{K} is irreducible between L_i and R_{i+1} , $\tilde{K} = \Lambda_i \cup \Lambda_{i+1}$.
4. Suppose that $\{D_i, D_{i+1}, \dots, D_{i+m}\}$ is a subchain of \underline{D} each link of which is an adjoining preconnection link. Then for each $i \leq j < j+1 \leq i+m$, there is a continuum K'_j in $D_j \cup D_{j+1}$ which intersects both L_j and R_{j+1} , but does not contain two disjoint preconnections. Then there is an irreducible subcontinuum $\Lambda_{j,j}$ of K'_j in D_j which intersects both L_j and R_j , and there is an irreducible subcontinuum $\Lambda_{j,j+1}$ in D_{j+1} which intersects both L_{j+1} and R_{j+1} . Then $\Lambda_{j,j} \cap \Lambda_{j,j+1} \neq \emptyset$, for otherwise K'_j does contain two disjoint preconnections, and $\Lambda_{j,u} \cup \Lambda_{j,j+1}$ is irreducible between L_j and R_{j+1} . Since each D_j^c is connected and locally connected, we can choose $K_{\underline{D},j}$ to be a continuum irreducible between L_j and R_{j+1} such that $K_{\underline{D},j} \cap L_j = \Lambda_{j,j} \cap L_j$, $K_{\underline{D},j} \cap R_j = \Lambda_{j,j} \cap R_j$, and $K_{\underline{D},j} \cup K_{\underline{D},j+1}$ does not contain two disjoint preconnections, and we can do this so that $\bigcup_{j=i}^{i+m} K_{\underline{D},j}$ is a continuum irreducible between L_i and R_{m+i} .
5. If for some i , D_i is a preconnection link which is not an adjoining preconnection link, then either no link that intersects D_i is a preconnection link, or

D_{i+1} is a preconnection link, but no continuum K in $D_i \cup D_{i+1}$ which intersects both L_i and R_{i+1} fails to contain two disjoint preconnections or D_{i-1} is a preconnection link, but no continuum K in $D_i \cup D_{i-1}$ which intersects both L_{i-1} and R_i fails to contain two disjoint preconnections. There is an irreducible continuum $K_{\underline{D},i}$ in D_i which intersects both L_i and R_i . Further, since each D_i° is connected and locally connected, we can choose each $K_{\underline{D},i}$ so that if D_j is also a preconnection link and D_j is adjacent to D_i , then $K_{\underline{D},i} \cap K_{\underline{D},j} \neq \emptyset$.

6. If D_i is not a preconnection link, choose $K_{\underline{D},i}$ to be an irreducible continuum from L_i to R_i contained in D_i which is not a preconnection. (This can happen in one of two ways. Either for $j = 0$ or $j = 1$ both L_i and R_i are contained in $f^{-1}(end_j)$, or $D_i \not\subseteq f^{-1}(Q)$.) Again, since each D_i° is connected and locally connected, we can choose each $K_{\underline{D},i}$ so that if D_j is adjacent to D_i , then $K_{\underline{D},i} \cap K_{\underline{D},j} \neq \emptyset$.

Thus, $K_{\underline{D}}$ is a connection in $\bigcup \underline{D}$ which has the property that no connection K' has fewer mutually disjoint preconnections than does $K_{\underline{D}}$. Since each preconnection of $K_{\underline{D}}$ is contained in a link of \underline{D} , but no link of \underline{D} contains more than one preconnection, it follows from Lemma 11 that the crossing number $M_{\underline{D}}$ for $\bigcup \underline{D}$ must be greater than or equal to M , and thus $\bigcup \underline{D}$ contains at least M preconnection links such that the continua $K_{\underline{D},i}$ corresponding to the $M_{\underline{D}}$ links in this set are mutually disjoint. More precisely, suppose that $\underline{D}' = \{D_{\beta_1}, D_{\beta_2}, \dots, D_{\beta_{N''}}\}$ denotes a maximal subcollection of preconnection links of \underline{D} having the property that $\{K_{\underline{D},\beta_i} : D_{\beta_i} \in \underline{D}'\}$ is a collection of $M_{\underline{D}}$ disjoint continua. If \underline{D} contains at least one subchain of at least 4 links consisting entirely of adjoining preconnection links, there may be more than one such set \underline{D}' for \underline{D} . However, this lack of uniqueness occurs only when \underline{D} contains such a subchain (1) containing an even number greater than 2 links, and (2) this subchain is not properly contained in another subchain of preconnection links in \underline{D} . In this case, we choose as members of \underline{D}' the first, third, fifth, etc., members of the subchain of adjoining links rather than the second, fourth, etc., links.

Because distinct tiling chains can share links, though some of the preconnection links in the sets \underline{D}' may not “count”: If $\underline{D} = \{D_0, D_1, \dots, D_{N'}\}$ and $\underline{E} = \{E_0, E_1, \dots, E_{N''}\}$ are irreducible end_0 - end_1 \mathcal{C}_{Com} tiling chains, $D_i = E_j$, and no link of $\{D_0, D_1, \dots, D_{i-1}\}$ is a link of $\{E_{j+1}, \dots, E_{N''}\}$, then $\{D_0, D_1, \dots, D_{k-1}, D_i, E_{j+1}, \dots, E_{N''}\}$ is also an irreducible end_0 - end_1 \mathcal{C}_{Com} tiling chain. Thus if $\underline{D} = \{D_0, D_1, \dots, D_{N'}\}$ is an irreducible end_0 - end_1 \mathcal{C}_{Com} tiling chain, we define $D_{\alpha(D)_1}$ to be the last link of \underline{D} which is the *first* preconnection link of some irreducible end_0 - end_1 \mathcal{C}_{Com} tiling chain E . (Since \underline{D} has a first preconnection link, \underline{D} has only finitely many preconnection links, and there are only finitely many irreducible end_0 - end_1 \mathcal{C}_{Com} tiling chains, $D_{\alpha(D)_1}$ exists.) We say that $D_{\alpha(D)_1}$ is the *first essential preconnection link* of \underline{D} . Next, we define $D_{\alpha(D)_2}$ to be the last link of \underline{D} after $D_{\alpha(D)_1}$ which is the *first nonadjoining* preconnection link following the first essential preconnection link $E_{\alpha(E)_1}$ in some irreducible end_0 - end_1 \mathcal{C}_{Com} tiling chain \underline{E} . Again, $D_{\alpha(D)_2}$ must exist (because otherwise we violate our basic assumption that each connection contains at least M mutually disjoint preconnections), and we call it the *second essential* preconnection link of \underline{D} . This finite process continues until we choose the M th essential preconnection link of \underline{D} , $D_{\alpha(D)_M}$ to be the last

preconnection link of \underline{D} after the link $D_{\alpha(D)_{M-1}}$ which is the first nonadjoining preconnection link after the link $E_{\alpha(D)_{M-1}}$ in some irreducible end_0 - end_1 \mathcal{C}_{Com} tiling chain E .

Then, for $1 \leq i \leq M$, $S_{Com,i} = \bigcup \{D_{\alpha(D)_i} : D \text{ is an irreducible } end_0\text{-}end_1 \mathcal{C}_{Com} \text{ tiling chain}\}$. Finally, define for $1 \leq i \leq M$, $S_i = \bigcup \{S_{Com,i} : Com \text{ is a component of } Q\}$. Checking that this collection has the properties claimed is straightforward. \square

Outline of the proof of the Chaos Lemma. Suppose that each member $i_* = (i_j)_{j=0}^\infty$ is in $\{1, \dots, M\}$. Let $K_*^\infty = \{i_* : i_* \text{ is a sequence of which each member is in } \{1, \dots, M\}\}$, and for each $n \geq 0$, let $K_*^{n+1} = \{(i_0, i_1, i_2, \dots, i_n) : (i_0, i_1, i_2, \dots, i_n) \text{ is an } n + 1\text{-member finite sequence of which each member is in } \{1, \dots, M\}\}$. Define inductively $S_{(i_0, i_1)} = S_{i_0} \cap f^{-1}(S_{i_1})$, $S_{(i_0, i_1, i_2)} = S_{(i_0, i_1)} \cap f^{-2}(S_{i_2}), \dots, S_{(i_0, i_1, i_2, \dots, i_n)} = S_{(i_0, i_1, \dots, i_{n-1})} \cap f^{-n}(S_{i_n}), \dots$. It can be proved that $S_{i_*} = \bigcap_{j=0}^\infty S_{(i_0, i_1, i_2, \dots, i_n)} \neq \emptyset$. Next, let $P = \bigcup_{i_* \in K_*^\infty} S_{i_*}$. Then P is a compact subset of Q , and it can be written as $P = \bigcup_{i_* \in K_*^\infty} (\bigcap_{n=0}^\infty S_{(i_0, i_1, i_2, \dots, i_n)}) = \bigcap_{n=0}^\infty (\bigcup_{(i_0, i_1, i_2, \dots, i_n) \in K_*^{n+1}} S_{(i_0, i_1, i_2, \dots, i_n)})$. Now $f(P) \subset P$. To get an invariant set, define $Y = \bigcap_{m=0}^\infty f^m(P)$ (i.e., Y is a closed subset of P such that $f(Y) = Y$). Since for each i_* , $Y_{i_*} = S_{i_*} \cap Y \neq \emptyset$ and $Y = \bigcup_{i_* \in K_*^\infty} Y_{i_*}$, there is a natural one-to-one correspondence between Y and K_*^∞ defined by $\alpha(Y_{i_*}) = i_*$. Let $g = f|_Y$. Then $g: Y \rightarrow Y$ is continuous, $\alpha: Y \rightarrow K_*^\infty$, and it is not difficult to show that $\sigma \circ \alpha = \alpha \circ g$, where σ denotes the one-sided shift on M symbols. \square

The main theorem follows immediately from Lemma 12 and the Chaos Lemma.

5. RELAXING THE HYPOTHESES

We do not need the strength of all the horseshoe hypotheses Ω , and we give lemmas below that enable the application of our results in some cases in which the ambient space X is not locally compact. (This might occur, for example, when partial differential equations are studied.) Also, the last lemma shows that an extremely careful choice of Q is not necessary, as long as it is precompact; in particular, it need not be the case that all components of Q are connections. Any components that are not connections play no role in the choice of the symbol sets \mathcal{S} , but they do not cause problems either.

Suppose that Y is a space, Q is a closed, locally connected subset of X , and $f: X \rightarrow X$ is continuous. If end_0 and end_1 are disjoint closed subsets of Q , then we say that a closed, connected subset K of Q which intersects both end_0 and end_1 is a end_0 - end_1 connection, or just a connection, if no confusion results. If K' is a closed connected subset of $Q \cap f^{-1}(Q)$ that intersects both $f^{-1}(end_0)$ and $f^{-1}(end_1)$, then K' is an end_0 - end_1 preconnection, or just a preconnection. We sometimes write $end(0)$ and $end(1)$ for end_0 and end_1 . If Q contains no end_0 - end_1 connection or if each end_0 - end_1 connection contains no preconnection, then the crossing number of f on Q is 0. If there is an end_0 - end_1 connection in Q and each end_0 - end_1 connection contains at least one preconnection, then the crossing number of f on Q is the largest positive integer such that each connection K in Q contains at least M mutually disjoint end_0 - end_1 preconnections.

In this section we make the following five assumption:

- Ω'_X : X is a separable metric space.
- Ω'_Q : $Q \subset X$ is locally connected and closed.

Ω'_f : The map $f: Q \rightarrow X$ is continuous, and $\overline{f(Q)}$ is a compact (i.e., Q is precompact).

Ω'_E : The sets $end_0 \subset Q$ and $end_1 \subset Q$ are disjoint and compact, and at least one component of Q intersects both end_0 and end_1 .

Ω'_M : Q has a crossing number (defined below) $M \geq 2$.

We refer to the above five hypotheses $\Omega'_X, \Omega'_Q, \Omega'_f, \Omega'_E$, and Ω'_M collectively as the *weak horseshoe hypotheses* Ω' . A *connection* Γ is a closed connected subset of Q that intersects both end_0 and end_1 . Note that by hypothesis Ω'_E , at least one component of Q is a connection. An example of a connection is a path in Q intersecting both end_0 and end_1 . A *preconnection* γ is a closed connected subset of Q for which $f(\gamma)$ is a connection. We define the crossing number M to be the largest number such that every connection contains at least M mutually disjoint preconnections.

The last lemmas below argue that Q contains a compact subset Q' , all of whose connected components are connections, and furthermore that Q' has a crossing number at least as big as the crossing number M of Q . When we later construct a compact invariant set in Q' that factors over an M -shift, the same will automatically be true of Q . Hence these two lemmas argue that without loss of generality, we can assume that Q is compact and all its connected components are connections, since the weak horseshoe hypotheses Ω' on Q with crossing number M imply the horseshoe hypotheses Ω on Q' with crossing number $M' \geq M$.

Lemma 13. *Suppose that $f: X \rightarrow X$ is continuous, and f satisfies the weak horseshoe hypotheses Ω' on Q with crossing number M relative to ends end_0 and end_1 . If $Q' = Q \cap \overline{f(Q)}$, then f satisfies Ω on the compact set Q' with crossing number $M' \geq M$ relative to ends $end'_0 = end_0 \cap \overline{f(Q)}$ and $end'_1 = end_1 \cap \overline{f(Q)}$.*

Proof. Since $\overline{f(Q)}$ is compact, and f satisfies Ω' on the closed set Q , there is some connection K in Q . If K is a connection in Q , the collection $\mathcal{K}_p = \{K_\alpha: K_\alpha \text{ is a preconnection in } K\}$ has at least M mutually disjoint members. Let $Q' = Q \cap \overline{f(Q)}$. Then Q' is compact, and Q' contains a connection that is also a continuum. (For example, if K_α is a preconnection in the connection K in Q , then $\overline{f(K_\alpha)}$ is a compact connection in $Q \cap \overline{f(Q)}$.) Furthermore, for each connection K' in Q' , $\mathcal{K}'_p = \{K'_\alpha: K'_\alpha \text{ is a preconnection in } K'\}$ has at least M mutually disjoint members. Thus, Q' is a compact subset of Q on which f satisfies Ω relative to ends $end'_0 = end_0 \cap Q'$ and $end'_1 = end_1 \cap Q'$. □

Lemma 14. *Suppose that $f: X \rightarrow X$ is continuous, and f satisfies the horseshoe hypotheses Ω on the compact set Q with crossing number M relative to ends end_0 and end_1 . Then $Q' = \bigcup\{K: K \text{ is a connection in } Q \text{ and } K \text{ is a connected component of } Q\}$ is a compact subset of Q and f satisfies Ω on Q' with crossing number $M' \geq M$ relative to ends $end'_0 = end_0 \cap Q'$ and $end'_1 = end_1 \cap Q'$.*

Proof. This is straightforward, and the proof is therefore omitted. □

With Lemmas 13 and 14 we have reduced our problem to one of working with a compact set Q of which each component is a connection. Thus, f satisfies the horseshoe hypotheses Ω on a closed subset Q' of Q with crossing number $M' \geq M$ as long as Q is precompact and contains at least one connection. Thus, we get the following slightly more general result:

Theorem 15. *More general version of the main result.* Assume the weak horseshoe hypotheses Ω' on the set Q . Then there is a closed invariant set $Q_I \subset Q$ for which $f|_{Q_I}$ is semiconjugate to a one-sided M -shift. (If f is a homeomorphism, then $f|_{Q_I}$ is also semiconjugate to a two-sided M -shift).

REFERENCES

- [BW] K. Burns and H. Weiss, *A geometric criterion for positive topological entropy*, Commun. Math. Phys. **172** (1995), 95–118. MR **96e**:58120
- [CKM] M. Carbinatto, J. Kwapisz, and K. Mischaikow, *Horseshoes and the Conley index spectrum*, Ergodic Theory Dynam. Systems **20** (2000), 365–377. CMP 2000:12
- [CM] M. Carbinatto and K. Mischaikow, *Horseshoes and the Conley index spectrum II. The theorem is sharp*, Discrete Contin. Dynam. Systems **5** (1999), 599–616. MR **2000i**:37009
- [KKY] J. Kennedy, S. Koçak, and J. Yorke, *A Chaos Lemma*, to appear, Amer. Math. Monthly.
- [KSYG] J. Kennedy, M. Sanjuan, J. Yorke, and C. Grebogi, *The topology of fluid flow past a sequence of cylinders*, Topology and its Applications **94** (1999), 207–242. CMP 99:14
- [Ku] C. Kuratowski, *Topology II*, Academic Press, New York, 1968. MR **41**:4467
- [MM] K. Mischaikow and M. Mrozek, *Isolating Neighborhoods and Chaos*, Japan Journal of Industrial and Applied Mathematics **12** (1995), 205–236. MR **96e**:58104
- [S] A. Szymczak, *The Conley index and symbolic dynamics*, Topology **35** (1996), 287–299. MR **97b**:58054
- [SKGY] M. Sanjuan, J. Kennedy, C. Grebogi, and J. Yorke, *Indecomposable continua in dynamical systems with noise: fluid flow dynamics past a spatial array of cylinders*, Chaos **7** (1997), 125–138. MR **98a**:76027

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF DELAWARE, NEWARK, DELAWARE 19716

E-mail address: jkennedy@math.udel.edu

INSTITUTE FOR PHYSICAL SCIENCE AND TECHNOLOGY, UNIVERSITY OF MARYLAND, COLLEGE PARK, MARYLAND 20742

E-mail address: yorke@ipst.umd.edu