

SEMI-LINEAR HOMOLOGY G -SPHERES AND THEIR EQUIVARIANT INERTIA GROUPS

ZHI LÜ

ABSTRACT. This paper introduces an abelian group $H\Theta_V^G$ for all semi-linear homology G -spheres, which corresponds to a known abelian group Θ_V^G for all semi-linear homotopy G -spheres, where G is a compact Lie group and V is a G -representation with $\dim V^G > 0$. Then using equivariant surgery techniques, we study the relation between both $H\Theta_V^G$ and Θ_V^G when G is finite. The main result is that under the conditions that G -action is semi-free and $\dim V - \dim V^G \geq 3$ with $\dim V^G > 0$, the homomorphism $T : \Theta_V^G \rightarrow H\Theta_V^G$ defined by $T([\Sigma]_G) = \langle \Sigma \rangle_G$ is an isomorphism if $\dim V^G \neq 3, 4$, and a monomorphism if $\dim V^G = 4$. This is an equivariant analog of a well-known result in differential topology. Such a result is also applied to the equivariant inertia groups of semi-linear homology G -spheres.

1. INTRODUCTION

Let Θ_n (resp. $H\Theta_n$) be the set of consisting of the h -cobordism (resp. homology cobordism) classes of all homotopy (resp. homology) spheres. It is well known that both Θ_n and $H\Theta_n$ form abelian groups under connected sum operation, and for $n \neq 3$, Θ_n is finite (see [KM]). Note that the group structure of Θ_3 is still open (this is associated to Poincaré conjecture in dimension three), but $H\Theta_3$ must be infinite as noticed by Fintushel and Stern [FS] (see also [Fu]). With respect to the relation between Θ_n and $H\Theta_n$, there is a natural homomorphism $T_n : \Theta_n \rightarrow H\Theta_n$ defined by $T_n([\Sigma^n]) = \langle \Sigma^n \rangle$ for $[\Sigma^n] \in \Theta_n$. Then the following result follows from [HH] and [Ke].

Theorem 1.1. *The homomorphism $T_n : \Theta_n \rightarrow H\Theta_n$ is an isomorphism for $n \neq 3$.*

As a consequence of Theorem 1.1, one has

Corollary 1.2. *For any $\langle M^n \rangle \in H\Theta_n$ with $n \neq 3$, then the inertia group $I(M^n)$ of M^n is trivial, i.e., $I(M^n) = (0)$.*

Note that the inertia group $I(M^n)$ of a closed oriented manifold M^n is defined as the group of those homotopy spheres Σ^n satisfying the condition that the connected

Received by the editors July 3, 2000.

2000 *Mathematics Subject Classification.* Primary 57S15, 57S17, 57R91, 57R55, 57R67.

Key words and phrases. Semi-linear homology G -sphere, equivariant inertia group, G -action, representation, surgery.

This work was supported by the Japanese Government Scholarship, and partially supported by the research fund of the Ministry of Education in China and the JSPS Postdoctoral Fellowship (No. P02299).

sum $M^n \sharp \Sigma^n$ is orientation preserving diffeomorphic to M^n . Such a group $I(M^n)$ may be used to classify the distinct differentiable structures almost diffeomorphic to M^n , or in other words, it may be used to measure the nonuniqueness of connected sums of M^n with exotic spheres. The inertia groups for some special manifolds have been studied (see, for example [Ko], [Sc1], [Wi1], [Wi2], [Win]).

The purpose of this paper is to discuss the equivariant analogs of the above results. There are mainly three kinds of equivariant analogs, Θ_V^G , $\Theta_V^{G,s}$, and Γ_V^G , of Θ_n , where G is a compact Lie group and V is a representation of G . These three equivariant analogs are isomorphic to each other in most cases, and are often infinite (see [MSc1], [MSc2]). Here we are only concerned with Θ_V^G (where Θ_V^G , which consists of G - h -cobordism classes of all semi-linear homotopy G -spheres such that the tangential spaces at fixed points are equivalent to V , forms an abelian group under the equivariant connected sum if $\dim V^G > 0$). In contrast to Θ_V^G , we will construct an equivariant analog $H\Theta_V^G$ of $H\Theta_n$. Then we will study the relation between Θ_V^G and $H\Theta_V^G$ when G is finite. It should be pointed out that it is possible to work very successfully with the other two groups $\Theta_V^{G,s}$ and Γ_V^G and to get corresponding comparative results. The argument would require some additional input involving Rothenberg's work on classifying semi-linear actions on disks by equivariant Whitehead torsion, but this will be beyond what one wants to consider at this point.

Θ_V^G works in homotopy sense, while $H\Theta_V^G$ will work in homology sense. Hence, in order to construct $H\Theta_V^G$, we need to give the definitions of semi-linear homology G -spheres, G -equivariant homology cobordism and G -homology equivalence. These notations correspond to those known notations of semi-linear homotopy G -spheres, G - h -cobordism and G -homotopy equivalence, respectively. Then $H\Theta_V^G$ is defined as the set of G -equivariant homology cobordism classes of all semi-linear homology G -spheres whose tangential spaces at fixed points are equivalent to V . We shall see that $H\Theta_V^G$ also forms an abelian group under the equivariant connected sum if $\dim V^G > 0$. Furthermore, by making use of equivariant surgery techniques, the relation between both Θ_V^G and $H\Theta_V^G$ is discussed under the assumption that G is finite and G -action is semi-free (i.e., G -action with two orbit types—fixed points and free orbits). The result is stated as follows.

Theorem 1.3. *Suppose G is a finite Lie group, and V is a semi-free G -representation with $\dim V - \dim V^G \geq 3$ and $\dim V^G > 0$. Then the homomorphism*

$$T : \Theta_V^G \longrightarrow H\Theta_V^G$$

defined by $T([\Sigma]_G) = \langle \Sigma \rangle_G$ is an isomorphism if $\dim V^G \neq 3, 4$, and a monomorphism if $\dim V^G = 4$.

Remark. It is possible that the discussions for the relation between Θ_V^G and $H\Theta_V^G$ may be carried out if G -actions are not semi-free. Such an example may be provided when $G = \mathbb{Z}_{p^r}$ for p a prime and $r \geq 2$ and if G -action is assumed to be not semi-free. Since the subgroups of \mathbb{Z}_{p^r} are always linearly ordered and the isotropy subgroups of \mathbb{Z}_{p^r} -action are also linearly ordered, in this case there is a realization result as in Theorem 1.3. However, it seems difficult to determine to what extent Theorem 1.3 extends to non-semi-free actions.

The equivariant inertia group of a closed, equivariantly oriented smooth G -manifold has been introduced and studied by Masuda and Schultz; for details,

see [Ma], [MSc1], [MSc2]. Applying Theorem 1.3 to the equivariant inertia group immediately gives the following result.

Corollary 1.4. *Suppose G is a finite Lie group and V is a semi-free G -representation with $\dim V - \dim V^G \geq 3$ and $\dim V^G > 0$. For $\langle M \rangle_G \in H\Theta_V^G$, the equivariant inertia group $I^G(M; M^G)$ is trivial if $\dim V^G \neq 3$.*

The arrangement of this paper is as follows. In Section 2 an equivariant analog $H\Theta_V^G$ of $H\Theta_n$ is constructed for G to be a compact Lie group. In Section 3 we discuss the relationship between both $H\Theta_V^G$ and Θ_V^G when G is finite, and give the proof of Theorem 1.3. The methods that we use are mainly equivariant spherical modification and equivariant handle addition. Throughout this paper, all G -manifolds, with or without boundary, are to be compact, smooth and G -oriented. The G -manifold M with G -orientation reversed is denoted by $-M$.

2. CONSTRUCTION OF $H\Theta_V^G$

Throughout this section, suppose that G is a compact Lie group (here G is not necessarily restricted to be finite) and V is a G -representation. The task of this section is to construct an analog $H\Theta_V^G$ of $H\Theta_n$. As stated in Section 1, we wish that $H\Theta_V^G$ is comparable with Θ_V^G . For this, we first review some standard notations with respect to Θ_V^G .

Recall from [Sc2] and [Sc3] that a semi-linear homotopy G -sphere is a manifold for which (1) the fixed point set of all subgroups are homotopy spheres, (2) if the fixed point set of one subgroup is a codimension 2 submanifold of the fixed point set of another subgroup, then the former is homotopically unknotted in the latter. An equivariant h -cobordism between such objects is an h -cobordism with group action such that the fixed point sets of all subgroups are h -cobordisms and there is an analog of the homotopy unknotting condition. Then Θ_V^G is defined as the set of the G - h -cobordism classes of all those G -manifolds that are G -homotopically equivalent to $S(V \oplus \mathbb{R})$ and whose tangent spaces at fixed points are equivalent to V , where the G -action on \mathbb{R} is trivial and $S(V \oplus \mathbb{R})$ is a linear G -sphere. Θ_V^G forms an abelian group under the equivariant connected sum operation if $\dim V^G > 0$.

Based on the above, the construction of $H\Theta_V^G$ needs the following notations. Remember that $H\Theta_V^G$ should work in homology sense.

Definition 2.1. A homology G -sphere M is a *semi-linear homology G -sphere* if M^H is also a homology sphere for each subgroup H of G . Note that Alexander duality ensures that a semi-linear homology G -sphere M always possesses the homology unknottedness, i.e., for $H \leq L \leq G$ and $\dim M^H - \dim M^L \geq 1$, $M^H - M^L$ is homology equivalent to S^l where $l = \dim M^H - \dim M^L - 1$.

Clearly, a semi-linear homotopy G -sphere is a semi-linear homology G -sphere. For a semi-linear homology G -sphere M on which G acts semi-freely, since M^G is also a homology sphere, M^G is either connected (when $\dim M^G > 0$) or S^0 (when $\dim M^G = 0$). Then by [AB], [Mi2], the tangent space at each point $x \in M^G$ is equivalent to V' , where V' is a given semi-free G -representation.

Definition 2.2. Two closed oriented smooth G -manifolds M_1, M_2 are *G -equivariantly homologically cobordant* if there exists an oriented compact smooth G -manifold W such that the boundary $\partial W^H = M_1^H \sqcup -M_2^H$ and the inclusions $i_j : M_j \rightarrow$

$W, j = 1, 2$, induce isomorphisms $H_*(M_j^H; \mathbb{Z}) \cong H_*(W^H; \mathbb{Z})$ ($j = 1, 2$) for each subgroup H of G .

It is obvious that G -equivariant homology cobordism is an equivalence relation, and for two oriented G -manifolds M_1, M_2 , if M_1 and M_2 are G -equivariantly h -cobordant, then they are G -equivariantly homologically cobordant. In homotopy category, it is well known that a G -map $f : X \rightarrow Y$ is a G -homotopy equivalence if and only if $f^H : X^H \rightarrow Y^H$ is an ordinary homotopy equivalence for each subgroup H of G . However, in homology category there does not exist the above analogous result in general. We say that a G -map $f : X \rightarrow Y$ is a G -homology equivalence (or both G -spaces X and Y are G -homologically equivalent) if f induces an isomorphism $H_*(X^H; \mathbb{Z}) \cong H_*(Y^H; \mathbb{Z})$ for each subgroup H of G .

With the above understood, we define $H\Theta_V^G$ as the set of the G -equivariant homology cobordism classes of all those G -manifolds that are G -homologically equivalent to $S(V \oplus \mathbb{R})$ and whose tangent spaces at fixed points are equivalent to V . $H\Theta_V^G$ can form an abelian group with respect to the equivariant connected sum if $\dim V^G > 0$. In fact, it is easy to see that the equivariant connected sum of two semi-linear homology G -spheres is also a semi-linear homology G -sphere, and the equivariant connected sum operation is associative and commutative up to equivariantly orientation preserving diffeomorphism. The sphere $S(V \oplus \mathbb{R})$ serves as the identity element. Note that the condition $\dim V^G > 0$ makes sure that the equivariant connected sum is commutative in $H\Theta_V^G$. It remains to check that each element of $H\Theta_V^G$ has an inverse.

Definition 2.3. A compact G -manifold W is said to be *equivariantly acyclic* if W^H is acyclic for any subgroup H of G .

In nonequivariant topology, there are two well-known results with respect to h -cobordism, contractible manifolds and homotopy spheres: (1) a simply connected manifold M^n is h -cobordant to the sphere S^n if and only if M^n bounds a contractible manifold; (2) if M^n is a homotopy sphere, then $M^n \sharp (-M^n)$ bounds a contractible manifold (see [KM, Lemmas 2.3 and 2.4]). Those methods used to prove the above two results can be carried out to obtain the analogous results in homology sense, and especially can still work in equivariant cases. The proofs of the following results are straightforward. We would like to leave them as an exercise to the reader.

Lemma 2.1. *Suppose M is a closed oriented G -manifold such that the tangential spaces at all points of M^G are equivalent to V . Then M equivariantly bounds an equivariantly acyclic G -manifold W if and only if M is G -equivariantly homologically cobordant to $S(V \oplus \mathbb{R})$, i.e., M is a representative of the identity element of $H\Theta_V^G$.*

Lemma 2.2. *If M is a semi-linear homology G -sphere, then $M \sharp (-M)$ equivariantly bounds an equivariant acyclic G -manifold.*

Now let $\langle M \rangle_G$ be an element in $H\Theta_V^G$. By Lemmas 2.1 and 2.2, $M \sharp (-M)$ is G -equivariantly homologically cobordant to $S(V \oplus \mathbb{R})$, and thus $\langle -M \rangle_G$ is the inverse element of $\langle M \rangle_G$.

Combining the above arguments, we have

Proposition 2.2. *If $\dim V^G > 0$, then $H\Theta_V^G$ forms an abelian group under the equivariant connected sum operation.*

3. RELATION BETWEEN $H\Theta_V^G$ AND Θ_V^G

This section is devoted to complete the proof of Theorem 1.3. The proof relies on the following two lemmas.

Lemma 3.1. *Let G be a finite Lie group, and let V be a semi-free n -dimensional G -representation such that $\dim V - \dim V^G \geq 3$ and $\dim V^G = k \geq 5$. For any $\langle M \rangle_G \in H\Theta_V^G$, there exists a semi-linear homotopy G -sphere Σ that is G -equivariantly homologically cobordant to M .*

Lemma 3.2. *Let G be a finite Lie group, and let V be a semi-free n -dimensional G -representation such that $\dim V - \dim V^G \geq 3$ and $\dim V^G = k \geq 4$. Let $\langle M \rangle_G \in H\Theta_V^G$; if $\langle M \rangle_G = 0$, i.e., M bounds an equivariantly acyclic G -manifold W , then it bounds an equivariantly contractible G -manifold.*

We shall use equivariant spherical modification and equivariant handle addition to prove Lemmas 3.1 and 3.2. In our case, since G -actions are assumed to be semi-free, i.e., the isotropy subgroups of G only are $\{1\}$ and itself G , the equivariant surgery on M in Lemma 3.1 (resp. W in Lemma 3.2) is divided into two stages. The first one is to do surgery on the fixed point set M^G (resp. W^G) and then on the G free part $M - M^G$ (resp. $W - W^G$). Fortunately we have not encountered any surgery obstruction because all Stiefel-Whitney classes of M^H (resp. W^H) are trivial for $H = \{1\}, G$, and in particular, only 1-dimensional and 2-dimensional surgeries are needed. Note that the triviality of the normal bundle of imbedded 1-dimensional (resp. 2-dimensional) sphere in ambient manifold is detected by the first (resp. second) Stiefel-Whitney class of ambient manifold. Also, according to [KM], when doing 1-dimensional surgery on a spin manifold, the resulting manifold may still preserve the spin structure. These facts will not be mentioned again in the proofs of Lemmas 3.1 and 3.2.

Proof of Lemma 3.1. Without loss of generality we assume that M^H is not simply connected for $H = \{1\}, G$. Note that by [Ke], the fundamental group of M^H has a finite presentation. Let $\Gamma = M \times I$, where I is the unit interval $[0, 1]$ with trivial G -action, and let $M \times \{1\}$ identify with M . The argument proceeds as follows.

Step I. *1-dimensional surgery on M^G .*

Let $\alpha_1, \dots, \alpha_l$ be a finite set of generators of $\pi_1(M^G)$. Then there exist G -imbeddings f_1, \dots, f_l of $S^1 \times D^{k-1} \times D(U)$ into M with disjoint images representing $\alpha_1, \dots, \alpha_l$, respectively, where G -actions on S^1 and D^{k-1} are trivial and U is the G -representation on the normal fibre $\mu(M, M^G)_x$ at a point $x \in M^G$ in M such that $U^G = \{0\}$. (Note that f_i should be expressed as the G -imbedding of $G \times_G (S^1 \times D^{k-1} \times D(U))$ into M . Since $G \times_G (S^1 \times D^{k-1} \times D(U))$ only contains a copy of $S^1 \times D^{k-1} \times D(U)$, we may identify $G \times_G (S^1 \times D^{k-1} \times D(U))$ with $S^1 \times D^{k-1} \times D(U)$. See [Br], [PR]). Let \mathbb{S}^1 (resp. $\mathbb{D}^2, \mathbb{D}^{k-1}, \mathbb{D}(U)$) denote the disjoint union of l copies of S^1 (resp. $D^2, D^{k-1}, D(U)$). We use the convention that $\mathbb{S}^1 \times \mathbb{D}^{k-1} \times \mathbb{D}(U)$ (resp. $\mathbb{D}^2 \times \mathbb{D}^{k-1} \times \mathbb{D}(U)$) means the disjoint union of l copies of $S^1 \times D^{k-1} \times D(U)$ (resp. $D^2 \times D^{k-1} \times D(U)$).

Now, using $\mathbf{f} = \{f_1, \dots, f_l\}$ to attach l equivariant handles $\mathbb{D}^2 \times \mathbb{D}^{k-1} \times \mathbb{D}(U)$ of type 2 to Γ along $M \times \{1\}$ simultaneously, we obtain a new G -manifold

$$\Gamma_1 = (M \times I) \cup_{\mathbf{f}} (\mathbb{D}^2 \times \mathbb{D}^{k-1} \times \mathbb{D}(U)).$$

The boundary $\partial\Gamma_1$ of Γ_1 consists of two connected components. One of them is equivariantly diffeomorphic to M . We denote the other one by M_1 . It is easy to

see that

$$(*) \quad M_1 = (M - \text{Intf}(\mathbb{S}^1 \times \mathbb{D}^{k-1} \times \mathbb{D}(U))) \cup_{\mathbf{f}} (\mathbb{D}^2 \times \partial(\mathbb{D}^{k-1} \times \mathbb{D}(U))).$$

First, let us observe Γ_1^G and M_1^G in this surgery process. We see that

$$\Gamma_1^G = (M^G \times I) \cup_{\mathbf{f}|_{\mathbb{S}^1 \times \mathbb{D}^{k-1}}} (\mathbb{D}^2 \times \mathbb{D}^{k-1})$$

and

$$M_1^G = (M^G - \text{Intf}(\mathbb{S}^1 \times \mathbb{D}^{k-1})) \cup_{\mathbf{f}|_{\mathbb{S}^1 \times \mathbb{S}^{k-2}}} (\mathbb{D}^2 \times \mathbb{S}^{k-2}),$$

and this surgery process on Γ_1^G and M_1^G just belongs to that of the nonequivariant case. Thus from [Ke] we know that Γ_1^G and M_1^G have the following properties:

- (i) $\pi_1(M_1^G) \cong \pi_1(\Gamma_1^G) \cong 0$.
- (ii) $H_i(M_1^G) \cong 0$ if $i \neq 0, 2, k-2, k$; $H_i(M_1^G) \cong \mathbb{Z}$ if $i = 0, k$. The groups $H_2(M_1^G)$ and $H_{k-2}(M_1^G)$ are free abelian of rank l .
- (iii) $H_i(\Gamma_1^G) \cong 0$ if $i \neq 0, 2, k$; $H_i(\Gamma_1^G) \cong \mathbb{Z}$ if $i = 0, k$. $H_2(\Gamma_1^G)$ is free abelian of rank l .

Next, we look at Γ_1 and M_1 . It is easy to see that

$$\pi_1(\Gamma_1) \cong \pi_1(\Gamma)/A$$

and

$$\pi_1(M_1) \cong \pi_1(M)/B$$

where A (resp. B) is the normal subgroup of $\pi_1(\Gamma)$ (resp. $\pi_1(M)$) generated by $f_i|_{\mathbb{S}^1}, i = 1, \dots, l$. Now we compute the homology groups of Γ_1 and M_1 .

Claim 1. If $i \neq 0, 2, n$, then $H_i(\Gamma_1) \cong 0$. $H_0(\Gamma_1) \cong H_n(\Gamma_1) \cong \mathbb{Z}$. $H_2(\Gamma_1)$ is free abelian of rank l .

Since Γ_1 has

$$O_1 = (M \times I) \cup_{\mathbf{f}|_{\mathbb{S}^1}} \mathbb{D}^2$$

as an G -deformation retract, applying the Mayer-Vietoris exact sequence of the triad $(O_1; M \times I, \mathbb{D}^2)$, it is easy to see that Claim 1 follows.

Claim 2. If $i \neq 0, 2, n-2, n$, then $H_i(M_1) \cong 0$. $H_0(M_1) \cong H_n(M_1) \cong \mathbb{Z}$. The groups $H_2(M_1)$ and $H_{n-2}(M_1)$ are free abelian of rank l .

The proof of Claim 2 can be completed by using the expression $(*)$ to compute the homology group of M_1 . This is basically a standard Quillenization argument combined with a *top hat* construction that was much used in the 1960's. We would like to leave it as an exercise to the reader.

Step II. *2-dimensional surgery on M_1^G .*

Let β_1, \dots, β_l and $\gamma_1, \dots, \gamma_l$ be bases of $H_2(M_1^G)$ and $H_{k-2}(M_1^G)$, respectively, such that the intersection numbers $\alpha_i \cdot \beta_j = \delta_{i,j}$, where $\delta_{i,j} = 1$ if $i = j$ and 0 if $i \neq j$. Then there exist disjoint G -imbeddings g_1, \dots, g_l of

$$S^2 \times D^{k-2} \times D(U_1)$$

into M_1 representing the classes β_1, \dots, β_l , where G -actions on S^2 and D^{k-2} are trivial, and U_1 is the G -representation on the normal fibre $\mu(M_1^G, M_1)_x$ at a point $x \in M_1^G$ in M_1 such that $U_1^G = \{0\}$. Now by simultaneously attaching l equivariant handles $\mathbb{D}^3 \times \mathbb{D}^{k-2} \times \mathbb{D}(U_1)$ of type 3 to Γ_1 along M_1 via $\mathbf{g} = \{g_1, \dots, g_l\}$, we obtain a new G -manifold

$$\Gamma_2 = \Gamma_1 \cup_{\mathbf{g}} (\mathbb{D}^3 \times \mathbb{D}^{k-2} \times \mathbb{D}(U_1)).$$

In two boundary components of Γ_2 , by M_2 we denote the boundary component of $\partial\Gamma_2$ which is not equivariantly diffeomorphic to M . Then M_2 may be expressed as

$$(**) \quad M_2 = (M_1 - \text{Intg}(\mathbb{S}^2 \times \mathbb{D}^{k-2} \times \mathbb{D}(U_1))) \cup_{\mathbf{g}} (\mathbb{D}^3 \times \partial(\mathbb{D}^{k-2} \times \mathbb{D}(U_1))).$$

Note that the above $\mathbb{D}^3 \times \mathbb{D}^{k-2} \times \mathbb{D}(U_1)$, $\mathbb{S}^2 \times \mathbb{D}^{k-2} \times \mathbb{D}(U_1)$ and $\mathbb{D}^3 \times \partial(\mathbb{D}^{k-2} \times \mathbb{D}(U_1))$ fit in with the convention of Step I. The same consideration will also be used in the following discussions.

The above 2-dimensional surgery on the fixed point set is actually done in the nonequivariant case. So we know from the work of Kervaire [Ke] that M_2^G is a homotopy sphere.

Claim 3. Γ_2^H has the same homology as M^H for $H = \{1\}, G$. Furthermore, Γ_2^G is a homology cobordism between M_2^G and M^G .

Since Γ_2 has $O_2 = \Gamma_1 \cup_{\mathbf{g}|_{\mathbb{S}^2}} \mathbb{D}^3$ as its G -deformation retract, applying the Mayer-Veitoris exact sequence of the triad $(O_2; \Gamma_1, \mathbb{D}^3)$ and using the result (iii) and Claim 1 in Step I, it at once follows that for $H = \{1\}, G$,

$$H_i(\Gamma_2^H) \cong H_i(M^H).$$

Next, let us observe the structure of $H_i(M_2)$. Similarly to the argument of Claim 2, by using (**) we can still carry out the Quillenization argument, and obtain

Claim 4. M_2 is a homology sphere.

Up to now, we have finished the 1-dimensional and 2-dimensional surgeries on the fixed point set of M , so that the fixed point set has become a homotopy sphere. If $\pi_1(M_2)$ is trivial, then we have completed the proof. In fact, M_2 is exactly the desired semi-linear homotopy G -sphere Σ , and Γ_2 is a G -equivariant homology cobordism between M and Σ . If $\pi_1(M_2)$ is nontrivial, we continue our discussion as follows.

Step III. *Surgeries on $M_2 - M_2^G$.*

Now we begin with doing surgeries on free orbits. Since M_2 is a homology sphere, the fundamental group of M_2 is finitely presented. Let ξ_1, \dots, ξ_t be a finite set of generators for $\pi_1(M_2)$. Consider the fibration

$$G \longrightarrow M_2 - M_2^G \longrightarrow (M_2 - M_2^G)/G.$$

Since G is finite, we easily see that the fundamental groups of $M_2 - M_2^G$ and $(M_2 - M_2^G)/G$ are isomorphic from the homotopy exact sequence of the above fibration, and there are liftings for embeddings of generators of π_1 of the orbit space. Thus, we may take G -imbeddings h_1, \dots, h_t of

$$G \times_{\{1\}} (S^1 \times D^{n-1})$$

into M_2 with disjoint images representing the generators ξ_1, \dots, ξ_t , where $S^1 \times D^{n-1}$ is a $\{1\}$ -space. Then, as in the arguments of Step I, we may simultaneously glue t equivariant handles

$$G \times_{\{1\}} \mathbb{D}^2 \times \mathbb{D}^{n-1}$$

of type 2 to Γ_2 along M_2 via $\mathbf{h} = \{h_1, \dots, h_t\}$, such that the resulting G -manifold denoted by Γ_3 has the following properties:

- (1) $\Gamma_3^G = \Gamma_2^G$ and $M_3^G = M_2^G$, where $M_3 = \partial\Gamma_3 - (M \times \{0\})$. Note that in this surgery process, the fixed point sets Γ_3^G and M_3^G have not been touched since $\dim V - \dim V^G \geq 3$.

- (2) $\pi_1(M_3) \cong 0$. If $i \neq 0, 2, n-2, n$, then $H_i(M_3) \cong 0$. $H_0(M_3) \cong H_n(M_3) \cong \mathbb{Z}$. $H_2(M_3)$ and $H_{n-2}(M_3)$ are free $\mathbb{Z}[G]$ -module of rank t .
- (3) If $i \neq 0, 2, n$, then $H_i(\Gamma_3) \cong 0$. $H_0(\Gamma_3) \cong H_n(\Gamma_3) \cong \mathbb{Z}$. The group $H_2(\Gamma_3)$ is free $\mathbb{Z}[G]$ -module of rank t .

Finally, we may do 2-dimensional equivariant surgery on Γ_3 along M_3 to kill the basis of $H_2(M_3)$ without touching the fixed point set. It is easily verified that the resulting G -manifold Γ_4 possesses the properties that $\partial\Gamma_4 - (M \times \{0\})$ denoted by Σ is exactly a semi-linear homotopy G -sphere as desired (note that Σ always possesses the homotopy unknottedness since the codimension $\dim \Sigma - \dim \Sigma^G$ is assumed to be greater than or equal to 3), and Γ_4 is a G -equivariant homology cobordism between Σ and M . This completes the proof. \square

Next, we will directly use equivariant spherical modifications to prove Lemma 3.2. Since G is assumed to be finite, this equivariant surgery can still be carried out well. Based upon this, we will omit the detailed computations in the proof, and only give an outline.

Proof of Lemma 3.2. Without loss of generality we assume that W^H is non-simply connected for $H = \{1\}, G$.

We first do surgery on the fixed point set W^G to kill the generators of $\pi_1(W^G)$. This surgery on W^G is done in the nonequivariant case. There always exist disjoint G -imbeddings $\mathbf{f} = \{f_1, \dots, f_l\}$ representing all the l generators of $\pi_1(W^G)$ since W is equivariantly acyclic. By doing 1-dimensional equivariant spherical modification on W via G -imbeddings \mathbf{f} to kill $\pi_1(W^G)$, we obtain a new G -manifold W_1 with the following properties:

- (i) $\pi_1(W_1^G) \cong 0$. If $i \neq 0, 2, k-1$, then $H_i(W_1^G) \cong 0$. The groups $H_2(W_1^G)$ and $H_{k-1}(W_1^G)$ are free \mathbb{Z} -module of rank l .
- (ii) If $i \neq 0, 2, n-1$, then $H_i(W_1) \cong 0$. $H_2(W_1) \cong H_{n-1}(W_1)$ is free \mathbb{Z} -module of rank l .
- (iii) $\partial W_1^H = \partial W^H = M^H$ for $H = \{1\}, G$.

Next, we do a 2-dimensional equivariant modification to kill the basis of $H_2(W_1^G)$. Then the resulting G -manifold W_2 has the properties that W_2^G becomes a contractible manifold with boundary M^G , and W_2 is still an acyclic manifold with boundary M .

If W_2 is not simply connected, since G is finite and $\dim V - \dim V^G \geq 3$, then we may do 1-dimensional and 2-dimensional equivariant surgeries on the free part $W_2 - W_2^G$ without touching the boundary and the fixed point set, creating a new contractible G -manifold as desired. \square

Now we begin with the proof of Theorem 1.3.

Proof of Theorem 1.3. Lemma 3.1 implies that T is an epimorphism if $\dim V^G \geq 5$, and Lemma 3.2 implies that T is a monomorphism if $\dim V^G \geq 4$ (note that for $\langle \Sigma \rangle_G \in \Theta_V^G$, $\langle \Sigma \rangle_G = 0$ if and only if Σ bounds an equivariantly contractible G -manifold). Therefore, T is an isomorphism if $\dim V^G \geq 5$, and a monomorphism if $\dim V^G = 4$. Next, we also need to consider the low-dimensional cases, i.e., $\dim V^G = 1, 2$. Let $\langle M \rangle_G \in H\Theta_V^G$. If $\dim V^G = 1$ (or 2), then M^G is diffeomorphic to S^1 (or S^2). Hence, in the cases $\dim V^G = 1, 2$, when we need to do surgery on M as an G -manifold, the surgery on the fixed point set M^G need not be considered.

Without loss of generality we assume that M is nonsimply connected. We proceed as follows:

- (I) *The case in which either $\dim V^G = 2$ or $\dim V^G = 1$ and $\dim V \geq 5$.* In this case, we claim that T is an isomorphism. Since $\dim V \geq 5$, we may change M into a semi-linear homotopy G -sphere Σ by doing surgeries without touching the fixed point set, such that M is G -equivariantly homologically cobordant to Σ , i.e., Σ may be used as a representative of the class $\langle M \rangle_G$. This means that if either $\dim V^G = 2$ or $\dim V^G = 1$ and $\dim V \geq 5$, then T is an epimorphism. If $\langle M \rangle_G = 0$, i.e., M bounds an equivariantly acyclic G -manifold W , then in a same way as the proof of Lemma 3.2, it is easy to check that M bounds an equivariantly contractible G -manifold. This means that if either $\dim V^G = 2$ or $\dim V^G = 1$ and $\dim V \geq 5$, then T is a monomorphism.
- (II) *The case in which $\dim V^G = 1$ and $\dim V = 4$.* In this case, we may directly compute the groups $H\Theta_V^G$ and Θ_V^G . By [Ke, Theorem 3], we know that every 4-dimensional homology sphere bounds a contractible manifold. Thus if $\dim V^G = 1$ and $\dim V = 4$, then the semi-linear homology G -sphere M bounds an equivariantly contractible G -manifold. This immediately deduces that

$$H\Theta_V^G \cong \Theta_V^G \cong 0$$

if $\dim V^G = 1$ and $\dim V = 4$.

Combining the above discussions, we complete the proof. \square

Remark. (1) If $\dim V^G = 4$, the above argument tells us that T is nothing but a monomorphism. Let $\dim V^G = 4$ and let the G -action be semi-free. Consider the homomorphism

$$K_1 : H\Theta_V^G \longrightarrow H\Theta_{\dim V}$$

defined by mapping $\langle M \rangle_G \in H\Theta_V^G$ to $\langle M \rangle \in H\Theta_{\dim V}$ by forgetting the G -action. If

$$K_1(\langle M \rangle_G) = \langle M \rangle = 0$$

in $H\Theta_{\dim V}$, then M is homologically cobordant to $S^{\dim V}$ so M bounds an acyclic manifold. By [Ke, Theorem 3], every 4-dimensional homology sphere always bounds an acyclic manifold, thus M^G as a homology 4-sphere bounds an acyclic manifold. Since the G -action is assumed to be semi-free, we conclude that M equivariantly bounds an equivariant acyclic G -manifold. Furthermore, by Lemma 2.1 it follows that M is G -equivariantly homologically cobordant to $S(V \oplus \mathbb{R})$, i.e., $\langle M \rangle_G = 0$. Therefore, K_1 is a monomorphism. Since $H\Theta_{\dim V}$ is finite, $H\Theta_V^G$ is finite if $\dim V^G = 4$. In a same way, we may also define the homomorphism

$$K_2 : \Theta_V^G \longrightarrow \Theta_{\dim V}$$

by forgetting the G -action, and obtain that K_2 is a monomorphism too, so Θ_V^G is finite if $\dim V^G = 4$. Obviously, if either K_1 and K_2 are epimorphisms or $\text{Im}K_1$ and $\text{Im}K_2$ are isomorphic, then we can conclude that T is an isomorphism if $\dim V^G = 4$. However, we do not know whether each homology (resp. homotopy) sphere of dimension ≥ 7 must admit a semi-free G -action such that the fixed point

set is a homology (resp. homotopy) 4-sphere. Even so, it is extremely tempting to conjecture that $T : \Theta_V^G \rightarrow H\Theta_V^G$ is an isomorphism if $\dim V^G = 4$.

(2) If G -actions are not assumed to be semi-free and $\dim V^G$ is not restricted to be 4, and G is not necessarily finite, by forgetting G -actions we can still obtain two homomorphisms like K_1 and K_2 , so that the following commutative diagram can be obtained:

$$\begin{array}{ccc} \Theta_V^G & \longrightarrow & H\Theta_V^G \\ \downarrow & & \downarrow \\ \Theta_{\dim V} & \longrightarrow & H\Theta_{\dim V} \end{array}$$

This gives a connection between ordinary cases and equivariant cases.

ACKNOWLEDGMENTS

The author expresses his gratitude to Professors T. Tsuboi and M. Masuda for their advice and suggestions in this work. The author also expresses his gratitude to the referee, who did an extremely careful reading of this paper and detected some flaws in the proof of the main result in earlier versions. The many suggestions and comments made by him considerably improved the presentation of this paper, leading to the present final version.

REFERENCES

- [AB] M. F. Atiyah and R. Bott, *A Lefschetz fixed point formula for elliptic complexes. II: Applications*, Ann. of Math. (2) **88** (1968), 451-491. MR **38**:731
- [Br] G. E. Bredon, *Introduction to compact transformation groups*, Academic Press, 1972. MR **54**:1265
- [FS] R. Fintushel and R. Stern, *Pseudofree orbifolds*, Ann. of Math. **122** (1985), 335-364. MR **87a**:57027
- [Fu] M. Furuta, *Homology cobordism group of homology 3-spheres*, Invent. Math. **100** (1990), 339-355. MR **91c**:57039
- [HH] W. C. Hsiang and W. Y. Hsiang, *Differentiable actions of compact connected classical groups I*, Amer. J. Math. **89** (1967), 705-786. MR **36**:304
- [Ke] M. Kervaire, *Smooth homology spheres and their fundamental groups*, Trans. Amer. Math. Soc. **144** (1969), 67-72. MR **40**:6562
- [KM] M. Kervaire and J. Milnor, *Groups of homotopy spheres*, Ann. of Math. **77** (1963), 504-537. MR **26**:5584
- [Ko] A. Kosiński, *On the inertia groups of π -manifolds*, Amer. J. Math. **89** (1967), 227-248. MR **35**:4936
- [Ma] M. Masuda, *A product formula for connected sum*, Transformation Groups (Proceedings, Osaka, 1987), Lecture Notes in Math. Vol. 1375, Springer, Berlin-Heidelberg-New York-Tokyo, 1989, 231-239. MR **90k**:57044
- [MSc1] M. Masuda and R. Schultz, *Invariants of Atiyah-Singer type, classifications up to finite ambiguity, and equivariant inertia groups*, Indiana Univ. Math. J. **45** (1996), 545-581. MR **97i**:57032
- [MSc2] M. Masuda and R. Schultz, *On the nonuniqueness of equivariant connected sums*, J. Math. Soc. Japan **51** (1999), 411-435. MR **2000b**:57049
- [Mi1] J. Milnor, *Lecture on the h-cobordism theorem*, Princeton University Press, 1965. MR **32**:8352
- [Mi2] J. Milnor, *Whitehead torsion*, Bull. Amer. Math. Soc. **72** (1966), 358-426. MR **33**:4922
- [PR] T. Petrie and J. D. Randall, *Transformation groups on manifolds*, Monographs and textbooks in pure and applied mathematics, 1984. MR **85m**:57026
- [Sc1] R. Schultz, *On the inertia group of a product of spheres*, Trans. Amer. Math. Soc. **156** (1971), 137-153. MR **43**:1209

- [Sc2] R. Schultz, *Differentiable group actions on homotopy spheres II: Ultrasemi-free actions*, Trans. Amer. Math. Soc. **268** (1981), 255-297. MR **83a**:57055
- [Sc3] R. Schultz, *Nonlinear analogs of linear group actions on spheres*, Bull. Amer. Math. Soc. (2) **11** (1984), 263-285. MR **86i**:57049
- [Wi1] D. L. Wilkens, *On the inertia groups of certain manifolds*, J. London Math. Soc. (2) **9** (1975), 537-548. MR **52**:4316
- [Wi2] D. L. Wilkens, *On inertia groups and bordism*, Michigan Math. J. **23** (1976), 105-106. MR **53**:14511
- [Win] H. E. Winkelkemper, *On the action of Θ^n* , Trans. Amer. Math. Soc. **206** (1975), 339-346. MR **54**:1257

INSTITUTE OF MATHEMATICS, FUDAN UNIVERSITY, SHANGHAI, 200433, PEOPLE'S REPUBLIC OF CHINA

E-mail address: zlu@fudan.edu.cn

Current address: Department of Mathematics, Osaka City University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan