

## VALUE GROUPS, RESIDUE FIELDS, AND BAD PLACES OF RATIONAL FUNCTION FIELDS

FRANZ-VIKTOR KUHLMANN

**ABSTRACT.** We classify all possible extensions of a valuation from a ground field  $K$  to a rational function field in one or several variables over  $K$ . We determine which value groups and residue fields can appear, and we show how to construct extensions having these value groups and residue fields. In particular, we give several constructions of extensions whose corresponding value group and residue field extensions are not finitely generated. In the case of a rational function field  $K(x)$  in one variable, we consider the relative algebraic closure of  $K$  in the henselization of  $K(x)$  with respect to the given extension, and we show that this can be any countably generated separable-algebraic extension of  $K$ . In the “tame case”, we show how to determine this relative algebraic closure. Finally, we apply our methods to power series fields and the  $p$ -adics.

### 1. INTRODUCTION

In this paper, we denote a valued field by  $(K, v)$ , its value group by  $vK$ , and its residue field by  $Kv$ . When we write  $(L|K, v)$  we mean a field extension  $L|K$  endowed with a valuation  $v$  on  $L$  and its restriction on  $K$ .

In many recent applications of valuation theory, valuations on algebraic function fields play a main role. To mention only a short and incomplete list of applications and references: local uniformization and resolution of singularities ([C], [CP], [S], [KU3], [KKU1], [KKU2]), model theory of valued fields ([KU1], [KU2], [KU4]), study of curves via constant reduction ([GMP1], [GMP2], [PL]), classification of all extensions of an ordering from a base field to a rational function field ([KUKMZ]), and Gröbner bases ([SW], [MOSW1], [MOSW2]).

In many cases, a basic tool is the classification of all extensions of a valuation from a base field to a function field. As the classification of all extensions of a valuation from a field to an algebraic extension is taken care of by general ramification theory (cf. [E], [KU2]), a crucial step in the classification is the case of rational function fields. Among the first papers describing valuations on rational function fields systematically were [M] and [MS]. Since then, an impressive number of papers have been written about the construction of such valuations and about their properties; the following list is by no means exhaustive: [AP], [APZ1]–[APZ3], [KH1]–[KH10], [KHG1]–[KHG6], [KHPR], [MO1], [MO2], [MOSW1], [O1]–[O3], [PP], [V]. From the paper [APZ3] the reader may get a good idea of how MacLane’s original approach has been developed further. Since then, the notion of “minimal pairs” has been

---

Received by the editors July 12, 2002 and, in revised form, July 15, 2003.

2000 *Mathematics Subject Classification.* Primary 12J10; Secondary 12J15, 16W60.

adopted and studied by several authors (see, e.g., [KHPR]). In the present paper, we will develop a new approach to this subject. It serves to determine in full generality which value groups and which residue fields can possibly occur. This question has recently played a role in two other papers:

1) In [KU3], we prove the existence of “bad places” on rational function fields of transcendence degree  $\geq 2$ . These are places whose value group is not finitely generated, or whose residue field is not finitely generated over the base field. The existence of such places has been shown by MacLane and Schilling ([MS]) and by Zariski and Samuel ([ZS], Chapter VI, §15, Examples 3 and 4). However, our approach in [KU3] using Hensel’s Lemma seems to be new, and the present paper contains a further refinement of it. The following theorem of [MS] and [ZS] is a special case of a result which we will prove by this refinement:

**Theorem 1.1.** *Let  $K$  be any field. Take  $\Gamma$  to be any non-trivial ordered abelian group of finite rational rank  $\rho$ , and  $k$  to be any countably generated extension of  $K$  of finite transcendence degree  $\tau$ . Choose any integer  $n > \rho + \tau$ . Then the rational function field in  $n$  variables over  $K$  admits a valuation whose restriction to  $K$  is trivial, whose value group is  $\Gamma$  and whose residue field is  $k$ .*

*In particular, every additive subgroup of  $\mathbb{Q}$  and every countably generated algebraic extension of  $K$  can be realized as value group and residue field of a place of the rational function field  $K(x, y)|K$  whose restriction to  $K$  is the identity.*

The rational rank of an abelian group  $\Gamma$  is the dimension of the  $\mathbb{Q}$ -vector space  $\mathbb{Q} \otimes_{\mathbb{Z}} \Gamma$ . We denote it by  $\text{rr } \Gamma$ . It is equal to the cardinality of any maximal set of rationally independent elements in  $\Gamma$ .

Bad places on function fields are indeed bad: the value group or residue field not being finitely generated constitutes a major hurdle for the attempt to prove local uniformization or other results which are related to resolution of singularities (cf. [CP]). Another hurdle is the phenomenon of **defect** which can appear when the residue characteristic of a valued field is positive, even if the field itself seems to be quite simple. Indeed, we will prove in Section 3.3, and by a different method in Section 3.5:

**Theorem 1.2.** *Let  $K$  be any algebraically closed field of positive characteristic. Then there exists a valuation  $v$  on the rational function field  $K(x, y)|K$  whose restriction to  $K$  is trivial, such that  $(K(x, y), v)$  admits an infinite chain of immediate Galois extensions of degree  $p$  and defect  $p$ .*

An extension  $(L'|L, v)$  of valued fields is called **immediate** if the canonical embeddings of  $vL$  in  $vL'$  and of  $Lv$  in  $L'v$  are surjective (which we will express by writing  $vL' = vL$  and  $L'v = Lv$ ). For a finite immediate extension  $(L'|L, v)$ , its defect is equal to its degree if and only if the extension of  $v$  from  $L$  to  $L'$  is unique (or equivalently,  $L'|L$  is linearly disjoint from some (or every) henselization of  $(L, v)$ ).

One of the examples we shall construct for the proof of the above theorem is essentially the same as in Section 7 of [CP], but we use a different and more direct approach (while the construction in [CP] is more intricate, since it serves an additional purpose).

2) In [KUKMZ], the classification of all extensions of an ordering to a rational function field is considered in the context of power series fields, and the above

question is partially answered in this setting. In the present paper, we will consider the question without referring to power series fields (see Theorem 1.8 below).

During the preparation of [KUKMZ], we found that the construction of an extension of the valuation  $v$  from  $K$  to the rational function field  $K(x)$  with prescribed value group  $vK(x)$  and residue field  $K(x)v$  is tightly connected with the determination of the relative algebraic closure of  $K$  in a henselization  $K(x)^h$  of  $K(x)$  with respect to  $v$ . In earlier papers, we have introduced the name “henselian function field” for the henselizations of valued function fields (although these are not function fields, unless the valuation is trivial). In the same vein, one can view the relative algebraic closure as being the (exact) constant field of the henselian function field  $(K(x)^h|K, v)$ . We will call it the **implicit constant field of  $(K(x)|K, v)$**  and denote it by  $\text{IC}(K(x)|K, v)$ . Clearly, the henselization  $K(x)^h$  depends on the valuation which has been fixed on the algebraic closure  $\widetilde{K(x)}$ . So whenever we will talk about the implicit constant field, we will do it in a setting where the valuation on  $\widetilde{K(x)}$  has been fixed. However, since the henselization  $L^h$  of any valued field  $(L, v)$  is unique up to valuation-preserving isomorphism over  $L$ , the implicit constant field is unique up to valuation-preserving isomorphism over  $K$ . If  $L_0$  is a subfield of  $L$ , then  $L^h$  contains a (unique) henselization of  $L_0$ . Hence,  $\text{IC}(K(x)|K, v)$  contains a henselization of  $K$  and is itself henselian. Further,  $L^h|L$  is a separable-algebraic extension; thus,  $K(x)^h|K$  is separable. Therefore,  $\text{IC}(K(x)|K, v)$  is a separable-algebraic extension of  $K$ .

In the present paper, we answer the above question on value groups and residue fields by determining which prescribed separable-algebraic extensions of  $K$  can be realized as implicit constant fields. The following result shows in particular that every countably generated separable-algebraic extension of a henselian base field can be so realized:

**Theorem 1.3.** *Let  $(K_1|K, v)$  be a countably generated separable-algebraic extension of non-trivially valued fields. Then there is an extension of  $v$  from  $K_1$  to the algebraic closure  $\widetilde{K_1(x)} = \widetilde{K(x)}$  of the rational function field  $K(x)$  such that, upon taking henselizations in  $(\widetilde{K(x)}, v)$ ,*

$$(1.1) \quad K_1^h = \text{IC}(K(x)|K, v).$$

In Section 3.1 we will introduce a basic classification (“value-transcendental” – “residue-transcendental” – “valuation-algebraic”) of all possible extensions of  $v$  from  $K$  to  $K(x)$ . In Section 3.2 we introduce a class of extensions  $(K(x)|K, v)$  for which  $\text{IC}(K(x)|K, v) = K^h$  holds. Building on this, we prove Theorem 1.3 in Section 3.5. In fact, we prove a more detailed version: we show under which additional conditions the extension can be chosen in a prescribed class of the basic classification. This yields the following:

**Theorem 1.4.** *Take any valued field  $(K, v)$ , an ordered abelian group extension  $\Gamma_0$  of  $vK$  such that  $\Gamma_0/vK$  is a torsion group, and an algebraic extension  $k_0$  of  $Kv$ . Further, take  $\Gamma$  to be the abelian group  $\Gamma_0 \oplus \mathbb{Z}$  endowed with any extension of the ordering of  $\Gamma_0$ .*

*Assume first that  $\Gamma_0/vK$  and  $k_0|Kv$  are finite. If  $v$  is trivial on  $K$ , then assume in addition that  $k_0|Kv$  is simple. Then there is an extension of  $v$  from  $K$  to the rational function field  $K(x)$  which has value group  $\Gamma$  and residue field  $k_0$ . If  $v$  is*

non-trivial on  $K$ , then there is also an extension which has value group  $\Gamma_0$  and as residue field a rational function field in one variable over  $k_0$ .

Now assume that  $v$  is non-trivial on  $K$  and that  $\Gamma_0/vK$  and  $k_0|Kv$  are countably generated. Suppose that at least one of them is infinite or that  $(K, v)$  admits an immediate transcendental extension. Then there is an extension of  $v$  from  $K$  to  $K(x)$  which has value group  $\Gamma_0$  and residue field  $k_0$ .

Here is the converse:

**Theorem 1.5.** *Let  $(K(x)|K, v)$  be a valued rational function field. Then one and only one of the following three cases holds:*

- 1)  $vK(x) \simeq \Gamma_0 \oplus \mathbb{Z}$ , where  $\Gamma_0|vK$  is a finite extension of ordered abelian groups, and  $K(x)v|Kv$  is finite;
  - 2)  $vK(x)/vK$  is finite, and  $K(x)v$  is a rational function field in one variable over a finite extension of  $Kv$ ;
  - 3)  $vK(x)/vK$  is a torsion group and  $K(x)v|Kv$  is algebraic.
- In all cases,  $vK(x)/vK$  is countable and  $K(x)v|Kv$  is countably generated.

In 2), we use a fact which was proved by J. Ohm [O2] and is known as the “Ruled Residue Theorem”: *If  $K(x)v|Kv$  is transcendental, then  $K(x)v$  is a rational function field in one variable over a finite extension of  $Kv$ .* For the countability assertion, see Theorem 2.9.

In Section 3.4 we give an explicit description of all possible extensions of  $v$  from  $K$  to  $K(x)$  (Theorem 3.11).

Theorem 1.4 is used in the proof of our next theorem:

**Theorem 1.6.** *Let  $(K, v)$  be any valued field,  $n, \rho, \tau$  non-negative integers,  $n \geq 1$ ,  $\Gamma \neq \{0\}$  an ordered abelian group extension of  $vK$  such that  $\Gamma/vK$  is of rational rank  $\rho$ , and  $k|Kv$  a field extension of transcendence degree  $\tau$ .*

**Part A.** *Suppose that  $n > \rho + \tau$  and that*

- A1)  $\Gamma/vK$  and  $k|Kv$  are countably generated, and
- A2)  $\Gamma/vK$  or  $k|Kv$  is infinite.

*Then there is an extension of  $v$  to the rational function field  $K(x_1, \dots, x_n)$  in  $n$  variables such that*

$$(1.2) \quad vK(x_1, \dots, x_n) = \Gamma \quad \text{and} \quad K(x_1, \dots, x_n)v = k.$$

**Part B.** *Suppose that  $n \geq \rho + \tau$  and that*

- B1)  $\Gamma/vK$  and  $k|Kv$  are finitely generated,
- B2) if  $v$  is trivial on  $K$ ,  $n = \rho + \tau$  and  $\rho = 1$ , then  $k$  is a simple algebraic extension of a rational function field in  $\tau$  variables over  $Kv$  (or of  $Kv$  itself if  $\tau = 0$ ), or a rational function field in one variable over a finitely generated field extension of  $Kv$  of transcendence degree  $\tau - 1$ ,
- B3) if  $n = \tau$ , then  $k$  is a rational function field in one variable over a finitely generated field extension of  $Kv$  of transcendence degree  $\tau - 1$ , and
- B4) if  $\rho = 0 = \tau$ , then there is an immediate extension of  $(K, v)$  which is either infinite separable-algebraic or of transcendence degree at least  $n$ .

*Then again there is an extension of  $v$  to  $K(x_1, \dots, x_n)$  such that (1.2) holds.*

Theorem 1.1 is the special case of Part A for  $v$  trivial on  $K$ . The following converse holds.

**Theorem 1.7.** *Let  $n \geq 1$ , and let  $v$  be a valuation on the rational function field  $F = K(x_1, \dots, x_n)$ . Set  $\rho = \text{rr } vF/vK$  and  $\tau = \text{trdeg } Fv|Kv$ . Then  $n \geq \rho + \tau$ ,  $vF/vK$  is countable, and  $Fv|Kv$  is countably generated.*

*If  $n = \rho + \tau$ , then  $vF/vK$  is finitely generated and  $Fv|Kv$  is a finitely generated field extension. Assertions B2) and B3) of Theorem 1.6 hold for  $k = Fv$ , and if  $\rho = 0 = \tau$ , then there is an immediate extension of  $(\tilde{K}, v)$  of transcendence degree  $n$  (for any extension of  $v$  from  $K$  to  $\tilde{K}$ ).*

There is a gap between Theorem 1.6 and this converse for the case of  $\rho = 0 = \tau$ , as the former talks about  $K$  and the latter talks about the algebraic closure  $\bar{K}$  of  $K$ . This gap can be closed if  $(K, v)$  has residue characteristic 0 or is a Kaplansky field; because the maximal immediate extension of such fields is unique up to isomorphism, one can show that  $\bar{K}$  can be replaced by  $K$ . But in the case where  $(K, v)$  is not such a field, we do not know enough about the behaviour of maximal immediate extensions under algebraic field extensions. This question should be considered in future research.

A valuation on an ordered field is called **convex** if the associated valuation ring is convex. For the case of ordered fields with convex valuations, we can derive from Theorem 1.6 the existence of convex extensions of the valuation with prescribed value groups and residue fields in the frame given by Theorem 1.7, provided that a natural additional condition for the residue fields is satisfied:

**Theorem 1.8.** *In the setting of Theorem 1.6, assume in addition that  $K$  is ordered and that  $v$  is convex w.r.t. the ordering. Assume further that  $k$  is equipped with an extension of the ordering induced by  $<$  on  $Kv$ . Then this extension can be lifted through  $v$  to  $K(x_1, \dots, x_n)$  in such a way that the lifted ordering extends  $<$ . It follows that  $v$  is convex w.r.t. this lifted ordering on  $K(x_1, \dots, x_n)$ .*

In Section 5 we shall introduce “homogeneous sequences”. In the “tame case”, they can be used to determine the implicit constant field of a valued rational function field, and also to characterize this “tame case”. In Section 6 we shall show how to apply our results to power series, in the spirit of [MS] and [ZS] (Theorem 6.1). We will also use our approach to give proofs of two well known facts in  $p$ -adic algebra: that the algebraic closure of  $\mathbb{Q}_p$  is not complete and that its completion is not maximal.

Finally, let us mention that we use our criteria for  $\text{IC}(K(x)|K, v) = K^h$  in Section 3.2 to give an example for the following fact: *Suppose that  $K$  is relatively algebraically closed in a henselian valued field  $(L, v)$  such that  $vL/vK$  is a torsion group. Then it is not necessarily true that  $vL = vK$ , even if  $v$  has residue characteristic 0.*

I would like to thank Murray Marshall and Salma Kuhlmann for our very inspiring joint seminar; without this seminar, this paper would not have been written. I also feel very much indebted to Sudesh Kaur Khanduja for many ideas she has shared with me. Finally, I would like to thank Roland Auer for finding some mistakes and asking critical questions.

## 2. NOTATION AND VALUATION THEORETICAL PRELIMINARIES

For an arbitrary field  $K$ , we will denote by  $K^{\text{sep}}$  the separable-algebraic closure of  $K$ , and by  $\bar{K}$  the algebraic closure of  $K$ . By  $\text{Gal } K$  we mean the absolute Galois

group  $\text{Gal}(\tilde{K}|K) = \text{Gal}(K^{\text{sep}}|K)$ . For a valuation  $v$  on  $K$ , we let  $\mathcal{O}_K$  denote the valuation ring of  $v$  on  $K$ .

Every finite extension  $(L|K, v)$  of valued fields satisfies the **fundamental inequality** (cf. [E]):

$$(2.1) \quad n \geq \sum_{i=1}^g e_i f_i$$

where  $n = [L : K]$  is the degree of the extension,  $v_1, \dots, v_g$  are the distinct extensions of  $v$  from  $K$  to  $L$ ,  $e_i = (v_i L : vK)$  are the respective ramification indices and  $f_i = [Lv_i : Kv]$  are the respective inertia degrees. Note that  $g = 1$  if  $(K, v)$  is henselian.

In analogy to field theory, an extension  $\Gamma \subset \Delta$  of abelian groups will also be written as  $\Delta/\Gamma$ , and it will be called **algebraic** if  $\Delta/\Gamma$  is a torsion group. The fundamental inequality implies the following well known fact:

**Lemma 2.1.** *If  $(L|K, v)$  is finite, then so are  $vL/vK$  and  $Lv|Kv$ . If  $(L|K, v)$  is algebraic, then so are  $vL/vK$  and  $Lv|Kv$ .*

Given two subextensions  $M|K$  and  $L|K$  within a fixed extension  $N|K$ , the **field compositum**  $M.L$  is defined to be the smallest subfield of  $N$  which contains both  $M$  and  $L$ . If  $L|K$  is algebraic, the compositum is uniquely determined by taking  $N = \tilde{M}$  and specifying a  $K$ -embedding of  $L$  in  $\tilde{M}$ .

**Lemma 2.2.** *Let  $(M|K, v)$  be an immediate extension of valued fields, and  $(L|K, v)$  a finite extension such that  $[L : K] = (vL : vK)[Lv : Kv]$ . Then for every  $K$ -embedding of  $L$  in  $\tilde{M}$  and every extension of  $v$  from  $M$  to  $\tilde{M}$ , the extension  $(M.L|L, v)$  is immediate.*

*Proof.* Via the embedding, we identify  $L$  with a subfield of  $\tilde{M}$ . Pick any extension of  $v$  from  $M$  to  $\tilde{M}$ . This will also be an extension of  $v$  from  $L$  to  $\tilde{M}$ , because by the fundamental inequality, the extension of  $v$  from  $K$  to  $L$  is unique. We consider the extension  $(M.L|M, v)$ . It is clear that  $vL \subseteq v(M.L)$  and  $Lv \subseteq (M.L)v$ ; therefore,  $(v(M.L) : vK) \geq (vL : vK)$  and  $[(M.L)v : Kv] \geq [Lv : Kv]$ . Since  $(M|K, v)$  is immediate, we have

$$\begin{aligned} [M.L : M] &\geq (v(M.L) : vM)[(M.L)v : Mv] = (v(M.L) : vK)[(M.L)v : Kv] \\ &\geq (vL : vK)[Lv : Kv] = [L : K] \geq [M.L : M]. \end{aligned}$$

This shows that  $(v(M.L) : vK) = (vL : vK)$  and  $[(M.L)v : Kv] = [Lv : Kv]$ ; that is,  $v(M.L) = vL$  and  $(M.L)v = Lv$ .  $\square$

**2.1. Pseudo Cauchy sequences.** We assume the reader to be familiar with the theory of pseudo Cauchy sequences as presented in [KA]. Recall that a pseudo Cauchy sequence  $\mathbf{A} = (a_\nu)_{\nu < \lambda}$  in  $(K, v)$  (where  $\lambda$  is some limit ordinal) is **of transcendental type** if for every  $g(x) \in K(x)$ , the value  $vg(a_\nu)$  is eventually constant, that is, there is some  $\nu_0 < \lambda$  such that

$$(2.2) \quad vg(a_\nu) = vg(a_{\nu_0}) \quad \text{for all } \nu \geq \nu_0, \nu < \lambda.$$

Otherwise,  $\mathbf{A}$  is **of algebraic type**.

Take a pseudo Cauchy sequence  $\mathbf{A}$  in  $(K, v)$  of transcendental type. We define an extension  $v_{\mathbf{A}}$  of  $v$  from  $K$  to the rational function field  $K(x)$  as follows. For

each  $g(x) \in K[x]$ , we choose  $\nu_0 < \lambda$  such that (2.2) holds. Then we set

$$v_{\mathbf{A}} g(x) := v g(a_{\nu_0}).$$

We extend  $v_{\mathbf{A}}$  to  $K(x)$  by setting  $v_{\mathbf{A}}(g/h) := v_{\mathbf{A}} g - v_{\mathbf{A}} h$ . The following is Theorem 2 of [KA]:

**Theorem 2.3.** *Let  $\mathbf{A}$  be a pseudo Cauchy sequence in  $(K, v)$  of transcendental type. Then  $v_{\mathbf{A}}$  is a valuation on the rational function field  $K(x)$ . The extension  $(K(x)|K, v_{\mathbf{A}})$  is immediate, and  $x$  is a pseudo limit of  $\mathbf{A}$  in  $(K(x), v_{\mathbf{A}})$ . If  $(K(y), w)$  is any other valued extension of  $(K, v)$  such that  $y$  is a pseudo limit of  $\mathbf{A}$  in  $(K(y), w)$ , then  $x \mapsto y$  induces a valuation preserving  $K$ -isomorphism from  $(K(x), v_{\mathbf{A}})$  onto  $(K(y), w)$ .*

From this theorem we deduce

**Lemma 2.4.** *Suppose that in some valued field extension of  $(K, v)$ ,  $x$  is the pseudo limit of a pseudo Cauchy sequence in  $(K, v)$  of transcendental type. Then the extension  $(K(x)|K, v)$  is immediate and  $x$  is transcendental over  $K$ .*

*Proof.* Assume that  $(a_{\nu})_{\nu < \lambda}$  is a pseudo Cauchy sequence in  $(K, v)$  of transcendental type. Then by Theorem 2.3 there is an immediate extension  $w$  of  $v$  to the rational function field  $K(y)$  such that  $y$  becomes a pseudo limit of  $(a_{\nu})_{\nu < \lambda}$ ; moreover, if also  $x$  is a pseudo limit of  $(a_{\nu})_{\nu < \lambda}$  in  $(K(x), v)$ , then  $x \mapsto y$  induces a valuation-preserving isomorphism from  $K(x)$  onto  $K(y)$  over  $K$ . Hence,  $(K(x)|K, v)$  is immediate and  $x$  is transcendental over  $K$ .  $\square$

**Lemma 2.5.** *A pseudo Cauchy sequence of transcendental type in a valued field remains a pseudo Cauchy sequence of transcendental type in every algebraic valued field extension of that field.*

*Proof.* Assume that  $(a_{\nu})_{\nu < \lambda}$  is a pseudo Cauchy sequence in  $(K, v)$  of transcendental type and that  $(L|K, v)$  is an algebraic extension. If  $(a_{\nu})_{\nu < \lambda}$  were of algebraic type over  $(L, v)$ , then by Theorem 3 of [KA] there would be an algebraic extension  $L(y)|L$  and an immediate extension of  $v$  to  $L(y)$  such that  $y$  is a pseudo limit of  $(a_{\nu})_{\nu < \lambda}$  in  $(L(y), v)$ . But then,  $y$  is also a pseudo limit of  $(a_{\nu})_{\nu < \lambda}$  in  $(K(y), v)$ . Hence by the foregoing lemma,  $y$  must be transcendental over  $K$ . This is a contradiction to the fact that  $L(y)|L$  and  $L|K$  are algebraic.  $\square$

**2.2. Valuation independence.** For the easy proof of the following lemma, see [B], Chapter VI, §10.3, Theorem 1.

**Lemma 2.6.** *Let  $(L|K, v)$  be an extension of valued fields. Take elements  $x_i, y_j \in L$ ,  $i \in I$ ,  $j \in J$ , such that the values  $vx_i$ ,  $i \in I$ , are rationally independent over  $vK$ , and the residues  $y_j v$ ,  $j \in J$ , are algebraically independent over  $Kv$ . Then the elements  $x_i, y_j$ ,  $i \in I$ ,  $j \in J$ , are algebraically independent over  $K$ .*

Moreover, if we write

$$f = \sum_k c_k \prod_{i \in I} x_i^{\mu_{k,i}} \prod_{j \in J} y_j^{\nu_{k,j}} \in K[x_i, y_j \mid i \in I, j \in J]$$

in such a way that for every  $k \neq \ell$  there is some  $i$  such that  $\mu_{k,i} \neq \mu_{\ell,i}$  or some  $j$  such that  $\nu_{k,j} \neq \nu_{\ell,j}$ , then

$$(2.3) \quad vf = \min_k v c_k \prod_{i \in I} x_i^{\mu_{k,i}} \prod_{j \in J} y_j^{\nu_{k,j}} = \min_k v c_k + \sum_{i \in I} \mu_{k,i} v x_i.$$

That is, the value of the polynomial  $f$  is equal to the least of the values of its monomials. In particular, this implies

$$\begin{aligned} vK(x_i, y_j \mid i \in I, j \in J) &= vK \oplus \bigoplus_{i \in I} \mathbb{Z}vx_i, \\ K(x_i, y_j \mid i \in I, j \in J)v &= Kv(y_jv \mid j \in J). \end{aligned}$$

Moreover, the valuation  $v$  on  $K(x_i, y_j \mid i \in I, j \in J)$  is uniquely determined by its restriction to  $K$ , the values  $vx_i$  and the residues  $y_jv$ .

Conversely, if  $(K, v)$  is any valued field and we assign to the  $vx_i$  any values in an ordered group extension of  $vK$  which are rationally independent, then (2.3) defines a valuation on  $L$ , and the residues  $y_jv$ ,  $j \in J$ , are algebraically independent over  $Kv$ .

**Corollary 2.7.** *Let  $(L|K, v)$  be an extension of finite transcendence degree of valued fields. Then*

$$(2.4) \quad \text{trdeg } L|K \geq \text{trdeg } Lv|Kv + \text{rr}(vL/vK).$$

If in addition  $L|K$  is a function field and if equality holds in (2.4), then the extensions  $vL|vK$  and  $Lv|Kv$  are finitely generated.

*Proof.* Choose elements  $x_1, \dots, x_\rho, y_1, \dots, y_\tau \in L$  such that the values  $vx_1, \dots, vx_\rho$  are rationally independent over  $vK$  and the residues  $y_1v, \dots, y_\tau v$  are algebraically independent over  $Kv$ . Then by the foregoing lemma,  $\rho + \tau \leq \text{trdeg } L|K$ . This proves that  $\text{trdeg } Lv|Kv$  and the rational rank of  $vL/vK$  are finite. Therefore, we may choose the elements  $x_i, y_j$  such that  $\tau = \text{trdeg } Lv|Kv$  and  $\rho = \text{rr}(vL/vK)$  to obtain inequality (2.4).

Assume that equality holds in (2.4). This means that  $L$  is an algebraic extension of  $L_0 := K(x_1, \dots, x_\rho, y_1, \dots, y_\tau)$ . Since  $L|K$  is finitely generated, it follows that  $L|L_0$  is finite; hence by Lemma 2.1, also  $vL/vL_0$  and  $Lv|L_0v$  are finite. Since already  $vL_0|vK$  and  $L_0v|Kv$  are finitely generated by the foregoing lemma, it follows that also  $vL|vK$  and  $Lv|Kv$  are finitely generated.  $\square$

The algebraic analogue of the transcendental case discussed in Lemma 2.6 is the following lemma (see [R] or [E]):

**Lemma 2.8.** *Let  $(L|K, v)$  be an extension of valued fields. Take  $\eta_i \in L$  such that  $v\eta_i$ ,  $i \in I$ , belong to distinct cosets modulo  $vK$ . Further, take  $\vartheta_j \in \mathcal{O}_L$ ,  $j \in J$ , such that  $\vartheta_jv$  are  $Kv$ -linearly independent. Then the elements  $\eta_i\vartheta_j$ ,  $i \in I$ ,  $j \in J$ , are  $K$ -linearly independent, and for every choice of elements  $c_{ij} \in K$ , only finitely many of them nonzero, we have that*

$$v \sum_{i,j} c_{ij} \eta_i \vartheta_j = \min_{i,j} v c_{ij} \eta_i \vartheta_j = \min_{i,j} (v c_{ij} + v \eta_i).$$

If the elements  $\eta_i\vartheta_j$  form a  $K$ -basis of  $L$ , then  $v\eta_i$ ,  $i \in I$ , is a system of representatives of the cosets of  $vL$  modulo  $vK$ , and  $\vartheta_jv$ ,  $j \in J$ , is a basis of  $Lv|Kv$ .

The following is an application which is important for our description of all possible value groups and residue fields of valuations on  $K(x)$ . The result has been proved with a different method in [APZ3] (Corollary 5.2); cf. Remark 3.1 in Section 3.1.

**Theorem 2.9.** *Let  $K$  be any field and  $v$  any valuation of the rational function field  $K(x)$ . Then  $vK(x)/vK$  is countable, and  $K(x)v|Kv$  is countably generated.*



*Proof.* Since  $K(x)$  is the quotient field of  $K[x]$ , we have that  $vK(x) = vK[x] - vK[x]$ . Hence, to show that  $vK(x)/vK$  is countable, it suffices to show that the set  $\{\alpha + vK \mid \alpha \in vK[x]\}$  is countable. If this were not true, then by Lemma 2.8 (applied with  $J = \{1\}$  and  $\vartheta_1 = 1$ ), we would have that  $K[x]$  contains uncountably many  $K$ -linearly independent elements. But this is not true, as  $K[x]$  admits the countable  $K$ -basis  $\{x^i \mid i \geq 0\}$ .

Now assume that  $K(x)v|Kv$  is not countably generated. Then by Corollary 2.7,  $K(x)v|Kv$  must be algebraic. It also follows that  $K(x)v$  has uncountable  $Kv$ -dimension. Pick an uncountable set  $\kappa$  and elements  $f_i(x)/g_i(x)$ ,  $i \in \kappa$ , with  $f_i(x), g_i(x) \in K[x]$  and  $vf_i(x) = vg_i(x)$  for all  $i$ , such that their residues are  $Kv$ -linearly independent. As  $vK(x)/vK$  is countable, there must be some uncountable subset  $\lambda \subset \kappa$  such that for all  $i \in \lambda$ , the values  $vf_i(x) = vg_i(x)$  lie in the same coset modulo  $vK$ , say  $vh(x) + vK$  with  $h(x) \in K[x]$ . The residues  $(f_i(x)/g_i(x))v$ ,  $i \in \lambda$ , generate an algebraic extension of uncountable dimension. Choosing suitable elements  $c_i \in K$  such that

$$vc_i f_i(x) = vh(x) = vc_i g_i(x),$$

we can write

$$\frac{f_i(x)}{g_i(x)} = \frac{c_i f_i(x)}{h(x)} \cdot \frac{h(x)}{c_i g_i(x)} = \frac{c_i f_i(x)}{h(x)} \cdot \left( \frac{c_i g_i(x)}{h(x)} \right)^{-1}$$

for all  $i \in \lambda$ . Therefore,

$$\frac{f_i(x)}{g_i(x)} v = \left( \frac{c_i f_i(x)}{h(x)} v \right) \cdot \left( \frac{c_i g_i(x)}{h(x)} v \right)^{-1}$$

for all  $i \in \lambda$ . In order that these elements generate an algebraic extension of  $Kv$  of uncountable dimension, the same must already be true for the elements  $(c_i f_i(x)/h(x))v$ ,  $i \in \lambda$ , or for the elements  $(c_i g_i(x)/h(x))v$ ,  $i \in \lambda$ . It follows that at least one of these two sets contains uncountably many  $Kv$ -linearly independent elements. But then by Lemma 2.8 (applied with  $I = \{1\}$  and  $\eta_1 = 1$ ), there are uncountably many  $K$ -linearly independent elements in the set

$$\frac{1}{h(x)} K[x]$$

and hence also in  $K[x]$ , a contradiction.  $\square$

Finally, let us mention the following lemma, which combines the algebraic and the transcendental case. We leave its easy proof to the reader.

**Lemma 2.10.** *Let  $(L|K, v)$  be an extension of valued fields. Take  $x \in L$ . Suppose that for some  $e \in \mathbb{N}$  there exists an element  $d \in K$  such that  $vd x^e = 0$  and  $dx^e v$  is transcendental over  $Kv$ . Let  $e$  be minimal with this property. Then for every  $f = c_n x^n + \dots + c_0 \in K[x]$ ,*

$$vf = \min_{1 \leq i \leq n} vc_i x^i.$$

Moreover,  $K(x)v = Kv(dx^e v)$  is a rational function field over  $Kv$ , and we have

$$vK(x) = vK + \mathbb{Z}vx \quad \text{with} \quad (vK(x) : vK) = e.$$

**2.3. Construction of valued field extensions with prescribed value groups and residue fields.** In this section, we will deal with the following problem. Suppose that  $(K, v)$  is a valued field,  $\Gamma|vK$  is an extension of ordered abelian groups and  $k|Kv$  is a field extension. Does there exist an extension  $(L|K, v)$  of valued fields such that  $vL = \Gamma$  and  $Lv = k$ ? We include the case of  $(K, v)$  being trivially valued; this amounts to the construction of a valued field with given value group and residue field. Throughout, we use the well known fact that if  $(K, v)$  is any valued field and  $L$  is any extension field of  $K$ , then there is at least one extension of  $v$  to  $L$  (cf. [E], [R]).

Let us adjust the following notion to our purposes. Usually, when one speaks of an **Artin-Schreier extension** then one means an extension of a field  $K$  generated by a root of an irreducible polynomial of the form  $X^p - X - c$ , provided that  $p = \text{char } K$ . We will replace this by the weaker condition “ $p = \text{char } Kv$ ”. In fact, such extensions also play an important role in the mixed characteristic case, where  $\text{char } K = 0$ .

Every **Artin-Schreier polynomial**  $X^p - X - c$  is separable, since its derivative does not vanish. The following is a simple but very useful observation:

**Lemma 2.11.** *Let  $(K, v)$  be a valued field and  $c \in K$  such that  $vc < 0$ . If  $a \in \tilde{K}$  is such that  $a^p - a = c$ , then for every extension of  $v$  from  $K$  to  $K(a)$ ,*

$$0 > v(a^p - c) = va > pva = vc.$$

*Proof.* Take any extension of  $v$  from  $K$  to  $K(a)$ . Necessarily,  $va < 0$ , since otherwise  $\infty = v(a^p - a - c) = \min\{pva, va, vc\} = vc$ , a contradiction. It follows that  $va^p = pva < va$ , and thus

$$vc = \min\{v(a^p - a - c), v(a^p - a)\} = v(a^p - a) = \min\{pva, va\} = pva.$$

□

**Lemma 2.12.** *Let  $(K, v)$  be a non-trivially valued field,  $p$  a prime and  $\alpha$  an element of the divisible hull of  $vK$  such that  $p\alpha \in vK$ ,  $\alpha \notin vK$ . Choose an element  $a \in \tilde{K}$  such that  $a^p \in K$  and  $va^p = p\alpha$ . Then  $v$  extends in a unique way from  $K$  to  $K(a)$ . It satisfies*

$$(2.5) \quad va = \alpha, \quad [K(a) : K] = (vK(a) : vK), \quad vK(a) = vK + \mathbb{Z}\alpha \quad \text{and} \quad K(a)v = Kv.$$

*If  $\text{char } K = \text{char } Kv = p$ , then this extension  $K(a)|K$  is purely inseparable. On the other hand, if  $\text{char } Kv = p$ , then there is always an Artin-Schreier extension  $K(a)|K$  with properties (2.5); if  $\alpha < 0$ , then  $a$  itself can be chosen to be the root of an Artin-Schreier polynomial over  $K$ .*

*Proof.* Take  $c \in K$  such that  $vc = p\alpha$  and  $a \in \tilde{K}$  such that  $a^p = c$ . Choose any extension of  $v$  from  $K$  to  $K(a)$ . Then  $pva = vc = p\alpha$ . Consequently,  $va = \alpha$  and  $(vK(a) : vK) \geq (vK + \mathbb{Z}\alpha : vK) = p$ . On the other hand, the fundamental inequality (2.1) shows that

$$p = [K(a) : K] \geq (vK(a) : vK) \cdot [K(a)v : Kv] \geq (vK(a) : vK) \geq p.$$

Hence, equality holds everywhere, and we find that

$$(vK(a) : vK) = p \quad \text{and} \quad [K(a)v : Kv] = 1.$$

That is,  $vK(a) = vK + \mathbb{Z}\alpha$  and  $K(a)v = Kv$ . Further, the fundamental inequality implies that the extension of  $v$  from  $K$  to  $K(a)$  is unique.

Now suppose that  $\text{char } Kv = p$ . Choose  $c \in K$  such that  $vc = -|p\alpha| < 0$ . By the foregoing lemma, every root  $b$  of the Artin-Schreier polynomial  $X^p - X - c$  must satisfy  $pvb = vc$ . Now we set  $a = b$  if  $\alpha < 0$ , and  $a = 1/b$  if  $\alpha > 0$  (but note that then  $1/a$  is in general not the root of an Artin-Schreier polynomial). Then as before one shows that (2.5) holds.  $\square$

For  $f \in \mathcal{O}_K[X]$ , we define the **reduction**  $fv \in Kv[X]$  to be the polynomial obtained from  $f$  through replacing every coefficient by its residue.

**Lemma 2.13.** *Let  $(K, v)$  be a valued field and  $\zeta$  an element of the algebraic closure of  $Kv$ . Choose a monic polynomial  $f \in \mathcal{O}_K[X]$  whose reduction  $fv$  is the minimal polynomial of  $\zeta$  over  $Kv$ . Further, choose a root  $b \in \tilde{K}$  of  $f$ . Then there is a unique extension of  $v$  from  $K$  to  $K(b)$  and a corresponding extension of the residue map such that*

$$(2.6) \quad bv = \zeta, \quad [K(b) : K] = [Kv(\zeta) : Kv], \quad vK(b) = vK \quad \text{and} \quad K(b)v = Kv(\zeta).$$

*In all cases,  $f$  can be chosen separable, provided that the valuation  $v$  is non-trivial. On the other hand, if  $\text{char } K = \text{char } Kv = p > 0$  and  $\zeta$  is purely inseparable over  $Kv$ , then  $b$  can be chosen purely inseparable over  $K$ .*

*If  $v$  is non-trivial,  $\text{char } Kv = p > 0$  and  $\zeta^p \in Kv$ ,  $\zeta \notin Kv$ , then there is also an Artin-Schreier extension  $K(b)|K$  such that (2.6) holds and  $db$  is the root of an Artin-Schreier polynomial over  $K$ , for a suitable  $d \in K$ .*

*Proof.* We choose an extension of  $v$  from  $K$  to  $K(b)$ . Since  $f$  is monic with integral coefficients,  $b$  must also be integral for this extension, and  $bv$  must be a root of  $fv$ . We may compose the residue map with an isomorphism in  $\text{Gal } Kv$  which sends this root to  $\zeta$ . Doing so, we obtain a residue map (still associated with  $v$ ) that satisfies  $bv = \zeta$ . Now  $\zeta \in K(b)v$ , and consequently  $[K(b)v : Kv] \geq [Kv(\zeta) : Kv] = \deg fv = \deg f$ . On the other hand, the fundamental inequality shows that

$$\deg f = [K(b) : K] \geq (vK(b) : vK) \cdot [K(b)v : Kv] \geq [K(b)v : Kv] \geq \deg f.$$

Hence, equality holds everywhere, and we find that  $[K(b)v : Kv] = [Kv(\zeta) : Kv] = [K(b) : K]$  and  $(vK(b) : vK) = 1$ . That is,  $vK(b) = vK$  and  $K(b)v = Kv(\zeta)$ . Further, the uniqueness of  $v$  on  $K(a)$  follows from the fundamental inequality.

If  $fv$  is separable, then so is  $f$ . Even if  $fv$  is not separable but  $v$  is non-trivial on  $K$ , then  $f$  can still be chosen separable since we can add a summand  $cX$  with  $c \neq 0$ ,  $vc > 0$  (we use that  $v$  is non-trivial) without changing the reduction of  $f$ . On the other hand, if  $fv$  is purely inseparable and hence of the form  $X^{p^r} - cv$ , then we can choose  $f = X^{p^r} - c$ , which also is purely inseparable if  $\text{char } K = p$ .

Now suppose that  $\text{char } Kv = p > 0$  and  $\zeta^p \in Kv$ ,  $\zeta \notin Kv$ . Choose  $c \in K$  such that  $cv = \zeta^p$ . To construct an Artin-Schreier extension, choose any  $d \in K$  with  $vd < 0$ , and let  $b_0$  be a root of the Artin-Schreier polynomial  $X^p - X - d^p c$ . Since  $vd^p c = pvd < 0$ , Lemma 2.11 shows that  $v(b_0^p - d^p c) = vb_0 > vb_0^p$ . Consequently,  $v((b_0/d)^p - c) > v(b_0/d)^p = vc = 0$ , whence  $(b_0/d)^p v = cv$  and  $(b_0/d)v = (cv)^{1/p} = \zeta$ . We set  $b = b_0/d$ ; so  $K(b) = K(b_0)$ . As before, it follows that  $vK(b) = vK$  and  $K(b)v = Kv(\zeta)$ .  $\square$

**Theorem 2.14.** *Let  $(K, v)$  be an arbitrary valued field. For every extension  $\Gamma|vK$  of ordered abelian groups and every field extension  $k|Kv$ , there is an extension  $(L, v)$  of  $(K, v)$  such that  $vL = \Gamma$  and  $Lv = k$ . If  $\Gamma|vK$  and  $k|Kv$  are algebraic, then  $L|K$  can be chosen to be algebraic, with a unique extension of  $v$  from  $K$  to*

*L.* If  $\rho = \text{rr } \Gamma/vK$  and  $\tau = \text{trdeg } k|Kv$  are finite, then  $L|K$  can be chosen of transcendence degree  $\rho + \tau$ . If  $\Gamma \neq \{0\}$ , then  $L$  can be chosen to be a separable extension of  $K$ .

If both  $\Gamma|vK$  and  $k|Kv$  are finite, then  $L|K$  can be taken to be a finite extension such that  $[L : K] = (\Gamma : vK)[k : Kv]$ . If in addition  $v$  is non-trivial on  $K$ , then  $L|K$  can be chosen to be a simple separable-algebraic extension.

If  $\Gamma|vK$  is countable and  $k|Kv$  is countably generated, then  $L|K$  can be taken to be a countably generated extension.

*Proof.* For the proof, we assume that  $\Gamma \neq \{0\}$  (the other case is trivial). Let  $\alpha_i$ ,  $i \in I$ , be a maximal set of elements in  $\Gamma$  rationally independent over  $vK$ . Then by Lemma 2.6 there is an extension  $(K_1, v) := (K(x_i \mid i \in I), v)$  of  $(K, v)$  such that  $vK_1 = vK \oplus \bigoplus_{i \in I} \mathbb{Z}\alpha_i$  and  $K_1v = Kv$ . Next, choose a transcendence basis  $\zeta_j$ ,  $j \in J$ , of  $k|Kv$ . Then by Lemma 2.6 there is an extension  $(K_2, v) := (K_1(y_j \mid j \in J), v)$  of  $(K_1, v)$  such that  $vK_2 = vK_1$  and  $K_2v = Kv(\zeta_j \mid j \in J)$ . If  $\Gamma|vK$  and  $k|Kv$  are algebraic, then  $I = J = \emptyset$  and  $K_2 = K$ .

If we have an ascending chain of valued fields whose value groups are subgroups of  $\Gamma$  and whose residue fields are subfields of  $k$ , then the union over this chain is again a valued field whose value group is a subgroup of  $\Gamma$  and whose residue field is a subfield of  $k$ . So a standard argument using Zorn's Lemma together with the transitivity of separable extensions shows that there are maximal separable-algebraic extension fields of  $(K_2, v)$  with these properties. Choose one of them and call it  $(L, v)$ . We have to show that  $vL = \Gamma$  and  $Lv = k$ . Since already  $\Gamma|vK_2$  and  $k|K_2v$  are algebraic, the same holds for  $\Gamma|vL$  and  $k|Lv$ . If  $vL$  is a proper subgroup of  $\Gamma$ , then there are some prime  $p$  and some element  $\alpha \in \Gamma \setminus vL$  such that  $p\alpha \in vL$ . But then, Lemma 2.12 shows that there exists a proper separable-algebraic extension  $(L', v)$  of  $(L, v)$  such that  $vL' = vL + \mathbb{Z}\alpha \subset \Gamma$  and  $L'v = Lv \subset k$ , which contradicts the maximality of  $L$ . If  $Lv$  is a proper subfield of  $k$ , then there is some element  $\zeta \in k \setminus Lv$  algebraic over  $Lv$ . But then, Lemma 2.13 shows that there exists a proper separable-algebraic extension  $(L', v)$  of  $(L, v)$  such that  $vL' = vL \subset \Gamma$  and  $L'v = Lv(\zeta) \subset k$ , which again contradicts the maximality of  $L$  (here we have used  $\Gamma \neq \{0\}$ , which implies that  $v$  is not trivial on  $L$ ). This proves that  $vL = \Gamma$  and  $Lv = k$ , and  $(L, v)$  is the required extension of  $(K, v)$ . Since  $K_2$  is generated over  $K$  by a set of elements which are algebraically independent over  $K$ , we know that  $K_2|K$  is separable. Since also  $L|K_2$  is separable, we find that  $L|K$  is separable. Since  $L|K_2$  is algebraic,  $\{x_i, y_j \mid i \in I, j \in J\}$  is a transcendence basis of  $(K_2|K, v)$ . If  $\Gamma|vK$  and  $k|Kv$  are algebraic, then  $I = J = \emptyset$  and  $L$  is an algebraic extension of  $K = K_2$ .

If  $\Gamma|vK$  and  $k|Kv$  are finite, then  $L$  can be constructed by a finite number of applications of Lemma 2.12 and Lemma 2.13. Since extension degree, ramification index and inertia degree are multiplicative, we obtain

$$[L : K] = (vL : vK)[Lv : Kv] = (\Gamma : vK)[k : Kv].$$

If in addition  $v$  is non-trivial, then  $L|K$  can be chosen to be a separable extension. Since it is finite, it is simple.

If  $\Gamma|vK$  and  $k|Kv$  are countably generated algebraic, then they are unions over a countable chain of algebraic extensions. Hence also  $L$  can be constructed as a union over a countable chain of algebraic extensions, and will thus be a countably generated extension of  $K$ .

If  $\Gamma|vK$  and  $k|Kv$  are countably generated, then the sets  $I$  and  $J$  in our above construction are both countable, that is,  $K_2|K$  is countably generated. Moreover, the extensions  $\Gamma|vK_2$  and  $k|K_2v$  are countably generated algebraic. Hence by what we have just shown,  $L$  can be taken to be a countably generated extension of  $K_2$ , and thus also of  $K$ .  $\square$

Every ordered abelian group is an extension of the trivial group  $\{0\}$  as well as of the ordered abelian group  $\mathbb{Z}$ . Every field of characteristic 0 is an extension of  $\mathbb{Q}$ , and every field of characteristic  $p > 0$  is an extension of  $\mathbb{F}_p$ . Let  $\Gamma \neq 0$  be an ordered abelian group and  $k$  a field. If  $\text{char } k = 0$ , then  $\mathbb{Q}$  endowed with the trivial valuation  $v$  will satisfy  $v\mathbb{Q} = \{0\} \subset \Gamma$  and  $\mathbb{Q}v = \mathbb{Q} \subset k$ . If  $\text{char } k = p > 0$ , then we can choose  $v$  to be the  $p$ -adic valuation on  $\mathbb{Q}$  to obtain that  $v\mathbb{Q} = \mathbb{Z} \subset \Gamma$  and  $\mathbb{Q}v = \mathbb{F}_p \subset k$ . But also  $\mathbb{F}_p$  endowed with the trivial valuation  $v$  will satisfy  $v\mathbb{F}_p = \{0\} \subset \Gamma$  and  $\mathbb{F}_pv = \mathbb{F}_p \subset k$ . An application of the foregoing theorem now proves

**Corollary 2.15.** *Let  $\Gamma \neq 0$  be an ordered abelian group and  $k$  a field. Then there is a valued field  $(L, v)$  with  $vL = \Gamma$  and  $Lv = k$ . If  $\text{char } k = p > 0$ , then  $L$  can be chosen to be of characteristic 0 (mixed characteristic case) or of characteristic  $p$  (equal characteristic case).*

For the sake of completeness, we add the following information. From the fundamental inequality it follows that  $v\tilde{K}|vK$ ,  $vK^{\text{sep}}|vK$ ,  $\tilde{K}v|Kv$  and  $K^{\text{sep}}v|Kv$  are algebraic extensions. On the other hand, Lemma 2.12 shows that the value group of a separable-algebraically closed field must be divisible. Similarly, it follows from Lemma 2.13 that the residue field of a separable-algebraically closed non-trivially valued field must be algebraically closed. This proves

**Lemma 2.16.** *Let  $(K, v)$  be a non-trivially valued field and extend  $v$  to  $\tilde{K}$ . Then the value groups  $v\tilde{K}$  and  $vK^{\text{sep}}$  are equal to the divisible hull of  $vK$ , and the residue fields  $\tilde{K}v$  and  $K^{\text{sep}}v$  are equal to the algebraic closure of  $Kv$ .*

A valued field  $(K, v)$  of residue characteristic  $p > 0$  is called **Artin-Schreier closed** if every Artin-Schreier polynomial with coefficients in  $K$  admits a root in  $K$ . Recall that if  $\text{char } K = p$ , then this means that every Artin-Schreier polynomial with coefficients in  $K$  splits into linear factors over  $K$ . If  $K$  is Artin-Schreier closed, then so is  $Kv$ . As a corollary to Lemmas 2.12 and 2.13, we obtain

**Corollary 2.17.** *Every Artin-Schreier closed non-trivially valued field of residue characteristic  $p > 0$  has  $p$ -divisible value group and perfect Artin-Schreier closed residue field.*

**2.4. Orderings and valuations.** We will assume the reader to be familiar with the basic theory of convex valuations, which can be found in [L] and [PR].

**Proposition 2.18.** *Suppose that  $(K, <)$  is an ordered field with convex valuation  $v$ , and denote by  $<_r$  the ordering induced by  $<$  through  $v$  on  $Kv$ . Let  $(L|K, v)$  be an extension of valued fields. If  $2vL \cap vK = 2vK$ , then each extension  $<'_r$  of  $<_r$  to an ordering of  $Lv$  can be lifted through  $v$  to an ordering of  $L$  which extends the ordering  $<$  of  $K$ .*

*Proof.* We fix a section from  $vK/2vK$  to  $K^\times/K^{\times 2}$ . Since  $2vL \cap vK = 2vK$ , this section can be extended to a section from  $vL/2vL$  to  $L^\times/L^{\times 2}$ . Now there is a

bijection between the set of all liftings of  $<_r$  through  $v$  to orderings of  $K$ , and the set of all group characters of  $vK/2vK$ ; see [PR], (7.5) to (7.9). The same construction yields a bijection between the set of all liftings of  $<'_r$  through  $v$  to orderings of  $L$ , and the set of all group characters of  $vL/2vL$ . Since we use an extension of the section  $vK/2vK \rightarrow K^\times/K^{\times 2}$  to a section  $vL/2vL \rightarrow L^\times/L^{\times 2}$ , the bijection maps commute with the restriction from  $L$  to  $K$  of any lifting. That is, if a lifting  $<'$  of  $<'_r$  to  $L$  corresponds to a character  $\chi$  of  $vL/2vL$ , then the restriction of  $<'$  to  $K$  is the unique lifting of  $<_r$  to  $K$  which corresponds to the restriction of  $\chi$  to  $vK/2vK$ .

As the given ordering  $<$  of  $K$  is a lifting of  $<_r$ , it corresponds to a unique group character of  $vK/2vK$ . Since  $2vL \cap vK = 2vK$ , we can extend it to a group character of  $vL/2vL$ . Take the lifting of  $<'_r$  through  $v$  to an ordering  $<'$  of  $L$  which corresponds to this group character of  $vL/2vL$ . Then its restriction to  $K$  is  $<$ .  $\square$

The following was proved by Knebusch and Wright [KW] and by Prestel (cf. [PR]); see Theorem 5.6 of [L].

**Theorem 2.19.** *Let  $v$  be a convex valuation on the ordered field  $(K, <)$ ,  $<_r$  the ordering induced by  $<$  on  $Kv$ , and  $R$  a real closure of  $(K, <)$ . Then there exists a unique extension of  $v$  to a convex valuation of  $R$ . Denoting this extension again by  $v$ , we have that  $(R, v)$  is henselian,  $vR$  is divisible, and  $Rv$  with the ordering induced by  $<$  is a real closure of  $(Kv, <_r)$ .*

**Corollary 2.20.** *Let  $v$  be a convex valuation on the ordered field  $(K, <)$  and  $R$  a real closure of  $(K, <)$ , endowed with the unique convex extension of  $v$ . Further, let  $\Gamma|vK$  be an algebraic extension of ordered abelian groups, and  $k|Kv$  a subextension of some real closure of  $Kv$ . Then there is a (separable-algebraic) subextension  $(L|K, v)$  of  $(R|K, v)$  such that  $vL = \Gamma$  and  $Lv = k$ . If both  $\Gamma|vK$  and  $k|Kv$  are finite, then  $L|K$  can be taken to be a finite simple extension of the form  $K(a)|K$  such that  $[K(a) : K] = (\Gamma : vK)[k : Kv]$ .*

We leave the proof of the corollary as an exercise to the reader. It is a straightforward application of Hensel's Lemma, using the fact that  $(R, v)$  is henselian. One also uses the fact that all real closures of  $Kv$  are isomorphic over  $Kv$ , so by passing to an equivalent residue map (place), one passes from  $Rv$  to the real closure given in the hypothesis.

**2.5. A version of Krasner's Lemma.** Let  $(K, v)$  be any valued field. If  $a \in \tilde{K} \setminus K$  is not purely inseparable over  $K$ , we choose some extension of  $v$  from  $K$  to  $\tilde{K}$  and define

$$\text{kras}(a, K) := \max\{v(\tau a - \sigma a) \mid \sigma, \tau \in \text{Gal } K \text{ and } \tau a \neq \sigma a\} \in v\tilde{K}$$

and call it the **Krasner constant of  $a$  over  $K$** . Since all extensions of  $v$  from  $K$  to  $\tilde{K}$  are conjugate, this does not depend on the choice of the particular extension of  $v$ . For the same reason, over a henselian field  $(K, v)$  our Krasner constant  $\text{kras}(a, K)$  coincides with the Krasner constant

$$\max\{v(a - \sigma a) \mid \sigma \in \text{Gal } K \text{ and } a \neq \sigma a\}$$

as defined by S. K. Khanduja in [KH11], [KH12]. The following is a variant of the well-known Krasner's Lemma (cf. [R]).

**Lemma 2.21.** *Take  $K(a)|K$  to be a separable-algebraic extension, and  $(K(a, b), v)$  to be any valued field extension of  $(K, v)$  such that*

$$(2.7) \quad v(b - a) > \text{kras}(a, K) .$$

*Then for every extension of  $v$  from  $K(a, b)$  to its algebraic closure  $\widetilde{K(a, b)} = \widetilde{K(b)}$ , the element  $a$  lies in the henselization of  $(K(b), v)$  in  $(\widetilde{K(b)}, v)$ .*

*Proof.* Take any extension of  $v$  from  $K(a, b)$  to  $\widetilde{K(b)}$  and denote by  $K(b)^h$  the henselization of  $(K(b), v)$  in  $(\widetilde{K(b)}, v)$ . Since  $a$  is separable-algebraic over  $K$ , it is also separable-algebraic over  $K(b)^h$ . Since for every  $\rho \in \text{Gal } K(b)^h$  we have that  $\rho a = \rho|_{\widetilde{K}} a$  and  $\rho|_{\widetilde{K}} \in \text{Gal } K$ , we find that

$$\begin{aligned} & \{v(a - \rho a) \mid \rho \in \text{Gal } K(b)^h \text{ and } a \neq \rho a\} \\ & \subseteq \{v(a - \sigma a) \mid \sigma \in \text{Gal } K \text{ and } a \neq \sigma a\} \\ & \subseteq \{v(\tau a - \sigma a) \mid \sigma, \tau \in \text{Gal } K \text{ and } \tau a \neq \sigma a\} . \end{aligned}$$

This implies that

$$\text{kras}(a, K(b)^h) \leq \text{kras}(a, K) ,$$

and consequently,  $v(b - a) > \text{kras}(a, K(b)^h)$ . Now  $a \in K(b)^h$  follows from the usual Krasner's Lemma.  $\square$

### 3. VALUATIONS ON $K(x)$

**3.1. A basic classification.** In this section, we wish to classify all extensions of the valuation  $v$  of  $K$  to a valuation of the rational function field  $K(x)$ . As

$$(3.1) \quad 1 = \text{trdeg } K(x)|K \geq \text{rr } vK(x)/vK + \text{trdeg } K(x)v|Kv$$

holds by Lemma 2.6, there are the following mutually exclusive cases:

- $(K(x)|K, v)$  is **valuation-algebraic**:  
 $vK(x)/vK$  is a torsion group and  $K(x)v|Kv$  is algebraic,
- $(K(x)|K, v)$  is **value-transcendental**:  
 $vK(x)/vK$  has rational rank 1, but  $K(x)v|Kv$  is algebraic,
- $(K(x)|K, v)$  is **residue-transcendental**:  
 $K(x)v|Kv$  has transcendence degree 1, but  $vK(x)/vK$  is a torsion group.

We will combine the value-transcendental case and the residue-transcendental case by saying that

- $(K(x)|K, v)$  is **valuation-transcendental**:  
 $vK(x)/vK$  has rational rank 1, or  $K(x)v|Kv$  has transcendence degree 1.

A special case of the valuation-algebraic case is the following:

- $(K(x)|K, v)$  is **immediate**:  
 $vK(x) = vK$  and  $K(x)v = Kv$ .

*Remark 3.1.* It has been observed by several authors that a valuation-algebraic extension of  $v$  from  $K$  to  $K(x)$  can be represented as a limit of an infinite sequence of residue-transcendental extensions. See, e.g., [APZ3], where the authors also derive the assertion of our Theorem 2.9 from this fact. A “higher form” of this approach is found in [S]. The approach is particularly important because residue-transcendental extensions behave better than valuation-algebraic extensions: the corresponding extensions of value group and residue field are finitely generated (Corollary 2.7), and they do not generate a defect: see the Generalized Stability Theorem (Theorem 3.1) and its application in [KKU1].

If  $K$  is algebraically closed, then the residue field  $Kv$  is algebraically closed, and the value group  $vK$  is divisible. So we see that for an extension  $(K(x)|K, v)$  with algebraically closed  $K$ , there are only the following mutually exclusive cases:

$(K(x)|K, v)$  is **immediate**:  $vK(x) = vK$  and  $K(x)v = Kv$ ,

$(K(x)|K, v)$  is **value-transcendental**:  $\text{rr } vK(x)/vK = 1$ , but  $K(x)v = Kv$ ,

$(K(x)|K, v)$  is **residue-transcendental**:  $\text{trdeg } K(x)v|Kv = 1$ , but  $vK(x) = vK$ .

Let us fix an arbitrary extension of  $v$  to  $\tilde{K}$ . Every valuation  $w$  on  $K(x)$  can be extended to a valuation on  $\tilde{K}(x)$ . If  $v$  and  $w$  agree on  $K$ , then this extension can be chosen in such a way that its restriction to  $\tilde{K}$  coincides with  $v$ . Indeed, if  $w'$  is any extension of  $w$  to  $\tilde{K}(x)$  and  $v'$  is its restriction to  $\tilde{K}$ , then there is an automorphism  $\tau$  of  $\tilde{K}|K$  such that  $v'\tau = v$  on  $\tilde{K}$ . We choose  $\sigma$  to be the (unique) automorphism of  $\tilde{K}(x)|K(x)$  whose restriction to  $\tilde{K}$  is  $\tau$  and which satisfies  $\sigma x = x$ . Then  $w'\sigma$  is an extension of  $w$  from  $K(x)$  to  $\tilde{K}(x)$  whose restriction to  $\tilde{K}$  is  $v$ . We conclude:

**Lemma 3.2.** *Take any extension of  $v$  from  $K$  to its algebraic closure  $\tilde{K}$ . Then every extension of  $v$  from  $K$  to  $K(x)$  is the restriction of some extension of  $v$  from  $\tilde{K}$  to  $\tilde{K}(x)$ .*

Now extend  $v$  to  $\widetilde{K(x)}$ . We know that  $\widetilde{vK(x)}/vK(x)$  and  $v\tilde{K}/vK$  are torsion groups, and also  $v\tilde{K}(x)/vK(x) \subset \widetilde{vK(x)}/vK(x)$  is a torsion group. Hence,

$$\text{rr } v\tilde{K}(x)/v\tilde{K} = \text{rr } vK(x)/vK.$$

Since  $v\tilde{K}$  is divisible,  $vK(x)/vK$  is a torsion group if and only if  $v\tilde{K}(x) = v\tilde{K}$ .

Further, the extensions  $\widetilde{K(x)v}|K(x)v$  and  $\tilde{K}v|Kv$  are algebraic, and also the subextension  $\tilde{K}(x)v|K(x)v$  of  $\widetilde{K(x)v}|K(x)v$  is algebraic. Hence,

$$\text{trdeg } \tilde{K}(x)v|\tilde{K}v = \text{trdeg } K(x)v|Kv.$$

Since  $\tilde{K}v$  is algebraically closed,  $K(x)v|Kv$  is algebraic if and only if  $\tilde{K}(x)v = \tilde{K}v$ . We have proved

**Lemma 3.3.**  *$(K(x)|K, v)$  is valuation-algebraic if and only if  $(\tilde{K}(x)|\tilde{K}, v)$  is immediate.  $(K(x)|K, v)$  is valuation-transcendental if and only if  $(\tilde{K}(x)|\tilde{K}, v)$  is not immediate.  $(K(x)|K, v)$  is value-transcendental if and only if  $(\tilde{K}(x)|\tilde{K}, v)$  is value-transcendental.  $(K(x)|K, v)$  is residue-transcendental if and only if  $(\tilde{K}(x)|\tilde{K}, v)$  is residue-transcendental.*

The proof can easily be generalized to show:

**Lemma 3.4.** *Let  $(F|K, v)$  be any valued field extension. Then  $vF|vK$  and  $Fv|Kv$  are algebraic if and only if  $(F.\tilde{K}|\tilde{K}, v)$  is immediate, for some (or any) extension of  $v$  from  $F$  to  $F.\tilde{K}$ .*

**3.2. Pure and weakly pure extensions.** Take  $t \in K(x)$ . If  $vt$  is not a torsion element modulo  $vK$ , then  $t$  will be called a **value-transcendental element**. If  $vt = 0$  and  $tv$  is transcendental over  $Kv$ , then  $t$  will be called a **residue-transcendental element**. An element will be called a **valuation-transcendental element** if it is value-transcendental or residue-transcendental. We will call the extension  $(K(x)|K, v)$  **pure (in  $x$ )** if one of the following cases holds:

- for some  $c, d \in K$ ,  $d \cdot (x - c)$  is valuation-transcendental,
- $x$  is the pseudo limit of some pseudo Cauchy sequence in  $(K, v)$  of transcendental type.



We leave it as an exercise to the reader to prove that  $(K(x)|K, v)$  is pure in  $x$  if and only if it is pure in any other generator of  $K(x)$  over  $K$ ; we will not need this fact in the present paper.

If  $(K(x)|K, v)$  is pure in  $x$ , then it follows from Lemma 2.6 and Lemma 2.4 that  $x$  is transcendental over  $K$ . If  $d \cdot (x - c)$  is value-transcendental, then  $vK(x) = vK \oplus \mathbb{Z}v(x - c)$  and  $K(x)v = Kv$  by Lemma 2.6 (in this case, we may in fact choose  $d = 1$ ). If  $d \cdot (x - c)$  is residue-transcendental, then, again by Lemma 2.6, we have  $vK(x) = vK$ , and  $K(x)v = Kv(d(x - c)v)$  is a rational function field over  $Kv$ . If  $x$  is the pseudo limit of some pseudo Cauchy sequence in  $(K, v)$  of transcendental type, then  $(K(x)|K, v)$  is immediate by Lemma 2.4. This proves

**Lemma 3.5.** *If  $(K(x)|K, v)$  is pure, then  $vK$  is pure in  $vK(x)$  (i.e.,  $vK(x)/vK$  is torsion free), and  $Kv$  is relatively algebraically closed in  $K(x)v$ .*

Here is the “prototype” of pure extensions:

**Lemma 3.6.** *If  $K$  is algebraically closed and  $x \notin K$ , then  $(K(x)|K, v)$  is pure.*

*Proof.* Suppose that the set

$$(3.2) \quad v(x - K) := \{v(x - b) \mid b \in K\}$$

has no maximum. Then there is a pseudo Cauchy sequence in  $(K, v)$  with pseudo limit  $x$ , but without a pseudo limit in  $K$ . Since  $K$  is algebraically closed, Theorem 3 of [KA] shows that this pseudo Cauchy sequence must be of transcendental type. The extension therefore satisfies the third condition for being pure.

Now assume that the set  $v(x - K)$  has a maximum, say,  $v(x - c)$  with  $c \in K$ . If  $v(x - c)$  is a torsion element over  $vK$ , then  $v(x - c) \in vK$ , because  $vK$  is divisible as  $K$  is algebraically closed. It then follows that there is some  $d \in K$  such that  $vd(x - c) = 0$ . If  $d(x - c)v$  were algebraic over  $Kv$ , then it would be in  $Kv$ , since  $K$  and thus also  $Kv$  is algebraically closed. But then, there would be some  $b_0 \in K$  such that  $v(d(x - c) - b_0) > 0$ . Putting  $b := c + d^{-1}b_0$ , we would then obtain that  $v(x - b) = v((x - c) - d^{-1}b_0) > -vd = v(x - c)$ , a contradiction to the maximality of  $v(x - c)$ . So we see that either  $v(x - c)$  is non-torsion over  $vK$ , or there is some  $d \in K$  such that  $vd(x - c) = 0$  and  $d(x - c)v$  is transcendental over  $Kv$ . In both cases, this shows that  $(K(x)|K, v)$  is pure.  $\square$

We will call the extension  $(K(x)|K, v)$  **weakly pure (in  $x$ )** if it is pure in  $x$  or if there are  $c, d \in K$  and  $e \in \mathbb{N}$  such that  $vd(x - c)^e = 0$  and  $d(x - c)^e v$  is transcendental over  $Kv$ .

**Lemma 3.7.** *Assume that the extension  $(K(x)|K, v)$  is weakly pure. If we take any extension of  $v$  to  $\widetilde{K(x)}$  and take  $K^h$  to be the henselization of  $K$  in  $(\widetilde{K(x)}, v)$ , then  $K^h$  is the implicit constant field of this extension:*

$$K^h = \text{IC}(K(x)|K, v).$$

*Proof.* As noted already in the introduction,  $K^h$  is contained in  $\text{IC}(K(x)|K, v)$ . Since  $K(x)^h$  is the fixed field of the decomposition group  $G_x^d := G^d(K(x)^{\text{sep}}|K(x), v)$  in the separable-algebraic closure  $K(x)^{\text{sep}}$  of  $K(x)$ , we know that  $\text{IC}(K(x)|K, v)$  is the fixed field in  $K^{\text{sep}}$  of the subgroup

$$G_{\text{res}} := \{\sigma|_{K^{\text{sep}}} \mid \sigma \in G_x^d\}$$

of  $\text{Gal } K$ . In order to show our assertion, it suffices to show that  $\text{IC}(K(x)|K, v) \subseteq K^h$ , that is, that the decomposition group  $G^d := G^d(K^{\text{sep}}|K, v)$  is contained in

$G_{\text{res}}$ . So we have to show that if  $\tau$  is an automorphism of  $K^{\text{sep}}|K$  such that  $v\tau = v$  on  $K^{\text{sep}}$ , then  $\tau$  can be lifted to an automorphism  $\sigma$  of  $K(x)^{\text{sep}}|K(x)$  such that  $v\sigma = v$  on  $K(x)^{\text{sep}}$ . In fact, it suffices to show that  $\tau$  can be lifted to an automorphism  $\sigma$  of  $K^{\text{sep}}(x)|K(x)$  such that  $v\sigma = v$  on  $K^{\text{sep}}(x)$ . Indeed, then we take any extension  $\sigma'$  of  $\sigma$  from  $K^{\text{sep}}(x)$  to  $K(x)^{\text{sep}}$ . Since the extensions  $v\sigma'$  and  $v$  of  $v$  from  $K^{\text{sep}}(x)$  to  $K(x)^{\text{sep}}$  are conjugate, there is some  $\rho \in \text{Gal}(K(x)^{\text{sep}}|K^{\text{sep}}(x))$  such that  $v\sigma'\rho = v$  on  $K(x)^{\text{sep}}$ . Thus,  $\sigma := \sigma'\rho \in G^d$  is the desired lifting of  $\tau$  to  $K(x)^{\text{sep}}$ .

We take  $\sigma$  on  $K^{\text{sep}}(x)$  to be the unique automorphism which satisfies  $\sigma x = x$  and  $\sigma|_{K^{\text{sep}}} = \tau$ . Using that  $(K(x)|K, v)$  is weakly pure, we have to show that  $v\sigma = v$  on  $K^{\text{sep}}(x)$ . Assume first that for some  $c, d \in K$  and  $e \in \mathbb{N}$ ,  $d(x-c)^e$  is valuation-transcendental. Since  $K(x-c) = K(x)$ , we may assume w.l.o.g. that  $c = 0$ . Every element of  $K^{\text{sep}}(x)$  can be written as a quotient of polynomials in  $x$  with coefficients from  $K^{\text{sep}}$ . For every polynomial  $f(x) = a_n x^n + \dots + a_1 x + a_0 \in K^{\text{sep}}[x]$ ,

$$\begin{aligned} v\sigma f(x) &= v(\sigma(a_n)x^n + \dots + \sigma(a_1)x + \sigma(a_0)) \\ &= \min_i (v\sigma(a_i) + ivx) = \min_i (v\tau(a_i) + ivx) \\ &= \min_i (va_i + ivx) = vf(x), \end{aligned}$$

where the second equality holds by Lemma 2.6 and Lemma 2.10. This shows that  $v\sigma = v$  on  $K^{\text{sep}}(x)$ .

Now assume that  $x$  is the pseudo limit of a pseudo Cauchy sequence in  $(K, v)$  of transcendental type. By Lemma 2.5, this pseudo Cauchy sequence is also of transcendental type over  $(K^{\text{sep}}, v)$ . Observe that  $x$  is still a pseudo limit of this pseudo Cauchy sequence in  $(K^{\text{sep}}(x), v\sigma)$ , because  $v\sigma(x-a) = v(\sigma x - \sigma a) = v(x-a)$  for all  $a \in K$ . But  $v\sigma = v\tau = v$  on  $K^{\text{sep}}$ , and the extension of  $v$  from  $K^{\text{sep}}$  to  $K^{\text{sep}}(x)$  is uniquely determined by the pseudo Cauchy sequence (cf. Theorem 2.3). Consequently,  $v\sigma = v$  on  $K^{\text{sep}}(x)$ .  $\square$

**3.3. Construction of nasty examples.** We are now able to give the

*Proof of Theorem 1.2.* Let  $K$  be any algebraically closed field of characteristic  $p > 0$ . On  $K(x)$ , we take  $v$  to be the  $x$ -adic valuation. We work in the power series field  $K((\frac{1}{p^\infty}\mathbb{Z}))$  of all power series in  $x$  with exponents in  $\frac{1}{p^\infty}\mathbb{Z}$ , the  $p$ -divisible hull of  $\mathbb{Z}$ . We choose  $y$  to be a power series

$$(3.3) \quad y = \sum_{i=1}^{\infty} x^{-p^{-e_i}},$$

where  $e_i$  is any increasing sequence of natural numbers such that  $e_{i+1} \geq e_i + i$  for all  $i$ . We then restrict the canonical ( $x$ -adic) valuation of  $K((\frac{1}{p^\infty}\mathbb{Z}))$  to  $K(x, y)$  and call it again  $v$ . We show first that  $vK(x, y) = \frac{1}{p^\infty}\mathbb{Z}$ . Indeed, taking  $p^{e_j}$ -th powers and using that the characteristic of  $K$  is  $p$ , we find that

$$y^{p^{e_j}} - \sum_{i=1}^j x^{-p^{e_j-e_i}} = \sum_{i=j+1}^{\infty} x^{-p^{e_j-e_i}}.$$

Since  $e_j - e_i \geq 0$  for  $i \leq j$ , the left hand side is an element in  $K(x, y)$ . The right hand side has value

$$-p^{e_j-e_{j+1}}vx;$$

since  $e_j - e_{j+1} \leq -j$ , we see that  $\frac{1}{p^j}vx$  lies in  $vK(x, y)$ . Hence,  $\frac{1}{p^\infty}\mathbb{Z} \subseteq vK(x, y)$ . On the other hand,  $vK(x, y) \subseteq vK((\frac{1}{p^\infty}\mathbb{Z})) = \frac{1}{p^\infty}\mathbb{Z}$ , and therefore  $vK(x, y) = \frac{1}{p^\infty}\mathbb{Z}$ .

By definition,  $y$  is a pseudo limit of the pseudo Cauchy sequence

$$\left( \sum_{i=1}^{\ell} x^{-p^{-e_i}} \right)_{\ell \in \mathbb{N}}$$

in the field  $L = K(x^{1/p^i} \mid i \in \mathbb{N}) \subset K((\frac{1}{p^\infty}\mathbb{Z}))$ . Suppose it were of algebraic type. Then by [KA], Theorem 3, there would exist some algebraic extension  $(L(b)|L, v)$  with  $b$  a pseudo limit of the sequence. But then  $b$  would also be algebraic over  $K(x)$ , and hence the extension  $K(x, b)|K(x)$  would be finite. On the other hand, since  $b$  is a pseudo limit of the above pseudo Cauchy sequence, it can be shown as before that  $vK(x, b) = \frac{1}{p^\infty}\mathbb{Z}$ , and thus  $(vK(x, b) : vK(x)) = \infty$ . This contradiction to the fundamental inequality shows that the sequence must be of transcendental type. Hence by Lemma 3.7,  $L^h$  is relatively algebraically closed in  $L(y)^h$ . Since  $L^h = L.K(x)^h$  is a purely inseparable extension of  $K(x)^h$  and  $K(x, y)^h|K(x)^h$  is separable, this shows that  $K(x)^h$  is relatively algebraically closed in  $K(x, y)^h$ .

Now we set  $\eta_0 := \frac{1}{x}$ , and by induction on  $i$  we choose  $\eta_i \in \widetilde{K(x)}$  such that  $\eta_i^p - \eta_i = \eta_{i-1}$ . Then we have

$$v\eta_i = -\frac{1}{p^i}vx$$

for every  $i$ . Since  $vK(x)^h = vK(x) = \mathbb{Z}vx$ , this shows that  $K(x)^h(\eta_i)|K(x)^h$  has ramification index at least  $p^i$ . On the other hand, it has degree at most  $p^i$ , and therefore it must have degree and ramification index equal to  $p^i$ . Note that, for all  $i \geq 0$ ,  $K(x, \eta_i) = K(x, \eta_1, \dots, \eta_i)$  and every extension  $K(x, \eta_{i+1})|K(x, \eta_i)$  is a Galois extension of degree and ramification index  $p$ . By what we have shown, the chain of these extensions is linearly disjoint from  $K(x)^h|K(x)$ . Since  $K(x)^h$  is relatively algebraically closed in  $K(x, y)^h$  and the extensions are separable, it follows that the chain is also linearly disjoint from  $K(x, y)^h|K(x)$ .

We will now show that all extensions  $K(x, y, \eta_i)|K(x, y)$  are immediate. First, we note that  $K(x, y)v = K$ , since  $K \subset K(x, y) \subset K((\frac{1}{p^\infty}\mathbb{Z}))$  and  $K((\frac{1}{p^\infty}\mathbb{Z}))v = K$ . Since  $K$  is algebraically closed, the inertia degree of the extensions must be 1. Further, as the ramification index of a Galois extension is always a divisor of the extension degree, the ramification index of these extensions must be a power of  $p$ . But the value group of  $K(x, y)$  is  $p$ -divisible, which yields that the ramification index of the extensions is 1. By what we have proved above, they are linearly disjoint from  $K(x, y)^h|K(x, y)$ ; that is, the extension of the valuation is unique. This shows that the defect of each extension  $K(x, y, \eta_i)|K(x, y)$  is equal to its degree  $p^i$ .  $\square$

*Remark 3.8.* Instead of defining  $y$  as in (3.3), we could also use any power series

$$(3.4) \quad y = \sum_{i=1}^{\infty} x^{n_i p^{-e_i}},$$

where  $n_i \in \mathbb{Z}$  are prime to  $p$  and the sequence  $n_i p^{-e_i}$  is strictly increasing. The example in [CP] is of this form. But in this example, the field  $K(x, y)$  is an extension of degree  $p^2$  of a field  $K(u, v)$  such that the extension of the valuation from  $K(u, v)$  to  $K(x, y)$  is unique. Since the value group of  $K(x, y)$  is  $\frac{1}{p^\infty}\mathbb{Z}$ , it must be equal to that of  $K(u, v)$ . Since  $K$  is algebraically closed, both have the same residue field.

Therefore, the extension has defect  $p^2$ . This shows that we can also use subfields instead of field extensions to produce defect extensions, in quite the same way.

A special case of (3.4) is the power series

$$(3.5) \quad y = \sum_{i=1}^{\infty} x^{i-p^{-e_i}} = \sum_{i=1}^{\infty} x^i x^{-p^{-e_i}},$$

which now has a support that is cofinal in  $\frac{1}{p^\infty}\mathbb{Z}$ .

To conclude this section, we use Lemma 3.7 to construct an example about relatively closed subfields in henselian fields. The following fact is well known:

*Suppose that  $K$  is relatively closed in a henselian valued field  $(L, v)$  of residue characteristic 0 and that  $Lv|Kv$  is algebraic. Then  $vL/vK$  is torsion free.*

We show that the assumption “ $Lv|Kv$  is algebraic” is necessary.

**Example 3.9.** On the rational function field  $\mathbb{Q}(x)$ , we take  $v$  to be the  $x$ -adic valuation. We extend  $v$  to the rational function field  $\mathbb{Q}(x, y)$  in such a way that  $vy = 0$  and  $yv$  is transcendental over  $\mathbb{Q}(x)v = \mathbb{Q}$ . So by Lemma 2.6 we have  $v\mathbb{Q}(x, y) = v\mathbb{Q}(x) = \mathbb{Z}vx$  and  $\mathbb{Q}(x, y)v = \mathbb{Q}(yv)$ . We pick any integer  $n > 1$ . Then  $v\mathbb{Q}(x^n) = \mathbb{Z}nvx$  and  $\mathbb{Q}(x^n)v = \mathbb{Q}$ . Further,  $v\mathbb{Q}(x^n, xy) = \mathbb{Z}vx$ , since  $vx = vxy \in v\mathbb{Q}(x^n, xy) \subseteq \mathbb{Z}nvx$ . Also,  $\mathbb{Q}(x^n, xy)v = \mathbb{Q}(y^n v)$  by Lemma 2.10. From Lemma 3.7 we infer that  $\mathbb{Q}(x^n)^h$  is relatively algebraically closed in  $\mathbb{Q}(x^n, xy)^h$ . But

$$v\mathbb{Q}(x^n, xy)^h / v\mathbb{Q}(x^n)^h = v\mathbb{Q}(x^n, xy) / v\mathbb{Q}(x^n) = \mathbb{Z}vx / \mathbb{Z}nvx \simeq \mathbb{Z}/n\mathbb{Z}$$

is a non-trivial torsion group.

**3.4. All valuations on  $K(x)$ .** In this section, we will explicitly define extensions of a given valuation on  $K$  to a valuation on  $K(x)$ . First, we define valuation-transcendental extensions, using the idea of valuation independence. Let  $(K, v)$  be an arbitrary valued field, and  $x$  transcendental over  $K$ . Take  $a \in K$  and an element  $\gamma$  in some ordered abelian group extension of  $vK$ . We define a map  $v_{a,\gamma} : K(x)^\times \rightarrow vK + \mathbb{Z}\gamma$  as follows. Given any  $g(x) \in K[x]$  of degree  $n$ , we can write

$$(3.6) \quad g(x) = \sum_{i=0}^n c_i(x-a)^i.$$

Then we set

$$(3.7) \quad v_{a,\gamma} g(x) := \min_{0 \leq i \leq n} vc_i + i\gamma.$$

We extend  $v_{a,\gamma}$  to  $K(x)$  by setting  $v_{a,\gamma}(g/h) := v_{a,\gamma}g - v_{a,\gamma}h$ .

For example, the valuation  $v_{0,0}$  is called the **Gauß valuation** or the **functional valuation** and is given by

$$v_{0,0}(c_n x^n + \dots + c_1 x + c_0) = \min_{0 \leq i \leq n} vc_i.$$

**Lemma 3.10.**  $v_{a,\gamma}$  is a valuation which extends  $v$  from  $K$  to  $K(x)$ . It satisfies the following conditions:

- 1) If  $\gamma$  is non-torsion over  $vK$ , then  $v_{a,\gamma}K(x) = vK \oplus \mathbb{Z}\gamma$  and  $K(x)v_{a,\gamma} = Kv$ .
- 2) If  $\gamma$  is torsion over  $vK$ ,  $e$  is the smallest positive integer such that  $e\gamma \in vK$  and  $d \in K$  is some element such that  $vd = -e\gamma$ , then  $d(x-a)^e v_{a,\gamma}$  is transcendental over  $Kv$ ,  $K(x)v_{a,\gamma} = Kv(d(x-a)^e v_{a,\gamma})$  and  $v_{a,\gamma}K(x) = vK + \mathbb{Z}\gamma$ . In particular, if  $\gamma = 0$  then  $(x-a)v_{a,\gamma}$  is transcendental over  $Kv$ ,  $K(x)v_{a,\gamma} = Kv((x-a)v_{a,\gamma})$  and  $v_{a,\gamma}K(x) = vK$ .

*Proof.* It is a straightforward exercise to prove that  $v_{a,\gamma}$  is a valuation and that 1) and 2) hold. However, one can also deduce this from Lemma 2.6. It says that if we assign a non-torsion value  $\gamma$  to  $x - a$ , then we obtain a unique valuation which satisfies (3.7). Since this defines a unique map  $v_{a,\gamma}$  on  $K[x]$ , we see that  $v_{a,\gamma}$  must coincide with the valuation given by Lemma 2.6, which in turn satisfies assertion 1). Similarly, if  $\gamma \in vK$ ,  $d \in K$  with  $vd = -\gamma$  and we assign a transcendental residue to  $d(x - a)$ , then Lemma 2.6 gives us a valuation on  $K(x)$  which satisfies (3.7) and hence must coincide with  $v_{a,\gamma}$ . This shows that  $v_{a,\gamma}$  is a valuation and satisfies 2).

If  $e > 1$ , then we can first use Lemma 2.6 to see that  $v_{a,\gamma}$  is a valuation on the subfield  $K(d(x - a)^e)$  of  $K(x)$  and that

$$v_{a,\gamma}K(d(x - a)^e) = vK$$

and

$$K(d(x - a)^e)v_{a,\gamma} = Kv(d(x - a)^ev_{a,\gamma})$$

with  $d(x - a)^ev_{a,\gamma}$  transcendental over  $Kv$ . We know that there is an extension  $w$  of  $v_{a,\gamma}$  to  $K(x)$ . It must satisfy  $w(x - a) = -vd/e = \gamma$ . So  $0, w(x - a), w(x - a)^2, \dots, w(x - a)^{e-1}$  lie in distinct cosets modulo  $vK$ . From Lemma 2.8 it follows that  $w$  satisfies (3.7) on  $K(x)$ ; hence it must coincide with the valuation  $v_{a,\gamma}$  on  $K(x)$ . Assertion 2) for this case follows from Lemma 2.8.  $\square$

Now we are able to prove

**Theorem 3.11.** *Take any valued field  $(K, v)$ . Then all extensions of  $v$  to the rational function field  $\tilde{K}(x)$  are of the form*

- $\tilde{v}_{a,\gamma}$ , where  $a \in \tilde{K}$  and  $\gamma$  is an element of some ordered group extension of  $vK$ , or
- $\tilde{v}_{\mathbf{A}}$ , where  $\mathbf{A}$  is a pseudo Cauchy sequence in  $(\tilde{K}, \tilde{v})$  of transcendental type; here  $\tilde{v}$  runs through all extensions of  $v$  to  $\tilde{K}$ . The extension is of the form  $\tilde{v}_{a,\gamma}$  with  $\gamma \notin \tilde{v}\tilde{K}$  if and only if it is value-transcendental, and with  $\gamma \in \tilde{v}\tilde{K}$  if and only if it is residue-transcendental. The extension is of the form  $\tilde{v}_{\mathbf{A}}$  if and only if it is valuation-algebraic.

All extensions of  $v$  to  $K(x)$  are obtained by restricting the above extensions, already from just one fixed extension  $\tilde{v}$  of  $v$  to  $\tilde{K}$ .

*Proof.* By Lemma 3.10 and Theorem 2.3,  $\tilde{v}_{a,\gamma}$  and  $\tilde{v}_{\mathbf{A}}$  are extensions of  $\tilde{v}$  to  $\tilde{K}(x)$ . For the converse, let  $w$  be any extension of  $v$  to  $\tilde{K}(x)$  and set  $\tilde{w} = w|_{\tilde{K}}$ . From Lemma 3.6 we know that  $(\tilde{K}(x)|\tilde{K}, w)$  is always pure. Hence, either  $d(x - c)$  is valuation-transcendental for some  $c, d \in \tilde{K}$ , or  $x$  is the pseudo limit of some pseudo Cauchy sequence  $\mathbf{A}$  in  $(\tilde{K}, \tilde{v})$  of transcendental type. In the first case, Lemma 2.6 shows that

$$w \sum_{i=0}^n d_i(d(x - c))^i = \min_{0 \leq i \leq n} vd_i + iwd(x - c) = \min_{0 \leq i \leq n} vd_id^i + iw(x - c)$$

for all  $d_i \in \tilde{K}$ . This shows that  $w = \tilde{v}_{c,\gamma}$  for  $\gamma = w(x - c)$ . If this value is not in  $\tilde{v}\tilde{K}$ , then it is non-torsion over  $\tilde{v}\tilde{K}$ , and thus the extension of  $\tilde{v}$  to  $\tilde{K}(x)$ , and hence also the extension of  $v$  to  $K(x)$ , is value-transcendental. If it is in  $\tilde{v}\tilde{K}$ , then the residue of  $d(x - c)$  is not in  $\tilde{K}\tilde{v}$ , and the extension of  $\tilde{v}$  to  $\tilde{K}(x)$ , and hence also the extension of  $v$  to  $K(x)$ , is residue-transcendental.

In the second case, we know from Theorem 2.3 that  $\mathbf{A}$  induces an extension  $\tilde{v}_{\mathbf{A}}$  of  $\tilde{v}$  to  $\tilde{K}(x)$  such that  $x$  is a pseudo limit of  $\mathbf{A}$  in  $(\tilde{K}(x), \tilde{v}_{\mathbf{A}})$ . Since  $x$  is also a pseudo limit of  $\mathbf{A}$  in  $(\tilde{K}(x), w)$ , we can infer from Lemma 2.4 that  $w = \tilde{v}_{\mathbf{A}}$ . It also follows from Theorem 2.3 that  $(\tilde{K}(x)|\tilde{K}, \tilde{v}_{\mathbf{A}})$  is immediate, and consequently  $(\tilde{K}(x)|K, \tilde{v}_{\mathbf{A}})$  is valuation-algebraic.

For the last assertion, we invoke Lemma 3.2. Now it just remains to show that it suffices to take the restrictions of the valuations  $\tilde{v}_{a,\gamma}$  and  $\tilde{v}_{\mathbf{A}}$  for one fixed  $\tilde{v}$ . Suppose that  $\tilde{w}$  is another extension of  $v$  to  $\tilde{K}$ . Since all such extensions are conjugate, there is  $\sigma \in \text{Gal } K$  such that  $\tilde{w} = \tilde{v} \circ \sigma$ . Let  $g(x) \in K[x]$  be given as in (3.6). Extend  $\sigma$  to an automorphism of  $\tilde{K}(x)$  which satisfies  $\sigma x = x$ . Since  $g$  has coefficients in  $K$ , we then have

$$g(x) = \sigma g(x) = \sum_i \sigma c_i (x - \sigma a)^i,$$

and therefore

$$\tilde{w}_{a,\gamma} g(x) = \min_i (\tilde{w} c_i + i\gamma) = \min_i (\tilde{v} \sigma c_i + i\gamma) = \tilde{v}_{\sigma a, \gamma} g(x).$$

This shows that  $\tilde{w}_{a,\gamma} = \tilde{v}_{\sigma a, \gamma}$  on  $K(x)$ .

Given a pseudo Cauchy sequence  $\mathbf{A}$  in  $(\tilde{K}, \tilde{w})$ , we set  $\mathbf{A}_{\sigma} = (\sigma a_{\nu})_{\nu < \lambda}$ . This is a pseudo Cauchy sequence in  $(\tilde{K}, \tilde{v})$ , since  $\tilde{v}(\sigma a_{\mu} - \sigma a_{\nu}) = \tilde{v}\sigma(a_{\mu} - a_{\nu}) = \tilde{w}(a_{\mu} - a_{\nu})$ . For every polynomial  $f(x) \in \tilde{K}[x]$ , we have  $\tilde{v}f(\sigma a_{\nu}) = \tilde{w}\sigma^{-1}(f(\sigma a_{\nu})) = \tilde{w}(\sigma^{-1}(f))(a_{\nu})$ , where  $\sigma^{-1}(f)$  denotes the polynomial obtained from  $f(x)$  by applying  $\sigma^{-1}$  to the coefficients. So we see that  $\mathbf{A}_{\sigma}$  is of transcendental type if and only if  $\mathbf{A}$  is. If  $g(x) \in K[x]$ , then  $\sigma^{-1}(g) = g$ , and the above computation shows that  $\tilde{v}g(\sigma a_{\nu}) = \tilde{w}g(a_{\nu})$ . This implies that  $\tilde{w}_{\mathbf{A}} = \tilde{v}_{\mathbf{A}_{\sigma}}$  on  $K(x)$ .  $\square$

*Remark 3.12.* If  $v$  is trivial on  $K$ , hence  $Kv = K$  (modulo an isomorphism), and if we choose  $\gamma > 0$ , then the restriction  $w$  of  $\tilde{v}_{a,\gamma}$  to  $K(x)$  will satisfy  $xw = aw = a$ . It follows that  $K(x)w = K(a)$ . Further,  $wK(x) \subseteq \mathbb{Z}\gamma$ , and thus  $wK(x) \simeq \mathbb{Z}$ .

**3.5. Prescribed implicit constant fields.** If not stated otherwise, we will always assume that  $(K, v)$  is any valued field. The following is an immediate consequence of our version of Krasner's Lemma:

**Lemma 3.13.** *Assume that  $K(a)|K$  is a separable-algebraic extension. Assume further that  $K(x)|K$  is a rational function field and  $v$  is a valuation on  $\widehat{K(x)}$  such that*

$$(3.8) \quad v(x - a) > \text{kras}(a, K).$$

*Then  $K(a) \subseteq (K(x)|K, v)$ , and consequently,*

$$vK(a) \subseteq vK(x) \quad \text{and} \quad K(a)v \subseteq K(x)v.$$

*Proof.* By Lemma 2.21, condition (3.8) implies that  $K(a) \subseteq K(x)^h$ , the henselization being chosen in  $(\widehat{K(x)}, v)$ . Consequently,  $K(a) \subseteq \text{IC}(K(x)|K, v)$ ,  $vK(a) \subseteq vK(x)^h = vK(x)$  and  $K(a)v \subseteq K(x)^h v = K(x)v$ .  $\square$

**Proposition 3.14.** *Assume that  $(K(a)|K, v)$  is a separable-algebraic extension of valued fields. Further, take  $\Gamma$  to be the abelian group  $vK(a) \oplus \mathbb{Z}$  endowed with any extension of the ordering of  $vK(a)$ , and take  $k$  to be the rational function field in*

one variable over  $K(a)v$ . Then there exists an extension  $v_1$  of  $v$  from  $K(a)$  to  $\widetilde{K(x)}$  such that

$$(3.9) \quad v_1 K(x) = \Gamma \text{ and } K(x)v_1 = K(a)v \text{ and } K(a)^h = \text{IC}(K(x)|K, v_1).$$

If  $v$  is non-trivial on  $K$ , then there exists an extension  $v_2$  of  $v$  from  $K(a)$  to  $\widetilde{K(x)}$  such that

$$(3.10) \quad v_2 K(x) = vK(a) \text{ and } K(x)v_2 = k \text{ and } K(a)^h = \text{IC}(K(x)|K, v_2).$$

If  $(K(a), v)$  admits a transcendental immediate extension, then there is also an extension  $v_3$  such that

$$(3.11) \quad v_3 K(x) = vK(a) \text{ and } K(x)v_3 = K(a)v.$$

If in addition  $(K(a), v)$  admits a pseudo Cauchy sequence of transcendental type, then  $v_3$  can be chosen such that  $K(a)^h = \text{IC}(K(x)|K, v_3)$ .

Suppose that  $(K(a), <)$  is an ordered field and that  $v$  is convex. Denote by  $<_r$  the ordering induced by  $<$  on the residue field  $K(a)v$ . Then  $<_r$  can be lifted through  $v_1$  and through  $v_3$  to  $K(x)$  in such a way that the lifted orderings extend  $<$  (from  $K$ ). If  $<'_r$  is an extension of  $<_r$  to  $k$ , then  $<'_r$  can be lifted through  $v_2$  to  $K(x)$  in such a way that the lifted ordering extends  $<$  (from  $K$ ).

*Proof.* If  $K(a) = K$ , then we do the following. We take a generator  $\gamma$  of  $\Gamma$  over  $vK$  and let  $v_1$  be any extension of  $v_{0,\gamma}$  to  $\widetilde{K(x)}$ . Further, we take  $v_2$  to be any extension of  $v_{0,0}$  to  $\widetilde{K(x)}$ . Then the desired properties of  $v_1$  and  $v_2$  follow from Lemma 3.10 and Lemma 3.7. To construct  $v_3$ , we send  $x$  to some transcendental element in the given immediate extension of  $(K, v)$ . The embedding so obtained induces an extension  $v_3$  of  $v$  to  $K(x)$  such that  $(K(x)|K, v_3)$  is immediate. We extend  $v_3$  further to  $\widetilde{K(x)}$ . If  $(K, v)$  admits a pseudo Cauchy sequence of transcendental type, then we can use Theorem 2.3 to construct an immediate extension  $v_3$  of  $v$  to  $K(x)$  such that  $x$  is a pseudo limit of this pseudo Cauchy sequence. Then by Lemma 3.7,  $K^h = \text{IC}(K(x)|K, v_3)$ .

Now assume that  $a \notin K$ . If  $v'$  is any extension of  $v$  to  $\widetilde{K(x)}$  such that  $v'(x-a) > \text{kras}(a, K)$ , then Lemma 3.13 shows that  $a \in K(x)^h$ , where the henselization is taken in  $(\widetilde{K(x)}, v')$ . Thus,

$$(3.12) \quad K(x)^h = K(a, x)^h,$$

and consequently,

$$(3.13) \quad \begin{aligned} v'K(x) &= v'K(x)^h = v'K(a, x)^h = v'K(a, x), \\ K(x)v' &= K(x)^h v' = K(a, x)^h v' = K(a, x)v'. \end{aligned}$$

If in addition we know that

$$(3.14) \quad K(a)^h = \text{IC}(K(a, x)|K(a), v'),$$

then

$$K(a)^h \subseteq \text{IC}(K(x)|K, v') \subseteq \text{IC}(K(a, x)|K(a), v') = K(a)^h,$$

which yields that  $K(a)^h = \text{IC}(K(x)|K, v')$ .

As  $vK$  is cofinal in its divisible hull  $v\widetilde{K}$ , we can choose some  $\alpha \in vK$  such that  $\alpha \geq \text{kras}(a, K)$ .

To construct  $v_1$ , we choose any positive generator  $\beta$  of  $\Gamma$  over  $vK(a)$ . Then also  $\gamma := \alpha + \beta$  is a generator of  $\Gamma$  over  $vK(a)$ . Now we take  $v_1$  to be any extension of

$v_{a,\gamma}$  to  $\widetilde{K(x)}$ . From Lemma 3.10 we know that  $v_{a,\gamma}K(a, x) = vK(a) \oplus \mathbb{Z}\gamma = \Gamma$  and  $K(a, x)v_{a,\gamma} = K(a)v$ . Since

$$(3.15) \quad v_{a,\gamma}(x - a) = \gamma > \text{kras}(a, K),$$

equations (3.13) hold for  $v' = v_{a,\gamma}$ , and we obtain

$$\begin{aligned} v_1K(x) &= v_{a,\gamma}K(a, x) = vK(a) + \mathbb{Z}\gamma = \Gamma, \\ K(x)v_1 &= K(a, x)v_{a,\gamma} = K(a)v. \end{aligned}$$

From Lemma 3.7 we infer that (3.14) holds for  $v' = v_{a,\gamma}$ . Consequently,  $K(a)^h = \text{IC}(K(x)|K, v_1)$ .

To construct  $v_2$ , we make use of our assumption that  $v$  is non-trivial on  $K$  and choose some positive  $\beta \in vK$ . We set  $\gamma := \alpha + \beta$  and take  $v_2$  to be any extension of  $v_{a,\gamma}$  to  $\widetilde{K(x)}$ . From Lemma 3.10 we know that  $v_{a,\gamma}K(a, x) = vK(a) = \Gamma$  and  $K(a, x)v_{a,\gamma}$  is a rational function field in one variable over  $K(a)v$ ; in this sense,  $K(a, x)v_{a,\gamma} = k$ . Again we have (3.15), and equations (3.13) hold for  $v' = v_{a,\gamma}$ ; so we obtain

$$\begin{aligned} v_2K(x) &= v_{a,\gamma}K(a, x) = vK(a), \\ K(x)v_2 &= K(a, x)v_{a,\gamma} = k. \end{aligned}$$

From Lemma 3.7 we infer that (3.14) holds for  $v' = v_{a,\gamma}$ . Consequently,  $K(a)^h = \text{IC}(K(x)|K, v_2)$ .

To construct  $v_3$ , we take  $(L|K(a), v)$  to be the transcendental immediate extension which exists by hypothesis. We choose some  $y \in L$ , transcendental over  $K(a)$  and such that  $\gamma := vy > \text{kras}(a, K)$ . Then  $(K(a, y)|K(a), v)$  is immediate. Now the isomorphism  $K(a, y) \simeq K(a, x)$  induced by  $y \mapsto x - a$  induces on  $K(a, x)$  a valuation  $v'$  such that  $(K(a, x)|\widetilde{K(a)}, v')$  is immediate and  $v'(x - a) = \gamma$ . We take  $v_3$  to be any extension of  $v'$  to  $\widetilde{K(x)}$ . Again we have (3.13), and we obtain

$$\begin{aligned} v_3K(x) &= v'K(a, x) = vK(a), \\ K(x)v_3 &= K(a, x)v' = K(a)v. \end{aligned}$$

If  $(K(a), v)$  admits a pseudo Cauchy sequence of transcendental type, then we can use Theorem 2.3 to construct an immediate extension  $(K(a, z)|K(a), v)$ . Multiplying  $z$  with a suitable element from  $K$  will give us  $y$  such that  $vy > \text{kras}(a, K)$ . By our above construction, also  $x$  will be a pseudo limit of a pseudo Cauchy sequence of transcendental type. Then by Lemma 3.7,  $K(a)^h = \text{IC}(K(x)|K, v_3)$ .

Finally, suppose that  $(K(a), <)$  is an ordered field and that  $<_r$  is the ordering induced by  $<$  on  $K(a)v$ . We have that  $vK(a) = v_2K(a, x) = v_3K(a, x)$ , so it is trivially true that  $2v_2K(a, x) \cap vK(a) = 2vK(a) = 2v_3K(a, x) \cap vK(a)$ . Further,  $v_1K(a, x) = vK(a) \oplus \mathbb{Z}\gamma$ ; hence also in this case  $2v_1K(a, x) \cap vK(a) = 2vK(a)$ . Therefore, Proposition 2.18 shows that the given orderings on  $K(a)v$  and  $k$  can be lifted through  $v_1$ ,  $v_2$  and  $v_3$  respectively, to orderings on  $K(a, x)$  which extend the ordering  $<$  of  $K(a)$ .  $\square$

This proposition proves Theorem 1.3 in the case where  $K_1^h|K^h$  is finite, since then there is some  $a \in K_1$  such that  $K_1^h = K(a)^h$ . Further, we obtain

**Proposition 3.15.** *Take any finite ordered abelian group extension  $\Gamma_0$  of  $vK$  and any finite field extension  $k_0$  of  $Kv$ . Further, take  $\Gamma$  to be the abelian group  $\Gamma_0 \oplus \mathbb{Z}$  endowed with any extension of the ordering of  $\Gamma_0$ , and take  $k$  to be the rational*



function field in one variable over  $k_0$ . If  $v$  is trivial on  $K$ , then assume in addition that  $k_0|Kv$  is separable. Then there exists an extension  $v_1$  of  $v$  from  $K$  to  $K(x)$  such that

$$(3.16) \quad v_1 K(x) = \Gamma \quad \text{and} \quad K(x)v_1 = k_0.$$

If  $v$  is non-trivial on  $K$ , then there exists an extension  $v_2$  of  $v$  from  $K$  to  $K(x)$  such that

$$(3.17) \quad v_2 K(x) = \Gamma_0 \quad \text{and} \quad K(x)v_2 = k.$$

If  $(K, v)$  admits a transcendental immediate extension, then there is also an extension  $v_3$  such that

$$(3.18) \quad v_3 K(x) = \Gamma_0 \quad \text{and} \quad K(x)v_3 = k_0.$$

Suppose that  $(K, <)$  is an ordered field and that  $v$  is convex. Suppose further that  $k_0$  and  $k$  are equipped with extensions of the ordering induced by  $<$  on  $Kv$ . Then these extensions can be lifted through  $v_1, v_2, v_3$  to  $K(x)$  in such a way that the lifted orderings extend  $<$ .

*Proof.* We choose any finite separable extension  $(K(a)|K, v)$  such that  $vK(a) = \Gamma_0$ ,  $K(a)v = k_0$  and  $[K(a) : K] = (vK(a) : vK)[K(a)v : Kv]$ . If  $v$  is non-trivial on  $K$ , then such an extension exists by Theorem 2.14; otherwise,  $K(a)$  is just equal to  $k_0$ , up to the isomorphism induced by the residue map of the trivial valuation  $v$ . If  $(K, v)$  admits a transcendental immediate extension, then by Lemma 2.2, also  $(K(a), v)$  admits a transcendental immediate extension. Now the first part of our proposition follows from Proposition 3.14.

Suppose that  $(K, <)$  is an ordered field with  $v$  convex. Then by Corollary 2.20 we can choose the field  $K(a)$  to be a subfield of a real closure  $(R, <)$  of  $K$ , equipped with a convex extension of  $v$ , in such a way that the given ordering on  $k_0$  is induced by  $<$  through this extension of  $v$ . Now again, our assertion for the ordered case follows from Proposition 3.14.  $\square$

We turn to the realization of countably infinite separable-algebraic extensions as implicit constant fields.

**Proposition 3.16.** *Let  $(K_1|K, v)$  be a countably infinite separable-algebraic extension of non-trivially valued henselian fields. Then  $(K_1, v)$  admits a pseudo Cauchy sequence of transcendental type. In particular, there is an extension  $v_3$  of  $v$  to  $\widetilde{K(x)}$  such that  $(K_1(x)|K_1, v)$  is immediate, with  $x$  being the pseudo limit of this pseudo Cauchy sequence. Moreover,  $K_1 = \text{IC}(K(x)|K, v_3)$ .*

*Proof.*  $K_1|K$  is a countably infinite union of finite subextensions. Thus, we can choose a sequence  $a_i$ ,  $i \in \mathbb{N}$ , such that  $K(a_i)|K$  is separable-algebraic and  $K(a_i) \subsetneq K(a_{i+1})$  for all  $i$ , and such that  $K_1 = \bigcup_{i \in \mathbb{N}} K(a_i)$ . Through multiplication with elements from  $K$  it is possible to choose each  $a_i$  in such a way that

$$(3.19) \quad va_{i+1} > \max\{va_i, \text{kras}(a_i, K(a_1, \dots, a_{i-1}))\}.$$

We set

$$b_i := \sum_{j=1}^i a_j.$$

By construction,  $(b_i)_{i \in \mathbb{N}}$  is a pseudo Cauchy sequence. Suppose that  $x$  is a pseudo limit of it, for some extension of  $v$  from  $K_1$  to  $K_1(x)$ . Then by induction on  $i$ , we

show that  $a_i \in K(x)^h$ , where  $K(x)^h$  is chosen in some henselization of  $(K_1(x), v)$ . First, by Lemma 2.21,  $v(x - a_1) = v(x - b_1) = va_2 > \text{kras}(a_1, K)$  implies that  $a_1 \in K(x)^h$ . If we have already shown that  $a_1, \dots, a_i \in K(x)^h$ , then  $b_i \in K(x)^h$ , and  $v(x - b_i - a_{i+1}) = v(x - b_{i+1}) = va_{i+2} > \text{kras}(a_{i+1}, K(a_1, \dots, a_i))$  implies that  $a_{i+1} \in K(a_1, \dots, a_i)(x - b_i)^h = K(a_1, \dots, a_i)(x)^h = K(x)^h$ .

This proves that  $K_1 \subseteq K(x)^h$ . Since  $K_1|K$  is infinite, this also proves that  $x$  must be transcendental. As a pseudo Cauchy sequence of algebraic type would admit an algebraic pseudo limit ([KA], Theorem 3), this yields that  $(b_i)_{i \in \mathbb{N}}$  is a pseudo Cauchy sequence of transcendental type in  $(K_1, v)$ .

By Theorem 2.3 we can now extend  $v$  to  $K_1(x)$  so that  $(K_1(x)|K_1, v)$  is immediate and  $x$  is a pseudo limit of  $(b_i)_{i \in \mathbb{N}}$ . We choose any extension  $v_3$  of  $v$  from  $K_1(x)$  to  $\widetilde{K(x)}$ . By Lemma 3.7 we know that  $K_1 = \text{IC}(K_1(x)|K_1, v_3)$ . Hence,

$$K_1 \subseteq K(x)^h \subseteq \text{IC}(K(x)|K, v_3) \subseteq \text{IC}(K_1(x)|K_1, v_3) = K_1,$$

which shows that  $K_1 = \text{IC}(K(x)|K, v_3)$ .  $\square$

Since  $\text{IC}(K(x)|K, v) = \text{IC}(K^h(x)|K^h, v)$ , this proposition proves Theorem 1.3 in the case where  $K_1^h|K^h$  is countably infinite.

**Proposition 3.17.** *Take any non-trivially valued field  $(K, v)$ , any countably generated ordered abelian group extension  $\Gamma_0$  of  $vK$  such that  $\Gamma_0/vK$  is a torsion group, and any countably generated algebraic field extension  $k_0$  of  $Kv$ . Assume that  $\Gamma_0/vK$  or  $k_0|Kv$  is infinite. Then there exists an extension  $v_3$  of  $v$  from  $K$  to  $K(x)$  such that equations (3.18) hold.*

*Suppose that  $(K, <)$  is an ordered field and that  $v$  is convex. If  $k_0$  is equipped with an extension of the ordering induced by  $<$  on  $Kv$ , then this can be lifted through  $v_3$  to an ordering of  $K(x)$  which extends  $<$ .*

*Proof.* We fix an extension of  $v$  to  $\tilde{K}$ . By Theorem 2.14 there is a countably generated separable-algebraic extension  $K_1|K^h$  such that  $vK_1 = \Gamma_0$  and  $K_1v = k_0$ . Since at least one of the extensions  $\Gamma_0|vK$  and  $k_0|Kv$  is infinite,  $K_1^h|K^h$  is infinite too, taking the henselizations in  $(\tilde{K}, v)$ . Hence by Proposition 3.16 there is an extension  $v_3$  of  $v$  to  $K_1(x)$  such that  $K_1 \subset K(x)^h$  and  $(K_1(x)|K_1, v_3)$  is immediate. Consequently,  $K_1(x)^h = K(x)^h$ , which gives us

$$v_3K(x) = v_3K(x)^h = v_3K_1(x) = vK_1 = \Gamma_0$$

and

$$K(x)v_3 = K(x)^hv_3 = K_1(x)v_3 = K_1v = k_0.$$

Suppose that  $(K, <)$  is ordered,  $v$  is convex and  $k_0$  is equipped with an extension of the ordering induced by  $<$  on  $Kv$ . Take any real closure  $(R, <)$  of  $(K, <)$  with an extension of  $v$  to a convex valuation of  $R$ . By Corollary 2.20 we can choose the extension  $K_1|K$  as a subextension of  $R|K$ , with the ordering on  $k_0$  induced by  $<$  through  $v$ . Since  $(K_1(x)|K_1, v_3)$  is immediate, Proposition 2.18 shows that there is a lifting of the ordering of  $k_0$  through  $v$  to an ordering of  $K_1(x)$  which extends the ordering  $<$  of  $K_1$ . Its restriction to  $K(x)$  is an extension of the ordering  $<$  of  $K$ .  $\square$

To conclude this section, we now give an alternative

*Proof of Theorem 1.2.* Let  $K$  be an algebraically closed field of characteristic  $p > 0$ . On  $K(x)$ , we again take  $v$  to be the  $x$ -adic valuation. We assume that  $K$  contains

an element  $t$  which is transcendental over the prime field of  $K$ . Then it can be proved that  $K(x)^h$  admits two infinite linearly disjoint towers of Galois extensions of degree  $p$  and ramification index  $p$ . They can be defined as follows. For the first tower, we set  $\eta_0 = x^{-1}$ , take  $\eta_{i+1}$  to be a root of  $X^p - X - \eta_i$ , and set  $L_i := K(\eta_i)$ . For the second tower, we set  $\vartheta_0 = tx^{-1}$ , take  $\vartheta_{i+1}$  to be a root of  $X^p - X - t\vartheta_i$ , and set  $N_i := K(\vartheta_i)$ . Note that  $L_0 = N_0 = K(x)$ ,  $\eta_i \in L_{i+1}$  and  $\vartheta_i \in N_{i+1}$ . For each  $i \geq 0$ , we have that  $v\eta_i = v\vartheta_i = -\frac{1}{p^i}vx$ , and the extensions  $L_{i+1}|L_i$  and  $N_{i+1}|N_i$  are Galois of degree  $p$  with ramification index  $p$ . Consequently, the same is true for the extensions  $L_{i+1}^h|L_i^h$  and  $N_{i+1}^h|N_i^h$  (note that  $L_i^h = L_i.K(x)^h$  and  $N_i^h = N_i.K(x)^h$ ).

We set  $L := \bigcup_{i \in \mathbb{N}} L_i$  and  $N := \bigcup_{i \in \mathbb{N}} N_i$ . By the above,  $L$  and  $N$  are linearly disjoint from  $K(x)^h$  over  $K(x)$ . Thus,  $L^h = L.K(x)^h$  and  $N^h = N.K(x)^h$  are countably infinite separable-algebraic extensions of  $K(x)^h$ , and it can be proved that they are linearly disjoint over  $K(x)^h$ . We use Proposition 3.16 (with  $K$  replaced by  $K(x)^h$  and  $x$  replaced by  $y$ ) to obtain an extension of  $v$  from  $K(x)^h$  to  $K(x)^h(y)$  such that  $vK(x)^h(y) = vL^h = \frac{1}{p^\infty}\mathbb{Z}$  and  $K(x)^h(y)v = L^hv = K(x)^hv = K$ , and such that the extension  $(K(x)^h(y)|K(x)^h, v)$  has implicit constant field  $L^h$  (i.e.,  $L^h$  is relatively algebraically closed in  $K(x, y)^h = (K(x)^h(y))^h$ ). Since  $K(x)$  is relatively algebraically closed in  $K(x, y)$ , we see that  $N$  is linearly disjoint from  $K(x, y)$  over  $K(x)$ , and therefore  $N.K(x, y)|K(x, y)$  is again an infinite tower of Galois extensions of degree  $p$ . Since  $N$  is linearly disjoint from  $K(x)^h$  over  $K(x)$  and  $N^h = N.K(x)^h$  is linearly disjoint from  $L^h$  over  $K(x)^h$ , we see that  $N$  is linearly disjoint from  $L^h$  over  $K(x)$ . Since  $L^h$  is relatively algebraically closed in  $K(x, y)^h$ , this implies that  $N$  is linearly disjoint from  $K(x, y)^h$  over  $K(x)$ , and hence,  $N.K(x, y)$  is linearly disjoint from  $K(x, y)^h$  over  $K(x, y)$ . Therefore, the extension of  $v$  from  $K(x, y)$  to  $N.K(x, y)$  is unique. Since  $(K(x)^h(y))^h = K(x, y)^h$ , we see that  $(K(x)^h(y)|K(x, y), v)$  is immediate. So we have  $vK(x, y) = \frac{1}{p^\infty}\mathbb{Z}$ , which is  $p$ -divisible, and  $K(x, y)v = K$ , which is algebraically closed. Hence, the extension  $(N.K(x, y)|K(x, y), v)$  is immediate, and so it is an infinite tower of Galois extensions of degree  $p$  and defect  $p$ .  $\square$

*Remark 3.18.* For the above defined  $\eta_i$ , we have that the roots of  $X^p - X - \eta_{i-1}$  are  $\eta_i, \eta_i + 1, \dots, \eta_i + p - 1$ . Therefore,

$$\text{kas}(\eta_i, K(\eta_{i-1})) = 0$$

for all  $i$ . Setting  $a_i := x^i \eta_i$ , we obtain

$$\begin{aligned} \text{kas}(a_i, K(x)^h(a_1, \dots, a_{i-1})) &= \text{kas}(a_i, K(x)^h(\eta_{i-1})) \\ &= ivx + \text{kas}(\eta_i, K(x)^h(\eta_{i-1})) = ivx \end{aligned}$$

and  $va_{i+1} = (i+1)vx - \frac{1}{p^{i+1}}vx > ivx$ . This shows that  $va_{i+1} > va_i$  and that (3.19) is satisfied. So we can take  $y$  to be a pseudo limit of the Cauchy sequence  $(\sum_{j=1}^i a_j)_{i \in \mathbb{N}}$ . That is,

$$(3.20) \quad y = \sum_{i=1}^{\infty} x^i \eta_i.$$

Comparing this with (3.5), we see that we have replaced the term  $x^{-p^{-e_i}}$  by  $\eta_i$ , which has an infinite expansion in powers of  $x$ , starting with  $x^{-p^{-i}}$ .

## 4. RATIONAL FUNCTION FIELDS OF HIGHER TRANSCENDENCE DEGREE

This section is devoted to the proof of Theorems 1.6 and 1.8. We will make use of the following theorem, which we prove in [KU5]:

**Theorem 4.1.** *Let  $(L|K, v)$  be a valued field extension of finite transcendence degree  $\geq 0$ , with  $v$  non-trivial on  $L$ . Assume that one of the following two cases holds:*

*transcendental case:  $vL/vK$  has rational rank at least 1 or  $Lv|Kv$  is transcendental;*

*separable-algebraic case:  $L|K$  contains a separable-algebraic subextension  $L_0|K$  such that within some henselization of  $L$ , the corresponding extension  $L_0^h|K^h$  is infinite.*

*Then each maximal immediate extension of  $(L, v)$  has infinite transcendence degree over  $L$ .*

Note: The assertion need not be true for an infinite purely inseparable extension  $L|K$ .

Now we can give the

*Proof of Theorem 1.6.* Assume  $(K, v)$  is a valued field,  $n, \rho, \tau, \ell$  are non-negative integers,  $\Gamma$  is an ordered abelian group extension of  $vK$  such that  $\Gamma/vK$  is of rational rank  $\rho$ , and  $k|Kv$  is a field extension of transcendence degree  $\tau$ . We pick a maximal set of elements  $\alpha_1, \dots, \alpha_\rho$  in  $\Gamma$  rationally independent over  $vK$ , and a transcendence basis  $\zeta_1, \dots, \zeta_\tau$  of  $k|Kv$ .

We prove **Part A** of Theorem 1.6 first. So assume further that  $n > \rho + \tau$ ,  $\Gamma|vK$  and  $k|Kv$  are countably generated, and at least one of them is infinite. By Lemma 2.6 there is a unique extension of  $v$  from  $K$  to  $K(x_1, \dots, x_{\rho+\tau})$  such that  $vx_i = \alpha_i$  for  $1 \leq i \leq \rho$ , and  $x_{\rho+i}v = \zeta_i$  for  $1 \leq i \leq \tau$ . Since  $\Gamma|vK$  and  $k|Kv$  are countably generated, and since  $vK(x_1, \dots, x_{\rho+\tau})$  contains  $\alpha_1, \dots, \alpha_\rho$  and  $K(x_1, \dots, x_{\rho+\tau})v$  contains  $\zeta_1, \dots, \zeta_\tau$ , it follows that the extensions  $\Gamma|vK(x_1, \dots, x_{\rho+\tau})$  and  $k|K(x_1, \dots, x_{\rho+\tau})v$  are countably generated and algebraic.

Suppose first that at least one of these extensions is infinite. From our assumption that  $\Gamma \neq \{0\}$  it follows that  $v$  is non-trivial on  $K(x_1, \dots, x_{\rho+\tau})$ . Hence we can use Proposition 3.17 to find an extension of  $v$  from  $K(x_1, \dots, x_{\rho+\tau})$  to  $K(x_1, \dots, x_{\rho+\tau+1})$  such that

$$vK(x_1, \dots, x_{\rho+\tau+1}) = \Gamma \quad \text{and} \quad K(x_1, \dots, x_{\rho+\tau+1})v = k,$$

and such that the implicit constant field of  $(K(x_1, \dots, x_{\rho+\tau+1})|K(x_1, \dots, x_{\rho+\tau}), v)$  is an infinite separable-algebraic extension of  $K(x_1, \dots, x_{\rho+\tau})^h$ . That means that  $K(x_1, \dots, x_{\rho+\tau+1})^h|K(x_1, \dots, x_{\rho+\tau})^h$  contains an infinite separable-algebraic subextension. Hence by the separable-algebraic case of Theorem 4.1, each maximal immediate extension of  $K(x_1, \dots, x_{\rho+\tau+1})^h$  is of infinite transcendence degree. This shows that we can find an immediate extension of  $v$  from  $K(x_1, \dots, x_{\rho+\tau+1})^h$  to  $K(x_1, \dots, x_{\rho+\tau+1})^h(x_{\rho+\tau+2}, \dots, x_n)$ . Its restriction to  $K(x_1, \dots, x_n)$  is an immediate extension of  $v$  from  $K(x_1, \dots, x_{\rho+\tau+1})$ .

Now suppose that both extensions  $\Gamma|vK(x_1, \dots, x_{\rho+\tau})$  and  $k|K(x_1, \dots, x_{\rho+\tau})v$  are finite. Then from our hypothesis that at least one of the extensions  $\Gamma|vK$  and  $k|Kv$  is infinite, it follows that  $\rho > 0$  or  $\tau > 0$ . Hence by the transcendental case of Theorem 4.1, any maximal immediate extension of  $(K(x_1, \dots, x_{\rho+\tau}), v)$  is of infinite transcendence degree. In combination with Proposition 3.15, this shows

that there is an extension of  $v$  from  $K(x_1, \dots, x_{\rho+\tau})$  to  $K(x_1, \dots, x_{\rho+\tau+1})$  such that  $vK(x_1, \dots, x_{\rho+\tau+1}) = \Gamma$  and  $K(x_1, \dots, x_{\rho+\tau+1})v = k$ . By the transcendental case of Theorem 4.1, every maximal immediate extension of  $(K(x_1, \dots, x_{\rho+\tau+1}), v)$  is of infinite transcendence degree. So we can find an extension of  $v$  to  $K(x_1, \dots, x_n)$  as in the previous case.

In both cases we have that  $vK(x_1, \dots, x_n) = \Gamma$  and  $K(x_1, \dots, x_n)v = k$ , as required.

Now we prove **Part B**. So assume that  $n \geq \rho + \tau$ , and that  $\Gamma|vK$  and  $k|Kv$  are finitely generated.

**I)** First, we consider the case of  $\rho > 0$ . By Lemma 2.6 there is an extension of  $v$  from  $K$  to  $K(x_1, \dots, x_{\rho-1+\tau})$  such that  $vx_i = \alpha_i$  for  $1 \leq i \leq \rho - 1$ , and  $x_{\rho-1+i}v = \zeta_i$  for  $1 \leq i \leq \tau$ . Since  $\Gamma|vK$  and  $k|Kv$  are finitely generated, and since  $vK(x_1, \dots, x_{\rho-1+\tau})$  contains  $\alpha_1, \dots, \alpha_{\rho-1}$  and  $K(x_1, \dots, x_{\rho-1+\tau})v$  contains  $\zeta_1, \dots, \zeta_\tau$ , it follows that the extension  $\Gamma|vK(x_1, \dots, x_{\rho-1+\tau})$  is finitely generated of rational rank 1 (and thus  $\Gamma$  is of the form  $\Gamma_0 \oplus \mathbb{Z}$  with  $\Gamma_0/vK(x_1, \dots, x_{\rho-1+\tau})$  finite), and the extension  $k|K(x_1, \dots, x_{\rho-1+\tau})v$  is finite.

If  $v$  is not trivial on  $K(x_1, \dots, x_{\rho-1+\tau})$ , then we can apply Proposition 3.15 to obtain an extension of  $v$  from  $K(x_1, \dots, x_{\rho-1+\tau})$  to  $K(x_1, \dots, x_{\rho+\tau})$  such that  $vK(x_1, \dots, x_{\rho+\tau}) = \Gamma$  and  $K(x_1, \dots, x_{\rho+\tau})v = k$ . From the transcendental case of Theorem 4.1 we see that any maximal immediate extension of  $(K(x_1, \dots, x_{\rho+\tau}), v)$  is of infinite transcendence degree. This shows that we can find an immediate extension of  $v$  from  $K(x_1, \dots, x_{\rho+\tau})$  to  $K(x_1, \dots, x_n)$ .

Now suppose that  $v$  is trivial on  $K(x_1, \dots, x_{\rho-1+\tau})$ . Then  $\rho = 1$  and  $v$  is trivial on  $K$ . It follows that  $\Gamma \simeq \mathbb{Z}$ , and we pick a generator  $\gamma$  of  $\Gamma$ .

If  $n > 1 + \tau$ , we modify the above procedure in such a way that we use Lemma 2.6 to choose an extension of  $v$  from  $K(x_1, \dots, x_\tau)$  to  $K(x_1, \dots, x_{1+\tau})$  so that the latter has value group  $\mathbb{Z}\gamma = \Gamma$  and residue field  $Kv(\zeta_1, \dots, \zeta_\tau)$ . Then we use Proposition 3.15 in combination with the transcendental case of Theorem 4.1 to find an extension of  $v$  to  $K(x_1, \dots, x_n)$  with value group  $\Gamma$  and residue field  $k$ .

If  $n = 1 + \tau$ , then by our assumption B2),  $k$  is a simple algebraic extension of a rational function field  $k'$  in  $\tau$  variables over  $Kv$  (or of  $Kv$  itself if  $\tau = 0$ ), or a rational function field in one variable over a finitely generated field extension  $k_0$  of  $Kv$  of transcendence degree  $\tau - 1$ .

Assume the first case holds. The extension of  $v$  to the rational function field  $K(x_1, \dots, x_\tau)$  can be chosen such that  $k' = Kv(x_1v, \dots, x_\tau v) = K(x_1, \dots, x_\tau)v$ . Then the extension  $k|K(x_1, \dots, x_\tau)v$  is simple algebraic, and by Remark 3.12 there is an extension of  $v$  to the rational function field  $K(x_1, \dots, x_\tau)(x_{1+\tau})$  which satisfies  $vK(x_1, \dots, x_{1+\tau}) = \mathbb{Z}\gamma = \Gamma$  and  $K(x_1, \dots, x_{1+\tau})v = k$ .

In the second case, we know that  $\tau \geq 1$ . The elements  $\zeta_i \in k$  can be chosen in such a way that  $\zeta_1, \dots, \zeta_{\tau-1}$  form a transcendence basis of  $k_0|Kv$ . We pick  $\gamma \in \Gamma$ ,  $\gamma \neq 0$ . By Lemma 2.6 there is an extension of  $v$  from  $K$  to  $K(x_1, \dots, x_\tau)$  such that  $x_i v = \zeta_i$  for  $1 \leq i \leq \tau - 1$ , and  $vx_\tau = \gamma$ . Since  $K(x_1, \dots, x_\tau)v = Kv(\zeta_1, \dots, \zeta_{\tau-1})$ , we see that  $k$  is a rational function field in one variable over a finite extension of  $K(x_1, \dots, x_\tau)v$ . Further,  $vK(x_1, \dots, x_\tau) = \Gamma$ . Now we use Proposition 3.15 to find an extension of  $v$  to  $K(x_1, \dots, x_{1+\tau})$  with value group  $\Gamma$  and residue field  $k$ .

**II)** Second, suppose that  $\rho = 0$ . Then because of  $\Gamma \neq \{0\}$ , we know that  $v$  is non-trivial on  $K$ . If  $\tau > 0$ , then we proceed as follows. The case of  $n > \tau$  is covered by Part A. So we assume that  $n = \tau$ , and that  $k$  is a rational function

field in one variable over a finitely generated field extension  $k_0$  of  $Kv$  of transcendence degree  $\tau - 1$ . Again we choose  $\zeta_1, \dots, \zeta_{\tau-1}$  to form a transcendence basis of  $k_0|Kv$ , and use Lemma 2.6 to find an extension of  $v$  from  $K$  to  $K(x_1, \dots, x_{\tau-1})$  such that  $x_i v = \zeta_i$  for  $1 \leq i \leq \tau - 1$ . Again we obtain that  $k$  is a rational function field in one variable over a finite extension of  $K(x_1, \dots, x_{\tau-1})v$ . By assumption,  $\Gamma/vK(x_1, \dots, x_{\tau-1}) = \Gamma/vK$  is finite. Hence by Proposition 3.15 there is an extension of  $v$  from  $K(x_1, \dots, x_{\tau-1})$  to  $K(x_1, \dots, x_\tau)$  such that  $vK(x_1, \dots, x_\tau) = \Gamma$  and  $K(x_1, \dots, x_\tau)v = k$ .

Finally, suppose that  $\rho = 0 = \tau$  and that there is an immediate extension  $(K', v)$  of  $(K, v)$  which is either infinite separable-algebraic or of transcendence degree at least  $n$ . If the former holds, then we obtain from the separable-algebraic case of Theorem 4.1 that every maximal immediate extension of  $(K', v)$  has infinite transcendence degree. But a maximal immediate extension of  $(K', v)$  is also a maximal immediate extension of  $(K, v)$ . Thus in all cases, we have the existence of immediate extensions of  $(K, v)$  of transcendence degree at least  $n$ . Therefore, we can choose an immediate extension of  $v$  from  $K$  to  $K(x_1, \dots, x_{n-1})$  such that  $(K(x_1, \dots, x_{n-1}), v)$  still admits a transcendental immediate extension. The extensions  $\Gamma|vK = \Gamma|vK(x_1, \dots, x_{n-1})$  and  $k|Kv = k|K(x_1, \dots, x_{n-1})v$  are both finite since they are finitely generated and algebraic. Hence by Proposition 3.15 there is an extension of  $v$  from  $K(x_1, \dots, x_{n-1})$  to  $K(x_1, \dots, x_n)$  such that

$$vK(x_1, \dots, x_n) = \Gamma \quad \text{and} \quad K(x_1, \dots, x_n)v = k. \quad \square$$

Now we give the

*Proof of Theorem 1.7.* Corollary 2.7 shows that  $n \geq \rho + \tau$ . If  $n = \rho + \tau$ , then Corollary 2.7 tells us that  $vF|vK$  and  $Fv|Kv$  are finitely generated extensions. The fact that  $vF|vK$  and  $Fv|Kv$  are always countably generated follows from Theorem 2.9 by induction on the transcendence degree  $n$ .

Suppose that  $F$  admits a transcendence basis  $x_1, \dots, x_n$  such that the residue field extension  $Fv|K(x_1, \dots, x_{n-1})v$  is of transcendence degree 1. (This is in particular the case when  $n = \tau$ .) Then by Ohm's Ruled Residue Theorem,  $Fv$  is a rational function field in one variable over a finite extension of  $K(x_1, \dots, x_{n-1})v$ . Whenever equality holds in (2.4) for an extension  $L|K$  of finite transcendence degree, it will hold in the respective inequality for any subextension of  $L|K$ . Hence if  $n = \rho + \tau$ , then by Corollary 2.7,  $K(x_1, \dots, x_{n-1})v|Kv$  is a finitely generated field extension, and if  $n = \tau$ , this extension is of transcendence degree  $\tau - 1$ . This yields that B3) holds for  $k = Fv$ .

Now suppose that there does not exist such a transcendence basis, and that  $n = \rho + \tau$  with  $\rho = 1$  and  $v$  is trivial on  $K$ . We pick any transcendence basis  $x_1, \dots, x_n$ . Since  $Fv|K(x_1, \dots, x_{n-1})v$  is algebraic and  $n = \rho + \tau$ , the rational rank of  $vF/vK(x_1, \dots, x_{n-1})$  must be 1. Since  $\rho = 1$  and  $v$  is trivial on  $K$ , this means that  $v$  is also trivial on  $K(x_1, \dots, x_{n-1})$ . Hence,  $Kv = K$  and

$$K(x_1, \dots, x_{n-1})v = K(x_1, \dots, x_{n-1})$$

(modulo an isomorphism). Remark 3.12 now shows that  $Fv$  is a simple algebraic extension of  $K(x_1, \dots, x_{n-1})$ , which in turn is a rational function field in  $\tau$  variables over  $K = Kv$ , or equal to  $Kv$  if  $\tau = 0$ . Together with what we have shown before, this proves that B2) holds for  $k = Fv$ .

To conclude this proof, assume that  $\rho = 0 = \tau$ . Then by Lemma 3.4,  $(F.\tilde{K}|\tilde{K}, v)$  is immediate, for any extension of  $v$  from  $F$  to  $F.\tilde{K}$ . This shows that  $(\tilde{K}, v)$  admits an immediate extension of transcendence degree  $n$ .  $\square$

Finally, we give the

*Proof of Theorem 1.8.* We have to show that whenever we construct an extension of the form  $(L_2|L_1, v)$  with  $Kv \subseteq L_1v \subseteq L_2v \subseteq k$  in the previous proof, then the ordering of  $L_2v$  induced by the given ordering of  $k$  can be lifted to an extension of the lifting that we have already obtained on  $L_1$ . Whenever we apply Propositions 3.14, 3.15 and 3.17, we obtain this already from the assertions of these propositions. Whenever we apply Lemma 2.6, we obtain this from Proposition 2.18, because in this case we always have that  $vL_2$  is generated over  $vL_1$  by rationally independent values, and therefore  $2vL_2 \cap vL_1 = 2vL_1$ . Finally, whenever  $(L_2|L_1, v)$  is an immediate extension, we can also apply Proposition 2.18, because  $2vL_2 \cap vL_1 = 2vL_1 \cap vL_1 = 2vL_1$ .  $\square$

## 5. HOMOGENEOUS SEQUENCES

In this section, we will develop special sequences which under certain tameness conditions can be used to determine implicit constant fields.

**5.1. Homogeneous approximations.** Let  $(K, v)$  be any valued field and  $a, b$  elements in some valued field extension  $(L, v)$  of  $(K, v)$ . We will say that  $a$  is **strongly homogeneous over**  $(K, v)$  if  $a \in K^{\text{sep}} \setminus K$ , the extension of  $v$  from  $K$  to  $K(a)$  is unique (or equivalently,  $K(a)|K$  is linearly disjoint from all henselizations of  $(K, v)$ ), and

$$(5.1) \quad va = \text{kas}(a, K) .$$

Note that in this case,  $va = v(\sigma a - a)$  for all automorphisms  $\sigma$  such that  $\sigma a \neq a$ ; indeed, we have  $v\sigma a = va$ , and therefore  $va \geq v(\sigma a - a) \geq va$ .

We will say that  $a$  is **homogeneous over**  $(K, v)$  if there is some  $d \in K$  such that  $a - d$  is strongly homogeneous over  $(K, v)$ , i.e.,

$$v(a - d) = \text{kas}(a - d, K) = \text{kas}(a, K) .$$

We call  $a \in L$  a **homogeneous approximation of  $b$  over  $K$**  if  $a$  is homogeneous over  $K$  and  $v(b - a) > vb$  (and consequently,  $va = vb$ ). From Corollary 2.21 we obtain

**Lemma 5.1.** *If  $a \in L$  is a homogeneous approximation of  $b$ , then  $a$  lies in the henselization of  $K(b)$  w.r.t. every extension of the valuation  $v$  from  $K(a, b)$  to  $\widetilde{K(b)}$ .*

We will also exploit the following easy observation:

**Lemma 5.2.** *Let  $(K', v)$  be any henselian extension field of  $(K, v)$  such that  $a \notin K'$ . If  $a$  is homogeneous over  $(K, v)$ , then it is also homogeneous over  $(K', v)$ , and  $\text{kas}(a, K) = \text{kas}(a, K')$ . If  $a$  is strongly homogeneous over  $(K, v)$ , then it is also strongly homogeneous over  $(K', v)$ .*

*Proof.* Suppose  $a - d$  is separable-algebraic over  $K$  and  $v(a - d) = \text{kas}(a - d, K)$  for some  $d \in K$ . Then  $a - d$  is also separable-algebraic over  $K'$ . Further,

$$\text{kas}(a - d, K') \leq \text{kas}(a - d, K)$$

since restriction to  $\tilde{K}$  is a map sending  $\{\sigma \in \text{Gal } K' \mid \sigma a \neq a\}$  into  $\{\sigma \in \text{Gal } K \mid \sigma a \neq a\}$ . Hence,

$$v(a - d) \leq \text{kras}(a - d, K') \leq \text{kras}(a - d, K) = v(a - d),$$

which shows that equality holds everywhere. Thus,

$$\text{kras}(a, K') = \text{kras}(a - d, K') = \text{kras}(a - d, K) = \text{kras}(a, K).$$

Since  $(K', v)$  is henselian by assumption, the extension of  $v$  from  $K'$  to  $K'(a)$  is unique. This shows that  $a - d$  is strongly homogeneous over  $(K', v)$ , and concludes the proof of our assertions.  $\square$

The following gives the crucial criterion for an element to be (strongly) homogeneous over  $(K, v)$ :

**Lemma 5.3.** *Suppose that  $a \in \tilde{K}$  and that there is some extension of  $v$  from  $K$  to  $K(a)$  such that if  $e$  is the least positive integer for which  $eva \in vK$ , then*

- a)  $e$  is not divisible by  $\text{char } Kv$ , and*
- b) there exists some  $c \in K$  such that  $vca^e = 0$ ,  $ca^e v$  is separable-algebraic over  $Kv$ , and the degree of  $ca^e$  over  $K$  is equal to the degree  $f$  of  $ca^e v$  over  $Kv$ .*

*Then  $[K(a) : K] = ef$ , and if  $a \notin K$ , then  $a$  is strongly homogeneous over  $(K, v)$ .*

*Proof.* We have

$$\begin{aligned} ef &\geq [K(a) : K(a^e)] \cdot [K(a^e) : K] = [K(a) : K] \\ &\geq (vK(a) : vK) \cdot [K(a)v : Kv] \geq ef. \end{aligned}$$

So equality holds everywhere, and we obtain  $[K(a) : K] = ef$ ,  $(vK(a) : vK) = e$  and  $[K(a)v : Kv] = f$ . By the fundamental inequality, the latter implies that the extension of  $v$  from  $K$  to  $K(a)$  is unique.

Now assume that  $a \notin K$ . Take two distinct conjugates  $\sigma a \neq \tau a$  of  $a$  and set  $\eta := \sigma a / \tau a \neq 1$ . If  $\sigma a^e \neq \tau a^e$ , then  $c\sigma a^e = \sigma c a^e$  and  $c\tau a^e = \tau c a^e$  are distinct conjugates of  $ca^e$ . By hypothesis, their residues are also distinct, and therefore the residue of  $\sigma a^e / \tau a^e = \eta^e$  is not 1. It follows that the residue of  $\eta$  is not 1. If  $\sigma a^e = \tau a^e$ , then  $\eta$  is an  $e$ -th root of unity. Since  $e$  is not divisible by the residue characteristic, it again follows that the residue of  $\eta$  is not equal to 1. Hence in both cases, we obtain that  $v(\eta - 1) = 0$ , which shows that  $v(\sigma a - \tau a) = v\tau a = va$ . We have now proved (5.1).  $\square$

**Lemma 5.4.** *Assume that  $b$  is an element in some algebraically closed valued field extension  $(L, v)$  of  $(K, v)$ . Suppose that there is some  $e \in \mathbb{N}$  not divisible by  $\text{char } Kv$ , and some  $c \in K$  such that  $vcb^e = 0$  and  $cb^e v$  is separable-algebraic over  $Kv$ . If  $e > 0$  or  $cb^e v \notin Kv$ , then we can find in  $L$  a strongly homogeneous approximation of  $b$  over  $K$ .*

*Proof.* Take a monic polynomial  $g$  over  $K$  with  $v$ -integral coefficients whose reduction modulo  $v$  is the minimal polynomial of  $cb^e v$  over  $Kv$ . Then let  $d \in \tilde{K}$  be the root of  $g$  whose residue is  $cb^e v$ . The degree of  $d$  over  $K$  is the same as that of  $cb^e v$  over  $Kv$ . We have that  $v(\frac{d}{cb^e} - 1) > 0$ . So there exists  $a_0 \in \tilde{K}$  with residue 1 and such that  $a_0^e = \frac{d}{cb^e}$ . Then for  $a := a_0 b$ , we find that  $v(a - b) = vb + v(a_0 - 1) > vb$  and  $ca^e = d$ . By the foregoing lemma, this shows that  $a$  is a strongly homogeneous approximation of  $b$ .  $\square$



**5.2. Homogeneous sequences.** Let  $(K(x)|K, v)$  be any extension of valued fields. We fix an extension of  $v$  to  $\widehat{K(x)}$ .

Let  $S$  be an initial segment of  $\mathbb{N}$ , that is,  $S = \mathbb{N}$  or  $S = \{1, \dots, n\}$  for some  $n \in \mathbb{N}$  or  $S = \emptyset$ . A sequence

$$\mathfrak{S} := (a_i)_{i \in S}$$

of elements in  $\tilde{K}$  will be called a **homogeneous sequence for  $x$**  if the following conditions are satisfied for all  $i \in S$  (where we set  $a_0 := 0$ ):

**(HS)**  $a_i - a_{i-1}$  is a homogeneous approximation of  $x - a_{i-1}$  over  $K(a_0, \dots, a_{i-1})$ .

Recall that then by definition of “homogeneous”,  $a_i \notin K(a_0, \dots, a_{i-1})^h$ . We call  $S$  the **support** of the sequence  $\mathfrak{S}$ . We set

$$K_{\mathfrak{S}} := K(a_i \mid i \in S).$$

If  $\mathfrak{S}$  is the empty sequence, then  $K_{\mathfrak{S}} = K$ .

From this definition, we obtain

**Lemma 5.5.** *If  $1 \leq i < j \in S$ , then*

$$(5.2) \quad v(x - a_j) > v(x - a_i) = v(a_{i+1} - a_i).$$

*If  $S = \mathbb{N}$ , then  $(a_i)_{i \in S}$  is a pseudo Cauchy sequence with pseudo limit  $x$ .*

*Proof.* If  $1 \leq i \in S$ , then  $a_i - a_{i-1}$  is a homogeneous approximation of  $x - a_{i-1}$ . Hence, by definition,

$$v(x - a_i) = v(x - a_{i-1} - (a_i - a_{i-1})) > v(x - a_{i-1}),$$

whence  $v(a_i - a_{i-1}) = \min\{v(x - a_i), v(x - a_{i-1})\} = v(x - a_{i-1})$ . If  $i < j \in S$ , then by induction,  $v(x - a_j) > v(x - a_i)$ .

Suppose that  $S = \mathbb{N}$ . Then it follows by induction that for all  $k > j > i \geq 1$ ,

$$v(x - a_k) > v(x - a_j) > v(x - a_i),$$

and therefore

$$\begin{aligned} v(a_k - a_j) &= \min\{v(x - a_k), v(x - a_j)\} = v(x - a_j) > v(x - a_i) \\ &= \min\{v(x - a_j), v(x - a_i)\} = v(a_j - a_i). \end{aligned}$$

This shows that  $(a_i)_{i \in S}$  is a pseudo Cauchy sequence. The equality in (5.2) shows that  $x$  is a pseudo limit of this sequence.  $\square$

Let us also observe the following:

**Lemma 5.6.** *Take  $x, x' \in L$ . If  $a \in L$  is a homogeneous approximation of  $x$  over  $K$  and if  $v(x - x') > vx$ , then  $a$  is also a homogeneous approximation of  $x'$  over  $K$ . If  $(a_i)_{i \in S}$  is a homogeneous sequence for  $x$  over  $K$  and if  $v(x - x') > v(x - a_k)$  for all  $k \in S$ , then  $(a_i)_{i \in S}$  is also a homogeneous sequence for  $x'$  over  $K$ .*

*In particular, for each  $k \in S$  such that  $k > 1$ ,  $(a_i)_{i < k}$  is a homogeneous sequence for  $a_k$  over  $K$ .*

*Proof.* Suppose  $a$  is a homogeneous approximation of  $x$  over  $K$ . Then  $v(x - a) > vx$ . If also  $v(x - x') > vx$ , then  $vx' = vx$  and  $v(x' - a) \geq \max\{v(x - x'), v(x - a)\} > vx = vx'$ . This yields the first assertion.

Now assume that  $(a_i)_{i \in S}$  is a homogeneous sequence for  $x$  over  $K$  and that  $v(x - x') > v(x - a_k)$  for all  $k \in S$ . Then for all  $k \in S$ ,

$$v(x' - a_k) = \min\{v(x' - x), v(x - a_k)\} = v(x - a_k).$$

Hence,

$$\begin{aligned} v(x' - a_{k-1} - (a_k - a_{k-1})) &= v(x' - a_k) = v(x - a_k) \\ &> v(x - a_{k-1}) = v(x' - a_{k-1}), \end{aligned}$$

showing that  $a_k - a_{k-1}$  is also a homogeneous approximation of  $x' - a_{k-1}$ . Hence,  $(a_i)_{i \in S}$  is a homogeneous sequence for  $x'$  over  $K$ .  $\square$

What is special about homogeneous sequences is described by the following lemma:

**Lemma 5.7.** *Assume that  $(a_i)_{i \in S}$  is a homogeneous sequence for  $x$  over  $K$ . Then*

$$(5.3) \quad K_{\mathfrak{S}} \subset K(x)^h.$$

*For every  $k \in S$ , we have  $a_1, \dots, a_k \in K(a_k)^h$ . If  $S = \{1, \dots, n\}$ , then*

$$(5.4) \quad K_{\mathfrak{S}}^h = K(a_n)^h.$$

*Proof.* We use induction on  $k \in S$ . Suppose that we have already shown that  $a_{k-1} \in K(x)^h$ . As  $a_k - a_{k-1}$  is a homogeneous approximation of  $x - a_{k-1}$ , we know from Lemma 5.1 that

$$a_k - a_{k-1} \in K(x - a_{k-1})^h \subseteq K(x)^h.$$

This proves (5.3). Now all other assertions follow when we replace  $x$  by  $a_k$  in the above argument, using the fact that, by the previous lemma,  $(a_i)_{i < n}$  is a homogeneous sequence for  $a_n$  over  $K$ .  $\square$

**Proposition 5.8.** *Assume that  $\mathfrak{S} = (a_i)_{i \in S}$  is a homogeneous sequence for  $x$  over  $K$  with support  $S = \mathbb{N}$ . Then  $(a_i)_{i \in \mathbb{N}}$  is a pseudo Cauchy sequence of transcendental type in  $(K_{\mathfrak{S}}, v)$  with pseudo limit  $x$ , and  $(K_{\mathfrak{S}}(x)|K_{\mathfrak{S}}, v)$  is immediate and pure.*

*Proof.* By Lemma 5.5,  $(a_i)_{i \in \mathbb{N}}$  is a pseudo Cauchy sequence with pseudo limit  $x$ . Suppose it were of algebraic type. Then by [KA], Theorem 3, there would exist some algebraic extension  $(K_{\mathfrak{S}}(b)|K_{\mathfrak{S}}, v)$  with  $b$  a pseudo limit of the sequence. But then  $v(x - b) > v(x - a_k)$  for all  $k \in S$  and, by Lemma 5.6,  $(a_i)_{i \in S}$  is also a homogeneous sequence for  $b$  over  $K$ . Hence by Lemma 5.7,  $K_{\mathfrak{S}}^h \subset K(b)^h = K^h(b)$ . Since  $b$  is algebraic over  $K$ , the extension  $K^h(b)|K^h$  is finite. On the other hand,  $K_{\mathfrak{S}}^h|K^h$  is infinite since, by the definition of homogeneous elements,  $a_k \notin K(a_i \mid 1 \leq i < k)^h$  for every  $k \in \mathbb{N}$ , and therefore each extension  $K(a_i \mid 1 \leq i \leq k+1)^h|K(a_i \mid 1 \leq i \leq k)^h$  is non-trivial. This contradiction shows that the sequence is of transcendental type. Hence by definition,  $(K_{\mathfrak{S}}(x)|K_{\mathfrak{S}}, v)$  is pure. Further, it follows from Lemma 2.4 that  $(K_{\mathfrak{S}}(x)|K_{\mathfrak{S}}, v)$  is immediate.  $\square$

This proposition leads to the following definition. A homogeneous sequence  $\mathfrak{S}$  for  $x$  over  $K$  will be called **(weakly) pure homogeneous** if  $(K_{\mathfrak{S}}(x)|K_{\mathfrak{S}}, v)$  is (weakly) pure in  $x$ . Hence if  $S = \mathbb{N}$ , then  $\mathfrak{S}$  is always a pure homogeneous sequence. The empty sequence is a (weakly) pure homogeneous sequence for  $x$  over  $K$  if and only if already  $(K(x)|K, v)$  is (weakly) pure in  $x$ .

**Theorem 5.9.** *Suppose that  $\mathfrak{S}$  is a (weakly) pure homogeneous sequence for  $x$  over  $K$ . Then*

$$K_{\mathfrak{S}}^h = \text{IC}(K(x)|K, v).$$

*Further,  $K_{\mathfrak{S}}v$  is the relative algebraic closure of  $Kv$  in  $K(x)v$ , and the torsion subgroup of  $vK(x)/vK_{\mathfrak{S}}$  is finite. If  $\mathfrak{S}$  is pure, then  $vK_{\mathfrak{S}}$  is the relative divisible closure of  $vK$  in  $vK(x)$ .*

*Proof.* The assertions follow from Lemma 3.5, Lemma 2.10 and Lemma 3.7, together with the fact that because  $K_{\mathfrak{S}}|K$  is algebraic, the same holds for  $vK_{\mathfrak{S}}|vK$  and  $K_{\mathfrak{S}}v|Kv$  by Lemma 2.1.  $\square$

**5.3. Conditions for the existence of homogeneous sequences.** Now we have to discuss for which extensions  $(K(x)|K, v)$  there exist homogeneous sequences.

An algebraic extension  $(L|K, v)$  of henselian fields is called **tame** if the following conditions hold:

- (TE1)  $Lv|Kv$  is separable,
- (TE2) if  $\text{char } Kv = p > 0$ , then the order of each element in  $vL/vK$  is prime to  $p$ , and
- (TE3)  $[K' : K] = (vK' : vK)[K'v : Kv]$  holds for every finite subextension  $K'|K$  of  $L|K$ .

Condition (TE3) means that equality holds in the fundamental inequality (2.1). If  $L'|K$  is any subextension of  $L|K$ , then  $(L|K, v)$  is a tame extension if and only if  $(L|L', v)$  and  $(L'|K, v)$  are (this is easy to prove if  $L|K$  is finite). Further, it is well known that for  $(K, v)$  henselian, the ramification field of the extension  $(K^{\text{sep}}|K, v)$  is the unique maximal tame extension of  $(K, v)$  (cf. [E]). A henselian valued field  $(K, v)$  is called a **tame field** if all its algebraic extensions are tame, or equivalently, the following conditions hold:

- (T1)  $Kv$  is perfect,
- (T2) if  $\text{char } Kv = p > 0$ , then  $vK$  is  $p$ -divisible, and
- (T3) for every finite extension  $K'|K$ ,  $[K' : K] = (vK' : vK)[K'v : Kv]$ .

Note that every valued field with a residue field of characteristic zero is tame; this is a consequence of the Lemma of Ostrowski (cf. [R]). It follows directly from the definition together with the multiplicativity of ramification index and inertia degree that every finite extension of a tame field is again a tame field. If  $(K, v)$  is a tame field, then condition (T3) shows that  $(K, v)$  does not admit any proper immediate algebraic extensions; hence by Theorem 3 of [KA], every pseudo Cauchy sequence in  $(K, v)$  without a pseudo limit in  $K$  must be of transcendental type.

If an element  $a \in \tilde{K}$  satisfies the conditions of Lemma 5.3, then  $(K(a)|K, v)$  is a tame extension. The following implication is also true, as was noticed by Sudesh K. Khanduja (cf. [KH11], Theorem 1.2):

**Proposition 5.10.** *Suppose that  $(K, v)$  is henselian. If  $a$  is homogeneous over  $(K, v)$ , then  $(K(a)|K, v)$  is a tame extension. If  $\mathfrak{S}$  is a homogeneous sequence over  $(K, v)$ , then  $K_{\mathfrak{S}}$  is a tame extension of  $K$ .*

*Proof.* Since  $K(a - d) = K(a)$  for  $d \in K$ , we may assume w.l.o.g. that  $a$  is strongly homogeneous over  $(K, v)$ . If  $(K(a)|K, v)$  were not a tame extension, then  $a$  would not lie in the ramification field of the extension  $(K^{\text{sep}}|K, v)$ . So there would exist an automorphism  $\sigma$  in the ramification group such that  $\sigma a \neq a$ . But by the definition of the ramification group,

$$v(\sigma a - a) > va = \text{kras}(a, K),$$

a contradiction.

The second assertion is proved using the first assertion and the fact that a (possibly infinite) tower of tame extensions is itself a tame extension.  $\square$

In fact, it can also be shown that  $va = \text{kras}(a, K)$  implies that  $a$  satisfies the conditions of Lemma 5.3.

We can give the following characterization of elements in tame extensions:

**Proposition 5.11.** *An element  $b \in \tilde{K}$  belongs to a tame extension of the henselian field  $(K, v)$  if and only if there is a finite homogeneous sequence  $a_1, \dots, a_k$  for  $a$  over  $(K, v)$  such that  $b \in K(a_k)$ .*

*Proof.* Suppose that such a sequence exists. By the foregoing proposition,  $K_{\mathfrak{S}}$  is a tame extension of  $K$ . Since  $K(a_1, \dots, a_k) = K(a_k)$  by Lemma 5.7, it contains  $b$ .

For the converse, let  $b$  be an element in some tame extension of  $(K, v)$ . Then  $b$  satisfies the assumptions of Lemma 5.4, and hence there is a homogeneous approximation  $a_1 \in \tilde{K}$  of  $b$  over  $K$ . By the foregoing proposition,  $K(a_1)$  is a tame extension of  $K$ , and therefore, by the general facts we have noted following the definition of tame extensions,  $K(a_1, b - a_1)$  is a tame extension of  $K(a_1)$ . We repeat this step, replacing  $b$  by  $b - a_1$ . By induction, we build a homogeneous sequence for  $b$  over  $K$ . It cannot be infinite, since  $b$  is algebraic over  $K$  (cf. Proposition 5.8). Hence it stops with some element  $a_k$ . Our construction shows that this can only happen if  $b \in K(a_1, \dots, a_k) = K(a_k)$ .  $\square$

**Proposition 5.12.** *Assume that  $(K, v)$  is a henselian field. Then  $(K, v)$  is a tame field if and only if for every element  $x$  in any extension  $(L, v)$  of  $(K, v)$  there exists a weakly pure homogeneous sequence for  $x$  over  $K$ , provided that  $x$  is transcendental over  $K$ .*

*Proof.* First, let us assume that  $(K, v)$  is a tame field and that  $x$  is an element in some extension  $(L, v)$  of  $(K, v)$ , transcendental over  $K$ . We set  $a_0 = 0$ . We assume that  $k \geq 0$  and that  $a_i$  for  $i \leq k$  are already constructed. Like  $K$ , also the finite extension  $K_k := K(a_0, \dots, a_k)$  is a tame field. Therefore, if  $x$  is the pseudo limit of a pseudo Cauchy sequence in  $K_k$ , then this pseudo Cauchy sequence must be of transcendental type, and  $K_k(x)|K_k$  is pure and hence weakly pure in  $x$ .

If  $K_k(x)|K_k$  is weakly pure in  $x$ , then we take  $a_k$  to be the last element of  $\mathfrak{S}$  if  $k > 0$ , and  $\mathfrak{S}$  to be empty if  $k = 0$ .

Assume that this is not the case. Then  $x$  cannot be the pseudo limit of a pseudo Cauchy sequence without pseudo limit in  $K_k$ . So the set  $v(x - a_k - K_k)$  must have a maximum, say  $x - a_k - d$  with  $d \in K_k$ . Since we assume that  $K_k(x)|K_k$  is not weakly pure in  $x$ , there exist  $e \in \mathbb{N}$  and  $c \in K_k$  such that  $vc(x - a_k - d)^e = 0$  and  $c(x - a_k - d)^e$  is algebraic over  $K_kv$ . Conditions (T1) and (T2) tell us that  $e$  can be chosen to be prime to  $\text{char } K_v$  and that  $c(x - a_k - d)^e v$  is separable-algebraic over  $K_kv$ . Since  $v(x - a_k - d)$  is maximal in  $v(x - a_k - K_k)$ , we must have that  $e > 1$  or  $c(x - a_k - d)^e v \notin K_kv$ .

Now Lemma 5.4 shows that there exists a homogeneous approximation  $a \in \tilde{K}$  of  $x - a_k - d$  over  $K_k$ ; so  $a + d$  is a homogeneous approximation of  $x - a_k$  over  $K_k$ , and we set  $a_{k+1} := a_k + a + d$ . This completes our induction step. If our construction stops at some  $k$ , then  $K_k(x)|K_k$  is weakly pure in  $x$  and we have obtained a weakly pure homogeneous sequence. If the construction does not stop, then  $S = \mathbb{N}$  and the obtained sequence is pure homogeneous.

For the converse, assume that  $(K, v)$  is not a tame field. We choose an element  $b \in \tilde{K}$  such that  $K(b)|K$  is not a tame extension. On  $K(b, x)$  we take the valuation  $v_{b, \gamma}$  with  $\gamma$  an element in some ordered abelian group extension such that  $\gamma > vK$ . Choose any extension of  $v$  to  $\tilde{K}(x)$ . Since  $vK$  is cofinal in  $v\tilde{K}$ , we have that  $\gamma > v\tilde{K}$ . Since  $b \in \tilde{K}$ , we have  $\gamma \in v\tilde{K}(x)$ . Hence,  $(\tilde{K}(x)|\tilde{K}, v)$  is value-transcendental.

Now suppose that there exists a weakly pure homogeneous sequence  $\mathfrak{S}$  for  $x$  over  $K$ . By Lemma 3.3, also  $(K_{\mathfrak{S}}(x)|K_{\mathfrak{S}}, v)$  is value-transcendental. Since  $(K_{\mathfrak{S}}(x)|K_{\mathfrak{S}}, v)$  is also weakly pure, it follows that there must be some  $c \in K_{\mathfrak{S}}$  such that  $x - c$  is a value-transcendental element (all other cases in the definition of “weakly pure” lead to immediate or residue-transcendental extensions). But if  $c \neq b$  then  $v(b - c) \in v\tilde{K}$ , and thus  $v(c - b) < \gamma$ . This implies  $v(x - c) = \min\{v(x - b), v(b - c)\} = v(b - c) \in v\tilde{K}$ , a contradiction. This shows that  $b = c \in K_{\mathfrak{S}}$ . On the other hand,  $K_{\mathfrak{S}}$  is a tame extension of  $K$  by Proposition 5.10 and cannot contain  $b$ . This contradiction shows that there cannot exist a weakly pure homogeneous sequence for  $x$  over  $K$ .  $\square$

## 6. APPLICATIONS

Let us show how to apply our results to power series fields. We denote by  $k((G))$  the field of power series with coefficients in the field  $k$  and exponents in the ordered abelian group  $G$ .

**Theorem 6.1.** *Let  $(K, v)$  be a henselian subfield of a power series field  $k((G))$  such that  $v$  is the restriction of the canonical valuation of  $k((G))$ . Suppose that  $K$  contains all monomials of the form  $ct^\gamma$  for  $c \in Kv \subseteq k$  and  $\gamma \in vK \subseteq \Gamma$ . Consider a power series*

$$z = \sum_{i \in \mathbb{N}} c_i t^{\gamma_i},$$

where  $(\gamma_i)_{i \in \mathbb{N}}$  is a strictly increasing sequence in  $G$ , all  $c_i \in k$  are separable-algebraic over  $Kv$ , and for each  $i$  there is an integer  $e_i > 0$  prime to the characteristic of  $Kv$  and such that  $e_i \gamma_i \in vK$ . Then, upon taking the henselization of  $K(z)$  in  $k((G))$ , we obtain that  $K(c_i t^{\gamma_i} \mid i \in \mathbb{N}) \subseteq K(z)^h$ . Consequently,  $vK(z)$  contains all  $\gamma_i$ , and if  $\gamma_i \in vK$  for all  $i \in \mathbb{N}$ , then  $K(z)v$  contains all  $c_i$ .

If  $vK + \sum_{i=1}^{\infty} \mathbb{Z}\gamma_i/vK$  or  $Kv(c_i \mid i \in \mathbb{N})|Kv$  is infinite, then  $z$  is transcendental over  $K$ , we have  $\text{IC}(K(z)|K, v) = K(c_i t^{\gamma_i} \mid i \in \mathbb{N})$ , and

- a)  $vK(z)$  is the group generated over  $vK$  by the elements  $\gamma_i$ ,
- b) if  $\gamma_i \in vK$  for all  $i \in \mathbb{N}$ , then  $K(z)v = Kv(c_i \mid i \in \mathbb{N})$ .

*Proof.* We derive a homogeneous sequence  $\mathfrak{S}$  from  $z$  as follows. We set  $a_0 = 0$ . If all  $c_i$  are in  $Kv$  and all  $\gamma_i$  are in  $vK$ , then we take  $\mathfrak{S}$  to be the empty sequence. Otherwise, having chosen  $i_j \in \mathbb{N}$  and defined

$$a_j := \sum_{1 \leq i \leq i_j} c_i t^{\gamma_i}$$

for all  $j < m$ , we proceed as follows. We let  $\ell$  be the first index in the power series  $z - a_{m-1}$  for which  $c_\ell t^{\gamma_\ell} \notin K(a_1, \dots, a_{m-1})$ ; if such an index does not exist, we let  $a_{m-1}$  be the last element of  $\mathfrak{S}$ . Otherwise, we set

$$i_m := \ell \quad \text{and} \quad a_m := \sum_{1 \leq i \leq i_m} c_i t^{\gamma_i}.$$

We have

$$c_\ell t^{\gamma_\ell} = a_m - a_{m-1} - d \quad \text{with} \quad d = \sum_{i_{m-1} < i < \ell} c_i t^{\gamma_i} \in K(a_1, \dots, a_{m-1}).$$

By assumption,  $e_\ell v c_\ell t^{\gamma_\ell} = e_\ell \gamma_\ell \in vK$ , and hence  $c := t^{-e_\ell \gamma_\ell} \in K$ . We have that  $c(c_\ell t^{\gamma_\ell})^{e_\ell} = c_\ell^{e_\ell}$ . Since  $c_\ell$  is separable-algebraic over  $Kv$ , the same holds for  $c_\ell^{e_\ell}$ .

Since  $v$  is trivial on  $k$ , the degree of  $c_\ell^{\text{e}_\ell} v$  over  $Kv$  is equal to that of  $c_\ell^{\text{e}_\ell}$  over  $K$ . It now follows by Lemma 5.3 that  $c_\ell t^{\gamma_\ell}$  is strongly homogeneous over  $K$ . By Lemma 5.2 it follows that it is also strongly homogeneous over the henselian field  $K(a_1, \dots, a_{m-1})$ . Therefore,  $a_m - a_{m-1}$  is homogeneous over  $K(a_1, \dots, a_{m-1})$ . Further,  $v(z - a_{m-1} - (a_m - a_{m-1})) = v(z - a_m) = \gamma_{\ell+1} > \gamma_\ell \geq v(z - a_{m-1})$ . This proves that  $a_m - a_{m-1}$  is a homogeneous approximation of  $z - a_{m-1}$  over  $K(a_0, \dots, a_{m-1})$ . By induction, we obtain a homogeneous sequence  $\mathfrak{S}$  for  $z$  in  $k((G))$ . It now follows from Lemma 5.7 that  $K(c_i t^{\gamma_i} \mid i \in \mathbb{N}) = K(a_j \mid j \in S) \subseteq K(z)^h$ ; thus,  $\gamma_i = v c_i t^{\gamma_i} \in vK(z)$  for all  $i \in \mathbb{N}$ . If  $\gamma_i \in vK$  and hence  $t^{\gamma_i} \in K$ , then  $c_i \in K(z)^h$ ; since the residue map is the identity on elements of  $k$ , this implies that  $c_i \in K(z)v$ .

Now assume that  $vK + \sum_{i=1}^{\infty} \mathbb{Z}\gamma_i/vK$  or  $Kv(c_i \mid i \in \mathbb{N})|Kv$  is infinite. Then  $\mathfrak{S}$  must be infinite, and it follows from Proposition 5.8 that  $z$  is a pseudo limit of the pseudo Cauchy sequence  $(a_i)_{i \in \mathbb{N}}$  of transcendental type. Thus,  $z$  is transcendental over  $K$  by Lemma 2.4. Theorem 5.9 now shows that

$$\text{IC}(K(z)|K, v) = K(c_i t^{\gamma_i} \mid i \in \mathbb{N})$$

and

$$vK(z) = vK(c_i t^{\gamma_i} \mid i \in \mathbb{N}), \quad K(z)v = K(c_i t^{\gamma_i} \mid i \in \mathbb{N})v.$$

Since  $K(c_i t^{\gamma_i} \mid i \in \mathbb{N}) \subseteq K(c_i, t^{\gamma_i} \mid i \in \mathbb{N})$ , for the proof of assertion a) it now suffices to show that the value group of the latter field is generated over  $vK$  by the  $\gamma_i$ , and that its residue field is generated over  $Kv$  by the  $c_i$ . As  $t^\gamma \in K$  for every  $\gamma \in vK$ , we have

$$(vK + \sum_{i=1}^{\ell} \mathbb{Z}\gamma_i : vK) = [K(t^{\gamma_i} \mid 1 \leq i \leq \ell) : K].$$

On the other hand,

$$[K(t^{\gamma_i} \mid 1 \leq i \leq \ell) : K] \geq (vK(t^{\gamma_i} \mid 1 \leq i \leq \ell) : vK) \geq (vK + \sum_{i=1}^{\ell} \mathbb{Z}\gamma_i : vK),$$

where the last inequality holds since  $\gamma_i \in vK(t^{\gamma_i} \mid 1 \leq i \leq \ell)$  for  $i = 1, \dots, \ell$ . We obtain that for all  $\ell \in \mathbb{N}$

$$vK(t^{\gamma_i} \mid 1 \leq i \leq \ell) = vK + \sum_{i=1}^{\ell} \mathbb{Z}\gamma_i$$

and

$$K(t^{\gamma_i} \mid 1 \leq i \leq \ell)v = Kv.$$

This implies that  $vK(t^{\gamma_i} \mid i \in \mathbb{N}) = vK + \sum_{i=1}^{\infty} \mathbb{Z}\gamma_i$  and  $K(t^{\gamma_i} \mid i \in \mathbb{N})v = Kv$ .

Set  $K' := K(t^{\gamma_i} \mid i \in \mathbb{N})$ . As  $K'v = Kv \subset K \subset K'$ , we have

$$[K'v(c_i \mid 1 \leq i \leq \ell) : K'v] \geq [K'(c_i \mid 1 \leq i \leq \ell) : K'].$$

On the other hand,

$$[K'(c_i \mid 1 \leq i \leq \ell) : K'] \geq [K'(c_i \mid 1 \leq i \leq \ell)v : K'v] \geq [K'v(c_i \mid 1 \leq i \leq \ell) : K'v],$$

where the last inequality holds since  $c_i \in K'(c_i \mid 1 \leq i \leq \ell)v$  for  $i = 1, \dots, \ell$ . We obtain  $K'(c_i \mid 1 \leq i \leq \ell)v = K'v(c_i \mid 1 \leq i \leq \ell)$  and  $vK'(c_i \mid 1 \leq i \leq \ell) = vK'$ .

This implies that

$$vK(c_i, t^{\gamma_i} \mid i \in \mathbb{N}) = vK'(c_i \mid i \in \mathbb{N}) = vK' = vK + \sum_{i=1}^{\infty} \mathbb{Z}\gamma_i$$

and

$$K(c_i, t^{\gamma_i} \mid i \in \mathbb{N})v = K'(c_i \mid i \in \mathbb{N})v = K'v(c_i \mid i \in \mathbb{N}) = Kv(c_i \mid i \in \mathbb{N}).$$

This proves assertions a) and b).  $\square$

*Remark 6.2.* Assertions a) and b) of the previous theorem will also hold if  $K = Kv((vK))$ . Indeed, if  $G_0$  denotes the subgroup of  $G$  generated by the  $\gamma_i$  over  $Kv$ , and  $k_0 = Kv(c_i \mid i \in \mathbb{N})$ , then  $K(z) \subset k_0((G_0))$ , and therefore  $vK(z) \subseteq G_0$  and  $K(z)v \subseteq k_0$ .

Our methods also yield an alternative proof of the following well known fact:

**Theorem 6.3.** *The algebraic closure of the field  $\mathbb{Q}_p$  of  $p$ -adic numbers is not complete.*

*Proof.* Choose any compatible system of  $n$ -th roots  $p^{1/n}$  of  $p$ , that is, such that  $(p^{1/mn})^m = p^{1/n}$  for all  $m, n \in \mathbb{N}$ . For  $i \in \mathbb{N}$ , choose any  $n_i \in \mathbb{N}$  not divisible by  $p$ , and set  $\gamma_i := i + \frac{1}{n_i}$  if  $n_i > 1$ , and  $\gamma_i = i$  otherwise. Further, choose  $c_i$  in some fixed set of representatives in  $\widetilde{\mathbb{Q}_p}$  of its residue field  $\widetilde{\mathbb{F}_p}$  such that the degree of  $c_i^{n_i}$  over  $\mathbb{Q}_p$  is equal to the degree of  $c_i^{n_i}v_p$  over  $\mathbb{Q}_p v_p = \mathbb{F}_p$ . Then set

$$(6.1) \quad b_i := \sum_{1 \leq j \leq i} c_j p^{\gamma_j}.$$

Since  $\gamma_i < \gamma_{i+1}$  for all  $i$  and the sequence  $(\gamma_i)_{i \in \mathbb{N}}$  is cofinal in the value group  $\mathbb{Q}$  of  $\widetilde{\mathbb{Q}_p}$ , the sequence  $(b_i)_{i \in \mathbb{N}}$  is a Cauchy sequence in  $(\widetilde{\mathbb{Q}_p}, v_p)$ . If this field were complete, it would contain a pseudo limit  $z$  to every such Cauchy sequence. On the other hand, as in the proof of Theorem 6.1 one shows that  $\mathbb{Q}_p(c_i p^{\gamma_i} \mid i \in \mathbb{N}) \subseteq \mathbb{Q}_p(z)^h$ , and that

- a)  $v\mathbb{Q}_p(z)$  is the group generated over  $\mathbb{Z}$  by the elements  $\gamma_i$ ,
- b) if  $\gamma_i \in \mathbb{Z}$  for all  $i \in \mathbb{N}$ , then  $\mathbb{Q}_p(z)v = \mathbb{F}_p(c_i v_p \mid i \in \mathbb{N})$ , and
- c) if  $v\mathbb{Q}_p(z)/\mathbb{Z}$  or  $\mathbb{F}_p(c_i v_p \mid i \in \mathbb{N})/\mathbb{F}_p$  is infinite, then  $z$  is transcendental over  $\mathbb{Q}_p$ .

Hence, if we choose  $(n_i)_{i \in \mathbb{N}}$  to be a strictly increasing sequence and  $c_i = 1$  for all  $i$ , or if we choose  $n_i = 1$  for all  $i$  and the elements  $c_i$  of increasing degree over  $\mathbb{Q}_p$ , then  $z$  will be transcendental over  $\mathbb{Q}_p$ . Since  $z$  lies in the completion of  $\widetilde{\mathbb{Q}_p}$ , we have now proved that this completion is transcendental over  $\mathbb{Q}_p$ .  $\square$

With the same method, we can also prove another well known result:

**Theorem 6.4.** *The completion  $\mathbb{C}_p$  of  $\widetilde{\mathbb{Q}_p}$  admits a pseudo Cauchy sequence without a pseudo limit in  $\mathbb{C}_p$ . Hence,  $\mathbb{C}_p$  is not maximal and not spherically complete.*

*Proof.* In the same setting as in the foregoing proof, we now choose  $c_i = 1$  for all  $i$ . Further, we choose  $(n_i)_{i \in \mathbb{N}}$  to be a strictly increasing sequence and set  $\gamma_i := 1 - \frac{1}{n_i}$ . Then  $(b_i)_{i \in \mathbb{N}}$  is a pseudo Cauchy sequence. Suppose it admitted a pseudo limit  $y$  in  $\mathbb{C}_p$ . Then, using that  $\widetilde{\mathbb{Q}_p}$  is dense in  $\mathbb{C}_p$ , we could choose  $z \in \widetilde{\mathbb{Q}_p}$  such that  $v_p(y - z) \geq 1$ . Since  $1 > \gamma_i$  for all  $i$ , it would follow that also  $z$  is a pseudo limit

of  $(b_i)_{i \in \mathbb{N}}$ . But as in the foregoing proof one shows that  $z$  must be transcendental over  $\mathbb{Q}_p$ . This contradiction shows that  $(b_i)_{i \in \mathbb{N}}$  cannot have a pseudo limit in  $\mathbb{C}_p$ . By the results of [KA], this implies that  $\mathbb{C}_p$  admits a proper immediate extension, which shows that  $\mathbb{C}_p$  is not maximal. With the elements  $b_i$  defined as in (6.1), we also find that the intersection of the nest  $(\{a \in \mathbb{C}_p \mid v(a - b_i) \geq \gamma_{i+1}\})_{i \in \mathbb{N}}$  of balls is empty. Hence,  $\mathbb{C}_p$  is not spherically complete.  $\square$

## REFERENCES

- [AP] Alexandru, V. – Popescu, N.: *Sur une classe de prolongements à  $K(X)$  d'une valuation sur une corps  $K$* , Rev. Roumaine Math. Pures Appl. **33** (1988), 393–400 MR 89h:12015
- [APZ1] Alexandru, V. – Popescu, N. – Zaharescu, A.: *A theorem of characterization of residual transcendental extensions of a valuation*, J. of Math. Kyoto Univ. **28** (1988), 579–592 MR 90c:12011
- [APZ2] Alexandru, V. – Popescu, N. – Zaharescu, A.: *Minimal pairs of definition of a residual transcendental extension of a valuation*, J. of Math. Kyoto Univ. **30** (1990), 207–225 MR 92b:12014
- [APZ3] Alexandru, V. – Popescu, N. – Zaharescu, A.: *All valuations on  $K(X)$* , J. of Math. Kyoto Univ. **30** (1990), 281–296 MR 92c:12011
- [B] Bourbaki, N.: *Commutative algebra*, Paris (1972)
- [C] Cutkosky, D.: *Local factorization and monomialization of morphisms*, Astérisque **260** (1999) MR 2001c:14027
- [CP] Cutkosky, D. – Piltant, O.: *Ramification of valuations*, to appear in: Advances in Mathematics
- [E] Endler, O.: *Valuation theory*, Springer, Berlin (1972) MR 50:9847
- [GMP1] Green, B. – Matignon, M. – Pop, F.: *On valued function fields I*, manuscripta math. **65** (1989), 357–376 MR 91g:12010
- [GMP2] Green, B. – Matignon, M. – Pop, F.: *On valued function fields II, III*, J. reine angew. Math. **412** (1990), 128–149; **432** (1992), 117–133
- [KA] Kaplansky, I.: *Maximal fields with valuations I*, Duke Math. Journ. **9** (1942), 303–321 MR 3:264d
- [KH1] Khanduja, S. K.: *Value groups and simple transcendental extensions*, Mathematika **38** (1991), 381–385 MR 93c:12016
- [KH2] Khanduja, S. K.: *Prolongations of valuations to simple transcendental extensions with given residue field and value group*, Mathematika **38** (1991), 386–390 MR 93c:12017
- [KH3] Khanduja, S. K.: *On valuations of  $K(X)$* , Proc. Edinburgh Math. Soc. **35** (1992), 419–426 MR 94a:12008
- [KH4] Khanduja, S. K.: *A uniqueness problem in simple transcendental extensions of valued fields*, Proc. Edinburgh Math. Soc. **37** (1993), 13–23 MR 95b:12010
- [KH5] Khanduja, S. K.: *On value groups and residue fields of some valued function fields*, Proc. Edinburgh Math. Soc. **37** (1994), 445–454 MR 95h:12006
- [KH6] Khanduja, S. K.: *A note on residually transcendental prolongations with uniqueness property*, J. Math. Kyoto Univ. **36** (1996), 553–556 MR 97j:12007
- [KH7] Khanduja, S. K.: *On residually transcendental valued function fields of conics*, Glasgow Math. J. **38** (1996), 137–145 MR 98a:12007
- [KH8] Khanduja, S. K.: *On extensions of valuations with prescribed value groups and residue fields*, J. Indian Math. Soc. **62** (1996), 57–60 MR 98c:12013
- [KH9] Khanduja, S. K.: *Valued function fields with given genus and residue field*, J. Indian Math. Soc. **63** (1997), 115–121 MR 99c:12008
- [KH10] Khanduja, S. K.: *An independence theorem in simple transcendental extensions of valued fields*, J. Indian Math. Soc. **63** (1997), 243–248 MR 99g:12008
- [KH11] Khanduja, S. K.: *Tame fields and tame extensions*, J. Alg. **201** (1998), 647–655 MR 99a:12010



- [KH12] Khanduja, S. K.: *The minimum property for Krasner's constant*, in: Valuation Theory and its Applications, Proceedings of the Valuation Theory Conference Saskatoon 1999, Volume I, eds. F.-V. Kuhlmann, S. Kuhlmann and M. Marshall, Fields Institute Communications **32**, Amer. Math. Soc. (2002) MR 2003c:12001
- [KHG1] Khanduja, S. K. – Garg, U.: *On extensions of valuations to simple transcendental extensions*, Proc. Edinburgh Math. Soc. **32** (1989), 147–156 MR 90e:12019
- [KHG2] Khanduja, S. K. – Garg, U.: *Rank 2 valuations of  $K(x)$* , Mathematika **37** (1990), 97–105 MR 91j:12016
- [KHG3] Khanduja, S. K. – Garg, U.: *On the rank of extensions of valuations*, Colloq. Math. **59** (1990), 25–29 MR 92c:12008
- [KHG4] Khanduja, S. K. – Garg, U.: *On residually generic prolongations to a simple transcendental extension*, J. Indian Math. Soc. **57** (1990), 1–8 MR 93g:12010
- [KHG5] Khanduja, S. K. – Garg, U.: *Residue fields of valued function fields of conics*, Proc. Edinburgh Math. Soc. **36** (1993), 469–478 MR 94i:12002
- [KHG6] Khanduja, S. K. – Garg, U.: *Prolongations of a Krull valuation to a simple transcendental extension*, J. Indian Math. Soc. **59** (1993), 13–21 MR 94j:12012
- [KHPR] Khanduja, S. K. – Popescu, N. – Roggenkamp, K. W.: *On minimal pairs and residually transcendental extensions of valuations*, to appear in: Mathematika
- [KHS] Khanduja, S. K. – Saha, J.: *A uniqueness problem in valued function fields of conics*, Bull. London Math. Soc. **28** (1996), 455–462 MR 98a:12008
- [KKU1] Knaf, H. – Kuhlmann, F.-V.: *Abhyankar places admit local uniformization in any characteristic*, submitted, available at:  
<http://math.usask.ca/fvk/Valth.html>
- [KKU2] Knaf, H. – Kuhlmann, F.-V.: *Every place admits local uniformization in a finite extension of the function field*, preprint available at:  
<http://math.usask.ca/fvk/Valth.html>
- [KU1] Kuhlmann, F.-V.: *Henselian function fields and tame fields*, preprint (extended version of Ph.D. thesis), Heidelberg (1990)
- [KU2] Kuhlmann, F.-V.: *Valuation theory of fields, abelian groups and modules*, to appear in the “Algebra, Logic and Applications” series (formerly Gordon and Breach, eds. A. Macintyre and R. Göbel). Preliminary versions of several chapters available at:  
<http://math.usask.ca/~fvk/Fvkbook.htm>
- [KU3] Kuhlmann, F.-V.: *Valuation theoretic and model theoretic aspects of local uniformization*, in: Resolution of Singularities - A Research Textbook in Tribute to Oscar Zariski. Herwig Hauser, Joseph Lipman, Frans Oort, Adolfo Quiros (eds.), Progress in Mathematics Vol. **181**, Birkhäuser Verlag Basel (2000), 381–456 MR 2001c:14001
- [KU4] Kuhlmann, F.-V.: *The model theory of tame valued fields*, in preparation
- [KU5] Kuhlmann, F.-V.: *Algebraic independence of elements in completions and maximal immediate extensions of valued fields*, in preparation
- [KUKMZ] Kuhlmann, F.-V. – Kuhlmann, S. – Marshall, M. – Zekavat, M.: *Embedding ordered fields in formal power series fields*, J. Pure Appl. Algebra **169** (2002), 71–90 MR 2002m:12003
- [KUPR] Kuhlmann, F.-V. – Pank, M. – Roquette, P.: *Immediate and purely wild extensions of valued fields*, manuscripta mathematica **55** (1986), 39–67 MR 87d:12012
- [KW] Knebusch, M. – Wright, M.: *Bewertungen mit reeller Henselisierung*, J. reine angew. Math. **286/287** (1976), 314–321 MR 54:7440
- [L] Lam, T. Y.: *The theory of ordered fields*, in: Ring Theory and Algebra III (ed. B. McDonald), Lecture Notes in Pure and Applied Math. **55**, Dekker, New York (1980), 1–152 MR 82e:12033
- [M] MacLane, S.: *A construction for absolute values in polynomial rings*, Trans. Amer. Math. Soc. **40** (1936), 363–395
- [MO1] Matignon, M. – Ohm, J.: *A structure theorem for simple transcendental extensions of valued fields*, Proc. Amer. Math. Soc. **104** (1988), 392–402 MR 90h:12011
- [MO2] Matignon, M. – Ohm, J.: *Simple transcendental extensions of valued fields III: The uniqueness property*, J. of Math. Kyoto Univ. **30** (1990), 347–366 MR 91j:12017
- [MS] MacLane, S. – Schilling, O.F.G.: *Zero-dimensional branches of rank 1 on algebraic varieties*, Annals of Math. **40** (1939), 507–520 MR 1:26c

- [MOSW1] Mosteig, E. – Sweedler, M.: *Well-ordered valuations on characteristic zero rational function fields of transcendence degree two*, manuscript (2001), available at: <http://math.usask.ca/fvk/Valth.html>
- [MOSW2] Mosteig, E. – Sweedler, M.: *Valuations and Filtrations*, J. Symbolic Comput. **34** (2002), 399–435 MR 2003j:12008
- [N] Nagata, M.: *A theorem on valuation rings and its applications*, Nagoya Math. J. **29** (1967), 85–91 MR 34:7503
- [O1] Ohm, J.: *Simple transcendental extensions of valued fields*, J. of Math. Kyoto Univ. **22** (1982), 201–221 MR 84a:13005
- [O2] Ohm, J.: *The Ruled Residue Theorem for simple transcendental extensions of valued fields*, Proc. Amer. Math. Soc. **89** (1983), 16–18 MR 85b:12010
- [O3] Ohm, J.: *Simple transcendental extensions of valued fields II: A fundamental inequality*, J. of Math. Kyoto Univ. **25** (1985), 583–596 MR 87c:12013
- [PL] Polzin, M.: *Prolongement de la valeur absolue de Gauss et problème de Skolem*, Bull. Soc. Math. France **116** (1988), 103–132 MR 90e:12020
- [PP] Popescu, L. – Popescu, N.: *On the residual transcendental extensions of a valuation. Key polynomials and augmented valuation*, Tsukuba J. Math. **1** (1991), 57–78 MR 92h:12008
- [PR] Prestel, A.: *Lectures on Formally Real Fields*, Lecture Notes in Math. **1093**, Springer, Berlin–Heidelberg–New York–Tokyo (1984) MR 86h:12013
- [R] Ribenboim, P.: *Théorie des valuations*, Les Presses de l’Université de Montréal, Montréal, 1st ed. (1964), 2nd ed. (1968) MR 40:2670
- [S] Spivakovsky, M.: *Valuations on function fields of surfaces*, Amer. J. Math. **112** (1990), 107–156 MR 91c:14037
- [SW] Sweedler, M.: *Ideal Bases and Valuation Rings*, manuscript (1986), available at: <http://myweb.lmu.edu/faculty/emosteig/>
- [V] Vaquié, M.: *Extension d’une valuation*, preprint (2002), available at: <http://math.usask.ca/fvk/Valth.html>
- [ZS] Zariski, O. – Samuel, P.: *Commutative Algebra*, Vol. II, Springer, New York–Heidelberg–Berlin (1960) MR 22:11006

MATHEMATICAL SCIENCES GROUP, UNIVERSITY OF SASKATCHEWAN, 106 WIGGINS ROAD, SASKATOON, SASKATCHEWAN, CANADA S7N 5E6  
*E-mail address:* [fvk@math.usask.ca](mailto:fvk@math.usask.ca)