

COMPLETE NONORIENTABLE MINIMAL SURFACES IN A BALL OF \mathbb{R}^3

F. J. LÓPEZ, FRANCISCO MARTIN, AND SANTIAGO MORALES

ABSTRACT. The existence of complete minimal surfaces in a ball was proved by N. Nadirashvili in 1996. However, the construction of such surfaces with nontrivial topology remained open. In 2002, the authors showed examples of complete orientable minimal surfaces with arbitrary genus and one end. In this paper we construct complete bounded nonorientable minimal surfaces in \mathbb{R}^3 with arbitrary finite topology. The method we present here can also be used to construct *orientable* complete minimal surfaces with arbitrary genus and *number of ends*.

1. INTRODUCTION

This work is part of a series of papers about the admissible topological type of a complete bounded minimal surface in \mathbb{R}^3 . The first example of such a surface, due to Nadirashvili [5], had the topology of a disk. His proof consisted of successive modifications of the Weierstrass representation of a minimal surface that involved the use of the classic Runge's theorem in the plane. Since Nadirashvili's result, examples with more complicated topological type were constructed: annuli in [4], and orientable surfaces with finite genus and one end in [2]. The second family of examples was obtained by combining Nadirashvili's ideas with a technique that solves period problems for minimal surfaces with arbitrary genus. In this paper we extend this kind of result to the nonorientable case. Actually, we have proved:

Theorem. *For any genus $\sigma \geq 1$ and any $k \in \mathbb{N}$, there exists a complete bounded nonorientable minimal surface in \mathbb{R}^3 with genus σ and k ends.*

Similar techniques to those developed in [5] and [2] have been applied to the orientable double covering of our nonorientable surfaces. However, in this case the Weierstrass have to satisfy a compatibility condition with the reversing orientable involution. This fact introduces additional difficulties in the treatment of the period problem. To overcome these obstacles, we have had to use, among other things, a general version of Runge's theorem on Riemann surfaces [8].

We would like to point out that the present method can also be used to construct *orientable* complete minimal surfaces with arbitrary genus and *number of ends*.

Received by the editors May 24, 2004.

2000 *Mathematics Subject Classification.* Primary 53A10; Secondary 49Q05, 49Q10, 53C42.

Key words and phrases. Complete bounded minimal surfaces, nonorientable minimal immersions.

This research was partially supported by MEC-FEDER Grant No. MTM2004-00160.

©2006 American Mathematical Society
Reverts to public domain 28 years from publication

2. BACKGROUND AND NOTATION

In this section, we provide some definition and notation that we will use throughout the paper.

2.1. Metrics and divisors on a Riemann surface. Let \mathcal{N} and $d\hat{s}^2$ be a Riemann surface and a Riemannian conformal metric on \mathcal{N} , respectively. Given a curve α in \mathcal{N} , by $\text{length}(\alpha, d\hat{s})$ we mean the length of α with the metric $d\hat{s}^2$. Given a subset $W \subset \mathcal{N}$, we define:

- $\text{dist}_{(d\hat{s},W)}(p, q) = \inf\{\text{length}(\alpha, d\hat{s}) \mid \alpha : [0, 1] \rightarrow W, \alpha(0) = p, \alpha(1) = q\}$, for any $p, q \in W$,
- $\text{dist}_{(d\hat{s},W)}(T_1, T_2) = \inf\{\text{dist}_{(d\hat{s},W)}(p, q) : p \in T_1, q \in T_2\}$, for any $T_1, T_2 \subset W$,
- $\text{diam}_{d\hat{s}}(W) = \sup\{\text{dist}_{(d\hat{s},W)}(p, q) : p, q \in W\}$.

The concepts of the (multiplicative) divisor on \mathcal{N} , the integral divisor on \mathcal{N} , and the natural partial ordering, \geq , on divisors can be found in [1]. Let ω be a meromorphic function or 1-form on \mathcal{N} . Let $W \subset \mathcal{N}$ and suppose that ω has a finite number of zeroes, z_1, \dots, z_n , and a finite number of poles, p_1, \dots, p_m , in W . We denote by $(\omega|_W)_0 = z_1 \cdots z_n$, $(\omega|_W)_\infty = p_1 \cdots p_m$, and $(\omega|_W) = (\omega|_W)_0 / (\omega|_W)_\infty$, the zero divisor, the polar divisor, and the divisor of ω on W , respectively. When $W = \mathcal{N}$, we simply write (ω) , $(\omega)_0$, and $(\omega)_\infty$, respectively.

2.2. The conformal structure. In this subsection, we describe the family of Riemann surfaces that we will deal with in the next sections.

Let $\sigma \geq 0$ be an integer number, let $d_1, \dots, d_{\sigma+1}$ be a sequence of pairwise distinct complex numbers satisfying $\text{Im } d_i \neq 0$, $i = 1, \dots, \sigma + 1$, and let \overline{M} be the algebraic hyperelliptic curve of genus σ given by

$$\overline{M} = \left\{ (z, w) \in \overline{\mathbb{C}}^2 : w^2 = \prod_{i=1}^{\sigma+1} (z - d_i)(z - \overline{d}_i) \right\}.$$

Let $I : \overline{M} \rightarrow \overline{M}$ be the antiholomorphic involution without fixed points on \overline{M} , given by $I(z, w) = (\overline{z}, -\overline{w})$, and label $A(z, w) = (z, -w)$ the hyperelliptic involution.

In what follows Δ will be an open bounded simply-connected domain in \mathbb{C} whose boundary is a Jordan curve, and such that $\{d_1, \dots, d_{\sigma+1}\} \subset \Delta$. Observe that $z^{-1}(\overline{\mathbb{C}} \setminus \overline{\Delta}) = D_+ \cup D_-$, where D_\pm is a conformal disk around $\infty^\pm = (\infty, \pm\infty)$, and $z|_{D_\pm}$ is one to one. We denote $f_\pm = z|_{D_\pm}^{-1}$. Let $\{\beta_1, \dots, \beta_{k-1}\} \subset \mathbb{C} \setminus \overline{\Delta}$ be a set of real numbers and label $\beta_i^\pm = f_\pm(\beta_i)$, $i = 1, \dots, k - 1$. We denote $M = \overline{M} - \{\beta_1^\pm, \dots, \beta_{k-1}^\pm, \infty^\pm\}$.

Now, we are going to describe the domains of M that we will deal with. Given α as a Jordan curve in \mathbb{C} , we denote $\text{Int } \alpha$ as the bounded connected component of $\mathbb{C} \setminus \alpha$.

Definition 1. By a multigon, we mean a finite family of simple closed polygonal curves in \mathbb{C} , $\mathfrak{P} = \{P_1, \dots, P_k\}$, pairwise disjoint, such that:

- $\overline{\Delta} \subset \text{Int } P_k$,
- $\overline{\text{Int } P_i} \subset \text{Int } P_k \setminus \overline{\Delta}$, $i = 1, \dots, k - 1$,
- $\beta_i \in \text{Int } P_i$, $i = 1, \dots, k - 1$,
- $\overline{\text{Int } P_1}, \dots, \overline{\text{Int } P_{k-1}}$ are pairwise disjoint.

We denote $\text{Int } \mathfrak{P}$ as the domain of \mathbb{C} whose boundary consists of the curves in \mathfrak{P} . See Figure 1.

If \mathfrak{P} is a multigon, we write $D(\mathfrak{P}) \subset M$ as the domain whose boundary coincides with the union of the Jordan curves $f_+(\mathfrak{P})$ and $I(f_+(\mathfrak{P}))$.

Given $\xi > 0$, small enough, we define a new multigon $\mathfrak{P}^\xi = \{P_1^\xi, \dots, P_k^\xi\}$, where $P_i^\xi, i = 1, \dots, k$, are the parallel polygonal curves to P_i , satisfying that the distance between parallel sides is equal to ξ , and $\overline{\text{Int } \mathfrak{P}^\xi} \subset \text{Int } \mathfrak{P}$. Whenever we write \mathfrak{P}^ξ in the paper we are assuming that ξ is small enough to \mathfrak{P}^ξ were a multigon. Observe that $\overline{D(\mathfrak{P}^\xi)} \subset D(\mathfrak{P})$.

2.3. The Weierstrass representation of a nonorientable minimal surface.

Finally, we introduce the Weierstrass representation of nonorientable minimal surfaces.

Given $D \subset \overline{M}$ a domain, we will say that a function, or a 1-form, is harmonic, holomorphic, meromorphic, ... on \overline{D} , if it is harmonic, holomorphic, meromorphic, ... on a domain containing \overline{D} .

Let $D \subset M$ be a domain invariant under I , and let $\Phi = (\Phi_1, \Phi_2, \Phi_3)$ be the Weierstrass representation of a minimal immersion $X : \overline{D} \rightarrow \mathbb{R}^3$ such that $I^*\Phi = \overline{\Phi}$. As $X \circ I = X$, then it induces a minimal immersion $\mathcal{X} : \overline{\mathcal{D}} \rightarrow \mathbb{R}^3$, where $\overline{\mathcal{D}} = \overline{D}/\langle I \rangle$ is a nonorientable surface with boundary. Obviously $X = \mathcal{X} \circ \pi$, where $\pi : \overline{D} \rightarrow \overline{\mathcal{D}}$ is the canonical projection.




FIGURE 1. The multigon \mathfrak{P} and the set Δ .

We can write $\Phi_j = \varphi_j(z, w) \frac{dz}{w}$, where φ_j is a holomorphic function on \overline{D} , $j = 1, 2, 3$, and denote $\varphi \stackrel{\text{def}}{=} (\varphi_1, \varphi_2, \varphi_3)$. Note that $\varphi \circ I = -\overline{\varphi}$. With this notation, if we write the Riemannian metric induced by X as $ds_X^2 = \lambda_X^2 \|\frac{dz}{w}\|^2$, then

$$(1) \quad \lambda_X = \frac{1}{\sqrt{2}} \|\varphi\| = \frac{1}{\sqrt{2}} \sqrt{|\varphi_1|^2 + |\varphi_2|^2 + |\varphi_3|^2} > 0 \quad \text{on } \overline{D}.$$

For the sake of simplicity, given $W \subset M$, $p, q \in W$ and $T \subset W$, we write $\text{dist}_{(X,W)}(p, q)$ and $\text{dist}_{(X,W)}(p, T)$ instead of $\text{dist}_{(ds_X,W)}(p, q)$ and $\text{dist}_{(ds_X,W)}(p, T)$, respectively.

3. PREVIOUS LEMMAS

Using the previous notations, a precise statement of our main result is the following.

Theorem 1. *For any $\sigma \geq 0$ and $k \geq 1$, there exists a complete bounded minimal immersion $\mathcal{X} : \mathcal{D} \rightarrow \mathbb{R}^3$, where $D \subset M$, $I(D) = D$, $\mathcal{D} = D/\langle I \rangle$ and \mathcal{D} is homeomorphic to a nonorientable compact surface of genus $\sigma + 1$ minus k points.*

In order to get this theorem, we need the following two lemmas.

Lemma 1. *Consider a multigon \mathfrak{P} and a minimal immersion $F : \overline{D(\mathfrak{P})} \rightarrow \mathbb{R}^3$ whose Weierstrass representation Φ satisfies $I^*(\Phi) = \overline{\Phi}$. Let K_1, K_2 be compact subsets in $D(\mathfrak{P})$ satisfying the facts that K_1, K_2 and $I(K_1)$ are pairwise disjoint, $I(K_2) = K_2$, $D(\mathfrak{P}^\vartheta) \subset K_2$, for some $\vartheta > 0$, and the set $\{\beta_1^\pm, \dots, \beta_{k-1}^\pm, \infty^\pm\}$ has a (unique) point in each connected component of $\overline{M} \setminus (K_1 \cup K_2 \cup I(K_1))$.*

Then for any $\alpha > 0$ there exists $h : \overline{D(\mathfrak{P})} \rightarrow \mathbb{C}$, a holomorphic function without zeroes, such that:

1. $h \circ I = 1/\overline{h}$,
2. $|h - \alpha| < 1/\alpha$ in K_1 ,
3. $|h - 1| < 1/\alpha$ in K_2 ,
4. the Weierstrass data $\tilde{\Phi}$ given by $\tilde{g} = g/h$ and $\tilde{\Phi}_3 = \Phi_3$ satisfy $I^*(\tilde{\Phi}) = \overline{\tilde{\Phi}}$, and the associated minimal immersion $\tilde{F} : \overline{D(\mathfrak{P})} \rightarrow \mathbb{R}^3$ is well defined.

Lemma 2. *Let \mathfrak{P} be a multigon and let $r > 0$. Consider a minimal immersion $X : \overline{D(\mathfrak{P})} \rightarrow \mathbb{R}^3$ satisfying:*

1. $X = \text{Re} \left(\int_{p_0} \Phi \right)$, where $p_0 = (d_1, 0)$ and $I^*\Phi = \overline{\Phi}$;
2. $\|X\| < r$ in $\overline{D(\mathfrak{P})}$.

Then, for any $\varepsilon, s > 0$ such that \mathfrak{P}^ε is a multigon, there exist a multigon $\tilde{\mathfrak{P}}$ and a conformal minimal immersion $Y : D(\tilde{\mathfrak{P}}) \rightarrow \mathbb{R}^3$, such that:

1. $\overline{D(\mathfrak{P}^\varepsilon)} \subset D(\tilde{\mathfrak{P}}) \subset \overline{D(\tilde{\mathfrak{P}})} \subset D(\mathfrak{P})$,
2. $Y = \text{Re} \left(\int_{p_0} \tilde{\Phi} \right)$, where $\tilde{\Phi}$ satisfies $I^*(\tilde{\Phi}) = \overline{\tilde{\Phi}}$,
3. $\text{dist}_{(Y, D(\tilde{\mathfrak{P}}))} \left(\partial(D(\tilde{\mathfrak{P}})), \partial(D(\mathfrak{P}^\varepsilon)) \right) > s$,
4. $Y(D(\tilde{\mathfrak{P}})) \subset B_R$, $R = \sqrt{r^2 + (2s)^2} + \varepsilon$,
5. $\|Y - X\| < \varepsilon$ in $D(\mathfrak{P}^\varepsilon)$.

3.1. Proof of Lemma 1. First, we construct a basis $\mathcal{B} = \{\gamma_1, \dots, \gamma_{\sigma+k}, \Gamma_1, \dots, \Gamma_{\sigma+k-1}\}$ of $H_1(D(\mathfrak{P}), \mathbb{R})$ contained in K_2 , and satisfying $I_*(\gamma_j) = \gamma_j$ and $I_*(\Gamma_j) = -\Gamma_j$.

To do this, consider Jordan curves $\hat{c}_1, \dots, \hat{c}_\sigma, c_1, \dots, c_{\sigma+k}$ in the z -plane satisfying:

- The winding number of \hat{c}_j around d_j and d_{j+1} is 1, and the winding number around $\beta_l, \overline{d_h}$ and d_i is 0, $l = 1, \dots, k - 1$, $h = 1, \dots, \sigma + 1$, and $i = 1, \dots, j - 1, j + 2, \dots, \sigma + 1$.
- If $j \leq \sigma$ the winding number of c_j around d_i and $\overline{d_i}$ is $\delta_{i,j}$, $i = 1, \dots, \sigma + 1$, and the winding number around β_l is 0, $l = 1, \dots, k - 1$.

- If $\sigma < j < \sigma+k$ the winding number of c_j around β_i is $\delta_{i,j-k}$, $i = 1, \dots, k-1$, and the winding number around d_l, \bar{d}_l is 0, $l = 1, \dots, \sigma+1$.
- The winding number of $c_{\sigma+k}$ around β_i, d_j, \bar{d}_j is -1 , $i = 1, \dots, k-1, j = 1, \dots, \sigma+1$.

Take $\hat{\alpha}_j$ and α_j as curves in K_2 obtained by lifting \hat{c}_j and c_j , respectively, for any j . Then, we define $\gamma_j \stackrel{\text{def}}{=} \frac{1}{2}(\alpha_j + I_*(\alpha_j))$, for $j = 1, \dots, \sigma+k$, and

$$\Gamma_j = \begin{cases} \frac{1}{2}(\hat{\alpha}_j - I_*(\hat{\alpha}_j)), & j = 1, \dots, \sigma, \\ \frac{1}{2}(\alpha_j - I_*(\alpha_j)), & j = \sigma+1, \dots, \sigma+k-1. \end{cases}$$

Note that $\gamma_j = \alpha_j, j = 1, \dots, \sigma$. It is not hard to check that \mathcal{B} is indeed a basis of $H_1(D(\mathfrak{P}), \mathbb{R})$, satisfying the required conditions.




FIGURE 2. The z -projection of the curves in \mathcal{B} .

To prove our theorem we will need to close the period problem for holomorphic 1-forms on $D(\mathfrak{P}), \Phi$, satisfying $I^*(\Phi) = \bar{\Phi}$. In this case we have the following claim.

Claim 1. *If τ is a holomorphic differential on $D(\mathfrak{P})$ satisfying $I^*(\tau) = \bar{\tau}$, then $\text{Re} \left(\int_{\gamma} \tau \right) = 0, \forall \gamma \in H_1(D(\mathfrak{P}), \mathbb{R})$ if, and only if, $\int_{\gamma_j} \tau = 0, j = 1, \dots, \sigma+k$.*

In particular, if in addition τ is holomorphic in \overline{M} , then $\tau = 0$ if, and only if, $\int_{\gamma_j} \tau = 0, j = 1, \dots, \sigma$.

Proof. Observe that $\int_{\gamma} \tau = \overline{\int_{I_*(\gamma)} \tau}$. From the choice of \mathcal{B} , the first part of the claim trivially holds. For the second part, take into account that a holomorphic differential on \overline{M} vanishes if, and only if, it has imaginary periods. \square

We denote $\mathcal{V} = \{\omega \text{ holomorphic differentials on } M\}$. It is clear that $\omega \in \mathcal{V}$ if, and only if, $\overline{I^*\omega} \in \mathcal{V}$. Define the following real subspaces of \mathcal{V} :

$$\mathcal{V}^+ = \{\tau \in \mathcal{V} / I^*\tau = \bar{\tau}\}, \quad \mathcal{V}^- = \{\tau \in \mathcal{V} / I^*\tau = -\bar{\tau}\}.$$

Claim 2. *Let p be a point in M and $\omega \in \mathcal{V}^-$ satisfying $A^*\omega = -\omega$. Then it is possible to construct a holomorphic function $H : M \rightarrow \mathbb{C}$ such that:*

- (i) $H \circ I = -\overline{H}$, $H \circ A = -H$,
- (ii) $(\omega + dH)_0 \geq (\omega)_0 \cdot p \cdot I(p) \cdot A(p) \cdot I(A(p))$.

Proof. From the hypotheses of the claim we know that $\omega = \frac{P(z)}{Q(z)} \frac{dz}{w}$, where $P(z)$ and $Q(z)$ are polynomial with real coefficients and the roots of Q belong to the set $\{\beta_1, \dots, \beta_{k-1}\}$. Assume that $(\omega)_0 = p^n \cdot I(p)^n \cdot A(p)^n \cdot I(A(p))^n \cdot E$, $n \geq 0$, where E is an integral divisor not containing the points $p, I(p), A(p)$ and $I(A(p))$. Define:

$$J(z, w) = \begin{cases} \frac{P(z)^2}{(z-z(p))^{n-1}(z-z(p))^{n-1}} w, & \text{if } p \neq A(p) \text{ and } p \neq I(A(p)), \\ \frac{P(z)^2}{(z-z(p))^{n-1}} w, & \text{if } p \neq A(p) \text{ and } p = I(A(p)), \\ \frac{P(z)^2}{(z-z(p))^n(z-z(p))^n} w, & \text{if } p = A(p). \end{cases}$$

It is clear that J satisfies (i) and that it has zeroes at $p, I(p), A(p)$ and $I(A(p))$ of order $n + 1$. Furthermore the divisor E^2 is contained in $(J)_0$. Then it is possible to find $\lambda \in \mathbb{R}$ such that $(\omega + \lambda dJ)_0 \geq (\omega)_0 \cdot p \cdot I(p) \cdot A(p) \cdot I(A(p))$. This completes the proof. □

Claim 3. *Consider $(b_1, \dots, b_{\sigma+k}) \in \mathbb{R}^{\sigma+k} - \{(0, \dots, 0)\}$ and $c = \sum_{j=1}^{\sigma+k} b_j \gamma_j$. Then there exists $\tau \in \mathcal{V}^-$ satisfying $\text{Im}(\int_c \tau) \neq 0$.*

Furthermore, if L is an integral divisor invariant under I and whose support is in $D(\mathfrak{P})$, then τ can be chosen in such a way that $(\tau)_0 \geq L$.

Proof. We consider the linear map $f : \mathcal{V}^- \rightarrow \mathbb{R}^{\sigma+k}$, $f(\omega) = \left(\int_{\gamma_j} -i\omega \right)_{j=1, \dots, \sigma+k}$. Note that the following differentials are linearly independent and are not in the kernel of f :

$$\left\{ z^m \frac{dz}{w}, \quad m = 0, \dots, \sigma \right\} \cup \left\{ \frac{dz}{(z - \beta_m)w}, \quad m = 1, \dots, k - 1 \right\}.$$

Using Claim 1 we get that the matrix

$$\left(f\left(z^m \frac{dz}{w}\right) \right)_{m=0, \dots, \sigma-1} = (a_0^m, \dots, a_{\sigma-1}^m, 0, \dots, 0)_{m=0, \dots, \sigma-1}$$

has range σ . Moreover, it is easy to check that $f\left(\frac{dz}{(z-\beta_j)w}\right) = (a_0^j, \dots, a_{\sigma-1}^j, 0, \dots, 0, 2^j \pi b_j, 0, \dots, 0)$, and $f\left(z^\sigma \frac{dz}{w}\right) = (a_0^k, \dots, a_{\sigma-1}^k, 0, \dots, 0, 2\pi b_k)$, where $b_j \in \mathbb{R}^*$, $j = 1, \dots, k$. Both facts imply that f is onto. Thanks to this there exists $\tau_0 \in \mathcal{V}^-$ such that $\text{Im}(\int_c \tau_0) \neq 0$.

Moreover, taking into account our construction, we know that $A^*(\tau_0) = -\tau_0$. The second part of the lemma is a consequence of a successive use of Claim 2. □

Claim 4. *Let $H^-\left(\overline{D(\mathfrak{P})}\right)$ be the real vector space of the holomorphic functions $t : \overline{D(\mathfrak{P})} \rightarrow \mathbb{C}$ satisfying $t \circ I = -\bar{t}$. Then the linear map $F : H^-\left(\overline{D(\mathfrak{P})}\right) \rightarrow \mathbb{R}^{2(\sigma+k)}$,*

given by

$$F(t) = \left(\int_{\gamma_j} t \Phi_3 \left(\frac{1}{g} + g \right), -i \int_{\gamma_j} t \Phi_3 \left(\frac{1}{g} - g \right) \right)_{j=1, \dots, \sigma+k},$$

is surjective.

Proof. We proceed by contradiction. Assume F is not onto. Then, there is

$$(\vartheta_1, \dots, \vartheta_{\sigma+k}, \mu_1, \dots, \mu_{\sigma+k}) \in \mathbb{R}^{2(\sigma+k)} - \{(0, \dots, 0)\},$$

such that

$$(2) \quad \sum_{j=1}^{\sigma+k} \left[\vartheta_j \int_{\gamma_j} t \Phi_3 \left(\frac{1}{g} + g \right) - i \mu_j \int_{\gamma_j} t \Phi_3 \left(\frac{1}{g} - g \right) \right] = 0 \quad \forall t \in H^-(\overline{D(\mathfrak{P})}).$$

Claim 3 guarantees the existence of a differential $\tau \in \mathcal{V}^-$ satisfying

- (i) $(\tau)_0 \geq \left(\left(\left(\frac{1}{g} + g \right) \Phi_3 \right)_{|\overline{\Omega}} \right)_0^2 \left(\left(d \left(\frac{1-g^2}{1+g^2} \right) \right)_{|\overline{\Omega}} \right)_0$,
- (ii) $-i \sum_{j=1}^{\sigma+k} \mu_j \int_{\gamma_j} \tau \neq 0$.

Let us define $y \stackrel{\text{def}}{=} \frac{\tau}{d \left(\frac{1-g^2}{1+g^2} \right)}$ and $t \stackrel{\text{def}}{=} \frac{d(y)}{\left(\frac{1}{g} + g \right) \Phi_3}$. Taking the choice of τ into

account the function t belongs to $H^-(\overline{D(\mathfrak{P})})$. In this case, and integrating by parts, (2) becomes $-i \sum_{j=1}^{\sigma+k} \mu_j \int_{\gamma_j} \tau = 0$, which is absurd. This contradiction proves the claim. \square

Using the previous claim we infer the existence of $\{t_1, \dots, t_{2(\sigma+k)}\} \subset H^-(\overline{D(\mathfrak{P})})$ such that $\det(F(t_1), \dots, F(t_{2(\sigma+k)})) \neq 0$. Up to changing $t_i \leftrightarrow t_i/x$, $x > 0$, large enough, we can assume that

$$(3) \quad \left| \exp \left(\sum_{i=1}^{2(\sigma+k)} x_i t_i(p) \right) - 1 \right| < 1/(2\alpha),$$

$$\forall (x_1, \dots, x_{2(\sigma+k)}) \in \mathbb{R}^{2(\sigma+k)}, |x_i| < 1, i = 1, \dots, 2(\sigma+k), \quad \forall p \in \overline{D(\mathfrak{P})}.$$

Claim 5. For each $n \in \mathbb{N}$, there is $t_0^n \in H^-(\overline{D(\mathfrak{P})})$ such that:

- (i) $|t_0^n - n \log \alpha| < 1/n$ in K_1 (and so $|t_0^n + n \log \alpha| < 1/n$ in $I(K_1)$),
- (ii) $|t_0^n| < 1/n$ in K_2 .

Proof. Given $n \in \mathbb{N}$, we apply a Runge-type theorem on \overline{M} (see [8, Theorem 10]) and obtain a holomorphic function $T_0^n : \overline{D(\mathfrak{P})} \rightarrow \mathbb{C}$ satisfying

- $|T_0^n - n \log \alpha| < 1/n$ in K_1 ,
- $|T_0^n + n \log \alpha| < 1/n$ in $I(K_1)$,
- $|T_0^n| < 1/n$ in K_2 .

We take $t_0^n = \frac{1}{2}(T_0^n - \overline{T_0^n \circ I})$. From this, it is trivial to check properties (i) and (ii). \square

For $\Theta = (\lambda_0, \dots, \lambda_{2(\sigma+k)}) \in \mathbb{R}^{2(\sigma+k)+1}$, we define

$$h^{\Theta,n}(p) \stackrel{\text{def}}{=} \exp \left[\lambda_0 t_0^n(p) + \sum_{j=1}^{2(\sigma+k)} \lambda_j t_j(p) \right], \quad \forall p \in \overline{D(\mathfrak{P})}.$$

Label $g^{\Theta,n} = g/h^{\Theta,n}$ and $\Phi_3^{\Theta,n} = \Phi_3$. As $\{t_0^n|_{K_2}\}_{n \in \mathbb{N}}$ is uniformly bounded, then, up to a subsequence, we have $\{t_0^n|_{K_2}\} \rightarrow t_0^\infty \equiv 0$, uniformly on K_2 . We also define on K_2 the Weierstrass data $g^{\Theta,\infty} = g/h^{\Theta,\infty}$, $\Phi_3^{\Theta,\infty} = \Phi_3$, where

$$h^{\Theta,\infty}(p) \stackrel{\text{def}}{=} \exp \left[\sum_{j=1}^{2(\sigma+k)} \lambda_j t_j(p) \right], \quad \forall p \in K_2.$$

Observe that the third Weierstrass differential of the aforementioned holomorphic data has no real periods. Then we must only consider the period problem associated to $\Phi_j^{\Theta,n}$, $j = 1, 2$. To do this, we define the period map $\mathcal{P}_n : \mathbb{R}^{2(\sigma+k)+1} \rightarrow \mathbb{R}^{2(\sigma+k)}$, $n \in \mathbb{N} \cup \{\infty\}$:

$$\mathcal{P}_n(\Theta) = \left(\int_{\gamma_j} \Phi_1^{\Theta,n}, \int_{\gamma_j} \Phi_2^{\Theta,n} \right)_{j=1, \dots, \sigma+k}.$$

Since the initial immersion X is well defined, one has $\mathcal{P}_n(0) = 0$, $\forall n \in \mathbb{N} \cup \{\infty\}$. Moreover, it is not hard to check that

$$\text{Jac}_{\lambda_1, \dots, \lambda_{2(\sigma+k)}}(\mathcal{P}_n)(0) = \det(F(t_1), \dots, F(t_{2(\sigma+k)})) \neq 0, \quad \forall n \in \mathbb{N} \cup \{\infty\}.$$

Applying the Implicit Function Theorem to the map \mathcal{P}_n at $0 \in [-\epsilon, \epsilon] \times \overline{B}(0, r)$, we get a smooth function $L_n : I_n \rightarrow \mathbb{R}^{2(\sigma+k)}$ satisfying $\mathcal{P}_n(\lambda_0, L_n(\lambda_0)) = 0$, $\forall \lambda_0 \in I_n$, where I_n is a maximal open interval containing 0 (here, maximal means that L_n cannot be regularly extended beyond I_n).

Let us see that the supremum ϵ_n of the connected component of $L_n^{-1}(\overline{B}(0, r)) \cap [0, \epsilon]$ containing $\lambda_0 = 0$ belongs to I_n . Indeed, take a sequence $\{\lambda_0^k\}_{k \in \mathbb{N}} \nearrow \epsilon_n$. As $\{L_n(\lambda_0^k)\} \subset \overline{B}(0, r)$, then, up to a subsequence, $\{L_n(\lambda_0^k)\}_{k \in \mathbb{N}} \rightarrow \Lambda_n \in \overline{B}(0, r)$. Taking into account that $\text{Jac}_{\lambda_1, \dots, \lambda_{2(\sigma+k)}}(\mathcal{P}_n)(\epsilon_n, \Lambda_n) \neq 0$, the local unicity of the curve $(\lambda_0, L_n(\lambda_0))$ around the point (ϵ_n, Λ_n) , and the maximality of I_n , we infer that $\epsilon_n \in I_n$. Therefore, either $\epsilon_n = \epsilon$ or $L_n(\epsilon_n) = \Lambda_n \in \partial(B(0, r))$.

We will now see that $\epsilon_0 \stackrel{\text{def}}{=} \liminf \{\epsilon_n\} > 0$. Otherwise, there would be a subsequence $\{\epsilon_n\} \rightarrow 0$. Without loss of generality, $\epsilon_n < \epsilon$, $\forall n \in \mathbb{N}$, and so $\Lambda_n \in \partial(B(0, r))$, $\forall n \in \mathbb{N}$. Up to a subsequence, $\{\Lambda_n\} \rightarrow \Lambda_\infty \in \partial(B(0, r))$. The fact $\mathcal{P}_\infty(0, 0) = \mathcal{P}_\infty(0, \Lambda_\infty) = 0$ would contradict the injectivity of $\mathcal{P}_\infty(0, \cdot)$ in $\overline{B}(0, r)$. Hence the function $L_n : [0, \epsilon_0] \rightarrow \overline{B}(0, r)$ is well defined, $\forall n \geq n_0$, $n_0 > 1/\epsilon_0$, n_0 large enough.

Label $(\lambda_1^n, \dots, \lambda_{2(\sigma+k)}^n) = L_n(1/n)$. From (3) we have $|\exp[\sum_{j=1}^{2(\sigma+k)} \lambda_j^n t_j] - 1| < 1/(2\alpha)$ on $D(\mathfrak{P})$. Hence, if $n (\geq n_0)$ is large enough, the function

$$h(z) \stackrel{\text{def}}{=} \exp \left[\frac{1}{n} t_0^n(z) + \sum_{j=1}^{2(\sigma+k)} \lambda_j^n t_j(z) \right]$$

satisfies statements 1, 2 and 3 in Lemma 1. As the period function \mathcal{P}_n vanishes at $\Theta_n = (1/n, \lambda_1^n, \dots, \lambda_{2(\sigma+k)}^n)$, then the minimal immersion \tilde{F} associated to the Weierstrass data $g^{\Theta_n, n}, \Phi_3^{\Theta_n, n} = \Phi_3$ is well defined. This proves statement 4 in the lemma.

Proof of Lemma 2. Consider \mathfrak{P} , the multigon given in the statement of the lemma, $\mathfrak{P} = \{P_1, P_2, \dots, P_k\}$. As usual in this kind of construction, we are going to follow [2] to describe a labyrinth on $D(\mathfrak{P}) \subset M$ depending on \mathfrak{P} and a positive integer N . Then, we use Lemma 1 following Nadirashvili's ideas [5].

From now on, N will represent a positive multiple of k , and we will always assume that N is large enough to satisfy all the inequalities where it appears. In what follows, the symbol *const* will denote a family of positive real constants that do not depend on N . These constants only depend on $X, \mathfrak{P}, r, \varepsilon$ and s . So, we will use this symbol to refer distinct constants that appear in different parts of the proof.

Let $v_{1,j}, \dots, v_{\frac{2N}{k},j}$ be a set of points in the polygon P_j (containing the vertices of P_j) that divide each side of the polygon P_j into $2N/(k l_j)$ segments of equal length, where l_j is the number of sides of P_j . We are supposing that N is a multiple of l_1, \dots, l_k . We transfer this partition to the parallel polygon $P_j^{2/N}: v'_{1,j}, \dots, v'_{2N/k,j}$. Define the following sets:

- $L_{i,j}$ = the segment that joins $v_{i,j}$ and $v'_{i,j}$, $i = 1, \dots, 2N/k$;
- $P_{i,j} = P_j^{i/N^3}$, $i = 0, \dots, 2N^2$ (recall that P_j^{i/N^3} means the parallel polygon to P_j , in $\text{Int}(\mathfrak{P})$, such that the distance between parallel sides is i/N^3);
- **If** $j < k$, $\mathcal{A}_j = \bigcup_{i=0}^{N^2-1} \overline{\text{Int}(P_{2i+1,j}) \setminus \text{Int}(P_{2i,j})}$
and $\tilde{\mathcal{A}}_j = \bigcup_{i=1}^{N^2} \overline{\text{Int}(P_{2i,j}) \setminus \text{Int}(P_{2i-1,j})}$;
- **If** $j = k$, $\mathcal{A}_k = \bigcup_{i=0}^{N^2-1} \overline{\text{Int}(P_{2i,k}) \setminus \text{Int}(P_{2i+1,k})}$
and $\tilde{\mathcal{A}}_k = \bigcup_{i=1}^{N^2} \overline{\text{Int}(P_{2i-1,k}) \setminus \text{Int}(P_{2i,k})}$;
- $\mathcal{R}_j = \bigcup_{i=0}^{2N^2} P_{i,j}$;
- $L_j = \bigcup_{i=1}^{N/k} L_{2i,j} \cap \mathcal{A}_j$, $\tilde{L}_j = \bigcup_{i=0}^{N/k-1} L_{2i+1,j} \cap \tilde{\mathcal{A}}_j$, and $H_j = \mathcal{R}_j \cup L_j \cup \tilde{L}_j$;
- **If** $j < k$, $\Omega_{N,j} = \{z \in \text{Int}(P_{2N^2,j}) \setminus \text{Int}(P_{0,j}) : \text{dist}_{(ds_0, \mathbb{C})}(z, H_j) \geq \frac{1}{4N^3}\}$;
- **If** $j = k$, $\Omega_{N,k} = \{z \in \text{Int}(P_{0,k}) \setminus \text{Int}(P_{2N^2,k}) : \text{dist}_{(ds_0, \mathbb{C})}(z, H_k) \geq \frac{1}{4N^3}\}$;
where ds_0 is the Euclidean metric on \mathbb{C} ;
- $\Omega_N = \bigcup_{j=1}^k \Omega_{N,j}$.

We define $V_{i,j}$ as the union of the segment $L_{i,j}$ and those connected components of $\Omega_{N,j}$ that have nonempty intersection with $L_{i,j}$ for $i = 1, \dots, 2N/k$. We define the sets ω_l , $l = 1, \dots, 2N$, as $\omega_{(j-1)\frac{2N}{k}+i} = V_{i,j}$, $i = 1, \dots, 2N/k$, $j = 1, \dots, k$. We label $\varpi_i = \{z \in \mathbb{C} : \text{dist}_{(ds_0, \mathbb{C})}(z, \omega_i) < \delta(N)\}$, where $i = 1, \dots, 2N$, and $\delta(N) > 0$ is chosen in such a way that the sets $\overline{\varpi_i}$ ($i = 1, \dots, 2N$) are pairwise disjoint (see Figure 3). Finally, we denote $\omega'_i \stackrel{\text{def}}{=} f_+(\omega_i)$, $\varpi'_i \stackrel{\text{def}}{=} f_+(\varpi_i)$, $\omega_i^* \stackrel{\text{def}}{=} \omega'_i \cup I(\omega'_i)$, and $\varpi_i^* = \varpi'_i \cup I(\varpi'_i)$.

Now, we consider the Riemannian metric $ds^2 = \|dz/w\|^2$ on M . Observe that the metrics ds^2 and $\|dz\|$ are equivalent in $\overline{D(\mathfrak{P})} \setminus D(\mathfrak{P}^\varepsilon)$, i.e.,

$$\text{const} \left\| \frac{dz}{w} \right\| \leq \|dz\| \leq \text{const} \left\| \frac{dz}{w} \right\| \quad \text{in } \overline{D(\mathfrak{P})} \setminus D(\mathfrak{P}^\varepsilon).$$




FIGURE 3. The labyrinth.

Thanks to this relation, we can translate some metric properties of the sets ϖ_i to the sets ϖ_i^* . If N is large enough, it is clear that we can guarantee the following properties:

Claim A. $\text{diam}_{ds}(\varpi_i^j) \leq \frac{\text{const}}{N}$, $j = 1, 2$.

Claim B. If $\lambda^2(z, w)ds^2$ is a conformal metric in $\overline{D(\mathfrak{P})}$ and $\Upsilon \in \mathbb{R}^+$ satisfying

$$\lambda(z, w) \geq \begin{cases} \Upsilon & \text{in } \overline{D(\mathfrak{P})}, \\ \Upsilon N^4 & \text{in } \bigcup_{j=1}^{2N} \omega_j^*, \end{cases}$$

and if α is a curve in $D(\mathfrak{P})$ connecting $\partial(D(\mathfrak{P}^\epsilon))$ and $\partial(D(\mathfrak{P}))$, then $\text{length}(\alpha, \lambda ds) \geq \text{const} \Upsilon N$, where const does not depend on Υ .

Our next stage involves a modification of the immersion X in $2N$ steps, where each step is related to each set ϖ_i^* . This modification consists of constructing a sequence of $2N$ conformal minimal immersions $F_i : \overline{D(\mathfrak{P})} \rightarrow \mathbb{R}^3$, $i = 0, 1, \dots, 2N$, with $F_0 = X$, in a recursive way. The immersion obtained in the last step, F_{2N} , will be the immersion that proves the lemma. In order to get this, the sequence of immersions must satisfy the following properties:

- (P1_{*i*}) $F_i(p) = \text{Re} \left(\int_{p_0}^p \Phi^i \right)$, where $I^* \Phi^i = \overline{\Phi^i}$,
 $\Phi^i = (\varphi_1^i(z, w), \varphi_2^i(z, w), \varphi_3^i(z, w)) \frac{dz}{w}$;
- (P2_{*i*}) $\|\varphi^i - \varphi^{i-1}\| \leq 1/N^2$ in $D(\mathfrak{P}) \setminus \varpi_i^*$;
- (P3_{*i*}) $\|\varphi^i\| \geq N^{7/2}$ in ω_i^* ;
- (P4_{*i*}) $\|\varphi^i\| \geq \frac{\text{const}}{\sqrt{N}}$ in ϖ_i^* ;
- (P5_{*i*}) $\text{dist}_{(ds_1, S^2)}(\mathcal{G}_i(z, w), \mathcal{G}_{i-1}(z, w)) < \frac{1}{N\sqrt{N}}$ in $D(\mathfrak{P}) \setminus \varpi_i^*$, where ds_1 is the usual Riemannian metric in S^2 and \mathcal{G}_i represents the Gauss map of the immersion F_i ;

(P6_i) there exists an orthogonal frame $S_i = \{e_1, e_2, e_3\}$ in \mathbb{R}^3 such that:

(P6.1_i) if $p \in \overline{\varpi_i^*}$ and $\|F_{i-1}(p)\| \geq 1/\sqrt{N}$, then

$$\|(F_{i-1}(p))_*\| < \frac{\text{const}}{\sqrt{N}} \|F_{i-1}(p)\|,$$

(P6.2_i) $(F_i(p))_3 = (F_{i-1}(p))_3$ for all $p \in \overline{D(\mathfrak{P})}$,

where $(\cdot)_k$ is the k th coordinate function and $(\cdot)_* = ((\cdot)_1, (\cdot)_2)$, with respect to $S_i = \{e_1, e_2, e_3\}$.

Now, we construct the above sequence of immersions in a recursive way. Suppose that we have F_0, \dots, F_{j-1} satisfying the above claims. We must define F_j .

Observe that for an N large enough, the immersion F_{j-1} satisfies the following properties:

(L1) $\|\varphi^{j-1}\| \leq \text{const}$ in $D(\mathfrak{P}) \setminus \bigcup_{l=1}^{j-1} \varpi_l^*$.

(L2) $\|\varphi^{j-1}\| \geq \text{const}$ in $D(\mathfrak{P}) \setminus \bigcup_{l=1}^{j-1} \varpi_l^*$.

(L3) $F_{j-1} \circ I = F_{j-1}$ and $\mathcal{G}_{j-1} \circ I = -\mathcal{G}_{j-1}$.

(L4) The diameter in \mathbb{S}^2 of $\mathcal{G}_{j-1}(\varpi_j')$ is less than $\frac{\text{const}}{\sqrt{N}}$, and the diameter in \mathbb{R}^3 of $F_{j-1}(\varpi_j')$ is less than $\frac{\text{const}}{N}$.

(L5) There is a set of orthogonal coordinates $S_j = \{e_1, e_2, e_3\}$ in $\mathbb{R}^3 > 0$ such that:

(L5.a) if $p \in \varpi_j^*$ and $\|F_{j-1}(p)\| \geq \frac{1}{\sqrt{N}}$, then $\angle(e_3, F_{j-1}(p)) \leq \frac{\text{const}}{\sqrt{N}}$,

(L5.b) $\angle(\pm e_3, \mathcal{G}_{j-1}) \geq \frac{\text{const}}{\sqrt{N}}$ in ϖ_j^* .

Here, $\angle(a, b) \in [0, \pi[$ is the angle formed by a and b in \mathbb{R}^3 .

Claims (L1)–(L4) easily deduce from the properties (P₋). The choice of the coordinates S_j in (L5) is possible as a consequence of the bounds of the diameters of $F_{j-1}(\varpi_j^l)$ and $\mathcal{G}_{j-1}(\varpi_j^l)$ given in the statement of (L3) and property (L4).

We shall now construct F_j . Let (Φ_3^{j-1}, g^{j-1}) be the Weierstrass data of the immersion F_{j-1} in the coordinate system S_j . Let h_α be the function given by Lemma 1 for $K_1 = \omega_j'$, $K_2 = \overline{D(\mathfrak{P})} \setminus \varpi_j^*$ and α large enough in terms of N . We define the minimal immersion F_j in the frame S_j as $F_j(p) = \text{Re} \left(\int_{p_0}^p \Phi^j \right)$, where Φ^j is determined by $\Phi_3^j = \Phi_3^{j-1}$ and $g^j = g^{j-1}/h_\alpha$. We denote $\Phi^j = \varphi^j \frac{dz}{w}$. Observe that F_j is well defined because Lemma 1 tells us that the Weierstrass data Φ^j has no real periods.

We shall now see that F_j satisfies properties (P1_j)–(P6_j). Claim (P1_j) easily holds. Claims (P2_j) and (P5_j) are a consequence of the fact that $h_\alpha \rightarrow 1$ uniformly on $K_2 = \overline{D(\mathfrak{P})} \setminus \varpi_j^*$ as $\alpha \rightarrow \infty$. To satisfy (P3_j), we take into account that $h_\alpha \rightarrow \infty$ uniformly on $K_1 = \omega_j'$ as $\alpha \rightarrow \infty$. So, $\|\Phi^j\| \geq N^{7/2}$ in ω_j' , for α large enough. Property (P3_j) is a consequence of $I^* \Phi_j = \overline{\Phi_j}$. To deduce property (P4_j), we need the next inequalities:

$$(4) \quad \frac{\sin\left(\frac{\text{const}}{\sqrt{N}}\right)}{1 + \cos\left(\frac{\text{const}}{\sqrt{N}}\right)} \leq |g^{j-1}| \leq \frac{\sin\left(\frac{\text{const}}{\sqrt{N}}\right)}{1 - \cos\left(\frac{\text{const}}{\sqrt{N}}\right)} \quad \text{in } \varpi_j^*.$$

These inequalities are due to (L5.b). Then using (L2) and (4) (for a large enough N), one has

$$\|\varphi^j\| \geq |\varphi_3^j| = |\varphi_3^{j-1}| \geq \sqrt{2} \|\varphi^{j-1}\| \frac{|g^{j-1}|}{1 + |g^{j-1}|^2} \geq \text{const} \sin\left(\frac{\text{const}}{\sqrt{N}}\right) \geq \frac{\text{const}}{\sqrt{N}} \quad \text{in } \varpi_j^*.$$

To obtain (P6.1_j) and (P6.2_j), use that $\Phi_3^{j-1} = \Phi_3^j$ in the frame S_j , and (L5.a).

Hence, we have constructed the immersions F_0, F_1, \dots, F_{2N} satisfying claims (P1_j)–(P6_j) for $j = 1, \dots, 2N$.

In the following proposition, we study the properties of the immersion F_{2N} .

Proposition 1. *If N is large enough, then F_{2N} satisfies that:*

- (i) $2s < \text{dist}_{(F_{2N}, \overline{D(\mathfrak{P})})}(\partial(D(\mathfrak{P})), \partial(D(\mathfrak{P}^\varepsilon)))$;
- (ii) $\|F_j - F_{j-1}\| \leq \frac{\text{const}}{N^2}$ in $D(\mathfrak{P}) \setminus \varpi_j^*$;
- (iii) $\|F_{2N} - X\| \leq \frac{\text{const}}{N}$ in $D(\mathfrak{P}) \setminus \bigcup_{j=1}^{2N} \varpi_j^*$;
- (iv) *there is a multigon $\tilde{\mathfrak{P}}$ satisfying:*
 - (iv.1) $\overline{\text{Int}(\mathfrak{P}^\varepsilon)} \subset \text{Int}(\tilde{\mathfrak{P}}) \subset \text{Int}(\mathfrak{P}) \subset \text{Int}(\mathfrak{P})$;
 - (iv.2) $s < \text{dist}_{(F_{2N}, \overline{D(\tilde{\mathfrak{P}})})}(p, \partial(D(\mathfrak{P}^\varepsilon))) < 2s, \forall p \in \partial(D(\tilde{\mathfrak{P}}))$;
 - (iv.3) $F_{2N}(D(\mathfrak{P})) \subset B_R$, where $R = \sqrt{r^2 + (2s)^2} + \varepsilon$.

Proof. Properties (L2), (P2_i), (P3_i) and (P4_i) give us the following bound of the conformal metric $\lambda_{F_{2N}}^2(z) \|\frac{dz}{w}\|^2$:

$$\lambda_{F_{2N}}(z, w) \geq \begin{cases} \Upsilon & \text{in } \overline{D(\mathfrak{P})}, \\ \Upsilon N^4 & \text{in } z^{-1}(\Omega_N), \end{cases}$$

for $\Upsilon = \frac{\text{const}}{\sqrt{N}}$. Then we conclude the proof of (i) thanks to Claim B.

Now we shall prove (ii). Observe that there is a constant, depending only on $D(\mathfrak{P})$, such that

$$\sup\{\text{dist}_{(ds, \Xi_N^j)}(p_0, p) : N \in \mathbb{N}, j \in \{1, \dots, 2N\}, p \in \Xi_N^j\} \leq \text{const},$$

where $\Xi_N^j = \overline{D(\mathfrak{P})} \setminus \varpi_j^*$. Bearing in mind the above and using (P2_j), we can obtain (ii). From (ii), it is not hard to deduce (iii).

Now, we deal with (iv). We will construct the multigon $\tilde{\mathfrak{P}}$. Let

$$\mathcal{S} = \{p \in D(\mathfrak{P}) \setminus \overline{D(\mathfrak{P}^\varepsilon)} : s < \text{dist}_{(F_{2N}, \overline{D(\mathfrak{P})})}(p, \partial(D(\mathfrak{P}^\varepsilon))) < 2s\}.$$

Note that \mathcal{S} is a nonempty open subset of $D(\mathfrak{P}) \setminus \overline{D(\mathfrak{P}^\varepsilon)}$, satisfying $I(\mathcal{S}) = \mathcal{S}$. As a consequence of (i), we deduce that $z(\mathcal{S})$ contains k Jordan curves $\Gamma = \{\Gamma_1, \dots, \Gamma_k\}$ satisfying that $f_+(\Gamma_1), \dots, f_+(\Gamma_k)$ and $I(f_+(\Gamma_1)), \dots, I(f_+(\Gamma_k))$ bound a domain \mathcal{D} such that $\overline{D(\mathfrak{P}^\varepsilon)} \subset \mathcal{D} \subset \overline{\mathcal{D}} \subset D(\mathfrak{P})$. Then we can approximate Γ by a multigon $\tilde{\mathfrak{P}} \subset z(\mathcal{S})$ satisfying statements (iv.1) and (iv.2).

Finally, we prove assertion (iv.3), or equivalently $F_{2N}(\partial(D(\tilde{\mathfrak{P}}))) \subset B_R$. Take $p \in \partial(D(\tilde{\mathfrak{P}}))$. If $p \in D(\mathfrak{P}) \setminus \bigcup_{j=1}^{2N} \varpi_j^*$, we conclude that

$$\|F_{2N}(p)\| \leq \|F_{2N}(p) - X(p)\| + \|X(p)\| \leq \frac{\text{const}}{N} + r \leq R.$$

Suppose now $p \in \varpi_j^*, j \in \{1, \dots, 2N\}$. From (iv.2), it is possible to find a curve $\gamma : [0, 1] \rightarrow D(\mathfrak{P})$ such that $\gamma(0) \in \partial(D(\mathfrak{P}^\varepsilon)), \gamma(1) = p$, and $\text{length}(\gamma, ds_{F_{2N}}) \leq 2s$. Let $\bar{p} = \gamma(\bar{t})$ be a point in the curve γ where

$$\bar{t} = \sup\{t \in [0, 1] : \gamma(t) \in \partial(\varpi_j^*)\}.$$

Using the former, we deduce that $\|F_{2N}(\bar{p}) - F_{2N}(p)\| \leq 2s$. Then, taking into account (ii), one has

$$(5) \quad \|F_j(\bar{p}) - F_j(p)\| \leq \frac{\text{const}}{N} + 2s.$$

At this point, we are able to get bounds for coordinates of $F_j(p)$ in the frame S_j . Henceforth, we will use the frame S_j . We deal with the coordinates $(F_j(p))_3$ and $(F_j(p))_* = ((F_j(p))_1, (F_j(p))_2)$. From (P6.2_j) and (ii), we obtain a bound of the third coordinate of $F_j(p)$:

$$|(F_j(p))_3| = |(F_{j-1}(p))_3| \leq |(F_{j-1}(p))_3 - (X(p))_3| + |(X(p))_3| \leq \frac{\text{const}}{N} + r.$$

On the other hand, using inequality (5), the fact that $\bar{p} \in D(\mathfrak{P}) \setminus \varpi_j^*$ and assertion (ii), one has

$$\begin{aligned} (6) \quad \|(F_j(p))_*\| &\leq \|(F_j(p))_* - (F_j(\bar{p}))_*\| + \|(F_j(\bar{p}))_* - (F_{j-1}(\bar{p}))_*\| + \|(F_{j-1}(\bar{p}))_*\| \\ &\leq \frac{\text{const}}{N} + 2s + \frac{\text{const}}{N^2} + \|(F_{j-1}(\bar{p}))_*\|. \end{aligned}$$

Observe that $\|(F_{j-1}(\bar{p}))_*\| \leq \frac{\text{const}}{\sqrt{N}}$. To check this, we follow the next arguments. If $\|F_{j-1}(\bar{p})\| \geq 1/\sqrt{N}$, then $\|(F_{j-1}(\bar{p}))_*\| \leq \frac{\text{const}}{\sqrt{N}} (\frac{\text{const}}{N} + r) \leq \frac{\text{const}}{\sqrt{N}}$, as a consequence of (P6.1_j) and (ii). If, instead, we have $\|F_{j-1}(\bar{p})\| < 1/\sqrt{N}$, then $\|(F_{j-1}(\bar{p}))_*\| \leq \frac{\text{const}}{\sqrt{N}}$ directly holds. In any case, following (6) and using the former, we can deduce that

$$\|(F_j(p))_*\| \leq \frac{\text{const}}{N} + 2s + \frac{\text{const}}{N^2} + \frac{\text{const}}{\sqrt{N}} \leq 2s + \frac{\text{const}}{\sqrt{N}}.$$

To finish, Pythagoras' theorem gives us

$$\begin{aligned} \|F_{2N}(p)\| &\leq \|F_{2N}(p) - F_j(p)\| + \|F_j(p)\| \\ &\leq \frac{\text{const}}{N} + \sqrt{|(F_j(p))_3|^2 + \|(F_j(p))_*\|^2} < \sqrt{r^2 + (2s)^2} + \varepsilon = R \end{aligned}$$

for an N large enough. □

It is easy to check that $Y = F_{2N}$ satisfies all claims in Lemma 2, and then this ends the proof of the lemma.

4. PROOF OF THE MAIN THEOREM

At this point, we prove our main result (Theorem 1).

Proof. Let $r_1 > 1$ and $\rho_1 > 0$, and define $r_n = \sqrt{r_{n-1}^2 + (2/n)^2} + 1/n^2$ and $\rho_n = \rho_1 + \sum_{i=2}^n 1/i$, $n \geq 2$. Our purpose consists of using Lemma 2 to define a sequence

$$\chi_n = (X_n : \overline{D(\mathfrak{P}_n)} \rightarrow \mathbb{R}^3, \mathfrak{P}_n, \varepsilon_n, \xi_n),$$

where X_n is a conformal minimal immersion, \mathfrak{P}_n is a multigon, $\{\varepsilon_n\}$, $\{\xi_n\}$ are decreasing sequences of nonvanishing terms satisfying $\varepsilon_n, \xi_n < 1/n^2$, and:

- (A_n) $\rho_n < \text{dist}_{(X_n, \overline{D(\mathfrak{P}_n^{\xi_n})})}(p_0, \partial(D(\mathfrak{P}_n^{\xi_n})))$,
- (B_n) $X_n(D(\mathfrak{P}_n)) \subset B_{r_n}$,
- (C_n) $X_n(p) = \text{Re} \left(\int_{p_0}^p \Phi^n \right)$, where $I^*(\Phi^n) = \bar{\Phi}^n$,
- (D_n) $\|X_n - X_{n-1}\| < \varepsilon_n$ in $D(\mathfrak{P}_{n-1}^{\xi_{n-1}})$,
- (E_n) $\lambda_{X_n} \geq \alpha_n \lambda_{X_{n-1}}$ in $D(\mathfrak{P}_{n-1}^{\xi_{n-1}})$, where $\{\alpha_i\}_{i \in \mathbb{N}}$ is a sequence of real numbers such that $0 < \alpha_i < 1$ and $\{\prod_{i=1}^n \alpha_i\}_n$ converges to $1/2$,
- (F_n) $\overline{D(\mathfrak{P}_{n-1}^{\xi_{n-1}})} \subset D(\mathfrak{P}_n^{\xi_n}) \subset \overline{D(\mathfrak{P}_n)} \subset D(\mathfrak{P}_{n-1})$.

The first element of the sequence is taken as follows. If $\sigma = 0$, we deal with $X_1 : M_0 \rightarrow \mathbb{R}^3$, the double oriented cover of Meeks' Möbius strip [6]. If $\sigma > 0$, then we take $X_1 : M_\sigma \rightarrow \mathbb{R}^3$ as the oriented double cover of the nonorientable surface of genus $\sigma + 1$ described in [3]. It is not hard to see that, in both cases, the closed Riemann surface can be written as

$$\overline{M} = \left\{ (z, w) \in \overline{\mathbb{C}}^2 : w^2 = \prod_{i=1}^{\sigma+1} (z - d_i)(z - \overline{d}_i) \right\}$$

for suitable $d_1, \dots, d_{\sigma+1}$, and the reversing orientation involution has the form $I(z, w) = (\overline{z}, -\overline{w})$. The simply connected domain Δ is chosen in such a way that $\{d_1, \dots, d_{\sigma+1}, \overline{d}_1, \dots, \overline{d}_{\sigma+1}\} \subset \Delta$, and $z^{-1}(\Delta) \subset M_\sigma$. The real numbers $\beta_1, \dots, \beta_{k-1}$ are chosen in $\mathbb{C} \setminus \Delta$ and outside the z -projection of the ends of X_1 . So, a convenient \mathfrak{P}_1, r_1 and ρ_1 , joint with $X_1 : D(\mathfrak{P}_1) \rightarrow \mathbb{R}^3$ completes the construction of the first element in our sequence. We also choose $\xi_1 < 1$ small enough satisfying (A_1) . The choice of $\varepsilon_1 < 1$ is irrelevant.

Lemma 2 allows us to construct the terms of the sequence in a recursive way. For more details see [2].

Now, we define $D = \bigcup_{n=1}^{\infty} D(\mathfrak{P}_n^{\xi_n})$. Observe that D is the limit of an expansive sequence of domains of \overline{M} with the same conformal type. Then, this conformal type remains at the limit, in other words, D has the conformal type of a compact Riemann surface of genus σ minus k closed disks. Thanks to properties (A_n) – (F_n) , it is not difficult to check that $\{X_n\}$ converges to a complete bounded minimal immersion $X : D \rightarrow \mathbb{R}^3$ satisfying $X \circ I = X$. So, it induces a complete nonorientable bounded minimal immersion $\mathcal{X} : \mathcal{D} \rightarrow \mathbb{R}^3$. This concludes the proof. \square

REFERENCES

- [1] H. M. Farkas, I. Kra, *Riemann surfaces*. Graduate Texts in Math., **72**, Springer-Verlag, Berlin, 1980. MR0583745 (82c:30067)
- [2] F.J. López, F. Martín, S. Morales, *Adding handles to Nadirashvili's surfaces*. J. Diff. Geom. **60** (2002), no. 3, 155-175. MR1924594 (2003f:53013)
- [3] Francisco J. López, Francisco Martín, *Complete nonorientable minimal surfaces and symmetries*. Duke Math. J. **79** (1995), no. 3, 667–686. MR1355180 (98b:53008)
- [4] F. Martín, S. Morales, *A complete bounded minimal cylinder in \mathbb{R}^3* . Michigan Math. J. **47** (2000), no. 3, 499-514. MR1813541 (2001m:53015)
- [5] N. Nadirashvili, *Hadamard's and Calabi-Yau's conjectures on negatively curved and minimal surfaces*. Invent. Math. **126** (1996), 457-465. MR1419004 (98d:53014)
- [6] W. H. Meeks, *The classification of complete minimal surfaces in \mathbb{R}^3 with total curvature greater than -8π* . Duke Mathematical Journal **48** (1981), 523-535. MR0630583 (82k:53009)
- [7] R. Osserman, *A survey of minimal surfaces*. Dover Publications, New York, second edition, 1986. MR0852409 (87j:53012)
- [8] H. L. Royden, *Function theory on compact Riemann surfaces*. J. Analyse Math. **18** (1967), 295–327. MR0214757 (35:5606)

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA, FACULTAD DE CIENCIAS, UNIVERSIDAD DE GRANADA, 18071 GRANADA, SPAIN

E-mail address: `fjlopez@ugr.es`

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA, FACULTAD DE CIENCIAS, UNIVERSIDAD DE GRANADA, 18071 GRANADA, SPAIN

E-mail address: `fmartin@ugr.es`

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA, FACULTAD DE CIENCIAS, UNIVERSIDAD DE GRANADA, 18071 GRANADA, SPAIN

E-mail address: `santimo@ugr.es`