

OPEN LOCI OF GRADED MODULES

CHRISTEL ROTTHAUS AND LIANA M. ŞEĞA

ABSTRACT. Let $A = \bigoplus_{i \in \mathbb{N}} A_i$ be an excellent homogeneous Noetherian graded ring and let $M = \bigoplus_{n \in \mathbb{Z}} M_n$ be a finitely generated graded A -module. We consider M as a module over A_0 and show that the (S_k) -loci of M are open in $\text{Spec}(A_0)$. In particular, the Cohen-Macaulay locus $U_{CM}^0 = \{\mathfrak{p} \in \text{Spec}(A_0) \mid M_{\mathfrak{p}} \text{ is Cohen-Macaulay}\}$ is an open subset of $\text{Spec}(A_0)$. We also show that the (S_k) -loci on the homogeneous parts M_n of M are eventually stable. As an application we obtain that for a finitely generated Cohen-Macaulay module M over an excellent ring A and for an ideal $I \subseteq A$ which is not contained in any minimal prime of M , the (S_k) -loci for the modules $M/I^n M$ are eventually stable.

INTRODUCTION

A well-known theorem of Grothendieck states that if M is a finitely generated module over an excellent Noetherian ring A , then for all $k \in \mathbb{N}$ the (S_k) -locus of M

$$U_{S_k}(M) = \{\mathfrak{p} \in \text{Spec}(A) \mid M_{\mathfrak{p}} \text{ satisfies } (S_k)\}$$

is an open subset of $\text{Spec}(A)$. As usual, (S_k) denotes the Serre condition, that is, $M_{\mathfrak{p}}$ satisfies (S_k) if for all $\mathfrak{q} \in \text{Spec}(A)$ with $\mathfrak{q} \subseteq \mathfrak{p}$ it holds that

$$\text{depth}_{A_{\mathfrak{q}}}(M_{\mathfrak{q}}) \geq \min(k, \dim(M_{\mathfrak{q}})).$$

It also follows that for such modules M the Cohen-Macaulay locus

$$U_{CM}(M) = \{\mathfrak{p} \in \text{Spec}(A) \mid M_{\mathfrak{p}} \text{ is Cohen-Macaulay}\}$$

is an open subset of $\text{Spec}(A)$.

Let $A = \bigoplus_{n \geq 0} A_n$ be a Noetherian graded excellent homogeneous ring and $M = \bigoplus_{i \in \mathbb{Z}} M_i$ a finitely generated graded A -module. Considered as a module over the base ring A_0 , M is a direct sum of finitely generated A_0 -modules. Moreover, if the base ring A_0 is local, the standard notion of depth is meaningful for the A_0 -module M and we may consider its (S_k) -loci

$$U_{S_k}^0(M) = \{\mathfrak{p} \in \text{Spec}(A_0) \mid M_{\mathfrak{p}} \text{ satisfies } S_k\},$$

where $M_{\mathfrak{p}}$ denotes the localization of M at the multiplicative set $A_0 \setminus \mathfrak{p}$. In this paper we prove that under these assumptions the (S_k) -loci of the A_0 -module M are open subsets of $\text{Spec}(A_0)$. In particular, the Cohen-Macaulay locus of M (as

Received by the editors March 23, 2004 and, in revised form, September 28, 2004.

2000 *Mathematics Subject Classification*. Primary 13A02, 13C15, 13F40; Secondary 13A30, 13C14.

©2006 American Mathematical Society
Reverts to public domain 28 years from publication

an A_0 -module)

$$U_{CM}^0(M) = \{\mathfrak{p} \in \text{Spec}(A_0) \mid M_{\mathfrak{p}} \text{ is Cohen-Macaulay}\}$$

is an open subset of $\text{Spec}(A_0)$.

The proof follows the main ideas of Grothendieck's proof. It is, however, not merely a copy of the proof in EGA and requires a number of modifications. For the benefit of the reader we have included complete proofs of the results. Our proof is based on the following two observations: First, if A is a polynomial ring over the base ring A_0 , then every graded resolution of M by finitely generated graded free A -modules provides a free resolution of the A_0 -module M which is finitely generated on the homogeneous parts. The second is a result by Hochster and Roberts which states for the A -module M that there is an element $a \in A_0 \setminus (0)$ so that M_a is a free $(A_0)_a$ -module provided that the ring A_0 is a domain.

The paper is organized as follows:

The first section contains basic facts about graded rings and modules which are relevant for the rest of the paper. As a main result we obtain that the Auslander-Buchsbaum formula holds for the A_0 -module M .

The second section shows that the codepth-loci of M are open in $\text{Spec}(A_0)$. This is the main step in proving the openness of the (S_k) -loci which we present in the next section.

In Section 4 we consider the homogeneous parts of the graded module M . We show that the codepth-loci and (S_k) -loci of the homogeneous parts of M are eventually stable. This is applied in the last section to the case of a finitely generated module M over an excellent Noetherian ring A . If $I \subseteq A$ is an ideal we recover a well-known result by Kodiyalam [7], namely that for $k \geq k_0$

$$\text{depth}(M/I^k M) = \text{depth}(M/I^{k_0} M).$$

We also show that if M is a Cohen-Macaulay module over A and if $I \subseteq A$ is not contained in a minimal prime of M , then the codepth- and (S_k) -loci of $M/I^n M$ are eventually stable.

1. BASIC FACTS

In this paper we assume that $A = \bigoplus_{i \in \mathbb{N}} A_i$ is a Noetherian homogeneous graded ring and that $M = \bigoplus_{i \in \mathbb{Z}} M_i$ is a finitely generated A -module. As usual, we let A_+ denote the irrelevant ideal of A , that is, $A_+ = \bigoplus_{i \geq 1} A_i$.

If $\mathfrak{p} \in \text{Spec}(A_0)$ is a prime ideal of A_0 , then $M_{\mathfrak{p}}$ denotes the localization $S^{-1}M$ where $S = A_0 \setminus \mathfrak{p}$. Note that $M_{\mathfrak{p}}$ is a graded module over the graded ring $A_{\mathfrak{p}}$.

Our goal is to show that if A is excellent, then the codepth-loci and the (S_k) -loci of M , considered as a module over the base ring A_0 , are open subsets of $\text{Spec}(A_0)$.

1.1. General remarks. We begin our investigation with some well-known facts about graded modules. Since these results are frequently used throughout the paper, we include them together with their (short) proofs in this introductory section.

1.1.1. Lemma. *There exists an integer t so that $\text{ann}_{A_0}(M_t) = \text{ann}_{A_0}(M_k)$ for all $k \geq t$.*

Proof. For all $k \in \mathbb{Z}$ set $J_k = \text{ann}_{A_0}(M_k)$. Since A is homogeneous and M is a finitely generated A -module, there exists $t_0 \in \mathbb{Z}$ such that

$$A_1 M_k = M_{k+1} \quad \text{for all } k \geq t_0.$$

We conclude $J_k \subseteq J_{k+1}$ for all $k \geq t_0$. Since A_0 is Noetherian, there then exists $t \geq t_0$ so that $J_k = J_t$ for all $k \geq t$. \square

1.1.2. Lemma. *The following two functions are well defined and surjective:*

- (1) *The function $\varphi: \text{Supp}_A(M) \rightarrow \text{Supp}_{A_0}(M)$ defined by $\varphi(P) = P \cap A_0$.*
- (2) *The function $\psi: \text{Ass}_A(M) \rightarrow \text{Ass}_{A_0}(M)$ defined by $\psi(P) = P \cap A_0$.*

Proof. (1) If $P \in \text{Supp}_A(M)$, then $M_P \neq 0$ and in particular $M_{\mathfrak{p}} \neq 0$, where $\mathfrak{p} = P \cap A_0$. This shows that φ is well defined. Let $\mathfrak{p} \in \text{Supp}_{A_0}(M)$. Then

$$M_{\mathfrak{p}} = \bigoplus_{i \in \mathbb{Z}} (M_i)_{\mathfrak{p}} \neq 0,$$

and we may consider $M_{\mathfrak{p}}$ as a graded module over the graded ring $A_{\mathfrak{p}}$. Note that $A_{\mathfrak{p}}$ is a *local ring with unique graded maximal ideal $\mathfrak{m} = \mathfrak{p}(A_0)_{\mathfrak{p}} \oplus (A_+)_{\mathfrak{p}}$. Since all minimal primes of $\text{Supp}_{A_{\mathfrak{p}}}(M_{\mathfrak{p}})$ are graded, $\mathfrak{m} \in \text{Supp}_{A_{\mathfrak{p}}}(M_{\mathfrak{p}})$. Thus there is a prime $P \in \text{Supp}_A(M)$ with $P \cap A_0 = \mathfrak{p}$.

(2) If $P \in \text{Ass}_A(M)$, then there exists $y \in M$ so that $\text{ann}_A(y) = P$. Thus $\text{ann}_{A_0}(y) = P \cap A_0 = \mathfrak{p}$ and $\mathfrak{p} \in \text{Ass}_{A_0}(M)$. Conversely, let $\mathfrak{p} \in \text{Ass}_{A_0}(M)$. Consider again the graded $A_{\mathfrak{p}}$ -module $M_{\mathfrak{p}}$. There exists $z \in M_{\mathfrak{p}}$ so that $\text{ann}_{(A_0)_{\mathfrak{p}}}(z) = \mathfrak{p}(A_0)_{\mathfrak{p}}$, and therefore

$$\mathfrak{p}(A_0)_{\mathfrak{p}} \subseteq \bigcup_{Q \in \text{Ass}_{A_{\mathfrak{p}}}(M_{\mathfrak{p}})} Q.$$

Since $M_{\mathfrak{p}}$ is a finitely generated $A_{\mathfrak{p}}$ -module, there exists $Q \in \text{Ass}_{A_{\mathfrak{p}}}(M_{\mathfrak{p}})$ with $\mathfrak{p}(A_0)_{\mathfrak{p}} \subseteq Q$. Since $A_{\mathfrak{p}}$ is *local with unique graded maximal ideal $\mathfrak{p}(A_0)_{\mathfrak{p}} \oplus (A_+)_{\mathfrak{p}}$, we obtain $Q \cap (A_0)_{\mathfrak{p}} = \mathfrak{p}(A_0)_{\mathfrak{p}}$, and a preimage $P \in \text{Spec}(A)$ of Q is an associated prime of the A -module M , with $P \cap A_0 = \mathfrak{p}$. \square

Lemma 1.1.2 shows in particular that M as an A_0 -module has a finite set of associated primes.

1.1.3. Lemma. *Let A and M be as above and set $I = \text{ann}_{A_0}(M)$. For any $\mathfrak{p} \in \text{Spec}(A_0)$ the following hold:*

- (1) *If $M_{\mathfrak{p}} = 0$, then there is an element $a \in A_0 \setminus \mathfrak{p}$ with $M_a = 0$.*
- (2) *$\text{ann}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) = I(A_0)_{\mathfrak{p}}$.*

Proof. (1) This is a basic fact about Noetherian modules using that M is a finitely generated module over A and $A_0 \setminus \mathfrak{p}$ is a multiplicative subset of A .

(2) Obviously, $I(A_0)_{\mathfrak{p}} \subseteq \text{ann}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}})$. Let $x \in \text{ann}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}})$ with $x = \frac{b}{s}$, where $b \in A_0$ and $s \in A_0 \setminus \mathfrak{p}$. Assume that m_1, \dots, m_r is a system of generators of the A -module M . Since $x \frac{m_i}{1} = 0$ for all $1 \leq i \leq r$ there is an element $t \in A_0 \setminus \mathfrak{p}$ with $t b m_i = 0$ for all $1 \leq i \leq r$. We have that $t b \in I$ and hence $x = \frac{b}{s} \in I(A_0)_{\mathfrak{p}}$. \square

1.2. The Auslander-Buchsbaum formula. Let $A = \bigoplus_{i \geq 0} A_i$ be a graded Noetherian homogeneous ring with (A_0, \mathfrak{m}_0) local and let $M = \bigoplus_{i \in \mathbb{Z}} M_i$ be a finitely generated A -module. Since M is (in general) not finitely generated as an A_0 -module, we need to verify that the classical definition of A_0 -depth works in the case of a finitely generated graded module. First note that an element $z \in \mathfrak{m}_0$ is regular on M if and only if z is regular on M_i for all $i \in \mathbb{Z}$ with $M_i \neq 0$. Let $x_1, \dots, x_s \in \mathfrak{m}_0$ and $y_1, \dots, y_t \in \mathfrak{m}_0$ be two maximal regular M -sequences (as an A_0 -module). Then for all $i \in \mathbb{Z}$ with $M_i \neq 0$ the two sequences are regular on the A_0 -module M_i , and the sets

$$\begin{aligned} \text{Ass}_{A_0}(M/(x_1, \dots, x_s)M) &= \bigcup_{i \in \mathbb{Z}} \text{Ass}_{A_0}(M_i/(x_1, \dots, x_s)M_i), \\ \text{Ass}_{A_0}(M/(y_1, \dots, y_t)M) &= \bigcup_{i \in \mathbb{Z}} \text{Ass}_{A_0}(M_i/(y_1, \dots, y_t)M_i) \end{aligned}$$

are finite by Lemma 1.1.2. The maximality of the first sequences yields that there is an $i \in \mathbb{Z}$ with $M_i \neq 0$ and $\mathfrak{m}_0 \in \text{Ass}_{A_0}(M_i/(x_1, \dots, x_s)M_i)$. Since the second sequence is also regular on M_i we have that $t \leq s$. A similar argument shows that $s \leq t$, and we obtain that two maximal regular sequences on M have the same length. Therefore the classical definition of depth is efficient and we put:

1.2.1. Definition. Let A and M be as above with (A_0, \mathfrak{m}_0) local. We define the *depth* of M as an A_0 -module to be the number

$$\text{depth}_{A_0}(M) := \sup\{n \in \mathbb{N} \mid \exists \text{ an } M\text{-sequence of length } n\}.$$

In general, for a (not necessarily finitely generated) module M over a Noetherian local ring A , the depth of M is defined by means of Koszul homology (see [2, Definition 9.1.1]). In our setting, the definition above coincides with the one in [2].

The aim of this section is to prove the Auslander-Buchsbaum theorem for finitely generated graded modules M over *local graded Noetherian rings A when M is considered a module over the base ring A_0 . There is a generalized version of the Auslander-Buchsbaum theorem which applies to our case (see [3, (12.2)] or [6, Theorem (2.1)]). For the convenience of the reader we include a proof of this theorem in the graded case, which only makes use of the classical definition of depth as given above.

1.2.2. Lemma. *Let A and M be as above and assume that (A_0, \mathfrak{m}_0) is local. Then:*

- (1) $\dim_{A_0}(M) = \sup\{\dim_{A_0}(M_i) \mid i \in \mathbb{Z}\}$,
- (2) $\text{depth}_{A_0}(M) = \inf\{\text{depth}_{A_0}(M_i) \mid i \in \mathbb{Z} \text{ with } M_i \neq 0\}$,
- (3) $\text{projdim}_{A_0}(M) = \sup\{\text{projdim}_{A_0}(M_i) \mid i \in \mathbb{Z}\}$.

Proof. (1) By Lemma 1.1.1 there is an integer $s \in \mathbb{Z}$ so that $\text{ann}_{A_0}(M_k) = \text{ann}_{A_0}(M_s)$ for all $k \geq s$. In particular, for all $k \geq s$, $\dim_{A_0}(M_k) = \dim_{A_0}(M_s)$ and

$$\dim_{A_0}(M) = \dim_{A_0}(M_r \oplus M_{r-1} \oplus \dots \oplus M_{s-1} \oplus M_s),$$

where $r \in \mathbb{Z}$ is the smallest integer j with $M_j \neq 0$. The dimension of a finite direct sum of A_0 -modules is the maximum of the dimensions of its summands.

(2) If $r_1, \dots, r_s \in A_0$ is a regular sequence on M , then r_1, \dots, r_s is a regular sequence on M_i for all $i \in \mathbb{Z}$ with $M_i \neq 0$. Thus $\text{depth}_{A_0}(M) \leq \text{depth}_{A_0}(M_i)$ for all $i \in \mathbb{Z}$ with $M_i \neq 0$, and hence

$$\text{depth}_{A_0}(M) \leq \inf\{\text{depth}_{A_0}(M_i) \mid i \in \mathbb{Z} \text{ with } M_i \neq 0\}.$$

In order to show the other inequality we proceed by induction on $t = \text{depth}_{A_0}(M)$.

Note that by Lemma 1.1.3, $\text{Ass}_{A_0}(M)$ is a finite set.

If $t = 0$, then $\mathfrak{m}_0 \in \text{Ass}_{A_0}(M)$ and there is an $i \in \mathbb{Z}$ so that $\mathfrak{m}_0 \in \text{Ass}_{A_0}(M_i)$. Thus

$$\inf\{\text{depth}_{A_0}(M_i) \mid i \in \mathbb{Z} \text{ with } M_i \neq 0\} = 0.$$

Now assume that $t = \text{depth}_{A_0}(M) > 0$. This implies that

$$\bigcup_{\mathfrak{p} \in \text{Ass}_{A_0}(M)} \mathfrak{p} \neq \mathfrak{m}_0.$$

Consider an element

$$r \in \mathfrak{m}_0 \setminus \bigcup_{\mathfrak{p} \in \text{Ass}_{A_0}(M)} \mathfrak{p}.$$

Since r is regular on M , and therefore is regular on M_i for all $i \in \mathbb{Z}$ with $M_i \neq 0$, we obtain

$$\text{depth}_{A_0}(M/rM) = \text{depth}_{A_0}(M) - 1,$$

and for all $i \in \mathbb{Z}$ with $M_i \neq 0$,

$$\text{depth}_{A_0}(M_i/rM_i) = \text{depth}_{A_0}(M_i) - 1.$$

By the induction hypothesis

$$\text{depth}_{A_0}(M/rM) = \inf\{\text{depth}_{A_0}(M_i/rM_i) \mid i \in \mathbb{Z} \text{ and } M_i/rM_i \neq 0\}.$$

The assertion follows.

(3) For all $i \in \mathbb{Z}$ let $F_\bullet^{(i)}$ be a finite free resolution of M_i . Then

$$F_\bullet = \bigoplus_{i \in \mathbb{Z}} F_\bullet^{(i)}$$

is a free resolution of the A_0 -module M yielding

$$\text{projdim}_{A_0}(M) \leq \sup\{\text{projdim}_{A_0}(M_i) \mid i \in \mathbb{Z}\}.$$

In order to show the other inequality, assume that $\text{projdim}_{A_0}(M) = r$ and consider for all $i \in \mathbb{Z}$ the r th syzygy $T_r^{(i)}$ of M_i and the exact sequence

$$0 \longrightarrow T_r^{(i)} \longrightarrow F_{r-1}^{(i)} \longrightarrow \dots \longrightarrow F_0^{(i)} \longrightarrow M_i \longrightarrow 0.$$

By taking direct sums we see that

$$\bigoplus_{i \in \mathbb{Z}} T_r^{(i)}$$

is an r th syzygy of M and thus projective. Therefore every $T_r^{(i)}$ is a projective finitely generated A_0 -module. Since A_0 is a local Noetherian ring, every $T_r^{(i)}$ is a free A_0 -module and thus for all $i \in \mathbb{Z}$

$$\text{projdim}_{A_0}(M_i) \leq r.$$

This shows (3). □

1.2.3. Proposition. *Let A and M be as above with (A_0, \mathfrak{m}_0) a local ring. Then the Auslander-Buchsbaum formula holds for M as an A_0 -module. That is, if $\text{projdim}_{A_0}(M)$ is finite, then*

$$\text{depth}_{A_0}(M) + \text{projdim}_{A_0}(M) = \text{depth}(A_0).$$

Proof. Let $\text{projdim}_{A_0}(M) = r < \infty$. Then by Lemma 1.2.2(2) there is an $i \in \mathbb{Z}$ with $\text{projdim}_{A_0}(M) = \text{projdim}_{A_0}(M_i)$, and for all $j \in \mathbb{Z}$

$$\text{projdim}_{A_0}(M_j) \leq r.$$

The Auslander-Buchsbaum formula holds for finitely generated A_0 -modules

$$\text{depth}_{A_0}(M_j) + \text{projdim}_{A_0}(M_j) = \text{depth}_{A_0}(A_0) \quad \text{for all } j \in \mathbb{Z},$$

and therefore

$$\text{depth}_{A_0}(M_j) \geq \text{depth}_{A_0}(M_i) \quad \text{for all } j \in \mathbb{Z}.$$

Using Lemma 1.2.2(1), we conclude $\text{depth}_{A_0}(M) = \text{depth}_{A_0}(M_i)$. The Auslander-Buchsbaum formula for M_i then gives the desired formula. \square

2. OPENNESS OF THE CODEPTH LOCUS

Throughout this section we assume that $A = \bigoplus_{i \in \mathbb{N}_0} A_i$ is a graded Noetherian homogeneous ring and that $M = \bigoplus_{i \in \mathbb{Z}} M_i$ is a finitely generated A -module. Our aim is to generalize and/or modify existing theorems for finitely generated modules over Noetherian rings to the graded case where the module M is considered a module over the base ring A_0 . We begin with a result on the flat locus of the A_0 -module M .

2.1. The flat locus of M . Our first result is a modification of [8, Theorem 24.3]. The proof follows the proof in Matsumura’s book. A key observation is that for a finitely generated graded module M the localizations $M_{\mathfrak{p}}$ are I -adically separated for every ideal $I \subseteq (A_0)_{\mathfrak{p}}$.

Proposition. *Let A and M be as above. The flat locus of M as an A_0 -module*

$$U^0(M) = \{\mathfrak{p} \in \text{Spec}(A_0) \mid M_{\mathfrak{p}} \text{ is flat over } A_0\}$$

is open in $\text{Spec}(A_0)$.

Proof. According to Nagata’s criterion on the openness of loci [8, Theorem 24.2] we have to show:

- (a) If $\mathfrak{p}, \mathfrak{q} \in \text{Spec}(A_0)$ with $\mathfrak{p} \in U^0(M)$ and $\mathfrak{q} \subseteq \mathfrak{p}$, then $\mathfrak{q} \in U^0(M)$.
- (b) If $\mathfrak{p} \in U^0(M)$, then $U^0(M)$ contains a nonempty open subset of $V^0(\mathfrak{p}) = \{\mathfrak{n} \in \text{Spec}(A_0) \mid \mathfrak{p} \subseteq \mathfrak{n}\}$.

(a) is trivial. Let $\mathfrak{p} \in U^0(M)$, that is, assume that $M_{\mathfrak{p}}$ is flat over A_0 . Set $\bar{A}_0 = A_0/\mathfrak{p}$. By [8, Theorem 22.3] for every $\mathfrak{q} \in V^0(\mathfrak{p})$ the module $M_{\mathfrak{q}}$ is flat over A_0 if and only if $(M/\mathfrak{p}M)_{\mathfrak{q}}$ is flat over \bar{A}_0 and $\text{Tor}_1^{A_0}(M_{\mathfrak{q}}, \bar{A}_0) = 0$. A similar argument as in the proof of [8, Theorem 23.2] shows that $\text{Tor}_1^{A_0}(M, \bar{A}_0)$ is a finitely generated module over A . Therefore there is an element $a \in A_0 \setminus \mathfrak{p}$ so that $(\text{Tor}_1^{A_0}(M, \bar{A}_0))_a = 0$. By applying [8, Theorem 24.1] to the \bar{A}_0 -module $M/\mathfrak{p}M$ we obtain an element $b \in A_0 \setminus \mathfrak{p}$ so that $(M/\mathfrak{p}M)_b$ is a free $(\bar{A}_0)_b$ -module. Set $D_{ab}^0 = \{\mathfrak{q} \in \text{Spec}(A_0) \mid ab \notin \mathfrak{q}\}$. Then for all $\mathfrak{q} \in V^0(\mathfrak{p}) \cap D_{ab}^0$ we have that $\text{Tor}_1^{A_0}(M_{\mathfrak{q}}, \bar{A}_0) = 0$ and that $(M/\mathfrak{p}M)_{\mathfrak{q}}$ is flat over $(\bar{A}_0)_{\mathfrak{q}}$. Thus by [8, Theorem 22.3] the module $M_{\mathfrak{q}}$ is flat over $(A_0)_{\mathfrak{q}}$ and $M_{\mathfrak{q}}$ is flat over A_0 . \square

2.2. A proposition by Auslander. As before, let A be a Noetherian graded homogeneous ring and let M be a finitely generated A -module. The following Proposition is an extension of a proposition in EGA [4, (6.11.1) and (6.11.2)] to the (not finitely generated) A_0 -module M .

Proposition. *The function $\gamma : \text{Spec}(A_0) \rightarrow \mathbb{N}$ defined by*

$$\gamma(\mathfrak{p}) = \text{projdim}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) \quad \text{for all } \mathfrak{p} \in \text{Spec}(A_0)$$

is upper semicontinuous. That is, for all $n \in \mathbb{N}$ the set

$$U_n^0(M) = \{\mathfrak{p} \in \text{Spec}(A_0) \mid \text{projdim}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) \leq n\}$$

is open in $\text{Spec}(A_0)$.

Proof. Note that the ring A is the homomorphic image of the polynomial ring $B = A_0[x_1, \dots, x_t]$, and that, with the standard grading on the polynomial ring B , the graded B -module M is finitely generated. We may replace A by B and assume that A is a graded polynomial ring over A_0 . Let $\mathfrak{p} \in \text{Spec}(A_0)$ with $\text{projdim}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) \leq n$.

Consider a graded finitely generated free resolution of the A -module M :

$$F_n \xrightarrow{\varphi_n} F_{n-1} \xrightarrow{\varphi_{n-1}} \dots \xrightarrow{\varphi_1} F_1 \xrightarrow{\varphi_0} M \rightarrow 0,$$

where the F_i are finitely generated graded free A -modules and the φ_i are homogeneous A -linear maps. Let T be the n th syzygy of M , yielding an exact sequence of graded A -modules:

$$(*) \quad 0 \rightarrow T \xrightarrow{\delta} F_{n-1} \xrightarrow{\varphi_{n-1}} \dots \xrightarrow{\varphi_1} F_1 \xrightarrow{\varphi_0} M \rightarrow 0.$$

Since all the homogeneous parts of F_i are free A_0 -modules and since T is a graded A -module, we obtain for all $k \in \mathbb{Z}$ an exact sequence of A_0 -modules

$$0 \rightarrow T_k \xrightarrow{(\delta)_k} (F_{n-1})_k \xrightarrow{(\varphi_{n-1})_k} \dots \xrightarrow{(\varphi_1)_k} (F_1)_k \xrightarrow{(\varphi_0)_k} M_k \rightarrow 0$$

with $(F_i)_k$ a finitely generated free A_0 -module. Therefore by considering $(*)$ as an exact sequence of A_0 -modules we obtain that every module F_i is free over A_0 and T is an n th syzygy of the A_0 -module M . Localization at \mathfrak{p} yields exact sequences:

$$0 \rightarrow T_{\mathfrak{p}} \xrightarrow{\delta_{\mathfrak{p}}} (F_{n-1})_{\mathfrak{p}} \xrightarrow{(\varphi_{n-1})_{\mathfrak{p}}} \dots \xrightarrow{(\varphi_1)_{\mathfrak{p}}} (F_1)_{\mathfrak{p}} \xrightarrow{(\varphi_0)_{\mathfrak{p}}} M_{\mathfrak{p}} \rightarrow 0.$$

Since $\text{projdim}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) \leq n$, it follows that $T_{\mathfrak{p}}$ is a projective $(A_0)_{\mathfrak{p}}$ -module. Therefore $T_{\mathfrak{p}}$ is a free $(A_0)_{\mathfrak{p}}$ -module. Since T is a finitely generated graded A -module, it follows from Proposition 2.1 that the set

$$U^0(T) = \{\mathfrak{q} \in \text{Spec}(A_0) \mid T_{\mathfrak{q}} \text{ is a flat over } (A_0)_{\mathfrak{q}}\}$$

is an open subset of $\text{Spec}(A_0)$. Since T is a finitely generated graded A -module,

$$T = \bigoplus_{i \in \mathbb{Z}} T_i,$$

we have for $\mathfrak{q} \in \text{Spec}(A_0)$

$$T_{\mathfrak{q}} = \bigoplus_{i \in \mathbb{Z}} (T_i)_{\mathfrak{q}}.$$

If $T_{\mathfrak{q}}$ is flat over $(A_0)_{\mathfrak{q}}$, then, by [1, chapter 1, §2.3, Proposition 2], for all $i \in \mathbb{Z}$, $(T_i)_{\mathfrak{q}}$ is flat over $(A_0)_{\mathfrak{q}}$. Since every $(T_i)_{\mathfrak{q}}$ is a finitely generated $(A_0)_{\mathfrak{q}}$ -module, each $(T_i)_{\mathfrak{q}}$ is a free $(A_0)_{\mathfrak{q}}$ -module and

$$U^0(T) = \{\mathfrak{q} \in \text{Spec}(A_0) \mid T_{\mathfrak{q}} \text{ is a free over } (A_0)_{\mathfrak{q}}\}.$$

This shows that $\mathfrak{p} \in U^0(T)$ and

$$U^0(T) \subseteq \{\mathfrak{q} \in \text{Spec}(A_0) \mid \text{projdim}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) \leq n\}.$$

The set $\{\mathfrak{q} \in \text{Spec}(A_0) \mid \text{projdim}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) \leq n\}$ is thus open in $\text{Spec}(A_0)$. □

2.3. A dimension formula.

Proposition. *Let A and M be as above. Assume that A_0 is catenary and let \mathfrak{p} be a prime ideal in A_0 with $\mathfrak{p} \in \text{Supp}_{A_0}(M)$. Then there is an open subset U in $\text{Spec}(A_0)$ such that $\mathfrak{p} \in U$, and for all $\mathfrak{q} \in U \cap V^0(\mathfrak{p})$ we have*

$$\dim(M_{\mathfrak{q}}) = \dim(M_{\mathfrak{p}}) + \dim((A_0/\mathfrak{p})_{\mathfrak{q}}).$$

Proof. Set $S = A_0/\text{ann}_{A_0}(M)$ and choose an element $a \in S \setminus \mathfrak{p}$ so that the following equality on the set of minimal primes holds:

$$\text{Min}(S_{\mathfrak{p}}) = \text{Min}(S_a).$$

Assume that $\dim(M_{\mathfrak{p}}) = \text{ht}(\mathfrak{p}S) = t$ and choose elements $y_1, y_2, \dots, y_t \in S$ so that

$$y_1 \text{ not in a minimal prime of } S_{\mathfrak{p}},$$

$$y_2 \text{ not in a minimal prime of } y_1 S_{\mathfrak{p}},$$

...

$$y_t \text{ not in a minimal prime of } (y_1, \dots, y_{t-1})S_{\mathfrak{p}}.$$

Then there is an element $b \in S \setminus \mathfrak{p}$ so that

$$y_1 \text{ not in a minimal prime of } S_b,$$

$$y_2 \text{ not in a minimal prime of } y_1 S_b,$$

...

$$y_t \text{ not in a minimal prime of } (y_1, \dots, y_{t-1})S_b.$$

Let a, b also denote preimages of a and b in A_0 and put $U = D_{ab} = \{\mathfrak{q} \in \text{Spec}(A_0) \mid ab \notin \mathfrak{q}\}$. Then for every $\mathfrak{q} \in U \cap V^0(\mathfrak{p})$ the elements y_1, \dots, y_t extend to a system of parameters of $S_{\mathfrak{q}}$. Since $S_{\mathfrak{p}}$ and $S_{\mathfrak{q}}$ have the same set of minimal primes and since S is catenary, we obtain that

$$\dim(S_{\mathfrak{q}}) = \dim(S_{\mathfrak{p}}) + \dim((S/\mathfrak{p})_{\mathfrak{q}}).$$

This is the same as

$$\dim(M_{\mathfrak{q}}) = \dim(M_{\mathfrak{p}}) + \dim((A_0/\mathfrak{p})_{\mathfrak{q}}). \quad \square$$

2.4. The special case of A_0 regular. Let (R, \mathfrak{m}) be a local Noetherian ring and M an R -module. Then we define

$$\text{codepth}_R(M) := \dim_R(M) - \text{depth}_R(M).$$

As usual the depth of the zero module is defined to be ∞ , and the dimension of the zero module is $-\infty$, implying that the codepth of the zero module is $-\infty$.

The following proposition extends a result by Auslander [4, (6.11.2)] to the graded case.

Proposition. *Let A and M be as above and assume that A_0 is a homomorphic image of a regular ring. The function $\varphi: \text{Spec}(A_0) \rightarrow \mathbb{N}$ defined by*

$$\varphi(\mathfrak{p}) = \text{codepth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) \quad \text{for all } \mathfrak{p} \in \text{Spec}(A_0)$$

is upper semicontinuous, that is, for all $n \in \mathbb{N}$, the set

$$U_{C_n}^0(M) = \{\mathfrak{p} \in \text{Spec}(A_0) \mid \text{codepth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) \leq n\}$$

is open in $\text{Spec}(A_0)$.

Proof. If A_0 is a homomorphic image of a regular ring R_0 , then the dimension and the depth of the R_0 -module M are identical to the dimension and depth of M considered as an R_0 -module. If we show that the set

$$\tilde{U}_{C_n}^0(M) = \{\mathfrak{q} \in \text{Spec}(R_0) \mid \text{codepth}_{(R_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) \leq n\}$$

is open in $\text{Spec}(R_0)$ (where M is considered an R_0 -module), then the corresponding set for the A_0 -module M is given by

$$U_{C_n}^0(M) = \tilde{U}_{C_n}^0(M) \cap V(J),$$

where $A_0 = R_0/J$. Thus we may assume that A_0 is a regular ring. We may also assume that A is a polynomial ring over A_0 equipped with the standard grading.

Let $\mathfrak{p} \in \text{Spec}(A_0)$. By Proposition 1.2.3, the Auslander-Buchsbaum formula holds:

$$\text{depth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) = \text{depth}((A_0)_{\mathfrak{p}}) - \text{projdim}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}).$$

Let $I = \text{ann}_{A_0}(M)$. By Lemma 1.1.3, $I_{\mathfrak{p}} = \text{ann}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}})$, and we have that

$$\dim_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) = \dim((A_0)_{\mathfrak{p}}) - \text{ht}(I(A_0)_{\mathfrak{p}}).$$

Suppose that $\mathfrak{p} \in \text{Spec}(A_0)$ is such that

$$\text{codepth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) \leq n.$$

If $M_{\mathfrak{p}} = 0$, then $\mathfrak{p} \not\supseteq I$. Take an element $a \in I \cap (A_0 \setminus \mathfrak{p})$. Then for all

$$\mathfrak{q} \in D_a = \{\mathfrak{w} \in \text{Spec}(A_0) \mid a \notin \mathfrak{w}\}$$

we have that $M_{\mathfrak{q}} = 0$ and $\text{codepth}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) = -\infty \leq n$.

If $M_{\mathfrak{p}} \neq 0$ pick an element $a_1 \in A_0 \setminus \mathfrak{p}$ so that $(A_0)_{\mathfrak{p}}$ and $(A_0)_{a_1}$ have the same minimal primes and put $U_1 = D_{a_1} = \{\mathfrak{w} \in \text{Spec}(A_0) \mid a_1 \notin \mathfrak{w}\}$. Then for all $\mathfrak{q} \in U_1 \cap V^0(I)$,

$$\text{ht}(I(A_0)_{\mathfrak{q}}) \geq \text{ht}(I(A_0)_{\mathfrak{p}}).$$

Let $\text{projdim}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) = t$. Then by Proposition 2.2 there is an open subset U_2 in $\text{Spec}(A_0)$ so that

$$\text{projdim}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) \leq t \quad \text{for all } \mathfrak{q} \in U_2.$$

Using the Auslander-Buchsbaum formula and the fact that A_0 is regular, we obtain for all $\mathfrak{q} \in U_2 \cap U_1 \cap V^0(I)$:

$$\begin{aligned} \text{codepth}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) &= \dim_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) - \text{depth}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) \\ &= \dim((A_0)_{\mathfrak{q}}) - \text{ht}(I(A_0)_{\mathfrak{q}}) - \dim((A_0)_{\mathfrak{q}}) + \text{projdim}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) \\ &= \text{projdim}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) - \text{ht}(I(A_0)_{\mathfrak{q}}). \end{aligned}$$

This implies that for all $\mathfrak{q} \in U = U_1 \cap U_2$,

$$\text{codepth}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) \leq \text{codepth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}),$$

and it follows that $U_{C_n}^0(M)$ is an open subset of $\text{Spec}(A_0)$. □

2.5. A local formula. Using the fact that a complete local Noetherian ring is the homomorphic image of a regular local ring, we obtain a result similar to [4, (6.11.5)]:

Lemma. *Let A be a Noetherian graded homogeneous ring and let M be a finitely generated graded A -module. Then for all prime ideals $\mathfrak{p}, \mathfrak{q} \in \text{Spec}(A_0)$ with $\mathfrak{p} \subseteq \mathfrak{q}$ we have that*

$$\text{codepth}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) \geq \text{codepth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}).$$

Proof. By replacing A_0 by $(A_0)_{\mathfrak{q}}$ (and A by $A_{\mathfrak{q}}$) we may assume that (A_0, \mathfrak{m}_0) is a local ring. Then we have to show

$$\text{codepth}_{A_0}(M) \geq \text{codepth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}).$$

Let $\widehat{\mathfrak{p}} \in \text{Spec}(\widehat{A}_0)$ be a minimal prime ideal over $\mathfrak{p}\widehat{A}_0$. Then $\widehat{\mathfrak{p}} \cap A_0 = \mathfrak{p}$ and $(\widehat{A}_0)_{\widehat{\mathfrak{p}}}$ is flat over $(A_0)_{\mathfrak{p}}$ with trivial special fiber. Moreover,

$$\begin{aligned} M_{\mathfrak{p}} \otimes_{(A_0)_{\mathfrak{p}}} (\widehat{A}_0)_{\widehat{\mathfrak{p}}} &= \left(\bigoplus_{i \in \mathbb{Z}} (M_i)_{\mathfrak{p}} \right) \otimes_{(A_0)_{\mathfrak{p}}} (\widehat{A}_0)_{\widehat{\mathfrak{p}}} \\ &= \bigoplus_{i \in \mathbb{Z}} ((M_i)_{\mathfrak{p}} \otimes_{(A_0)_{\mathfrak{p}}} (\widehat{A}_0)_{\widehat{\mathfrak{p}}}) \\ &\cong \bigoplus_{i \in \mathbb{Z}} (\widehat{M}_i)_{\widehat{\mathfrak{p}}}, \end{aligned}$$

where $\widehat{M}_i \cong M_i \otimes_{A_0} \widehat{A}_0$. We have that

$$\begin{aligned} \text{depth}_{A_0}(M) &= \inf\{\text{depth}_{A_0}(M_i) \mid M_i \neq 0\}, \\ \dim_{A_0}(M) &= \sup\{\dim_{A_0}(M_i) \mid i \in \mathbb{Z}\}. \end{aligned}$$

By [8, Theorem 23.3], for all $i \in \mathbb{Z}$,

$$\begin{aligned} \text{depth}_{(\widehat{A}_0)_{\widehat{\mathfrak{p}}}}((\widehat{M}_i)_{\widehat{\mathfrak{p}}}) &= \text{depth}_{(A_0)_{\mathfrak{p}}}((M_i)_{\mathfrak{p}}) + \text{depth}((\widehat{A}_0)_{\widehat{\mathfrak{p}}}/\mathfrak{p}(\widehat{A}_0)_{\widehat{\mathfrak{p}}}) \\ &= \text{depth}_{(A_0)_{\mathfrak{p}}}((M_i)_{\mathfrak{p}}), \end{aligned}$$

and by [8, Theorem 15.1],

$$\begin{aligned} \dim_{(\widehat{A}_0)_{\widehat{\mathfrak{p}}}}((\widehat{M}_i)_{\widehat{\mathfrak{p}}}) &= \dim_{(A_0)_{\mathfrak{p}}}((M_i)_{\mathfrak{p}}) + \dim((\widehat{A}_0)_{\widehat{\mathfrak{p}}}/\mathfrak{p}(\widehat{A}_0)_{\widehat{\mathfrak{p}}}) \\ &= \dim_{(A_0)_{\mathfrak{p}}}((M_i)_{\mathfrak{p}}). \end{aligned}$$

Let

$$\widetilde{M} := \bigoplus_{i \in \mathbb{Z}} \widehat{M}_i \cong M \otimes_{A_0} \widehat{A}_0,$$

and note that \widetilde{M} is a finitely generated graded module over the Noetherian homogeneous graded ring

$$\widetilde{A} := A \otimes_{A_0} \widehat{A}_0.$$

The computation above shows that

$$\text{codepth}_{(\widehat{A}_0)_{\widehat{\mathfrak{p}}}}(\widetilde{M}_{\widehat{\mathfrak{p}}}) = \text{codepth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) =: n.$$

Since \widehat{A}_0 is a homomorphic image of a regular local ring, by Proposition 2.3 the set $U_{C_{n-1}}^0(\widetilde{M})$ is open in $\text{Spec}(\widehat{A}_0)$. This implies that

$$\text{codepth}_{\widehat{A}_0}(\widetilde{M}) \geq \text{codepth}_{(\widehat{A}_0)_{\widehat{\mathfrak{p}}}}(\widetilde{M}_{\widehat{\mathfrak{p}}}).$$

The same argument as above shows that

$$\text{codepth}_{\widehat{A}_0}(\widetilde{M}) = \text{codepth}_{A_0}(M),$$

which proves the claim

$$\text{codepth}_{A_0}(M) \geq \text{codepth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}). \quad \square$$

2.6. Formulas for depth and codepth. In this section we make the same assumption as at the beginning, namely, A is a positively graded Noetherian homogeneous ring and M is a finitely generated graded A -module. The following proposition is the graded version of [4, (6.10.6)]:

2.6.1. Proposition. *Let A and M be as above and assume that A is excellent. Then for every $\mathfrak{p} \in \text{Spec}(A_0)$ there is an open subset $U^0 \subseteq \text{Spec}(A_0)$ with $\mathfrak{p} \in U^0$ so that for all $\mathfrak{q} \in U^0 \cap V^0(\mathfrak{p})$,*

$$\text{depth}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) = \text{depth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) + \text{depth}((A_0/\mathfrak{p})_{\mathfrak{q}}).$$

Proof. Let $\mathfrak{p} \in \text{Spec}(A_0)$. Then by Lemma 2.5 for all $\mathfrak{q} \in V^0(\mathfrak{p})$,

$$\text{codepth}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) \geq \text{codepth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}),$$

or equivalently,

$$(*) \quad \dim_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) - \text{depth}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) \geq \dim_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) - \text{depth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}).$$

According to Proposition 2.3 there is an open subset $U_1 \subseteq \text{Spec}(A_0)$ with $\mathfrak{p} \in U_1$ so that for all $\mathfrak{q} \in U_1 \cap V^0(\mathfrak{p})$,

$$\dim_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) = \dim_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) + \dim((A_0/\mathfrak{p})_{\mathfrak{q}}).$$

Since A_0 is excellent, there is an open subset $U_2 \subseteq \text{Spec}(A_0)$ so that $\mathfrak{p} \in U_2$, and for all $\mathfrak{q} \in U_2 \cap V^0(\mathfrak{p})$ the local ring

$$(A_0/\mathfrak{p})_{\mathfrak{q}} \text{ is Cohen-Macaulay.}$$

There is also an open subset $U_3 \subseteq \text{Spec}(A_0)$ so that $\mathfrak{p} \in U_3$, and for all $\mathfrak{q} \in U_3 \cap V^0(\mathfrak{p})$ we have equality on the set of minimal primes:

$$\text{Min}_{(A_0)_{\mathfrak{q}}}(I(A_0)_{\mathfrak{q}}) = \text{Min}_{(A_0)_{\mathfrak{p}}}(I(A_0)_{\mathfrak{p}}),$$

where $I := \text{ann}_{A_0}(M)$ denotes the A_0 -annihilator of M . In particular, for all $\mathfrak{q} \in U_3 \cap V^0(\mathfrak{p})$,

$$\text{ht}(I(A_0)_{\mathfrak{q}}) = \text{ht}(I(A_0)_{\mathfrak{p}}).$$

Put $\widetilde{U}_1 = U_1 \cap U_2 \cap U_3$; then for all $\mathfrak{q} \in \widetilde{U}_1 \cap V^0(\mathfrak{p})$,

$$\dim_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) = \dim((A_0/I)_{\mathfrak{q}}) \quad \text{and} \quad \dim_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) = \dim((A_0/I)_{\mathfrak{p}}).$$

Since A is excellent, the ring A_0 is universally catenary, and for all $\mathfrak{q} \in \widetilde{U}_1 \cap V^0(\mathfrak{p})$,

$$\dim((A_0/I)_{\mathfrak{q}}) - \dim((A_0/I)_{\mathfrak{p}}) = \dim((A_0/\mathfrak{p})_{\mathfrak{q}}) = \text{depth}((A_0/\mathfrak{p})_{\mathfrak{q}}).$$

From (*) we obtain

$$\text{depth}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) - \text{depth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) \leq \text{depth}((A_0/\mathfrak{p})_{\mathfrak{q}})$$

for all $\mathfrak{q} \in \widetilde{U}_1 \cap V^0(\mathfrak{p})$.

In order to prove the other inequality,

$$\text{depth}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) - \text{depth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) \geq \text{depth}((A_0/\mathfrak{p})_{\mathfrak{q}}),$$

assume that $\text{depth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) = t$ and let $f_1, \dots, f_t \in \mathfrak{p}$ be such that f_1, \dots, f_t is a regular sequence on $M_{\mathfrak{p}}$. A prime avoidance argument shows that there is an element $a \in A_0 \setminus \mathfrak{p}$ so that f_1, \dots, f_t is a regular sequence on M_a . (The argument again makes use of the fact that the sets $\text{Ass}_{A_0}(M)$ and $\text{Ass}_{A_0}(M/(f_1, \dots, f_i)M)$ for all $1 \leq i \leq t$ are finite.)

Put

$$\bar{M} := M/(f_1, \dots, f_t)M,$$

and consider the associated graded module

$$\text{gr}_{\mathfrak{p}}(\bar{M}) = \bigoplus_{i \in \mathbb{N}} \mathfrak{p}^i \bar{M} / \mathfrak{p}^{i+1} \bar{M}.$$

The module \bar{M} is finitely generated over A , and $\text{gr}_{\mathfrak{p}}(\bar{M})$ is a finitely generated $\text{gr}_{\mathfrak{p}}(A)$ -module. Also note that $\text{gr}_{\mathfrak{p}}(A)$ is a finitely generated algebra over $A/\mathfrak{p}A$ and that $A/\mathfrak{p}A$ is a finitely generated algebra over A_0/\mathfrak{p} . Thus $\text{gr}_{\mathfrak{p}}(A)$ is a finitely generated A_0/\mathfrak{p} -algebra. By [8, Theorem 24.1] there is an element $b \in A_0 \setminus \mathfrak{p}$ so that the $(A_0/\mathfrak{p})_b$ -module

$$\text{gr}_{\mathfrak{p}}(\bar{M})_b = \bigoplus_{i \in \mathbb{N}} (\mathfrak{p}^i \bar{M} / \mathfrak{p}^{i+1} \bar{M})_b$$

is free. Set $\tilde{U}_2 = D_b = \{\mathfrak{q} \in \text{Spec}(A_0) \mid b \notin \mathfrak{q}\}$ and fix a prime ideal $\mathfrak{q} \in \tilde{U}_2 \cap V^0(\mathfrak{p})$. Assume that

$$\text{depth}((A_0/\mathfrak{p})_{\mathfrak{q}}) = s,$$

and let $g_1, \dots, g_s \in \mathfrak{q}$ be such that g_1, \dots, g_s is a regular sequence on $(A_0/\mathfrak{p})_{\mathfrak{q}}$.

Claim 1. g_1 is a regular element on $\bar{M}_{\mathfrak{q}}$.

Claim 2. Set $N_1 := \bar{M}_{\mathfrak{q}}/g_1 \bar{M}_{\mathfrak{q}}$; then $\text{gr}_{\mathfrak{p}}(N_1) \cong \text{gr}_{\mathfrak{p}}(\bar{M}_{\mathfrak{q}})/g_1 \text{gr}_{\mathfrak{p}}(\bar{M}_{\mathfrak{q}})$.

Assuming the claims, we finish the proof. From the second claim it follows that $\text{gr}_{\mathfrak{p}}(N_1)$ is a free $(A_0/(g_1, \mathfrak{p})A_0)_{\mathfrak{q}}$ -module. Since g_2 is a regular element on $(A_0/(g_1, \mathfrak{p})A_0)_{\mathfrak{q}}$, we may apply Claims 1 and 2 to N_1 . Note that N_1 is also a finitely generated graded $A_{\mathfrak{q}}$ -module. This yields that g_2 is a regular element on N_1 and that with $N_2 = N_1/g_2 N_1$,

$$\text{gr}_{\mathfrak{p}}(N_2) \cong \text{gr}_{\mathfrak{p}}(N_1)/g_2 \text{gr}_{\mathfrak{p}}(N_1).$$

An induction argument yields that g_1, \dots, g_s is a regular sequence on $\bar{M}_{\mathfrak{q}}$, and we have that

$$\text{depth}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) \geq \text{depth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) + \text{depth}((A_0/\mathfrak{p})_{\mathfrak{q}}).$$

This inequality holds for all $\mathfrak{q} \in \tilde{U}_2 \cap V^0(\mathfrak{p})$. Assuming the claims the proposition is now proved with $U^0 = \tilde{U}_1 \cap \tilde{U}_2$.

In order to prove the claims, set $g = g_1$ and $N = N_1$.

Proof of Claim 1. Let $z \in \bar{M}_{\mathfrak{q}}$ with $gz = 0$. Consider the image \bar{z} of z in $\bar{M}_{\mathfrak{q}}/\mathfrak{p}\bar{M}_{\mathfrak{q}}$. Since $\bar{M}_{\mathfrak{q}}/\mathfrak{p}\bar{M}_{\mathfrak{q}}$ is a free module over $(A_0/\mathfrak{p})_{\mathfrak{q}}$ and since g is regular on $(A_0/\mathfrak{p})_{\mathfrak{q}}$, we obtain that $\bar{z} = 0$ and $z \in \mathfrak{p}\bar{M}_{\mathfrak{q}}$. Now consider the image of z in $\mathfrak{p}\bar{M}_{\mathfrak{q}}/\mathfrak{p}^2\bar{M}_{\mathfrak{q}}$ and repeat the argument. This yields

$$z \in \bigcap_{j=0}^{\infty} \mathfrak{p}^j \bar{M}_{\mathfrak{q}}.$$

Note that

$$\overline{M}_q = \bigoplus_{i \in \mathbb{Z}} (\overline{M}_i)_q \quad \text{with} \quad (\overline{M}_i)_q = (M_i)_q / (f_1, \dots, f_t)(M_i)_q.$$

In particular,

$$\mathfrak{p}^j \overline{M}_q = \bigoplus_{i \in \mathbb{Z}} \mathfrak{p}^j (\overline{M}_i)_q,$$

and every $(\overline{M}_i)_q$ is a finitely generated $(A_0)_q$ -module. This shows that $z = 0$.

Proof of Claim 2. By assumption, we have that $\text{gr}_{\mathfrak{p}}(\overline{M}_q)$ is a free $(A_0/\mathfrak{p})_q$ -module and $\mathfrak{p}^j \overline{M}_q / \mathfrak{p}^{j+1} \overline{M}_q$ is a direct summand of $\text{gr}_{\mathfrak{p}}(\overline{M}_q)$. Thus $\mathfrak{p}^j \overline{M}_q / \mathfrak{p}^{j+1} \overline{M}_q$ is a free $(A_0/\mathfrak{p})_q$ -module and g is regular on $(A_0/\mathfrak{p})_q$. Therefore

$$(**) \quad \mathfrak{p}^j \overline{M}_q \cap g \overline{M}_q = g \mathfrak{p}^j \overline{M}_q$$

and thus

$$\begin{aligned} \mathfrak{p}^j \overline{M}_q / g \mathfrak{p}^j \overline{M}_q &\cong \mathfrak{p}^j \overline{M}_q / (\mathfrak{p}^j \overline{M}_q \cap g \overline{M}_q) \\ &\cong \mathfrak{p}^j (\overline{M}_q / g \overline{M}_q). \end{aligned}$$

From the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathfrak{p}^{j+1} N & \longrightarrow & \mathfrak{p}^j N & \longrightarrow & \mathfrak{p}^j N / \mathfrak{p}^{j+1} N & \longrightarrow & 0 \\ & & \parallel & & \parallel & & \parallel & & \\ 0 & \longrightarrow & \mathfrak{p}^{j+1} (\overline{M}_q / g \overline{M}_q) & \longrightarrow & \mathfrak{p}^j (\overline{M}_q / g \overline{M}_q) & \longrightarrow & \mathfrak{p}^j (\overline{M}_q / g \overline{M}_q) / \mathfrak{p}^{j+1} (\overline{M}_q / g \overline{M}_q) & \longrightarrow & 0 \\ & & \downarrow \cong & & \downarrow \cong & & \downarrow & & \\ 0 & \longrightarrow & \mathfrak{p}^{j+1} \overline{M}_q / g \mathfrak{p}^{j+1} \overline{M}_q & \longrightarrow & \mathfrak{p}^j \overline{M}_q / g \mathfrak{p}^j \overline{M}_q & \longrightarrow & \mathfrak{p}^j \overline{M}_q / (g \mathfrak{p}^j \overline{M}_q + \mathfrak{p}^{j+1} \overline{M}_q) & \longrightarrow & 0 \end{array}$$

we obtain that

$$\begin{aligned} \text{gr}_{\mathfrak{p}}(N) &= \bigoplus_{j \in \mathbb{N}} \mathfrak{p}^j N / \mathfrak{p}^{j+1} N \\ &\cong \bigoplus_{j \in \mathbb{N}} \mathfrak{p}^j \overline{M}_q / (g \mathfrak{p}^j \overline{M}_q + \mathfrak{p}^{j+1} \overline{M}_q) \\ &\cong \bigoplus_{j \in \mathbb{N}} (\mathfrak{p}^j \overline{M}_q / \mathfrak{p}^{j+1} \overline{M}_q) / g (\mathfrak{p}^j \overline{M}_q / \mathfrak{p}^{j+1} \overline{M}_q) \\ &\cong \text{gr}_{\mathfrak{p}}(\overline{M}_q) / g (\text{gr}_{\mathfrak{p}}(\overline{M}_q)). \end{aligned}$$

This proves the claim, and finishes the proof. □

Similar to [4, (6.11.8.1)] we have in the graded case:

2.6.2. Corollary. *Let A and M be as above and assume that A is excellent. Then for every $\mathfrak{p} \in \text{Spec}(A_0)$ there is an open subset $U^0 \subseteq \text{Spec}(A_0)$ with $\mathfrak{p} \in U^0$, so that for all $\mathfrak{q} \in U^0 \cap V^0(\mathfrak{p})$,*

$$\text{codepth}_{(A_0)_q}(M_q) = \text{codepth}_{(A_0)_p}(M_p) + \text{codepth}((A_0)_q / \mathfrak{p}(A_0)_q).$$

Proof. Let $\mathfrak{p} \in \text{Spec}(A_0)$ and let U_1^0 be as in Proposition 2.6.1, so that $\mathfrak{p} \in U_1^0$, and for all $\mathfrak{q} \in U_1^0 \cap V^0(\mathfrak{p})$,

$$\text{depth}_{(A_0)_q}(M_q) = \text{depth}_{(A_0)_p}(M_p) + \text{depth}((A_0)_q / \mathfrak{p}(A_0)_q).$$

By Proposition 2.3 there is an open subset U_2^0 in $\text{Spec}(A_0)$, so that $\mathfrak{p} \in U_2^0$, and for all $\mathfrak{q} \in U_2^0 \cap V^0(\mathfrak{p})$,

$$\dim_{(A_0)_q}(M_q) = \dim_{(A_0)_p}(M_p) + \dim((A_0/\mathfrak{p})_q).$$

Thus with $U^0 = U_1^0 \cap U_2^0$ we have that $\mathfrak{p} \in U^0$, and for all $\mathfrak{q} \in U^0 \cap V^0(\mathfrak{p})$,

$$\text{codepth}_{(A_0)_q}(M_q) = \text{codepth}_{(A_0)_p}(M_p) + \text{codepth}((A_0)_q / \mathfrak{p}(A_0)_q). \quad \square$$

We are now ready to prove the graded version of [4, (6.11.2)(a)].

2.6.3. Theorem. *Let $A = \bigoplus_{i \in \mathbb{N}} A_i$ be an excellent graded homogeneous ring and let $M = \bigoplus_{i \in \mathbb{Z}} M_i$ be a finitely generated graded A -module. Then for all $n \in \mathbb{N}$ the set*

$$U_{C_n}^0(M) = \{\mathfrak{p} \in \text{Spec}(A_0) \mid \text{codepth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) \leq n\}$$

is open in $\text{Spec}(A_0)$.

Proof. According to Nagata’s criterion on openness of loci (see [8, Theorem 24.2]) we need to show:

- (a) If $\mathfrak{p}, \mathfrak{q} \in \text{Spec}(A_0)$ with $\mathfrak{q} \subseteq \mathfrak{p}$ and $\mathfrak{p} \in U_{C_n}^0(M)$, then $\mathfrak{q} \in U_{C_n}^0(M)$.
 - (b) If $\mathfrak{p} \in U_{C_n}^0(M)$, then $U_{C_n}^0(M)$ contains a nonempty open subset of $V(\mathfrak{p})$.
- (a) Let $\mathfrak{p}, \mathfrak{q} \in \text{Spec}(A_0)$ with $\mathfrak{q} \subseteq \mathfrak{p}$. By Lemma 2.5

$$\text{codepth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) \geq \text{codepth}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}),$$

and thus $\mathfrak{p} \in U_{C_n}^0(M)$ implies that $\mathfrak{q} \in U_{C_n}^0(M)$.

(b) Let $\mathfrak{p} \in U_{C_n}^0(M)$. By Corollary 2.6.2 there is an open subset U_1^0 in $\text{Spec}(A_0)$, so that $\mathfrak{p} \in U_1^0$, and for all $\mathfrak{q} \in U_1^0 \cap V^0(\mathfrak{p})$,

$$\text{codepth}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) = \text{codepth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}) + \text{codepth}((A_0)_{\mathfrak{q}}/\mathfrak{p}(A_0)_{\mathfrak{q}}).$$

Since A and A_0 are excellent, there is an open subset U_2^0 in $\text{Spec}(A_0)$, so that $\mathfrak{p} \in U_2^0$, and for all $\mathfrak{q} \in U_2^0 \cap V^0(\mathfrak{p})$, the ring $(A_0/\mathfrak{p})_{\mathfrak{q}}$ is Cohen-Macaulay. Therefore with $U^0 = U_1^0 \cap U_2^0$ we have that $\mathfrak{p} \in U^0$, and for all $\mathfrak{q} \in U^0 \cap V^0(\mathfrak{p})$,

$$\text{codepth}_{(A_0)_{\mathfrak{q}}}(M_{\mathfrak{q}}) = \text{codepth}_{(A_0)_{\mathfrak{p}}}(M_{\mathfrak{p}}).$$

This implies that $U^0 \cap V^0(\mathfrak{p}) \subseteq U_{C_n}^0(M)$, and the theorem is proved. □

2.6.4. Corollary. *Let A and M be as in Theorem 2.6.3. Then the Cohen-Macaulay locus of the A_0 -module M ,*

$$U_{CM}^0(M) = U_{C_0}^0(M) = \{\mathfrak{p} \in \text{Spec}(A_0) \mid M_{\mathfrak{p}} \text{ is a CM module over } (A_0)_{\mathfrak{p}}\},$$

is open in $\text{Spec}(A_0)$. □

3. OPENNESS OF THE (S_n) -LOCUS

Throughout this section we assume that $R = A_0$ is the base ring of a graded Noetherian homogeneous ring $A = \bigoplus_{i \geq 0} A_i$ and M is a finitely generated graded A -module. This includes the case of a finitely generated module M over a Noetherian ring R . For those modules we prove that the openness of the C_n -loci of M implies the openness of the (S_k) -loci of M . The argument is due to Grothendieck [4, (5.7.2) and (6.11.2)(b)], but we include it here for the convenience of the reader. The proof also shows that the (S_k) -loci of M only depend on the C_n -loci of M and on the annihilator of M , so that two R -modules M and N with the same annihilators and C_n -loci have identical (S_k) -loci.

Let M be an R -module and suppose that for all $n \in \mathbb{N}_0$, the set

$$U_{C_n}(M) = \{\mathfrak{p} \in \text{Spec}(R) \mid \text{codepth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) \leq n\}$$

is open in $\text{Spec}(R)$. Define

$$Z_n = V(\mathfrak{b}_n) = \text{Spec}(R) \setminus U_{C_n}(M),$$

where $\mathfrak{b}_n \subseteq R$ is a reduced ideal. Obviously, for all $n \in \mathbb{N}$,

$$U_{C_n}(M) \subseteq U_{C_{n+1}}(M),$$

and therefore

$$Z_{n+1} \subseteq Z_n \quad \text{and} \quad \mathfrak{b}_n \subseteq \mathfrak{b}_{n+1}.$$

Since R is Noetherian, there is an $m \in \mathbb{N}$ so that for all $t \in \mathbb{N}$,

$$\mathfrak{b}_m = \mathfrak{b}_{m+t} \quad \text{and} \quad Z_m = Z_{m+t}.$$

3.1. Lemma. *Let $m \in \mathbb{N}$ be as above. Then $Z_m = \emptyset$.*

Proof. If $\mathfrak{p} \in Z_m$, then $\mathfrak{p} \in Z_{m+t}$ for all $t \in \mathbb{N}$. By definition of Z_{m+t} ,

$$\text{codepth}_{(R)_{\mathfrak{p}}}(M_{\mathfrak{p}}) \geq m + t \quad \text{for all } t \in \mathbb{N}.$$

But $\text{codepth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) \leq \dim((R)_{\mathfrak{p}}) \leq \infty$, and therefore $Z_m = \emptyset$. □

Recall that the R -module M satisfies Serre's condition (S_k) if for all $\mathfrak{p} \in \text{Spec}(R)$,

$$(*) \quad \text{depth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) \geq \min(\dim(M_{\mathfrak{p}}), k).$$

From now on let m denote the minimal $m \in \mathbb{N}$ with $Z_m = \emptyset$.

3.2. Lemma. *With the assumptions as above put $\overline{R} = R/\text{ann}_R(M)$ and let $k \in \mathbb{N}$. Then the R -module M satisfies (S_k) if and only if for all $0 \leq n < m$,*

$$\text{ht}(\mathfrak{b}_n \overline{R}) > n + k.$$

Proof. Suppose that M satisfies (S_k) , and fix an integer n with $0 \leq n < m$. Let $\mathfrak{p} \in \text{Spec}(R)$ with $\mathfrak{b}_n \subseteq \mathfrak{p}$. Then $\mathfrak{p} \in Z_n$, and therefore

$$\text{codepth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) > n,$$

or equivalently,

$$\dim_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) - \text{depth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) > n.$$

Since M satisfies (S_k) , we obtain that whenever

$$\dim_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) - \text{depth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) \neq 0,$$

then

$$\text{depth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) \geq k.$$

Thus, if $\mathfrak{p} \in Z_n$, then

$$\dim_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) \geq n + k,$$

which implies that $\text{ht}(\mathfrak{b}_n \overline{R}) \geq n + k$.

Conversely, fix an integer k and assume that for all $0 \leq n < m$,

$$\text{ht}(\mathfrak{b}_n \overline{R}) > n + k.$$

Let $\mathfrak{p} \in \text{Spec}(R)$.

If $M_{\mathfrak{p}} = 0$, then $\text{depth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) = \infty$, and condition $(*)$ is satisfied.

Now assume $M_{\mathfrak{p}} \neq 0$. If $M_{\mathfrak{p}}$ is a Cohen-Macaulay R -module, then condition $(*)$ is satisfied. Now assume that

$$\text{codepth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) > 0,$$

and let $n \in \mathbb{N}_0$ with

$$\text{codepth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) = n + 1.$$

Thus $\mathfrak{p} \in Z_n$ and $\mathfrak{b}_n \subseteq \mathfrak{p}$. By assumption,

$$\text{ht}(\mathfrak{b}_n \overline{R}) > n + k \Rightarrow \text{ht}(\mathfrak{b}_n \overline{R}_{\mathfrak{p}}) > n + k \Rightarrow \dim(\overline{R}_{\mathfrak{p}}) > n + k.$$

This implies that

$$\begin{aligned} \text{codepth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) &= n + 1 \\ &= \dim(\overline{R}_{\mathfrak{p}}) - \text{depth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) \\ &\geq n + 1 + k - \text{depth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}), \end{aligned}$$

and therefore

$$\text{depth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) \geq k.$$

Thus $M_{\mathfrak{p}}$ satisfies condition $(*)$, and the R -module M satisfies Serre's condition (S_k) . \square

For all $0 \leq n < m$ consider the closed subset of $\text{Spec}(R)$,

$$Y_{n,k} = \{\mathfrak{q} \in V(\mathfrak{b}_n) \mid \text{ht}(\mathfrak{b}_n \overline{R}_{\mathfrak{q}}) \leq n + k\},$$

and its complement

$$V_{n,k} = \text{Spec}(R) - Y_{n,k}$$

an open subset of $\text{Spec}(R)$. By Lemma 3.2

$$U_{S_k}(M) = \bigcap_{0 \leq n < m} V_{n,k}$$

is an open subset of $\text{Spec}(R)$. We have shown:

3.3. Theorem. *Let M be an R -module as above. If for all $n \in \mathbb{N}_0$ the C_n -locus $U_{C_n}(M)$ is open in $\text{Spec}(R)$, then for all $k \in \mathbb{N}$, the (S_k) -locus*

$$U_{S_k}(M) = \{\mathfrak{p} \in \text{Spec}(R) \mid M_{\mathfrak{p}} \text{ satisfies } (S_k)\}$$

is open in $\text{Spec}(R)$.

In the graded case the theorem states:

3.4. Corollary. *Let $A = \bigoplus_{i \in \mathbb{N}} A_i$ be an excellent graded homogeneous ring and let $M = \bigoplus_{i \in \mathbb{Z}} M_i$ be a finitely generated graded A -module. Then for all $k \in \mathbb{N}$, the set*

$$U_{S_k}^0(M) = \{\mathfrak{p} \in \text{Spec}(A_0) \mid \text{the } (A_0)_{\mathfrak{p}}\text{-module } M_{\mathfrak{p}} \text{ satisfies } (S_k)\}$$

is open in $\text{Spec}(A_0)$.

The proof of the theorem also yields the following corollary:

3.5. Corollary. *Suppose that M and N are R -modules as above. Assume that $\text{ann}_R(M) = \text{ann}_R(N)$ and that for all $n \in \mathbb{N}_0$, the sets $U_{C_n}(M) = U_{C_n}(N)$ are open in $\text{Spec}(R)$. Then for all $k \in \mathbb{N}$,*

$$U_{S_k}(M) = U_{S_k}(N),$$

and the (S_k) -loci are open subsets of $\text{Spec}(R)$.

4. STABILITY ON THE HOMOGENEOUS PARTS

Let $A = \bigoplus_{i \in \mathbb{N}} A_i$ be an excellent graded homogeneous Noetherian ring and let $M = \bigoplus_{i \in \mathbb{Z}} M_i$ be a finitely generated graded A -module. In this section we prove that there is a $k \in \mathbb{N}$, so that for all $n \in \mathbb{N}$ and all $i \geq k$,

$$U_{C_n}^0(M_i) = U_{C_n}^0(M_k) \quad \text{and} \quad U_{S_n}^0(M_i) = U_{S_n}^0(M_k),$$

that is, the codepth and (S_n) -loci of the homogeneous parts of M are eventually stable (considered as an A_0 -module). As before we define for all $t \in \mathbb{Z}$

$$N_t = \bigoplus_{i \geq t} M_i,$$

and observe the following simple facts: Let $k_1 \in \mathbb{N}$ be an integer so that for all $t \geq k_1$, $\text{ann}_{A_0}(M_t) = \text{ann}_{A_0}(M_{k_1})$. Then for all $t \geq k_1$,

$$U_{C_n}^0(N_t) \supseteq U_{C_n}^0(N_{k_1}) \quad \text{and} \quad U_{S_n}^0(N_t) \supseteq U_{S_n}^0(N_{k_1}).$$

Since A_0 is Noetherian, there is an integer $k_2 \in \mathbb{Z}$, so that $k_2 \geq k_1$ and

$$U_{C_n}^0(N_t) = U_{C_n}^0(N_{k_2}) \quad \text{and} \quad U_{S_n}^0(N_t) = U_{S_n}^0(N_{k_2}).$$

We may also assume for large enough k_2 that

$$N_{k_2} = AM_{k_2},$$

which implies that for all $t \geq k_2$,

$$N_t = AM_t.$$

4.1. Lemma. *With the assumptions as above assume additionally that (A_0, \mathfrak{m}_0) is a local ring. Then there is a $k_3 \in \mathbb{Z}$, so that for all $t \geq k_3$,*

$$\text{depth}_{A_0}(M_t) = \text{depth}_{A_0}(M_{k_3}) = \text{depth}_{A_0}(N_{k_3}).$$

Proof. Let k_1 and k_2 be as above and take an integer k with $k > k_2$. Then $\text{codepth}_{A_0}(N_k) = n$ for some $n \in \mathbb{N}$, and therefore

$$\mathfrak{m}_0 \in U_{C_n}^0(N_k) \quad \text{and} \quad \mathfrak{m}_0 \notin U_{C_{n-1}}^0(N_k).$$

Since $k \geq k_2$, we have for all $t \geq k$

$$\text{codepth}_{A_0}(N_k) = n = \text{codepth}_{A_0}(N_t).$$

For all $t \geq k_1$ we also have that $\text{ann}_{A_0}(N_t) = \text{ann}_{A_0}(N_k)$, and therefore for all $t \geq k$,

$$\text{depth}_{A_0}(N_t) = s = \text{depth}_{A_0}(N_k).$$

Let r_1, \dots, r_s be a maximal regular sequence on N_k and put

$$\bar{N}_k = N_k / (r_1, \dots, r_s)N_k \quad \text{with homogeneous parts} \quad \bar{M}_i = M_i / (r_1, \dots, r_s)M_i$$

for $i \geq k$. Note that the torsion submodule $\Gamma_{A_+}(\bar{N}_k)$ is a finitely generated A -submodule of \bar{N}_k . This implies that there is an integer $k_3 \geq k$ so that $\Gamma_{A_+}(\bar{N}_k) \cap N_{k_3} = 0 = \Gamma_{A_+}(\bar{N}_{k_3})$. Thus for k_3 large enough the A -module \bar{N}_{k_3} is A_+ -torsion-free. Since by assumption $\text{depth}_{A_0}(N_k) = s = \text{depth}_{A_0}(N_{k_3})$, there is an integer $i \geq k_3$ and an element $\bar{x} \in \bar{M}_i$ so that $\bar{x} \neq 0$ and $\mathfrak{m}_0 \bar{x} = 0$. Since \bar{N}_{k_3} is A_+ -torsion-free, we obtain

$$(A_+)^l \bar{x} \neq 0 \quad \text{for all} \quad l \in \mathbb{N}.$$

Thus for $k_4 = i > k_3$ we have that $\text{depth}_{A_0}(\overline{M}_{k_4+l}) = 0$ for all $l \in \mathbb{N}_0$, and therefore for all $t \geq k_4$,

$$\text{depth}_{A_0}(M_t) = \text{depth}_{A_0}(M_{k_4}) = s. \quad \square$$

Choose an integer $k_0 \in \mathbb{Z}$ so that the following conditions are satisfied:

- (a) $N_{k_0} = AM_{k_0}$, that is, N_{k_0} is generated in the lowest nonvanishing degree.
- (b) For all $t \geq k_0$, $\text{ann}(M_{k_0}) = \text{ann}(M_t)$.
- (c) For all $n \in \mathbb{N}_0$ and all $t \geq k_0$,

$$U_{C_n}^0(N_t) = U_{C_n}^0(N_{k_0}) \quad \text{and} \quad U_{S_n}^0(N_t) = U_{S_n}^0(N_{k_0}).$$

As before put

$$Z_n = \text{Spec}(A_0) \setminus U_{C_n}^0(N_{k_0}) = V(\mathfrak{b}_n),$$

where $\mathfrak{b}_n \subseteq A_0$ is a reduced ideal. Then $\mathfrak{b}_n \subseteq \mathfrak{b}_{n+1}$, yielding an increasing sequence of ideals

$$\mathfrak{b}_0 \subseteq \mathfrak{b}_1 \subseteq \dots \subseteq \mathfrak{b}_{m-1} \subseteq \dots$$

We have seen before that the sequence stops with some $\mathfrak{b}_m = A_0$, and let m be minimal with this property, that is, let $\mathfrak{b}_m = A_0$ and $\mathfrak{b}_{m-1} \neq A_0$. For all $0 \leq j \leq m - 1$ we consider the set of minimal prime divisors of \mathfrak{b}_j :

$$\text{Min}(A_0/\mathfrak{b}_j) = \{\mathfrak{p}_{j1}, \dots, \mathfrak{p}_{jr_j}\}.$$

By Lemma 4.1, for all $0 \leq j \leq m - 1$ and all $r_j \geq h \geq 1$, there is an integer $k_{jh} \in \mathbb{N}$ with $k_{jh} \geq k_0$, so that for all $i \geq k_{jh}$,

$$\text{depth}_{(A_0)_{\mathfrak{p}_{jh}}}((M_i)_{\mathfrak{p}_{jh}}) = \text{depth}_{(A_0)_{\mathfrak{p}_{jh}}}((M_{k_{jh}})_{\mathfrak{p}_{jh}}) = \text{constant}.$$

Let $k = \max\{k_{jh} \mid 0 \leq j \leq m - 1; 1 \leq h \leq r_j\}$. Then for all $i \geq k$,

$$\text{depth}_{(A_0)_{\mathfrak{p}_{jh}}}((M_i)_{\mathfrak{p}_{jh}}) = \text{depth}_{(A_0)_{\mathfrak{p}_{jh}}}((M_k)_{\mathfrak{p}_{jh}}) = \text{depth}_{(A_0)_{\mathfrak{p}_{jh}}}((N_k)_{\mathfrak{p}_{jh}}).$$

By assumption on the annihilators we also have for all $i \geq k$

$$\dim_{(A_0)_{\mathfrak{p}_{jh}}}((M_i)_{\mathfrak{p}_{jh}}) = \dim_{(A_0)_{\mathfrak{p}_{jh}}}((M_k)_{\mathfrak{p}_{jh}}) = \dim_{(A_0)_{\mathfrak{p}_{jh}}}((N_k)_{\mathfrak{p}_{jh}}),$$

which implies that for all $i \geq k$ and all primes \mathfrak{p}_{jh} ,

$$\text{codepth}_{(A_0)_{\mathfrak{p}_{jh}}}((M_i)_{\mathfrak{p}_{jh}}) = \text{codepth}_{(A_0)_{\mathfrak{p}_{jh}}}((M_k)_{\mathfrak{p}_{jh}}) = \text{codepth}_{(A_0)_{\mathfrak{p}_{jh}}}((N_k)_{\mathfrak{p}_{jh}}).$$

We are now ready to prove:

4.2. Theorem. *Let k be as above. Then for all $i \geq k$ and all $\mathfrak{p} \in \text{Spec}(A_0)$,*

$$\text{codepth}_{(A_0)_{\mathfrak{p}}}((M_i)_{\mathfrak{p}}) = \text{codepth}_{(A_0)_{\mathfrak{p}}}((M_k)_{\mathfrak{p}}).$$

Proof. Let $\mathfrak{p} \in \text{Spec}(A_0)$. If $\mathfrak{b}_0 \not\subseteq \mathfrak{p}$, then $(N_k)_{\mathfrak{p}}$ is a Cohen-Macaulay module over $(A_0)_{\mathfrak{p}}$. It follows that $(M_i)_{\mathfrak{p}}$ is Cohen-Macaulay for all $i \geq k$.

Assume that $\mathfrak{b}_0 \subseteq \mathfrak{p}$ and let g be minimal so that $\mathfrak{b}_g \subseteq \mathfrak{p}$ and $\mathfrak{b}_{g+1} \not\subseteq \mathfrak{p}$. In this case $\text{codepth}_{(A_0)_{\mathfrak{p}}}((N_k)_{\mathfrak{p}}) = g + 1$, and there is an integer $1 \leq j \leq r_j$ so that $\mathfrak{p}_{gj} \subseteq \mathfrak{p}$. By [4, (6.11.5)], the nongraded version of Lemma 2.5, for all $i \geq k$,

$$\text{codepth}_{(A_0)_{\mathfrak{p}}}((M_i)_{\mathfrak{p}}) \geq \text{codepth}_{(A_0)_{\mathfrak{p}_{gj}}}((M_i)_{\mathfrak{p}_{gj}}) = \text{codepth}_{(A_0)_{\mathfrak{p}_{gj}}}((N_k)_{\mathfrak{p}_{gj}}) > g.$$

In order to verify the other inequality consider

$$\text{codepth}_{(A_0)_{\mathfrak{p}}}((N_k)_{\mathfrak{p}}) = g + 1 = \dim((N_k)_{\mathfrak{p}}) - \text{depth}_{(A_0)_{\mathfrak{p}}}((N_k)_{\mathfrak{p}}),$$

and assume that $\text{depth}_{(A_0)_{\mathfrak{p}}}((N_k)_{\mathfrak{p}}) = s$. Let x_1, \dots, x_s be a regular sequence on $(N_k)_{\mathfrak{p}}$. Then x_1, \dots, x_s is a regular sequence on $(M_i)_{\mathfrak{p}}$ for all $i \geq k$. Since N_k and M_i have the same annihilators, we obtain that

$$\text{codepth}_{(A_0)_{\mathfrak{p}}}((N_k)_{\mathfrak{p}}) = g + 1 \geq \text{codepth}_{(A_0)_{\mathfrak{p}}}((M_i)_{\mathfrak{p}})$$

for all $i \geq k$. This shows that for all $i \geq k$,

$$\text{codepth}_{(A_0)_{\mathfrak{p}}}((M_i)_{\mathfrak{p}}) = g + 1.$$

□

4.3. Corollary. *There is an integer $k \in \mathbb{N}$ so that for all $i \geq k$ and all $n \in \mathbb{N}$,*

$$U_{C_n}^0(M_i) = U_{C_n}^0(M_k) = U_{C_n}^0(N_k).$$

□

4.4. Corollary. *There is an integer $k \in \mathbb{N}$ so that for all $i \geq k$ and all $n \in \mathbb{N}$,*

$$U_{S_n}^0(M_i) = U_{S_n}^0(M_k) = U_{S_n}^0(N_k).$$

Proof. The second corollary follows from the first by using Corollary 3.5. □

5. APPLICATIONS

Let A be an excellent ring, let M be a finitely generated A -module, and let $I \subseteq A$ be an ideal of A . By applying the results of the previous section to the Rees algebra/module and to the associated graded ring/module, respectively, we see that there is an integer $k \in \mathbb{N}$, so that for all $i \geq k$ and all $n \in \mathbb{N}$,

$$\begin{aligned} U_{C_n}(I^i M) &= U_{C_n}(I^k M) & \text{and} & & U_{C_n}(I^i M/I^{i+1} M) &= U_{C_n}(I^k M/I^{k+1} M), \\ U_{S_n}(I^i M) &= U_{S_n}(I^k M) & \text{and} & & U_{S_n}(I^i M/I^{i+1} M) &= U_{S_n}(I^k M/I^{k+1} M). \end{aligned}$$

In the following we want to apply these results to the (S_n) - and codepth-loci of the modules $M/I^k M$. We want to show that these loci are again eventually stable, provided that M is a Cohen-Macaulay module over A .

5.1. Lemma. *Let A be any Noetherian ring, $I \subseteq A$ an ideal, and M a finitely generated A -module. Then for all $k \in \mathbb{N}$,*

$$\text{Supp}(M/I^k M) = \text{Supp}(M/IM).$$

Proof. It suffices to show that for all $k \in \mathbb{N}$,

$$\text{Supp}(M/I^k M) = \text{Supp}(M/I^{k+1} M).$$

Since $M/I^k M$ is a homomorphic image of $M/I^{k+1} M$, we have $\text{Supp}(M/I^k M) \subseteq \text{Supp}(M/I^{k+1} M)$. Consider the exact sequence:

$$0 \rightarrow I^k M/I^{k+1} M \rightarrow M/I^{k+1} M \rightarrow M/I^k M \rightarrow 0,$$

and let $\mathfrak{p} \in \text{Spec}(A)$ with $I \subseteq \mathfrak{p}$. The sequence stays exact after localization:

$$0 \rightarrow (I^k M/I^{k+1} M)_{\mathfrak{p}} \rightarrow (M/I^{k+1} M)_{\mathfrak{p}} \rightarrow (M/I^k M)_{\mathfrak{p}} \rightarrow 0.$$

If $(M/I^k M)_{\mathfrak{p}} = 0$ with $(M/I^{k+1} M)_{\mathfrak{p}} \neq 0$, then

$$(I^k M/I^{k+1} M)_{\mathfrak{p}} = (M/I^{k+1} M)_{\mathfrak{p}},$$

which implies by Nakayama that $(M/I^{k+1} M)_{\mathfrak{p}} = 0$, a contradiction. □

A more general version of the next result was proved, using different methods, by Kodiyalam [7, Corollary 9].

5.2. Theorem. *Suppose that (A, \mathfrak{m}) is a local Noetherian ring, let $I \subseteq A$ be an ideal of A , and let M be a finitely generated A -module. Then there is a $k \in \mathbb{N}$, so that for all $i \geq k$,*

$$\text{depth}_A(M/I^i M) = \text{depth}_A(M/I^k M).$$

Proof. Let \widehat{A} be the \mathfrak{m} -adic completion of A . Then for any finitely generated A -module T ,

$$\text{depth}_A(T) = \text{depth}_{\widehat{A}}(T \otimes_A \widehat{A}),$$

and we may replace A by \widehat{A} and M by $M \otimes_A \widehat{A}$, and assume that A is excellent. By Lemma 4.1 there is a $k_1 \in \mathbb{N}$, so that for all $t \geq k_1$,

$$\text{depth}_A(I^t M/I^{t+1} M) = \text{depth}_A(I^{k_1} M/I^{k_1+1} M) = g.$$

For all $t \geq k_1$ consider the exact sequence

$$0 \rightarrow I^t M/I^{t+1} M \rightarrow M/I^{t+1} M \rightarrow M/I^t M \rightarrow 0,$$

which leads to an exact sequence on the cohomology modules:

$$\begin{aligned} \dots \rightarrow H_{\mathfrak{m}}^i(M/I^{t+1} M) &\rightarrow H_{\mathfrak{m}}^i(M/I^t M) \rightarrow 0 \rightarrow \dots \rightarrow 0 \\ \rightarrow \dots \rightarrow H_{\mathfrak{m}}^{g-1}(M/I^{t+1} M) &\rightarrow H_{\mathfrak{m}}^{g-1}(M/I^t M) \rightarrow H_{\mathfrak{m}}^g(I^t M/I^{t+1} M) \\ &\rightarrow H_{\mathfrak{m}}^g(M/I^{t+1} M) \rightarrow H_{\mathfrak{m}}^g(M/I^t M) \rightarrow \dots, \end{aligned}$$

where g is minimal with $H_{\mathfrak{m}}^g(I^t M/I^{t+1} M) \neq 0$.

Case 1: There is an $i \leq g - 1$ and a $t_0 \geq k_1$, so that $H_{\mathfrak{m}}^i(M/I^{t_0} M) \neq 0$. Then for all $t \geq t_0$, $H_{\mathfrak{m}}^i(M/I^t M) \neq 0$. Let $h \leq g - 1$ be the minimal i with this property. Then

$$\text{depth}_A(M/I^t M) = h \quad \text{for all } t \geq t_0.$$

Case 2: For all $i \leq g - 1$ and all $t \geq k_1$,

$$H_{\mathfrak{m}}^i(M/I^t M) = 0.$$

This implies that $\text{depth}_A(M/I^t M) \geq g - 1$ for all $t \geq k_1$.

Case 2.1: There are infinitely many $t \geq k_1$, so that

$$H_{\mathfrak{m}}^{g-1}(M/I^t M) \neq 0.$$

From the long exact sequence we observe that $H_{\mathfrak{m}}^{g-1}(M/I^t M) \neq 0$ implies that $H_{\mathfrak{m}}^{g-1}(M/I^{t-1} M) \neq 0$ whenever $t - 1 \geq k_1$. Thus in this case there is a $t_1 \geq k_1$, so that for all $t \geq t_1$,

$$H_{\mathfrak{m}}^{g-1}(M/I^t M) \neq 0,$$

and therefore for all $t \geq t_1$, $\text{depth}_A(M/I^t M) = g - 1$.

Case 2.2: There is a $t_2 \geq k_1$, so that for all $t \geq t_2$, $H_{\mathfrak{m}}^{g-1}(M/I^t M) = 0$. Then for all $t \geq t_2$,

$$\text{depth}_A(M/I^t M) = g. \quad \square$$

5.3. Theorem. *Let A be an excellent ring and M a finitely generated Cohen-Macaulay A -module. Let $I \subseteq A$ be an ideal of A which is not contained in any minimal prime ideal of M . Then there is an integer $k \in \mathbb{N}$, so that for all $t \geq k$ and all $n \in \mathbb{N}_0$:*

- (1) $U_{C_n}(M/I^t M) = U_{C_n}(M/I^{k_0} M)$.
- (2) $U_{S_n}(M/I^t M) = U_{S_n}(M/I^{k_0} M)$.

Proof. (1) Fix $n \in \mathbb{N}$ and let $k \in \mathbb{N}$, so that for all $t \geq k$,

$$U_{C_n}(I^t M) = U_{C_n}(I^k M).$$

We claim that for all $i \geq k$ and all $\mathfrak{p} \in V(I)$,

$$\text{depth}_{A_{\mathfrak{p}}}(M/I^i M)_{\mathfrak{p}} = \text{depth}_{A_{\mathfrak{p}}}(M/I^k M)_{\mathfrak{p}}.$$

Obviously, for all $i \geq k$, $\dim((I^i M)_{\mathfrak{p}}) = \dim((I^k M)_{\mathfrak{p}})$, and thus because of the stability of the codepth-loci, we have for all $\mathfrak{p} \in V(I)$ and all $i \geq k$ that

$$\text{depth}_{A_{\mathfrak{p}}}((I^i M)_{\mathfrak{p}}) = \text{depth}_{A_{\mathfrak{p}}}((I^k M)_{\mathfrak{p}}).$$

Fix an integer $i \geq k$ and a prime ideal $\mathfrak{p} \in V(I)$, and consider the exact sequence

$$0 \rightarrow (I^i M)_{\mathfrak{p}} \rightarrow M_{\mathfrak{p}} \rightarrow (M/I^i M)_{\mathfrak{p}} \rightarrow 0.$$

With $d = \dim_{A_{\mathfrak{p}}}(M_{\mathfrak{p}}) = \text{depth}_{A_{\mathfrak{p}}}(M_{\mathfrak{p}})$ we obtain a long exact sequence of the local cohomology modules

$$\begin{aligned} \cdots \rightarrow 0 \rightarrow H_{\mathfrak{p}}^{i-1}((M/I^i M)_{\mathfrak{p}}) \rightarrow H_{\mathfrak{p}}^i((I^i M)_{\mathfrak{p}}) \rightarrow 0 \rightarrow \cdots \rightarrow 0 \\ \rightarrow H_{\mathfrak{p}}^{d-1}((M/I^i M)_{\mathfrak{p}}) \rightarrow H_{\mathfrak{p}}^d((I^i M)_{\mathfrak{p}}) \rightarrow H_{\mathfrak{p}}^d(M_{\mathfrak{p}}) \rightarrow 0 = H_{\mathfrak{p}}^d((M/I^i M)_{\mathfrak{p}}), \end{aligned}$$

where $H_{\mathfrak{p}}^d((M/I^i M)_{\mathfrak{p}}) = 0$, since $\dim_{A_{\mathfrak{p}}}((M/I^i M)_{\mathfrak{p}}) \leq d - 1$. This shows that

$$\text{depth}_{A_{\mathfrak{p}}}(M/I^i M)_{\mathfrak{p}} = \text{depth}_{A_{\mathfrak{p}}}((I^i M)_{\mathfrak{p}}) - 1 = \text{depth}_{A_{\mathfrak{p}}}((I^k M)_{\mathfrak{p}}) - 1,$$

and the claim is proven. For all $i \geq k$ and all $\mathfrak{p} \in V(I)$ we have

$$\begin{aligned} \text{depth}_{A_{\mathfrak{p}}}(M/I^i M)_{\mathfrak{p}} &= \text{depth}_{A_{\mathfrak{p}}}(M/I^k M)_{\mathfrak{p}}, \\ \dim((M/I^i M)_{\mathfrak{p}}) &= \dim((M/I^k M)_{\mathfrak{p}}). \end{aligned}$$

The last equation is obtained from Lemma 5.1. This yields that for all $n \in \mathbb{N}$ and for all $i \geq k$,

$$U_{C_n}(M/I^i M) = U_{C_n}(M/I^k M).$$

The second assumption follows with Corollary 3.5. □

5.4. Corollary. *Let A , M , and I be as in the theorem, and assume that $IM \neq M$. Then there is an element $a \in A$, so that for all $k \in \mathbb{N}$,*

- (1) $(M/I^k M)_a \neq 0$.
- (2) $(M/I^k M)_a$ is a Cohen-Macaulay module. □

5.5. Corollary. *Let A be an excellent ring and M a finitely generated A -module. Suppose that the ideal $I \subseteq A$ satisfies the following conditions:*

- (i) I is not contained in a minimal prime of M .
- (ii) If $\mathfrak{a} \subseteq A$ is the defining ideal of the non-Cohen-Macaulay locus of M , then $\mathfrak{a} \not\subseteq \sqrt{(IM : M)}$.

Then there is an element $a \in A$, so that for all $k \in \mathbb{N}$,

- (1) $(M/I^k M)_a \neq 0$.
- (2) $(M/I^k M)_a$ is a Cohen-Macaulay module.

Proof. Choose an element $b \in \mathfrak{a} \setminus \sqrt{(IM : M)}$. In order to prove the assertion apply the previous corollary to the Cohen-Macaulay A_b -module M_b . □

REFERENCES

- [1] N. Bourbaki, *Commutative algebra, chapters 1-7*, Springer Verlag, New York, 1989 MR0979760 (90a:13001)
- [2] W. Bruns, J. Herzog, *Cohen-Macaulay rings*, Cambridge Studies in Advanced Mathematics, Vol. 39, revised edition, Cambridge, 1998 MR1251956 (95h:13020)
- [3] H. B. Foxby, *Hyperhomological algebra and commutative rings*, in preparation
- [4] A. Grothendieck, *Éléments de géométrie algébrique IV*, Inst. Hautes Études Sci. Publ. Math **24** (1965) MR0199181 (33:7330)
- [5] M. Hochster, J. L. Roberts, *Rings of invariants of reductive groups acting on regular rings are Cohen-Macaulay*, Adv. Math. **13** (1974), 115–175 MR0347810 (50:311)
- [6] S. Iyengar, *Depth for complexes, and intersection theorems*, Math. Z. **230** (1999), 545–569 MR1680036 (2000a:13027)
- [7] V. Kodiyalam, *Homological invariants of powers of an ideal*, Proc. Amer. Math. Soc. **118** (1993), 757–764 MR1156471 (93i:13022)
- [8] H. Matsumura, *Commutative ring theory*, Cambridge Studies in Advanced Mathematics, Vol. 8, Cambridge, 1986 MR0879273 (88h:13001)

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MICHIGAN 48824

E-mail address: `rotthaus@math.msu.edu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MISSOURI, KANSAS CITY, MISSOURI 64110-2499

E-mail address: `segal@umkc.edu`