

A CLASS OF GARSIDE GROUPOID STRUCTURES ON THE PURE BRAID GROUP

DAAN KRAMMER

ABSTRACT. We construct a class of Garside groupoid structures on the pure braid groups, one for each function (called labelling) from the punctures to the integers greater than 1. The object set of the groupoid is the set of ball decompositions of the punctured disk; the labels are the perimeters of the regions. Our construction generalises Garside's original Garside structure, but not the one by Birman–Ko–Lee. As a consequence, we generalise the Tamari lattice ordering on the set of vertices of the associahedron.

1. INTRODUCTION

In [Gar69] Garside solved the word and conjugacy problems in the braid groups. Birman, Ko and Lee [BKL98] gave another solution, but it had something in common with Garside's approach. Indeed, both approaches were examples of what are now called *Garside structures* on the braid group. The general theory of Garside groups was developed mainly by Dehornoy; we mention three good overviews [DehPar99], [Deh00], [Deh02].

We propose a generalisation of the concept *Garside group* to *Garside groupoid*. It is not surprising, and the required proofs can easily be adapted from those in the literature covering the group case.

A short definition of Garside groupoids is as follows. A Garside groupoid is a group G acting freely on the left on a lattice L with the following properties.

- The orbit set $G \setminus L$ is finite.
- There exists an automorphism Δ (written on the right) of the lattice L commuting with the G -action.
- For any $x \in L$ the interval $[x, x\Delta] := \{y \in L : x \leq y \leq x\Delta\}$ is finite.
- The ordering on L is generated by $x \leq y$ whenever $y \in [x, x\Delta]$.

But this definition ought to be a theorem, and we shall use a different definition in the paper.

If the G -action on L is transitive it is a Garside group.

The main result of the paper is a class of new Garside (groupoid) structures on the pure braid group, one for each function, called labelling, from the set of punctures to $\mathbb{Z}_{\geq 2}$. The objects for the involved groupoid are the ball decompositions of a punctured disk; the labels are the perimeters of the regions.

Garside's original Garside structure on the braid group [Gar69] is a special case of our construction, but Birman–Ko–Lee's structure [BKL98] is not.

Received by the editors September 28, 2005 and, in revised form, March 27, 2006.

2000 *Mathematics Subject Classification*. Primary 20F36; Secondary 20F05, 20F60, 57M07.

©2008 American Mathematical Society
Reverts to public domain 28 years from publication

The set $[x, x\Delta]$ seems to be the vertex set of a polytope in a natural way, but we won't go into this. Among the polytopes obtained this way are the permutohedron (if all labels are 2) and the associahedron [Sta63], [Lee89] (if all labels are 3).

As a consequence of our main result we get many finite lattices $[x, x\Delta]$. They seem new even in the particular case where all labels are 3. In the more particular case where all triangles have a vertex in common, this was previously known by the work of Tamari [Tam51], [FriTam67], [Grä78, page 18]. See subsection 5.2 for more details.

The paper is written in the language of Garside groups, not least because it allows us to use the results from [Deh00], [Deh02], thus streamlining our proofs.

After the Garside structures by Garside and Birman–Ko–Lee there doesn't seem to be a need for any more of them. Our results are the byproducts of an investigation into surface mapping class groups. It would be interesting to know if mapping class groups are Garside; see [Par05].

In section 2 we construct a braid-like groupoid and a presentation for it. In section 3 we overview Garside groupoids in general. In section 4 we prove that the axioms for Garside groupoids are fulfilled by our braid groupoid. A short last section gives some more information about the finite lattices $[x, x\Delta]$ (there written $\Omega(xT)$) and especially the case where all labels are 3.

I would like to thank an anonymous referee who provided many suggestions for improvement.

Remark 1.1. Depending on the labels, some of our proofs can be shortened; that is, some case-by-case proofs can deal with fewer cases. This will be immediately clear from the text. If all labels are ≥ 3 , then:

- The elementary relations (ER2) and (ER4) don't exist (Figure 3).
- Cases 1 and 2 don't exist (Figures 7 and 8).
- Cases (f), (g), (h) in the proof of Lemma 4.4 (the cube condition) can be ignored.

If all labels are 3, then:

- The elementary relations (ER2) and (ER4) don't exist (Figure 3).
- Subsections 2.9 and 2.10 can be ignored (subsection 2.8 replaces them).
- Cases (a), (f), (g), (h) in the proof of Lemma 4.4 (the cube condition) can be ignored.

2. A BRAID-LIKE GROUPOID

2.1. Introduction. In this section we define a groupoid such that the automorphism groups of objects are finite index subgroups of the braid group. We give a presentation for it (generators and relations). It is a *complemented presentation* as required for Garside groupoids. The bulk of the section is there to prove that the presentation is correct.

2.2. A braid groupoid. Let D be a disk, that is, a topological space homeomorphic to $\{z \in \mathbb{C} : |z| \leq 1\}$. Let P be a finite set of n interior points of D called *punctures*. Let Q be a nonempty finite set of boundary points of D . Let ℓ be a map called *labelling* $\ell: P \rightarrow \mathbb{Z}_{\geq 2}$. For a reason which will become clear later we assume

$$(2.1) \quad |Q| - 2 = \sum_{p \in P} (\ell(p) - 2).$$




FIGURE 1. The punctured disk and an admissible decomposition.

Let M be the *mapping class group* of

$$(D, \text{orientation}, P, Q, \ell).$$

In other words, $M = H/H_0$; here H is the group of orientation preserving self-homeomorphisms g of D fixing P setwise and the boundary ∂D pointwise such that $\ell = \ell \circ g$, and H_0 is the component of the identity in H .

Our definition of the *braid group* B_n is precisely M provided all labels are equal. In general, M is isomorphic to the group of label preserving braids in B_n ; it is a subgroup of B_n of finite index

$$n! \left(\prod_{k \geq 2} n_k! \right)^{-1}$$

where n_k is the number of punctures of label k .

See Figure 1(a) for a picture of (D, P, Q, ℓ) ; each puncture p is labelled $\ell(p)$.

By a *ball complex* K we mean a CW complex whose cell attaching maps are injective. The cells of dimensions (respectively) 0,1,2 are called (respectively) *vertices*, *edges*, *regions*. By a *ball decomposition* of a topological space X we mean a ball complex whose underlying topological space is X , and whose cells are subsets of X , and whose cell attaching maps are the identity.

Definition 2.1. An *admissible decomposition* is a ball decomposition of D such that the following hold:

- The vertex set is Q .
- The 1-skeleton doesn't contain any punctures.
- Each region (= 2-dimensional cell) contains precisely one puncture p , and moreover, the label $\ell(p)$ equals the number of edges of that region.

See Figure 1(b) for a picture of an admissible decomposition. For example, the puncture in the middle of the region with 4 edges is labelled 4.

Remarks 2.2. (a). The boundary of D is in the 1-skeleton of any admissible decomposition and consists of $|Q|$ edges, which we shall call *boundary edges*. All other edges will be called *arcs*.

Every element of Q is a vertex, even if there isn't any arc emanating from it.

(b). Recall our assumption (2.1):

$$(2.2) \quad |Q| - 2 = \sum_{p \in P} (\ell(p) - 2).$$

Equation (2.2) is equivalent to saying that admissible decompositions exist. This follows because the number of even-dimensional cells minus the number of odd-dimensional cells is 1 (the Euler characteristic of the disk).

Definition 2.3. By L we will denote the set of isotopy classes relative to P (that is, H_0 -orbits) of admissible decompositions.

As is usual we shall abusively confuse admissible decompositions (or arcs) with their isotopy classes. Most of the time we are dealing with isotopy classes without saying we are.

The braid-like group M acts on L , by convention on the left. It is clear that the action is *free* ($gx = x \Rightarrow g = 1$ for all $g \in M, x \in L$) with a finite number of orbits.

Recall that a groupoid is a category all of whose morphisms are invertible. A group is then a groupoid with just one object. By $\text{Hom}_G(A, B)$ or $G(A, B)$ we shall denote the set of morphisms in a category G from an object A to an object B . Its elements are said to have *source* object A and *target* object B and are said to go from A to B . The order of multiplication will be

$$\begin{aligned} \text{Hom}_G(A, B) \times \text{Hom}_G(B, C) &\longrightarrow \text{Hom}_G(A, C), \\ (f, g) &\longmapsto fg. \end{aligned}$$

Definition 2.4. We define a groupoid G as follows. The object set is $G_0 = M \setminus L$.

We let M act on the Cartesian powers L^n diagonally:

$$g(x_1, \dots, x_n) = (gx_1, \dots, gx_n).$$

An M -orbit in L^n will be written $M(x_1, \dots, x_n)$.

The hom-set $\text{Hom}_G(A, B)$ is defined to be the set of M -orbits in $L \times L$ meeting (hence contained in) $A \times B$. Equivalently, $\text{Hom}_G(Mx, My) := \{M(x, gy) \mid g \in M\}$.

The multiplication in G is defined by $M(x, y) \circ M(y, z) := M(x, z)$. One readily checks that this is well defined. Note the order of multiplication.

Note that this construction of G could be applied whenever a group (in our case M) acts on a set (in our case L).

Remarks 2.5. (a). It is easier to think of the group M acting on the set L rather than the groupoids constructed in Definition 2.4. Groupoids are just a language and don't add anything to our reasoning. It is a good idea not to lose sight of the group M acting on the set L .

(b). Clearly, for $x, y \in L$, one has $Mx = My \iff x$ and y are isotopic *not* relative to P . In other words, the object set of G can be identified with the set of isotopy classes, *not* relative to P , of ball decompositions of D as follows.

- The vertex set is Q .
- For any $k \geq 2$, the number of k -gons equals the number of punctures with label k .

2.3. Elementary morphisms in G .

Definition 2.6 (Elementary morphisms). (See Figure 2 and ignore the w 's for the moment.) Let A be an arc of an admissible decomposition x . There is a




FIGURE 2. An elementary pair (x, y) . See Definition 2.7.

unique ball decomposition t of D whose edges are the edges of x except A . Let R denote the region of t containing A . Let B be the arc obtained from A by rotating both endpoints of A in the positive direction over one edge along the boundary of R , moving the rest of A along in a continuous way such that it never meets any punctures. There is a unique admissible decomposition y of D whose edges are the edges of t and B . We call (x, y) an *elementary pair* and $M(x, y) \in \text{Hom}_G(Mx, My)$ an *elementary morphism*. In diagrams we draw elementary morphisms by solid arrows.

For future reference we next define the diagonal morphisms, which generalise the elementary morphisms.


Definition 2.7 (Diagonal morphisms). Let x be an admissible decomposition. Let A_1, \dots, A_k be k distinct arcs of x . There is a unique ball decomposition t of D whose edges are the edges of x except A_1, \dots, A_k . For $1 \leq i \leq k$ let R_i denote the region of t containing A_i . Let B_i be the arc obtained from A_i by rotating both endpoints of A_i in the positive direction over one edge along the boundary of R_i , moving the rest of A_i along in a continuous way such that it never meets any punctures. There is a unique admissible decomposition y of D whose edges are the edges of t and B_1, \dots, B_k . The morphism $M(x, y) \in \text{Hom}_G(Mx, My)$ is called a *diagonal morphism*. If k is the maximal value $n - 1$, then we also call it a *full diagonal morphism*. We recover the elementary morphisms by taking $k = 1$.

For every $x \in L$ there are exactly $n - 1 = |P| - 1$ elementary pairs (x, y) (respectively, (y, x)), corresponding to the $n - 1$ arcs of x which are rotated in a positive (respectively, negative) direction to obtain y .

In Remark 2.5(b) we saw that for the mere purpose of drawing an object of G , one can omit the punctures. What about morphisms?

It often happens that a hom-set $\text{Hom}_G(Mx, My)$ contains at most one elementary morphism. This is at least the case if $Mx \neq My$. Moreover, if $Mx = My$, then the elementary morphisms from Mx to My are in bijection with those arcs of x which lie on two bigons in x . Rotating such an arc gives rise to the corresponding elementary morphism; see (2.5). Thanks to this observation we shall often simplify our diagrams in G by omitting the punctures, so that the arcs won't look like spaghetti.

Example 2.8. Consider the case where the labels are $(3, 3)$. Then there is a bijection $f: \mathbb{Z} \rightarrow L$. Here are some values of f .



We have $M \cong \mathbb{Z}$. A generator of M takes $f(n)$ to $f(n+2)$. So $M \setminus L$, which is also G_0 , the set of objects of G , has precisely two elements x_0, x_1 where $x_k = \{f(2n+k) \mid n \in \mathbb{Z}\}$.

In Remark 2.5(b) we saw that elements of G_0 are given by pictures without punctures. Such pictures of x_k are as follows.

$$x_0 = \text{Diagram with a horizontal line through the center} \quad x_1 = \text{Diagram with a vertical line through the center}$$

The elementary pairs are the pairs $(f(n), f(n+1))$. There are two elementary morphisms $g: x_0 \rightarrow x_1$ and $h: x_1 \rightarrow x_0$. Indeed g is represented by the elementary pairs $(f(2n), f(2n+1))$, and h by $(f(2n-1), f(2n))$. Note that g and h are not each other's inverses because $f(n) \neq f(n+2)$.

Consider $G(x_0, x_0)$, the set of morphisms in G from x_0 to x_0 . There is a bijection $r: \mathbb{Z} \rightarrow G(x_0, x_0)$ given by $r(n) = M(f(0), f(2n))$; that is, $r(n)$ is represented by the elementary pair $(f(0), f(2n))$. It restricts to a bijection $r: \mathbb{Z}_{\geq 0} \rightarrow G^+(x_0, x_0)$.

A flash forward. We define a relation \leq on L by putting $x \leq y$ if and only if there is a sequence $x = x_0, x_1, \dots, x_n = y$ such that (x_i, x_{i+1}) is an elementary pair for all i . We shall observe that \leq is an ordering on L . In Example 2.8, the ordering on L is given by $f(k) \leq f(\ell) \Leftrightarrow k \leq \ell$. One of our results, Corollary 4.24, states that (L, \leq) is a lattice.

2.4. Elementary relations in G .

Definition 2.9. An *elementary relation* is a relation in G as displayed in Figure 3.

Explanation. (a). A fat line in an object stands for any nonnegative number of edges. If the number of edges is zero, then the endpoints of the fat line are identical.

(b). In relations (ER2–4) any two objects of the relation differ only in a disk (a union of three regions), which is the only part depicted. In relation (ER1) two disks are involved, which may be disjoint or have a vertex or edge in common.

(c). All morphisms in elementary relations are elementary morphisms and are therefore depicted by solid arrows.

With the help of the punctures displayed in Figure 3, one observes that elementary relations really are relations in G ; that is, they are commuting diagrams.

Example 2.10. (See Figure 4.) If all labels are 3, then the elementary relation (ER3) simplifies to part (a) of Figure 4. On removing the punctures we obtain part (b) of the figure. Diagram (c) is obtained from diagram (b) by reversing the rightmost arrow and is therefore *not* a relation in G , certainly not an elementary relation. One toggles between right and wrong by either replacing each little 5-gon by its mirror image or by reversing all arrows.




FIGURE 3. Elementary relations. See Definition 2.9.




FIGURE 4. See Example 2.10.

2.5. Presentations of groupoids. There is a straightforward generalisation of the concept of group presentation (generators and relations) to presentations of categories and groupoids. We mention only that the object set is given in advance and is not affected by the presentation.

Recall that the braid group B_n is the mapping class group of (D, P, Q) . It can also be given by the *Artin presentation* with generators σ_i ($1 \leq i < n$) and relations

$$(2.3) \quad 4\text{-gon:} \quad \sigma_i \sigma_j = \sigma_j \sigma_i \quad \text{whenever } |i - j| > 1,$$

$$(2.4) \quad 6\text{-gon:} \quad \sigma_{i+1} \sigma_i \sigma_{i+1} = \sigma_i \sigma_{i+1} \sigma_i$$

for all appropriate indices.

Our aim is to prove that G is presented by elementary morphisms and elementary relations. Rather than building things from scratch we use the Artin presentation of the braid group as our starting point.

Let G^* be the groupoid presented by all elementary morphisms and elementary relations. So we want to prove $G^* = G$.


2.6. The case of only bigons. We now consider the special case where all labels are 2, so that every region of an admissible decomposition is a bigon. This makes a nice example and will also be used later on.

Lemma 2.11. *If all labels are 2, then $G^* = G$.*

Proof. Since all labels are equal, the automorphism group of any object of G is the braid group B_n .

By our assumption, all labels are 2, so G has only one object and is equal to the braid group.

An elementary morphism looks like this:



This is the pictorial interpretation of an Artin generator σ_i . Since B_n is generated by the Artin generators, we conclude that G is generated by the elementary morphisms.

One easily checks that the Artin 4-gons (2.3) are the elementary relations of the form (ER1) (Figure 3) and the Artin 6-gons (2.4) are the instances of (ER4). Relations (ER2) and (ER3) don't occur. The result follows. \square

2.7. Fans. From now on, and throughout the section, we fix a base vertex $q_0 \in Q$.

Definition 2.12. An admissible decomposition $x \in L$ is a *fan* if each arc of x contains q_0 . Its M -orbit $Mx \in G_0$ is also called a fan (confusion need not arise). Let F denote the full subgroupoid of G with the fans for objects (*full* means that every morphism of G between fans is a morphism of F).

Definition 2.13. By an *elementary fan morphism* we mean a morphism $M(x, y) \in \text{Hom}_F(Mx, My)$ between two fans Mx, My where an arc in x is rotated in the positive direction, not by one “click” as in elementary morphisms, but by the least positive number of clicks such that the result is again a fan (written y). In diagrams we denote elementary fan morphisms by dashed arrows.

Convention 2.14. From now on, with the exception of (2.6), we will follow the following convention for depicting vertices in objects in diagrams.

Of course: in a diagram, arrows go from objects to objects; in each object, there are vertices and edges.

In pictures of objects of G (rather than the objects themselves) we don't speak of edges but rather *apparent edges*, which stand for a (usually unspecified) nonnegative number of edges. In other words, not all vertices may be depicted. If an apparent edge stands for zero edges, then its endpoints coincide. Apparent arcs always stand for a single arc.

A black dot \bullet always stands for the base vertex q_0 , but we allow the possibility that q_0 is depicted by a different symbol or not at all.

Note also that this convention will become more restricted from section 4.3 on.




FIGURE 5. The 6-gon fan relation.

Let D' denote the quotient of D by contracting the union of all boundary edges not containing q_0 , to a point. Define P' , Q' , q'_0 in the obvious way, and let $\ell': P' \rightarrow \mathbb{Z}_{\geq 2}$ be the constant 2. Then (D', P', Q', ℓ') is again a collection of data as we started with, and indeed of the only-bigons form which we studied in subsection 2.6.

Define G' accordingly. The edge contraction defines an isomorphism $F \rightarrow G'$. But in Lemma 2.11 we have seen a presentation for G' ; on passing it through the isomorphism $F \cong G'$ we obtain a presentation for F . The generators in this presentation are precisely the elementary fan morphisms, and we shall call its relations the *fan relations*; they are just the Artin relations in disguise.

The 6-gon fan relation is depicted in Figure 5 using convention 2.14. The labels σ_i remind us that it is essentially an Artin relation. The perimeters of the regions are k, l, m .

Summarising:

Lemma 2.15. *The groupoid F is presented by the elementary fan morphisms and the fan relations.* \square

Lemma 2.16. *The groupoid G is generated by the elementary morphisms.*

Proof. It is easy and left to the reader to show that one can walk from any admissible decomposition to a fan using elementary morphisms. In other words, every object of G is isomorphic in G^* to a fan. It remains to show that every morphism in F is a product of elementary morphisms and their inverses (we say: it can be expressed in terms of elementary morphisms). Well, a morphism in F can be expressed in terms of elementary fan morphisms (by Lemma 2.15), which in turn can be expressed in terms of elementary morphisms. \square

We now prepare for the proof that $G = G^*$. There is a natural surjective functor $G^* \rightarrow G$. We don't yet know that it is injective, but we have at least a natural lifting of the elementary fan morphisms as follows. Every elementary fan morphism u can be uniquely written as a product $u_1 \cdots u_k$ of elementary morphisms where all arcs but one stay fixed. The expression $u_1 \cdots u_k$ defines a morphism in G^* which we write $a(u)$. The following is now clear.

Lemma 2.17. *The statement $G = G^*$ is equivalent to saying that for every fan relation the following holds. On replacing each arrow u in the fan relation by $a(u)$, one gets a relation in G^* (that is, a consequence of the elementary relations).* \square


2.8. The case of only triangles. Before finishing the proof that $G = G^*$ in the general case we do the case where all labels are 3, which is easier yet shows all steps.

Proposition 2.18. *If all labels are 3, then $G^* = G$.*

Proof. We prove the necessary and sufficient condition of Lemma 2.17.

A 4-gon fan relation can be ‘tessellated’ by four squares of the form (ER1); the details are easy and left to the reader.

Consider a 6-gon fan relation (see Figure 5). It necessarily looks like the dashed arrows in the following diagram.



The triangles in this diagram show how to replace each elementary fan morphism u of the fan relation by $a(u)$ consisting of two solid arrows. It is therefore our aim to prove that the twelve outer solid arrows form a relation in G^* . Well they do, because the six 5-gons are elementary relations as one can check. In particular there are no wrong 5-gons as in Figure 4(c). This settles the case of 6-gon fan relations. \square

2.9. Complementary relations.

Convention 2.19. In addition to convention 2.14 the present subsection 2.9 uses the following notation. If a vertex in an object is depicted by any symbol other than a white dot (namely, $\triangle, \bullet, 1, 2, 3, \dots$), this indicates that the vertex is being repeated in nearby objects (same place, same symbol). Knowing this helps understand the diagrams. White dots \circ denote other vertices, which we decide to depict.

Definition 2.20. Assume $n = 3$, so that every admissible decomposition has $n - 1 = 2$ arcs. We define two more special sorts of morphisms in G .

(a). Suppose the two arcs of $x \in L$ have a common vertex. Let $y \in L$ be the result of rotating both arcs of x in D over the same number ($\in \mathbb{Z}$) of clicks. The morphism $M(x, y) \in \text{Hom}_G(Mx, My)$ is called a *central morphism* and denoted by a solid arrow with a twiddle $\tilde{\rightarrow}$. Equivalently, a central morphism is any product of full diagonal morphisms, which were defined in Definition 2.6, and their inverses. See Figure 6 for pictures of central morphisms.

(b). Let $a, b, c \in Q$ be three distinct vertices in this cyclic order counterclockwise. Let $x \in L$ be such that one of its arcs A connects a, b and the other arc B connects b, c . Let $y \in L$ be obtained from x by rotating B counterclockwise by the least positive number of clicks such that it contains a . The morphism $M(x, y) \in \text{Hom}_G(Mx, My)$ is called a *parabolic morphism*. Like elementary morphisms, they will be denoted by solid arrows; some parabolic morphisms are indeed




FIGURE 6. Complementary relations. See Definition 2.21.

elementary morphisms. The two nontwiddled arrows in (CR2) in Figure 6 are examples of parabolic morphisms.

(c). The collective name for central and parabolic morphisms is *complementary morphisms*.

Definition 2.21. Assume $n = 3$. The *complementary relation* (CR1) reads $uv = w$ for any central morphisms u, v, w (provided it is meaningful and true in G).

The complementary relations (CR2–4) are defined in Figure 6. The fat apparent edge in (CR2) stands for $k \geq 1$ edges.

Finally, (CR5) are the relations $u = u_1 \cdots u_k$ where u is parabolic and u_i is elementary, provided the relation is meaningful and true in G and one arc is fixed along the path $u_1 \cdots u_k$ (recall that there are only two arcs).

Example 2.22. Here are some examples of reading diagrams by the rules we have introduced.

- (a) The nontwiddled arrows in (CR2) are not necessarily elementary.
- (b) The nontwiddled arrows in (CR3) are elementary. Three of the apparent boundary edges stand for single edges.
- (c) Consider the parabolic arrow $x \rightarrow w$ in (CR4). It is possible that the vertices are distinct and ordered (unexpectedly) $(1, 3, 2, 4)$ in this cyclic order. The vertex 2 is not necessarily repeated in x nor 3 in w .
- (d) The source and target of the twiddled arrow $z \rightarrow y$ in (CR4) are not necessarily equal. They may involve different sets of depicted vertices, and the pictures of the disk may differ by a rotation.

Lemma 2.23. Assume $n = 3$. The groupoid G^* is presented by elementary and complementary morphisms and elementary and complementary relations.

Proof. By definition, G^* is presented by elementary morphisms and relations. For each complementary morphism u we shall define a morphism $b(u)$ of G^* (with the same source and target objects as u). The result will then be proved by showing that replacing each complementary arrow u by $b(u)$ in the complementary relations yields relations in G^* (that is, consequences of the elementary relations).

The twiddled arrow in (CR2) with $k = 1$ is a central morphism u_0 , which for the moment we call a simple morphism; we define $b(u_0)$ and $b(u_0^{-1})$ by that diagram. Any other central morphism u is a product $u_1 \cdots u_\ell$ of simple morphisms and their inverses and we define $b(u) := b(u_1) \cdots b(u_\ell)$.

A parabolic morphism u can uniquely be written as a product $u_1 \cdots u_k$ of elementary morphisms where all arcs but one stay fixed, as in (CR5); we define $b(u) := u_1 \cdots u_k$.

Finally we prove that replacing $u \mapsto b(u)$ in complementary relations yields relations in G^* . For (CR1) and (CR5) this is trivial. For (CR2) this is an easy induction on k and left to the reader. For (CR3) this follows from the 5-gon (ER3) and (CR2) with $k = 1$. In order to prove (CR4), cut it into three pieces along the diagonals xy and xz and apply (CR2), (CR1) and (CR3) to the pieces (respectively, from top to bottom). \square

Remark 2.24. Note that (CR4) generalises (ER3). In order for (CR4) to be true we had to insert a central morphism into a 5-gon making it into a 6-gon. Inserting the central morphism can be done anywhere in the 5-gon (not necessarily the spot where (CR4) does) and we shall tacitly do so when necessary.

2.10. Presentation of G . We will use a variation of one of our lemmas as follows. Assume $n = 3$. Let u be an elementary fan morphism. We define a morphism $c(u)$ in G^* as follows.

- If u is an elementary morphism we put $c(u) = u$.
- Otherwise, in G^* one can write u uniquely as $u_1 u_2$ (with one arc fixed throughout) where u_1 or u_2 is an elementary morphism and the other is a parabolic morphism. We define $c(u) = u_1 u_2$.

The following mild variation of Lemma 2.17 is now clear.

Lemma 2.25. *Assume $n = 3$. The statement $G = G^*$ is equivalent to the following.*

- *For every 4-gon fan relation, the condition of Lemma 2.17 holds.*
- *For every 6-gon fan relation, the following holds. On replacing each arrow u in the fan relation by $c(u)$, one gets a relation in G^* (that is, a consequence of the elementary and complementary relations).* \square


Proposition 2.26. $G = G^*$.

Proof. As in the proof of Proposition 2.18, we verify the necessary and sufficient condition of Lemma 2.25.

First consider a 4-gon fan relation $uv = wx$ where u, v, w, x are elementary fan morphisms. Then there are $k, \ell \geq 1$ such that both $a(u), a(x)$ are a product of k elementary morphism while $a(v), a(w)$ consist of ℓ of them. Then the fan relation $uv = wx$ can be tessellated by $k\ell$ 2-cells of the form (ER1).

We turn to the 6-gon fan relation (see Figure 5). It entirely takes place in a subdisk of 3 regions, so without loss of generality we may assume $n = 3$. Then the definition of complementary morphisms and relations applies. Let $k, \ell, m \geq 2$ be the labels in the order defined by Figure 5. We consider three cases:

- Case 1: $\ell = 2$,
- Case 2: $\ell > 2, m = 2$,
- Case 3: $\ell > 2, m > 2$.

FIGURE 7. Case 1: $\ell = 2$.

The proofs of cases 1, 2, 3 can be found in (respectively) Figures 7, 8, 8. For convenience we have shaded one object of every fan 6-gon.

In order to be able to read the diagrams, we need to know which vertices are repeated from object to object. The rule, which is different from the previous subsection, is as follows.

If the value of a label is prescribed (namely, $\ell = 2$ in case 1 and $m = 2$ in case 2), then each apparent edge of a corresponding region stands for one edge.

Any other label only has a determined lower bound. These lower bounds are $k \geq 2$ (all cases), $\ell \geq 3$ (cases 2 and 3), $m \geq 2$ (case 1), $m \geq 3$ (case 3). Consider one of these lower bounds, say “label $\geq r$ ”. Every corresponding region will have $r + 1$ apparent edges, r of which stand for one edge, the remaining *exceptional* one standing for any nonnegative number of edges. In most cases, it is clear or it doesn’t matter which of the apparent edges is the exceptional one and we will not mark it to keep the diagrams simple. The exception is when both boundary edges containing q_0 are contained in the same region.¹ In that case and some other cases the exceptional apparent edge is drawn fat.

Consequently, the diagrams can almost be read as if every apparent edge stands for just one edge.

The main work is in checking that all bounded 2-cells in our diagrams are relations in G^* , so that we may conclude that so are the unbounded 2-cells. This cannot be done in print, but we mention the following. The triangles show how to express an elementary fan morphism u (a dashed arrow) as $c(u)$ (solid arrows), a product of an elementary and a parabolic morphism in some order. Two 2-cells in Figure 7 are (ER4) as indicated. The 4-gon in the middle of Figure 8 (second diagram) is not an elementary or complementary relation but is easily shown to be a consequence of such. All remaining (bounded) 2-cells are (CR4), but they may have the central morphism in a different place as explained in Remark 2.24. This finishes the proof. \square

Note that contraction of the central arrows in the second diagram of Figure 8 (they are enclosed in dashed ovals) recovers diagram (2.6), at least when taken as a planar directed graph.

¹It is rather flexible which vertex is depicted by a white dot, but not so q_0 .




FIGURE 8




FIGURE 9. Word reversing. See Example 3.1.

3. GARSIDE GROUPOIDS

3.1. Introduction. In this section we give a summary of Garside groupoids in general. It aims to be readable by people unfamiliar with Garside groups.

The literature is restricted to Garside groups rather than groupoids, but it is easy to generalise the assertions and proofs of all results pertaining to Garside groups to groupoids. If you want to learn about Garside groupoids, the easiest thing to do is to read about Garside groups first, because their notation is simpler. Conceptually there is hardly any difference. For this reason, we begin by summarising Garside groups. Subsequently we give a summary of Garside groupoids, which of course is almost identical to the group case.

Good overviews on Garside groups are [DehPar99], [Deh00], [Deh02]. Our summary is mainly based on [Deh00], but our terminology is mainly taken from [Deh02].

3.2. Garside groups.

Example 3.1. As an introduction to Garside groups we give an example. It is called the word reversing process and is at the basis of the theory of Garside groups.

Consider the braid group B_n presented by generators σ_i and relations (2.3), (2.4). The braid monoid B_n^+ is the submonoid of B_n generated by the σ_i . Define an ordering \leq on B_n by $x \leq z$ if and only if $z = xy$ for some $y \in B_n^+$. For $x, y \in B_n$, let $x \vee y$ denote the *join* of x, y , that is, the least common upper bound, provided it exists. It is known that the joins $\sigma_i \vee \sigma_j$ exist and are given by

$$(3.1) \quad \begin{aligned} \sigma_i \vee \sigma_j &= \sigma_i \sigma_j \sigma_i, & |i - j| &= 1, \\ \sigma_i \vee \sigma_j &= \sigma_i \sigma_j, & |i - j| &> 1. \end{aligned}$$

We write i for σ_i and aim to compute $2 \vee 13$ (and to prove that this join exists). See Figure 9. By (3.1) we have

$$g = a \vee f, \quad e = c \vee b, \quad h = g \vee d, \quad \ell = h \vee e.$$

(In particular, these joins exist.) Going backwards through these identities, we find

$$\ell = h \vee e = g \vee d \vee e = g \vee e = g \vee c \vee b = g \vee b = a \vee f \vee b = f \vee b.$$

We conclude $2 \vee 13 = 132312$. In algebraic notation the same process looks as follows:

$$\begin{aligned} \underline{2^{-1}1}3 &\rightarrow 121^{-1}\underline{2^{-1}3} \rightarrow 12\underline{1^{-1}3}23^{-1}2^{-1} \\ &\rightarrow 123\underline{1^{-1}2}3^{-1}2^{-1} \rightarrow 123212^{-1}1^{-1}3^{-1}2^{-1}. \end{aligned}$$

Thus, word reversing aims to remove all occurrences $a^{-1}b$ where a, b are (positive) letters.

3.2.1. Complemented presentations. Let A be a finite set. Let A^{-1} be a disjoint copy of A and let $a \mapsto a^{-1}$ denote a bijection from A to A^{-1} , whose inverse is also written $a \mapsto a^{-1}$. Let W^+ denote the free monoid on A and W the free monoid on $A \cup A^{-1}$. We call elements of A *letters*, elements of W *words* and elements of W^+ *positive words*. If $u \in W$ is a word, its inverse u^{-1} is defined by inverting all letters in u and reversing their order. Note that $u^{-1}u \neq 1$ unless $u = 1$.

A *complement* on A is a map f which takes any pair of distinct $a, b \in A$ to a positive word $f(a, b) \in W^+$. We also call (A, f) a complement.

The monoid G^+ (respectively, group G) associated with the above complement is defined as the monoid (respectively, group) presented by generators A and relations

$$(3.2) \quad a f(a, b) = b f(b, a)$$

for all distinct $a, b \in A$.

We have a natural map $W^+ \rightarrow G^+$. We write $u \equiv^+ v$ (where $u, v \in W^+$) if u, v have the same image in G^+ .

Example 3.2. We shall show that the Artin presentation of the braid group is a complemented presentation.

In the above notation, assume that $A = \{\sigma_1, \dots, \sigma_{n-1}\}$ and $f(\sigma_i, \sigma_j) = \sigma_j$ if $|i - j| > 1$ and $f(\sigma_i, \sigma_j) = \sigma_j \sigma_i$ if $|i - j| = 1$. Then the relations (3.2) are precisely the Artin relations (2.3), (2.4) as promised.

3.2.2. The cube condition. For letters a, b and words u, v write

$$u a^{-1} b v \rightarrow u f(a, b) f(b, a)^{-1} v.$$

This is what we called word reversing in Example 3.1. For words u, v write $u \curvearrowright v$ if there are $n \geq 0$ and u_i such that

$$u = u_0 \rightarrow u_1 \rightarrow \dots \rightarrow u_n = v.$$

Suppose u, v, u_0, v_0 are positive words such that $u^{-1}v \curvearrowright v_0u_0^{-1}$. It can be shown that v_0 depends only on u, v ; it will be written $u \backslash v$ (“ u under v ”). It may happen though that $u \backslash v$ is not defined if u_0, v_0 with the required properties don’t exist. Note $a \backslash b = f(a, b)$ for all $a, b \in A$.

We say that our complement satisfies the *cube condition*² if

$$(3.3) \quad (a \backslash b) \backslash (a \backslash c) \equiv^+ (b \backslash a) \backslash (b \backslash c)$$

for all $a, b, c \in A$; here, and henceforth, we take an equation such as (3.3) to mean that either both sides are not defined, or both sides are defined and equivalent as asserted.

If you find the cube condition mysterious, then you may like a more appealing version of it in terms of characteristic graphs, which is explained in Remark 4.6.

²In [Deh02] this is called the *weak cube condition on letters*.

3.2.3. Main theorem on Garside groups.

Definition 3.3. A monoid is called *atomic* if for each element x there is a natural number n such that x cannot be written as a product of n or more nontrivial elements.

Definition 3.4. An *automorphism* of a complement (A, f) (written on the right) is an automorphism ϕ of the associated monoid W^+ such that $A\phi = A$ and $(f(a, b))\phi = f(a\phi, b\phi)$ for all $a, b \in A$.

Definition 3.5. Retain the above notation. A *Garside element* is an element $\Delta \in G^+$ satisfying the following.

- (GE1) For all $a \in A$ we have $\Delta \in aG^+$.
- (GE2) There exists an automorphism ϕ of (A, f) such that for all $a \in A$ we have $a\Delta = \Delta(a\phi)$ (in G^+).

If $\Delta = uv$ (in G^+), then we call u a *simple element*.

Definition 3.6. A *Garside group tuple* consists of a complement (A, f) and a Garside element Δ such that G^+ is atomic, and the cube condition (3.3) is satisfied.

In this case, we call G a *Garside group* and G^+ a *Garside monoid* but technically one works with Garside group tuples.

The following theorem is a rather arbitrary collection of the many good properties Garside groups have, to give the reader an idea. For proofs we refer to [DehPar99], [Deh00], [Deh02] and the references cited therein.

Theorem 3.7. Let (A, f, Δ) be a Garside group tuple. Then the following hold.

- (G1) Let $[u]$ denote the image in G^+ of a morphism u in W^+ . If $u, v \in W^+$, then $[u \setminus v]$ depends only on $[u]$ and $[v]$ and will therefore be written $[u] \setminus [v]$.
- (G2) Let us put an ordering on G^+ by $a \leq c$ if and only if $c = ab$ for some $b \in G^+$. Then G^+ is a lattice. That is, any two elements have a join (a least common upper bound) and a meet (a greatest common lower bound). The join $u \vee v$ of $\{u, v\} \subset G^+$ is $u(u \setminus v) = v(v \setminus u)$.
- (G3) The natural map $G^+ \rightarrow G$ is injective.
- (G4) Every element of G is a quotient of two elements in the image of the map $G^+ \rightarrow G$.
- (G5) If $\Delta = uv$ in G^+ , then v is a simple element.
- (G6) (Greedy form). Every element $u \in G^+$ can uniquely be written $u = u_1 \cdots u_n$ (u_i simple and $u_n \neq 1$ unless $n = 0$) such that if also $u = u_1 \cdots u_{k-1} v_1 \cdots v_\ell$ (v_i also simple, $\ell \geq 1$), then $u_k = v_1 w$ for some w .
- (G7) The group G is automatic.³

3.3. Complemented presentations. We turn to Garside groupoids. Fix a finite set G_0 (which will be the object set of our categories). Let $A(x, y), \overline{A}(x, y)$ ($x, y \in G_0$) be pairwise disjoint sets (which will provide the generating morphisms from x to y). Let $a \mapsto a^{-1}$ denote a bijective map $A(x, y) \rightarrow \overline{A}(y, x)$ (note the orders of x and y); we shall denote its inverse by $a \mapsto a^{-1}$ too. The elements of $A(x, y)$ are called *letters*. Let W be the category with object set G_0 and presented by the generators $A(x, y) \cup \overline{A}(x, y)$ (consisting of morphisms from x to y) for all $x, y \in G_0$,

³See [Eps92] for a definition of automatic groups.

and no relations. Let W^+ be the subcategory of W with the same objects and generated by the letters. We call morphisms of W *words* and morphisms of W^+ *positive words*. If u is a word, then we define u^{-1} as the word obtained by inverting all letters occurring in u and reversing their order. Note that $u^{-1}u \neq 1$ except if u is a trivial morphism.

The (disjoint) union $\bigcup_y A(x, y)$ will be written $A(x, -)$. Let a positive word $f(a, b)$ be given for all $a, b \in A(x, -)$, $x \in G_0$. Suppose that, for all a, b , the positive words $a f(a, b)$ and $b f(b, a)$ are defined⁴ and have the same target object (they already have the same source object). The quotient of W^+ by the relations

$$(3.4) \quad \left\{ a f(a, b) = b f(b, a) \mid a, b \in A(x, -), x \in G_0 \right\}$$

is written G^+ . It is precisely the category presented by the same generators as W^+ and the relations (3.4). The *groupoid* of the same presentation is written G . So there is a natural functor $G^+ \rightarrow G$. We write $u \equiv^+ v$ (u, v morphisms in W^+) if the images in G^+ of u, v are equal.

We call

$$(G_0, \{A(x, y)\}_{xy}, f)$$

a *complemented presentation* or simply *complement*. In the rest of this section, we fix a complement and we use the above notation.

3.4. The cube condition.

For letters a, b and words u, v write

$$u a^{-1} b v \rightarrow u f(a, b) f(b, a)^{-1} v$$

provided the left hand side is defined (in which case so is the right hand side). For words u, v write $u \curvearrowright v$ if there are $n \geq 0$ and u_i such that

$$u = u_0 \rightarrow u_1 \rightarrow \cdots \rightarrow u_n = v.$$

Suppose u, v, u_0, v_0 are positive words such that $u^{-1}v \curvearrowright v_0 u_0^{-1}$. It can be shown that v_0 depends only on u, v ; it will be written $u \setminus v$ and may or may not exist. Note $a \setminus b = f(a, b)$ for all $a, b \in A(x, -)$.

We say that our complement satisfies the *cube condition* if

$$(3.5) \quad (a \setminus b) \setminus (a \setminus c) \equiv^+ (b \setminus a) \setminus (b \setminus c)$$

for all $a, b, c \in A(x, -)$ (and all $x \in G_0$).

3.5. Main theorem on Garside groupoids.

Definition 3.8. A category is called *atomic* if for each morphism x there is a natural number n such that x cannot be written as a product of n or more nontrivial morphisms.

Definition 3.9. By an *automorphism* of a complement (G_0, A, f) (written on the right) we mean an automorphism ϕ of the associated category W^+ of positive words, such that $(A(x, y))\phi = A(x\phi, y\phi)$ for all $x, y \in G_0$ and $(f(a, b))\phi = f(a\phi, b\phi)$ whenever $a, b \in A(x, -)$. Note that the action of ϕ on G_0 may be nontrivial.

⁴The product uv of two morphisms u, v in a category is *defined* if the target object of u is the source object of v .

Definition 3.10. A *Garside automorphism* is a pair

$$(\phi, \{\Delta_x \mid x \in G_0\})$$

where ϕ is an automorphism of the complement (G_0, A, f) and $\Delta_x \in G^+(x, x\phi)$ such that the following hold.

(GA1) Whenever $a \in A(x, y)$ there exists $u \in G^+(y, x\phi)$ such that $au = \Delta_x$ (in G^+).

(GA2) Whenever $a \in A(x, y)$ we have $a\Delta_y = \Delta_x(a\phi)$ (in G^+).

If $\Delta_x = uv$ (in G^+), then we call u a *simple morphism*.

Definition 3.11. A *Garside tuple* consists of a complement (G_0, A, f) and a Garside automorphism $(\phi, \{\Delta_x\}_x)$ such that G^+ is atomic, and the cube condition (3.5) is satisfied.

In this case, we call G a *Garside groupoid* and G^+ a *Garside category*.

The following is the generalisation of Theorem 3.7 to Garside groupoids, and the proof is easily adapted from the proof for the group case. We will use (P2) and (P3) later on.

Theorem 3.12. Let $(G_0, A, f, \phi, \{\Delta_x\}_x)$ be a Garside tuple. Then the following hold.

- (P1) Let $[u]$ denote the image in G^+ of a morphism u in W^+ . If $u, v \in W^+(x, -)$, then $[u \setminus v]$ depends only on $[u]$ and $[v]$ and will therefore be written $[u] \setminus [v]$.
- (P2) The category G^+ has finite limits and colimits. The limit $u \vee v$ of $\{u, v\} \subset G^+(x, -)$ is $u(u \setminus v) = v(v \setminus u)$.
- (P3) The natural functor $G^+ \rightarrow G$ is injective.
- (P4) Every morphism in G is a quotient of two morphisms in the image of the natural functor $G^+ \rightarrow G$.
- (P5) If $\Delta_x = uv$ in G^+ , then v is a simple morphism.
- (P6) (Greedy form). Every morphism u in G^+ can uniquely be written $u = u_1 \cdots u_n$ (u_i simple and $u_n \neq 1$ unless $n = 0$) such that whenever also $u = u_1 \cdots u_{k-1} v_1 \cdots v_\ell$ (v_i also simple, $\ell \geq 1$), then $u_k = v_1 w$ for some w .
- (P7) The groupoid G is automatic.

4. THE MAIN RESULT

4.1. Introduction. The elementary morphisms and elementary relations define a complemented groupoid presentation. In this section, we prove that it is part of a Garside tuple.

Let us recall what we already have. Let $G_0 = M \setminus L$ be the object set of G (Definition 2.4). Let $A(x, y)$ be the set of elementary morphisms from x to y ($x, y \in G_0$); see Definition 2.6. If $a_1, b_1 \in A(x, y)$, then there is a unique elementary relation of the form $a_1 \cdots a_k = b_1 \cdots b_\ell$; see section 2.4. The elementary morphisms and relations yield a complemented presentation as defined in subsection 3.3, which should not be confused with complementary generators and relations (subsection 2.9). The complement f is of course defined by $f(a_1, b_1) := a_2 \cdots a_k$.

We merge the notation of the two previous sections. Note that each of the two sections gave their own definition of G , but that the two definitions are equivalent by Proposition 2.26.

In subsection 4.2 we prove atomicity (3.8). In subsection 4.3 we prove the cube condition (3.4). In the remaining subsections we construct a Garside automorphism (4.17, 4.18) and prove that it is one.

4.2. Atomicity. As every group, \mathbb{Z} can be taken to be a groupoid. We will write \mathbb{Z}_G for it to remind us that it is a groupoid.

It is well known that there is a surjective homomorphism from the braid group to \mathbb{Z} . It follows that there is a surjective functor $w: G \rightarrow \mathbb{Z}_G$. In order to establish that it takes elementary morphisms to nonnegative integers, we construct w in an independent way.

Notation 4.1. If x is an admissible decomposition and p a puncture we will write $R(x, p)$ for the region of x containing p . We will write $\text{arcs}(x)$ for the set of arcs of x .

Lemma/Definition 4.2. (a). There exists a unique functor $w: G \rightarrow \mathbb{Z}_G$ with the following property. See Figure 2. Let (x, y) be an elementary pair. Write

$$\begin{aligned} \text{arcs}(x) \setminus \text{arcs}(y) &= \{A\}, \\ \text{arcs}(y) \setminus \text{arcs}(x) &= \{B\}, \\ A &= R(x, p) \cap R(x, q) \end{aligned}$$

so that also $B = R(y, p) \cap R(y, q)$. Then w takes the elementary morphism $M(x, y)$ to the number of boundary edges of D which are separated from p by both or neither of A, B . As an example, we have indicated the boundary edges with this property by w in Figure 2; there are 6 of them so w takes the elementary morphism of the figure to 6.

(b). The functor $w: G \rightarrow \mathbb{Z}_G$ takes elementary morphisms to nonnegative integers.

(c). The category G^+ is atomic.

Proof. (a). One proves existence of w by checking that w takes every elementary relation to a relation in \mathbb{Z}_G ; this can quickly be done using the diagrams for the elementary relations in Figure 3.

(b). This is obvious.

(c). Let G_0^+ denote the subcategory of G^+ generated by those elementary morphisms x satisfying $w(x) = 0$. By (b) it is enough to prove that G_0^+ is atomic.

Consider first the case where all labels are 2. Then $G_0^+ = G^+$. Also, G^+ is isomorphic to the positive braid monoid $B_n^+ \subset B_n$ generated by $\sigma_1, \dots, \sigma_{n-1}$, which is well known to be atomic; this follows for example from the existence of a homomorphism $B_n \rightarrow \mathbb{Z}$ which takes each σ_i to 1.

Finally we consider the general case. Notice that every morphism of G_0^+ is an endomorphism. It is therefore enough to prove the atomicity of $\text{End}(X)$, the monoid of endomorphisms in G_0^+ of any object X . Now $\text{End}(X)$ is isomorphic to a direct product of positive braid monoids, which is atomic by the foregoing. \square

4.3. Proof of the cube condition.

Convention 4.3. Recall our convention 2.14 on the meaning of apparent edges in objects. In addition, from now on, thin apparent edges will stand for precisely one edge.

Lemma 4.4. *The cube condition (3.4) is satisfied.*

Proof. Let us start with an example. See Figure 10(a). We will prove

$$(4.1) \quad (a \setminus b) \setminus (a \setminus c) \equiv^+ (b \setminus a) \setminus (b \setminus c),$$

where a, b, c are defined by the figure, and which is an instance of the cube condition (3.5). We have

$$\begin{aligned} a \setminus b &= d, \quad a \setminus c = k\ell, \quad b \setminus a = ef, \quad b \setminus c = g, \\ (a \setminus b)^{-1}(a \setminus c) &= \underline{d^{-1}} \underline{k} \ell \curvearrowright \underline{n} \underline{p^{-1}} \ell \curvearrowright \underline{n} \underline{q} \underline{r^{-1}}, \\ (b \setminus a)^{-1}(b \setminus c) &= f^{-1} \underline{e^{-1}} \underline{g} \curvearrowright \underline{f^{-1}} \underline{s} \underline{t^{-1}} \curvearrowright \underline{n} \underline{q} \underline{u^{-1}} \underline{t^{-1}} \end{aligned}$$

so

$$(b \setminus a) \setminus (b \setminus c) = nq = (a \setminus b) \setminus (a \setminus c).$$

It is not hard to do this calculation without writing much down, if you have Figure 10(a) in front of you. As the same diagram helps you verify two more equations (namely (4.1) with a, b, c permuted), drawing these diagrams is a useful step towards verifying the cube condition.

I haven't explained yet how the diagrams come about, even though we have rigorously used one already.

Definition 4.5. A commutative diagram over G^+ all of whose arrows are elementary morphisms is *closed* if the following holds for every elementary relation $a_1 \cdots a_k = b_1 \cdots b_\ell$. If the diagram contains a_1 and b_1 , then it contains the entire elementary relation.

Let $I \subset A(x, -)$ be any subset. The *characteristic graph spanned by I* is the (unique) closed diagram $\text{CG}(I)$ containing I and which is universal for this property. The *rank* of the characteristic graph is $|I|$ and its *initial object* is x .

Verifying the cube condition is done in two steps: one, drawing all rank 3 characteristic graphs; two, verifying the cube condition for each of them as we did in our example of Figure 10(a).

Remark 4.6. Assume for simplicity that every characteristic graph is finite. It can be shown that then the cube condition holds if and only if every characteristic graph has limits. We shall not use this.

Definition 4.7. Let A be an arc of an object $x \in G_0$. The arc A cuts the disk D into two pieces, say, D_1 and D_2 . Let $I \subset A(x, -)$ be such that A is not the rotating arc in any element of I . Then I is a disjoint union $I = I_1 \amalg I_2$ where I_i takes place in D_i .

The characteristic graph $\text{CG}(I)$ is said to be *reducible* if (for some choice of A) the above properties hold and I_1, I_2 are nonempty. We say it is *complete* if $I = A(x, -)$.

It is easy to prove the cube condition in reducible rank 3 characteristic graphs. We turn to the irreducible ones.




FIGURE 10. The irreducible rank 3 characteristic graphs if all labels are ≥ 3 .




FIGURE 11. Initial objects of characteristic graphs classified by shape.

By the *shape* of a characteristic graph we mean the underlying directed graph up to isomorphism. Figure 10(a) gives one shape in detail. Just as the elementary relations (ER2–4), it involves fat apparent edges, which stand for unspecified non-negative numbers of edges, the shape of the characteristic graph being independent of the exact numbers.

Checking the cube condition on a characteristic graph involves only the shape of that graph. Every irreducible characteristic graph has the shape of a complete one. So we need only look at complete rank 3 characteristic graphs from now on. They are completely determined by their initial objects, which are elements of G_0 of 3 arcs.

Figure 11 gives the initial objects of all (complete rank 3) characteristic graphs partitioned in eight *cases* (a)–(h) with the property that the characteristic graphs are of the same shape if (and only if) they are in the same case, as one finds when one draws these graphs explicitly, and which the reader should do. The figure also gives the number of objects in the characteristic graphs. Note that the boundary edges of the initial object can affect the shape of the graph; for example, by contracting two appropriate apparent boundary edges of (a) one obtains (b), which represents a different shape.

The characteristic graphs of cases (a)–(e) are drawn in Figure 10; these are precisely the cases which occur if all labels are ≥ 3 . For convenience, the initial and terminal objects are gray. Case (a) is the only rank 3 shape that does not occur if all labels are ≤ 3 . For simplicity the pictures of the characteristic graphs (b)–(e) assume that all labels are 3; that is, they avoid the fat apparent edges. Using Figure 10 and following our example of case (a) it is then straightforward and not-so-tedious to verify the cube condition for the case where all labels are at least 3.

As to cases (f), (g), (h), we leave it to the reader to draw their characteristic graphs and to verify the cube condition.

This finishes our proof of the cube condition. □

4.4. Shifted decompositions. The remainder of this section, which involves several subsections, is devoted to a proof of the existence of a Garside automorphism.

Let me begin by explaining what is difficult about it. Recall that we are aiming to construct $\Delta_x \in G^+(x, x\phi)$ satisfying two identities in the category G^+ stated in (GA1) and (GA2) in Definition 3.10. It is easy to establish these identities in G

rather than G^+ ; as the example $\langle a, b \mid ab = bb \rangle$ shows, this is not enough though. We need to prove the identities to hold in G^+ , and this is harder.

Let $D_0 = \{z \in \mathbb{C} : |z| \leq 1\}$. Fix a finite nonempty set P of interior points of D_0 and a labelling $\ell: P \rightarrow \mathbb{Z}_{\geq 2}$.

Define m by

$$m - 2 = \sum_{p \in P} (\ell(p) - 2);$$

compare (2.1). Define $S: \partial D_0 \rightarrow \partial D_0$ by $S(z) = z \exp(2\pi i/m)$. By an S -orbit we mean a set of the form $\{S^k(q) \mid k \in \mathbb{Z}\}$ for some $q \in \partial D_0$.

Definition 4.8. A *shifted decomposition* consists of a closed disk $D \subset D_0$ such that $D \cap \partial D_0$ is an S -orbit and such that the interior of D contains P , together with an admissible decomposition for $(D, P, D \cap \partial D_0, \ell)$.

We call $D \cap \partial D_0$ the vertex set of the shifted decomposition. We will soon be looking at pairs of shifted decompositions of distinct vertex sets. A *single* shifted decomposition is essentially the same as an admissible decomposition. Most of our conventions about admissible decompositions easily translate to shifted decompositions but we mention the following.

An arc of a shifted decomposition involving a disk $D \subset D_0$ is by definition an edge not contained in ∂D (it never is in ∂D_0).

Two shifted decompositions of the same vertex set are said to be *isotopic* if they are isotopic relative to ∂D_0 (pointwise) and P . In other words, two shifted decompositions of the same vertex set may be isotopic but not involve the same D .

Pictures of shifted decompositions can be simplified by not drawing the boundary of D . One only draws $D \cap \partial D_0$ and the arcs, that is, interior edges. See Figure 12 (d), (e) for an example.

If Q is an S -orbit we will write $L(Q)$ for the set of isotopy classes of shifted decompositions with vertex set Q . There is a natural bijection between L and $L(S^{\mathbb{Z}}(1))$ where $S^{\mathbb{Z}}(1) = \{S^k(1) \mid k \in \mathbb{Z}\}$ is the S -orbit of 1, which is the set of complex m -th roots of unity.

Every S -orbit Q yields $G(Q)$ and $G^+(Q)$, which are versions of G and G^+ in an obvious way.

4.5. Compatibility, part 1. Let x, y be two shifted decompositions of distinct vertex sets. After an isotopy on one of them we may assume that any edge of x intersects any edge of y transversally in a minimal number of points. The pair (x, y) is then said to be *tight*. The simultaneous isotopy class $[(x, y)]$ of a tight pair depends only on the individual isotopy classes $[x]$ and $[y]$.

Definition 4.9. (See Figure 12). Let x, y be two shifted decompositions of distinct vertex sets in tight position. We say that $[x], [y]$ are *compatible* if the following holds for all punctures p . Loosely speaking, some rotation with centre p takes $R(x, p)$ to $R(y, p)$. More precisely, the vertex sets of $R(x, p)$ and $R(y, p)$ alternate along the boundary, and there exists a homeomorphism $h: D_0 \rightarrow D_0$ such that $hR(x, p)$ and $hR(y, p)$ are convex. (If $\ell(p) > 2$, then an equivalent condition is that $hR(x, p)$ is the convex hull of $hR(x, p) \cap \partial D_0$ and the same for y .)

Definition 4.10. From now on we fix two distinct S -orbits Q_1, Q_2 . For every $y \in L(Q_2)$ we will write $\Omega(y)$ for the set of those $x \in L(Q_1)$ compatible with y . Elements of $L(Q_1)$ are drawn solid and those of $L(Q_2)$ dashed.




FIGURE 12. Compatibility. See Definition 4.9. Elements in $L(Q_1)$ are solid and those in $L(Q_2)$ dashed.

Example 4.11. If the number of arcs is $n - 1 = 2$, then the set of objects in an elementary relation (or rather their lifts to $L(Q_1)$) is of the form $\Omega(y)$. See Figure 12(c) where y is indicated by dashed arcs. Similarly, each of the diagrams in Figure 10 (or again rather their lifts to $L(Q_1)$) is of the form $\Omega(y)$, if $n - 1 = 3$.

The following two lemmas are easy and are left to the reader to prove.

Lemma 4.12. Let $([x], [y]) \in L(Q_1) \times L(Q_2)$ be compatible and suppose that x, y are tight.

- (1) Any edge of x meets any edge of y in at most one point.
- (2) The set $\Omega([y])$ is finite. □

From now on we will abuse language as usual and confuse shifted decompositions with their isotopy classes.

Lemma/Definition 4.13. Let $(x_2, y) \in L(Q_1) \times L(Q_2)$ be compatible. Let A be an arc of x_2 , and let $(x_1, x_2), (x_2, x_3)$ be the elementary pairs where A is moving. Suppose $A = R(x_2, p) \cap R(x_2, q)$. Due to the orientation of D_0 , one of two things can happen. In case 1, all arcs B of y separating p, q look as on the left in Figure 13. In case 2, all arcs B of y separating p, q look as on the right. There is always an arc




FIGURE 13. See Lemma/Definition 4.13.

B of y separating p, q . Case 1 is equivalent to (x_3, y) being compatible. Case 2 is equivalent to (x_1, y) being compatible. The triple (x_2, y, A) is said to be in *case 1* or *case 2* accordingly. \square

4.6. The universal coverings. In this intermezzo we introduce some more language. Let $Q \subset \partial D_0$ be an S -orbit. We shall define the “universal coverings” of our categories $G(Q)$, $G^+(Q)$ written $K(Q)$, $K^+(Q)$.

The object set of $K(Q)$ is $L(Q)$. All hom-sets $K(Q)(x, y)$ consist of one element.

The object set of $K^+(Q)$ is also $L(Q)$. We define $K^+(Q)(x, y)$ to be the set of those elements of $G^+(Mx, My)$ whose images in $G(Mx, My)$ equal $M(x, y)$. The definition of multiplication in $K^+(Q)$ is obvious. We have a commutative diagram as follows:

$$\begin{array}{ccc} K^+(Q) & \longrightarrow & K(Q) \\ \downarrow & & \downarrow \\ G^+(Q) & \longrightarrow & G(Q). \end{array}$$

Next we look at the presentation for $K^+(Q)$, which is of course closely related to the presentation for $G^+(Q)$.

Generators. The generators for $K^+(Q)$ are the elementary pairs in $L(Q) \times L(Q)$. The elementary pair (x, y) is a morphism from x to y .

A *K-word* (respectively, *positive K-word*) is a morphism in the groupoid (respectively, category) with object set $L(Q)$ freely generated by the elementary pairs.

Relations. We call $a_1 \cdots a_k = b_1 \cdots b_\ell$ a *K(Q)-relation* or *K-relation* (assuming that left and right hand sides are $K(Q)$ -words with the same source and target objects) if $(Ma_1) \cdots (Ma_k) = (Mb_1) \cdots (Mb_\ell)$ is an elementary relation (for $G(Q)$). The *K-words* on both sides of the *K-relation* are necessarily positive.

The foregoing taken as a category presentation presents $K^+(Q)$; taken as a groupoid presentation it presents $K(Q)$.

We define an ordering \leq on $L(Q)$ by $x \leq y$ if and only if $K^+(Q)(x, y)$ is nonempty. This is indeed an ordering by 4.2(b) (atomicity). We haven’t yet proved that every hom-set $K^+(Q)(x, y)$ has at most one element or, equivalently, that the natural functor $K^+(Q) \rightarrow K(Q)$ is injective, or again equivalently, that $G^+(Q) \rightarrow G(Q)$ is.

4.7. Compatibility, part 2. An elementary pair $(x, x') \in L(Q_1) \times L(Q_1)$ is said to be *compatible* with $y \in L(Q_2)$ if both x and x' are. Likewise, a *K-word* or *K-relation* is compatible with y if all involved elementary pairs are.

For $y \in L(Q_2)$ let \leq_y denote the ordering on $\Omega(y)$ defined by $x \leq x'$ if and only if there exists a sequence

$$x = z_0, z_1, \dots, z_k = x'$$




FIGURE 14. See the proof of Lemma 4.16.

such that $z_i \in \Omega(y)$ for all i and (z_{i-1}, z_i) is an elementary pair. Equivalently, there exists a positive $K(Q_1)$ -word from x to x' compatible with y .

It is clear that $x \leq_y x'$ implies $x \leq x'$; the converse will follow later on in Theorem 5.1(3).

Definition 4.14. We define a permutation T of $L(Q_1) \cup L(Q_2)$ written on the right. If $\{i, j\} = \{1, 2\}$ and $x \in L(Q_i)$, we define $xT \in L(Q_j)$ by rotating the endpoints of all edges of x in the positive direction along the boundary of the disk D_0 until they first meet Q_j .

Note that for all $y \in L(Q_2)$ one has $yT, yT^{-1} \in \Omega(y)$. These are special elements of $\Omega(y)$ as is shown in the following lemma.

Lemma 4.15. Let $y \in L(Q_2)$. Then $(\Omega(y), \leq_y)$ has a greatest⁵ element yT and a least element yT^{-1} .

Proof. We shall only be dealing with \leq_y , not the ordering on $L(Q)$. Let $x \in \Omega(y)$ be any maximal element. Then for all $A \in \text{arcs}(x)$ the triple (x, y, A) is in case 2 as defined in Lemma/Definition 4.13. A moment's thought now shows that $x = yT$, and we have shown that $\Omega(y)$ has at most one maximal element. Since $\Omega(y)$ is finite by Lemma 4.12(2) it has a greatest element yT . The case of the least element is similar. \square

Lemma 4.16. Let $y \in L(Q_2)$ and let $a_1 \cdots a_k = b_1 \cdots b_\ell$ be a $K(Q_1)$ -relation. If a_1 and b_1 are compatible with y , then so is the entire relation.

Proof. Recall that the $K(Q_1)$ -relations are versions of the elementary relations (ER1–4); see Figure 3. First consider the case where our relation corresponds to (ER3); see Figure 14(a). We assume that y is tight with the source object of a_1 . The fact that a_1 and b_1 are compatible with y implies by Lemma/Definition 4.13 that some of the arcs of y look like the dashed lines in the figure — more precisely, their intersections with the three regions that matter. The dashed lines in turn imply, again by Lemma/Definition 4.13, that the whole of the relation is compatible with y .

In the case (ER4) the fact that a_1 and b_1 are compatible with y implies that two arcs of y look like the dashed lines in Figure 14(b). We leave it to the reader to

⁵Let L be an ordered set (not necessarily totally ordered). A *greatest element* of L is an $x \in L$ such that $y \leq x$ for all $y \in L$. A *maximal element* of L is an $x \in L$ such that there is no $y \in L$ with $x < y$. Similarly for *least* and *minimal*.

conclude that the entire relation is compatible to y , as well as to handle the easier cases of (ER1) and (ER2). \square

Definition 4.17. We define an automorphism ϕ of the groupoid $G(Q_1)$. On objects it is defined by $(Mx)\phi = M(xT^2)$. On morphisms it is defined by $(M(x,y))\phi = M(xT^2, yT^2)$.

We turn to a definition of Δ_z . This is required to be a morphism in $G^+(Q_1)$, not $G(Q_1)$. The attempt $\Delta_{Mx} := M(x, xT^2)$ would define the correct morphism in $G(Q_1)$, but since we haven't yet proved the functor $G^+(Q_1) \rightarrow G(Q_1)$ to be injective, it wouldn't define Δ_z as an element of $G^+(Q_1)$. Our next lemma gets around this.

Lemma/Definition 4.18. Let $y \in L(Q_2)$, so that $x := yT^{-1}$ is the least element of $\Omega(y)$ and xT^2 the greatest by Lemma 4.15. Any positive $K(Q_1)$ -word from x to xT^2 compatible with y defines the same morphism in $K^+(Q_1)$. The image of this morphism in $G^+(Q_1)$ (from Mx to $Mx\phi$) will be written Δ_{Mx} .

Proof. We only deal with \leq_y , not the ordering on $L(Q_1)$. Let $u \in \Omega(y)$ be maximal such that there exist two positive words $a_1 \dots a_k, b_1 \dots b_\ell$ compatible with y from u to xT^2 defining *distinct* morphisms in $K^+(Q_1)$. We aim to deduce a contradiction. The maximality assumption implies $a_1 \neq b_1$. There is a (unique) K -relation $c_1 \dots c_p = d_1 \dots d_q$ with

$$(4.2) \quad c_1 = a_1 \text{ and } d_1 = b_1,$$

and we know the whole of this K -relation to be compatible with y by Lemma 4.16.

Extend the positive K -word $c_1 \dots c_p$ to a positive K -word $c_1 \dots c_r$ ($p \leq r$) compatible with y and which ends at xT^2 . This is possible because xT^2 is the greatest element of $\Omega(y)$. Write \equiv^+ for “defining the same morphism in $K^+(Q_1)$ ”. Then

$$(4.3) \quad \begin{aligned} a_1 \dots a_k &\equiv^+ (c_1 \dots c_p)(c_{p+1} \dots c_r) \\ &\equiv^+ (d_1 \dots d_q)(c_{p+1} \dots c_r) \equiv^+ b_1 \dots b_\ell \end{aligned}$$

where the middle \equiv^+ is obvious and the other two equivalences follow from (4.2) and the maximality of u and the fact that all K -words in (4.3) are compatible with y by Lemma 4.16. This contradiction proves the lemma. \square


Corollary 4.19. Let $a \in A(x, y)$ with $x, y \in G_0$. Then there exists $u \in G^+(y, x\phi)$ such that $au = \Delta_x$ in G^+ .

Proof. Write $z = xT$. Note that $y \in \Omega(z)$. Using Lemma 4.15, there is a positive K -word compatible with z from y to $\max \Omega(z) = x\Delta_x$. Its image u in G^+ satisfies $au = \Delta_x$. \square

Definition 4.20. Let $Y \subset L(Q_2)$. We define $\Omega(Y) := \bigcap_{y \in Y} \Omega(y)$ which we endow with an ordering \leq_Y defined by $x \leq_Y x'$ if and only if there exists a positive $K(Q_1)$ -word compatible with Y (that is, with all elements of Y) from x to x' . If $x \leq_Y x'$, then $x \leq_y x'$ for all $y \in Y$.

Lemma 4.21. *Let $(y, z) \in L(Q_2) \times L(Q_2)$ be an elementary pair and put $Y = \{y, z\}$. With respect to \leq_Y then, $\Omega(Y)$ has a greatest element yT and a least element zT^{-1} .*

Proof. Part (a) of



shows some common edges of y and z (solid) and a dashed arc belonging to y and a dotted arc belonging to z .

Let $x \in \Omega(Y)$ be maximal and suppose $x \neq yT$. Since $x \in \Omega(y)$ and $x \neq yT = \max \Omega(y)$ (by Lemma 4.15) there is an elementary pair (x, x') with $x' \in \Omega(y)$, say obtained by rotating the arc A of x . By Lemma/Definition 4.13, some arc B of y looks like the dashed line in (4.4b) where $A = R(x, p) \cap R(x, q)$.

Since x is maximal in $\Omega(Y)$, we have $x' \notin \Omega(Y)$ and therefore $x' \notin \Omega(z)$. Also $x \in \Omega(z)$, so by Lemma/Definition 4.13 some arc C of z looks like the dotted line in (4.4b).

But no pair $\in \text{arcs}(y) \times \text{arcs}(z)$ intersects the way B and C do as dictated by the figure. This is a contradiction. We have proved that $\Omega(Y)$ has at most one maximal element yT . Since $\Omega(Y)$ is finite it has a greatest element yT . The case of a least element is similar. \square

Lemma 4.22. *Let $a \in A(x, y)$ be an elementary morphism. Then $a\Delta_y = \Delta_x(a\phi)$.*

Proof. Let $(u, v) \in L(Q_1) \times L(Q_1)$ be an elementary pair such that $a = M(u, v)$. Then $(uT, vT) \in L(Q_2) \times L(Q_2)$ is an elementary pair too. Put $Y = \{uT, vT\}$. By Lemma 4.21, v is the least element of $\Omega(Y)$ (relative to \leq_Y) and uT^2 the greatest.

Choose any positive $K(Q_1)$ -word compatible with both elements of Y , from v (the least element) to uT^2 (the greatest), and let b be the morphism in $K^+(Q_1)$ defined by that word. Let $\pi: K^+(Q_1) \rightarrow G^+(Q_1)$ be the natural functor. Then $\pi((u, v)b) = \Delta_x$ and $\pi(b(uT^2, vT^2)) = \Delta_y$ by Lemma/Definition 4.18. Clearly we also have $\pi(u, v) = a$ and $\pi(uT^2, vT^2) = a\phi$. We get

$$\begin{aligned} a\Delta_y &= \pi(u, v) \cdot \pi(b(uT^2, vT^2)) \\ &= \pi((u, v)b) \cdot \pi(uT^2, vT^2) = \Delta_x(a\phi). \end{aligned}$$

Here is our main theorem.

Theorem 4.23. *The tuple $(G, \{A(x, y)\}_{xy}, f, \{\Delta_x\}_x, \phi)$ is a Garside tuple.*

Proof. This follows from Lemma/Definition 4.2(c) (atomicity), Lemma 4.4 (cube condition), Corollary 4.19 and Lemma 4.22. \square

A *lattice* is an ordered set (P, \leq) such that any two elements x, y have a least common upper bound (or *join*) and a greatest common lower bound (or *meet*).


An ordered set is the same as a category where every hom-set has at most one element, and no two objects are isomorphic. A lattice is then an ordered set in

which any two elements have a common upper bound and a common lower bound, and finite limits and colimits exist. This is the language we used in Theorem 3.12, (P2). It is clear that any two elements of L have a common upper bound and a common lower bound. So a corollary of Theorem 4.23 and (P2) and (P3) is the following.

Corollary 4.24. *The ordered set (L, \leq) is a lattice.* \square

It is clear from the proof of Lemma 2.11 that Garside's original Garside structure on the braid group [Gar69] is a special case of our construction, namely the case where all labels are 2.

Example 4.25. Let $x \in L$. If the orbit Mx is a sublattice of L (that is, closed under join and meet) we obtain a lattice ordering on M by putting $g \leq h \Leftrightarrow gx \leq hx$. But M -orbits in L are not sublattices of L in general, as is shown by



This diagram is an elementary relation, so $z = x \vee y$. Now x, y are in the same orbit but $x \vee y$ is in a different orbit.

5. FINITE LATTICES AND ASSOCIAHEDRA

Recall that in Corollary 4.24 we constructed a lattice $L(Q_1)$. The braid-like group M acts freely on it with finitely many orbits. We also have the finite subsets $\Omega(y) \subset L(Q_1)$ where $y \in L(Q_2)$. If all labels are 2, then $L(Q_1)$ can be identified with the braid group B_n and $\Omega(y)$ with the symmetric group S_n .

In Theorem 5.1 we prove that $\Omega(y)$ is a sublattice of $L(Q_1)$. In particular, it is a lattice in its own right. Most particular cases of these finite lattices seem to be new.

It seems that $\Omega(y)$ is the vertex set of a natural polytope (up to deformation); we shall not go into this. If all labels are 2, it is known as the permutohedron.

In subsection 5.2 we study the case where all labels are 3 more closely.

The *associahedron* is a certain polytope whose vertex set is the set of triangulations of a fixed n -gon such that the vertex set of the triangulation is the vertex set of the n -gon [Sta63], [Lee89].

Suppose now that all labels are 3. Recall that the object set of the groupoid $G = G(Q_1)$ is $G_0 = M \setminus L(Q_1)$. This is essentially the set of triangulations of the disk D_0 with vertex set Q_1 , that is, the vertex set of the associahedron.

In Proposition 5.2 we prove that $\Omega(y)$ is in bijection with G_0 . Combined with the lattice ordering on $\Omega(y)$ mentioned above this yields a family of lattice orderings on the vertex set of the associahedron. One of these orderings is known as the Tamari lattice.

We have already come across pictures of our orderings on the vertex set of the 3-dimensional associahedron in Figure 10 (b)–(d) when we were studying characteristic graphs. The 3-dimensional Tamari lattice is part (c) of the figure.


In contrast, we obtain essentially only one lattice ordering on the vertex set of the permutohedron, because if all labels are 2, then the M -action on L is transitive.

5.1. More on $\Omega(y)$.

Theorem 5.1. *Let $y \in L(Q_2)$.*

- (1) *Let $a_1 \cdots a_k = b_1 \cdots b_\ell$ be a $K(Q_1)$ -relation (in particular, the a_i and b_j are elementary pairs in $L(Q_1) \times L(Q_1)$). Then $a_1 \cdots a_k$ is compatible with y if and only if $b_1 \cdots b_\ell$ is.*
- (2) *We have $\Omega(y) = \{x \in L(Q_1) : yT^{-1} \leq x \leq yT\}$.*
- (3) *Let $x, x' \in \Omega(y)$. Then $x \leq_y x'$ is equivalent to $x \leq x'$.*
- (4) *The ordered set $(\Omega(y), \leq)$ is a lattice.*

Proof. (1) Suppose the relation is of the form (ER3) and more precisely reads $a_1a_2a_3 = b_1b_2$ as in Figure 3. We will show that if $a_1a_2a_3$ is compatible with y then so is b_1b_2 . We apply Lemma/Definition 4.13 three times as follows. Since a_1 is compatible with y , at least one of the dashed lines in part (a) of



describes an arc in y . Since a_3 is compatible with y , at least one of the dashed lines in part (b) also describes an arc in y . Since arcs of y don't intersect in the interior of the disk D_0 both dashed lines in part (c) describe arcs of y . They in turn imply that b_1b_2 is compatible with y . They also imply that Figure 14(a) matches our situation, including the dashed lines.

The reverse implication for (ER3) as well as the remaining elementary relations are similar and left to the reader. Note that (ER4) is slightly harder than (ER3) because the punctures cannot be omitted.

(2) The inclusion \subset follows from Lemma 4.21. We shall prove \supset . Let X be the set of positive K -words from yT^{-1} to yT . Let d be the largest metric (in a generalised sense, namely distances can be ∞) such that $d(uv_1w, uv_2w) = 1$ if " $v_1 = v_2$ " is a K -relation. In fact, any two elements of X have finite distance because the functor $G^+ \rightarrow G$ is injective by (P3) and our main theorem 4.23. But (1) states that if two elements of X have distance 1 and one is compatible with y , then so is the other. Since at least one element of X is compatible with y (by Lemma 4.21), all are as required.

(3) The implication \Rightarrow is trivial. We prove \Leftarrow . Let $x \leq x'$. Then there exists a positive K -word w from x to x' . By part (2), w is compatible with y . It follows that $x \leq_y x'$.

(4) This follows directly from (2) and the fact that $(L(Q_1), \leq)$ is a lattice by Corollary 4.24. \square

5.2. The case of only triangles. In this subsection, we assume all labels to be 3.

Proposition 5.2. *Suppose all labels are 3. For any $y \in L(Q_2)$, the natural map $\pi: \Omega(y) \rightarrow G_0$ is bijective.*

Proof. Injectivity is easy and left to the reader (and true even if all labels are ≥ 3).

We shall prove surjectivity. Let $x' \in G_0$, $y \in L(Q_2)$ be tight.

Let R be a region of y with vertices r_1, r_2, r_3 . Among all arcs of x' intersecting R and separating r_i from the other two vertices ($i = 1, 2, 3$),⁶ let A_i be the innermost one. By R' we shall denote the region of x' containing A_1, A_2, A_3 .

There exists a self-homeomorphism of D_0 fixing the boundary pointwise which preserves y and which takes the puncture in any region R of y to an interior point of $R \cap R'$. Then $x := h^{-1}x'$ is in $\Omega(y)$ (using the fact that all labels are 3); it is clear that $\pi x = x'$. This proves surjectivity. \square

Define π as in Proposition 5.2. For any $x \in L(Q_1)$ we define an ordering $\leq_{\pi x}$ on G_0 as follows. For $y, z \in \Omega(xT)$ we put

$$\pi y \leq_{\pi x} \pi z \iff y \leq z,$$

which is equivalent to $y \leq_{xT} z$ by Theorem 5.1(3). Clearly, $\leq_{\pi x}$ depends only on πx .

Notice that $x \leq_x y$ for all $x, y \in G_0$, and that \leq_x is a lattice ordering on G_0 , the vertex set of the associahedron. So we have many lattice orderings \leq_x on G_0 . If x is a fan, this ordering was discovered by Tamari [Tam51], [FriTam67], [Grä78, page 18]. If x is another triangulation of the n -gon it seems to be new.

We get the following amusing property, which we shall not prove. The braid-like groupoid G with object set G_0 is presented by a generator $[xy] \in G(x, y)$ (for all $x, y \in G_0$) and relations $[xx] = 1$ for all x and $[xy][yz] = [xz]$ whenever $y \leq_x z$.

REFERENCES

- [BKL98] Birman, Joan; Ko, Ki Hyoung; Lee, Sang Jin. A new approach to the word and conjugacy problems in the braid groups. *Adv. Math.* **139** (1998), no. 2, 322–353. MR1654165 (99m:20082)
- [Deh00] Dehornoy, Patrick. Chapter 2 in *Braids and self-distributivity*. Progress in Mathematics, 192. Birkhäuser-Verlag, Basel, 2000. MR1778150 (2001j:20057)
- [Deh02] Dehornoy, Patrick. Groupes de Garside. *Ann. Sci. École Norm. Sup. (4)* **35** (2002), no. 2, 267–306. MR1914933 (2003f:20068)
- [DehPar99] Dehornoy, Patrick; Paris, Luis. Gaussian groups and Garside groups, two generalisations of Artin groups. *Proc. London Math. Soc. (3)* **79** (1999), no. 3, 569–604. MR1710165 (2001f:20061)
- [Eps92] Epstein, D.B.A.; Cannon, J.W.; Holt, D.F.; Levy, S.V.F.; Paterson, M.S.; Thurston, W.P. Word processing in groups. Jones and Bartlett Publishers, Boston, MA, 1992. MR1161694 (93i:20036)
- [FriTam67] Friedman, Haya; Tamari, Dov. Problèmes d’associativité: Une structure de treillis finis induite par une loi demi-associative. *J. Combinatorial Theory* **2** (1967), 215–242. MR0238984 (39:344)
- [Gar69] Garside, F. A. The braid group and other groups. *Quart. J. Math. Oxford Ser. (2)* **20** (1969), 235–254. MR0248801 (40:2051)
- [Grä78] Grätzer, George. *General lattice theory*. Second edition. Birkhäuser-Verlag, Basel, 1998. (First edition published 1978). MR1670580 (2000b:06001)
- [Lee89] Lee, Carl W. The associahedron and triangulations of the n -gon. *European J. Combin.* **10** (1989), no. 6, 551–560. MR1022776 (90i:52010)
- [Par05] Paris, Luis. From braid groups to mapping class groups. *Proc. Sympos. Pure Math.* **74** (2006), 355–371.

⁶There is at least one such arc because of the boundary edges of x' .

- [Sta63] Stasheff, James. Homotopy associativity of H -spaces I. *Trans. Amer. Math. Soc.* **108** (1963), 275–292. MR0158400 (28:1623)
- [Tam51] Tamari, Dov. Monoïdes préordonnés et chaines de Malcev. Thesis, Paris, 1951. MR0051833 (14:532b)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WARWICK, COVENTRY CV4 7AL, UNITED KINGDOM

E-mail address: daan@maths.warwick.ac.uk