

CONSTRUCTION OF COMPLETE EMBEDDED SELF-SIMILAR SURFACES UNDER MEAN CURVATURE FLOW. PART I.

XUAN HIEN NGUYEN

ABSTRACT. We carry out the first main step towards the construction of new examples of complete embedded self-similar surfaces under mean curvature flow. An approximate solution is obtained by taking two known examples of self-similar surfaces and desingularizing the intersection circle using an appropriately modified singly periodic Scherk surface, called the core. Using an inverse function theorem, we show that for small boundary conditions on the core, there is an embedded surface close to the core that is a solution of the equation for self-similar surfaces. This provides us with an adequate central piece to substitute for the intersection.

1. INTRODUCTION

This paper is the first one of a series of three articles describing the construction of new examples of complete embedded self-similar surfaces under mean curvature flow. Our general strategy is inspired by Kapouleas' article [7]. His success in constructing complete embedded minimal surfaces motivates us to adapt his method towards finding self-similar surfaces under mean curvature flow.

Self-similar solutions are solutions to the mean curvature flow that do not change shape but are merely contracted (called self-shrinkers) or dilated (self-expanders) by it. Up to scaling, the self-shrinkers satisfy the equation

$$(1) \quad H + X \cdot \nu = 0,$$

where H is the mean curvature and ν is the normal vector so that the mean curvature vector is $\mathbf{H} = H\nu$. The sign of H is chosen so that the mean curvature of a convex surface is positive. These solutions are of special interest because they model the behavior of the mean curvature flow at singularities if the blow-up is of type I [5]. There are currently only four known embedded complete self-shrinkers: a plane, a sphere, a cylinder and a shrinking doughnut [1]. However, there exists numerical evidence of others [2], [3]. The availability of new examples will lead to a better understanding of the behavior of the flow near singularities; therefore, it is essential to find successful methods to construct them.

1.1. Strategy. To construct a new self-similar surface, we take two known examples and replace a neighborhood of their intersection with an appropriately bent

Received by the editors September 13, 2006.

2000 *Mathematics Subject Classification.* Primary 53C44.

Key words and phrases. Mean curvature flow, self-similar, singularities.

©2008 American Mathematical Society
Reverts to public domain 28 years from publication

scaled Scherk's singly periodic surface, called the core and denoted by $\tilde{\Sigma}_{1/N}^C$. In the case of a cylinder of radius one and a plane perpendicular to the axis of the cylinder, the desingularization process is shown in Figure 2 while Figure 1 shows a portion of the original unmodified Scherk surface. This idea also works if the cylinder is replaced by a sphere.

The resulting surface is not smooth; however it is a good approximate solution. The next task is to find a function whose graph over it satisfies the self-shrinker equation (1). Before considering graphs of functions on the entire surface, we have to work locally and study the Dirichlet problems with small boundary conditions corresponding to (1) for graphs of functions on the five different pieces: the core, the outer plane, the inner disk and the two cylindrical ends (or spherical caps, if we work with a sphere).

In this article, we give a precise description of the maps Φ_τ used to bend and the maps \mathcal{H}_τ used to scale the singly periodic Scherk surface as well as tackle the Dirichlet problem on the central core piece. The second article [9] shows that the Dirichlet problem corresponding to equation (1) for graphs of functions on the unbounded outer plane has a solution. These are the two more difficult Dirichlet problems. The results remain valid, whether we use a sphere or a cylinder in the construction. The third article [10] is to discuss the gluing of the solutions on the different pieces in a manner so as to obtain a smooth complete embedded self-similar surface.

1.2. Main result. Roughly speaking, our main result states that if the bent scaled Scherk surface used for desingularizing is scaled enough, then a small perturbation of it satisfies (1) and imposed boundary conditions, provided the latter are small enough. The statement is made more precise below.

Let us denote the bent scaled surface $\mathcal{H}_{1/N}(\Phi_{1/N}(S_C))$ by $\tilde{\Sigma}_{1/N}^C$ (this is the core in Figure 2). By our choice of bending and scaling, the surface $\tilde{\Sigma}_{1/N}^C$ has N handles. Moreover, it is invariant with respect to the rotation of 180 degrees about the x -axis and with respect to symmetries across the planes $\theta = \pi/2N + k\pi/N$, $k = 1, \dots, 2N$, where we used cylindrical coordinates.




FIGURE 1. Scherk surface




FIGURE 2. Removing a neighborhood of the intersection and inserting the core $\tilde{\Sigma}_{1/N}^C$

Theorem 1. *There is a constant C_1 so that, for any $C > C_1$, there exist an integer $N_0 > 0$ and a constant $\delta_0 > 0$ with the following properties: for every integer $N > N_0$ and for every function $\tilde{f} \in W^{2,p}(\tilde{\Sigma}_{1/N}^C)$ with $\|\tilde{f}\|_{W^{2,p}(\tilde{\Sigma}_{1/N}^C)} \leq \delta_0/(2N^2)$ and whose graph over $\tilde{\Sigma}_{1/N}^C$ satisfies the above symmetries, there exists a function $\tilde{h} \in W^{2,p}(\tilde{\Sigma}_{1/N}^C)$ such that*

$$\begin{aligned} & \text{the graph of } \tilde{h} \text{ over } \tilde{\Sigma}_{1/N}^C \text{ satisfies } H + X \cdot \nu = 0, \\ & \text{the graph of } \tilde{h} \text{ over } \tilde{\Sigma}_{1/N}^C \text{ enjoys the above symmetries,} \\ & \tilde{h} = \tilde{f} \text{ on } \partial\tilde{\Sigma}_{1/N}^C. \end{aligned}$$

Moreover, we can choose N_0 and δ_0 so that the graph of h over $\tilde{\Sigma}_{1/N}^C$ is an embedded surface.

Imposing symmetries on the solutions and boundary conditions that naturally mirror the symmetries of the Scherk surface greatly simplifies the problem. We can work on a piece of surface characterized by a single period and then reconstruct the complete surface using the symmetries. The central tool in the proof is the Inverse Function Theorem. The fact that the Scherk surface is minimal is also used strongly: the Gauss map is then conformal and allows us to transport partial differential equations to the sphere, where elliptic operators have been well studied.

Our methods for solving the Dirichlet problems differ from Kapouleas'. He was able to slightly change the relative position of the Scherk surface wings as well as the size and position of his catenoidal ends in order to compensate for the existence of an approximate kernel. Since in our case, the radius of the cylinder and position of the plane are fixed by equation (1), the Dirichlet problems have to be solved directly.

This paper is part of the author's thesis, written under the direction of Sigurd Angenent at the University of Wisconsin-Madison. The author is indebted to professor Angenent for invaluable discussion.

2. DEFINITIONS AND BENDING MAPS

In this section, we will describe in detail the transformations made on the Scherk surface S in order to obtain a suitable setting to define and solve a Dirichlet problem.

2.1. **Scherk surface.** Consider the Scherk surface S given by

$$(2) \quad \sin y = \sinh x \sinh z$$

and shown in Figure 1.

Lemma 2. *The surface S is a singly periodic embedded minimal surface which has the following properties:*

- S is asymptotic to the xy -plane and to the yz -plane,
- S is invariant under rotation of 180° about the axes $\{(x, k\pi, 0)\}$ ($k \in \mathbf{Z}$),
- S is invariant under reflections with respect to the planes $\{y = \pi/2 + k\pi\}$ ($k \in \mathbf{Z}$).

Proof. The symmetries can be readily derived from equation (2). A direct computation shows that the mean curvature vanishes. For more information on the Scherk surface, we refer to [4]. \square

We truncate our surface using the domain

$$(3) \quad \Omega = \{(x, y, z) \in \mathbf{R}^3 \mid |x| < C, \frac{-3\pi}{2} < y < \frac{\pi}{2}, |z| < C_0\},$$

for some large constant $C_0 > 100$ chosen later. We omit the dependence on C_0 in our notation for Ω and for the surfaces defined in this article, when we do not wish to emphasize it.

Fix a constant η so that $0 < \eta \leq \frac{1}{2C}$. For all $\tau \in (-2\eta, 2\eta)$, define the bending maps $\Phi_\tau : \Omega \rightarrow \mathbf{R}^3$ in the following manner:

$$(4) \quad \Phi_\tau(x, y, z) = \begin{cases} \left(\frac{1}{\tau}e^{\tau x} \cos(\tau y) - \frac{1}{\tau}, \frac{1}{\tau}e^{\tau x} \sin(\tau y), z\right), & \tau \neq 0, \\ (x, y, z), & \tau = 0. \end{cases}$$

The maps Φ_τ transform the yz -plane into cylinders of radius τ^{-1} around the axis $x = -\tau^{-1}, y = 0$. Therefore, bending the Scherk surface S using Φ_τ creates surfaces $\Phi_\tau(S)$ asymptotic to cylinders of radius τ^{-1} . In the end, we want our surfaces to be asymptotic to a cylinder of radius 1; however, working at this larger scale allows us to define a nonsingular surface for the case $\tau = 0$ and a smooth family of surfaces in a neighborhood of $\tau = 0$.

Denote by Σ_τ the bent truncated Scherk surface

$$(5) \quad \Sigma_\tau = \Phi_\tau(S \cap \Omega).$$

Lemma 3. *Let the maps Φ_τ be as in (4). Then the map*

$$\begin{aligned} \Phi : (-\eta, \eta) &\rightarrow C^5(\overline{\Omega}, \mathbf{R}^3), \\ \tau &\mapsto \Phi(\tau) = \Phi_\tau \end{aligned}$$

is continuously Fréchet differentiable.

Proof. From the explicit formula (4) defining the Φ_τ 's, it is immediate that Φ is continuously differentiable in all its variables up to order six and its derivatives are bounded on $(-\eta, \eta) \times \overline{\Omega}$. The argument is then completed by applying the Mean Value Theorem. \square

2.2. Tensor bundles on Σ_0 . Consider $\bar{\Sigma}_0$, the closure of Σ_0 as a smooth manifold with boundary. The metric on Σ_0 is induced by the Euclidean metric on \mathbf{R}^3 .

A tensor r times contravariant and s times covariant is called an (r, s) type tensor. Denote by $T_{r,s}$ the tensor bundle of type (r, s) over Σ_0 . This vector bundle possesses a metric induced by the metric of Σ_0 . An (r, s) type tensor f over Σ_0 is a section of $T_{r,s}$.

2.3. Definition of $\Sigma_{h,\tau}, X^{(h,\tau)}, a^{(h,\tau)}, A^{(h,\tau)}, \nu^{(h,\tau)}$ and $H^{(h,\tau)}$. We consider the surfaces $\bar{\Sigma}_\tau$ as smooth manifolds with boundary with respective metrics g_{Σ_τ} induced by the Euclidean metric in \mathbf{R}^3 . Let $p \in \Sigma_0$ and denote by $X^{(0,0)}(p)$ and $\nu^{(0,0)}(p)$ the position vector and the normal vector at p respectively.

The vectors $X^{(0,\tau)}(p)$, $\nu^{(0,\tau)}(p)$ and the function $H^{(0,\tau)}$ are pull backs of the position vector, normal vector and the mean curvature at $\Phi_\tau(p) \in \Sigma_\tau$ to $p \in \Sigma_0$, i.e.

$$\begin{aligned} X^{(0,\tau)}(p) &= \Phi_\tau(X^{(0,0)}(p)) = \Phi_\tau(p), \\ \nu^{(0,\tau)}(p) &= \text{unit normal to } \Sigma_\tau \text{ at } (\Phi_\tau(X^{(0,0)}(p))), \\ H^{(0,\tau)}(p) &= \text{mean curvature of } \Sigma_\tau \text{ at } (\Phi_\tau(X^{(0,0)}(p))). \end{aligned}$$

Similarly, the tensor $g^{(0,\tau)} \in T_{0,2}$ is the pull back to Σ_0 of the metric on Σ_τ , the tensor $a^{(0,\tau)} \in T_{0,2}$ is the pull back to Σ_0 of the second fundamental form on Σ_τ , and $A^{(0,\tau)} \in T_{1,1}$ is the pull back of the second fundamental tensor on Σ_τ to Σ_0 using the map Φ_τ : for $\xi, \rho \in T_p \Sigma_0$,

$$\begin{aligned} g^{(0,\tau)}(p)(\xi, \rho) &= (\Phi_\tau^* g_{\Sigma_\tau})(p)(\xi, \rho) = g_{\Sigma_\tau}(\Phi_\tau(p))(\Phi_{\tau*}\xi, \Phi_{\tau*}\rho), \\ a^{(0,\tau)}(p)(\xi, \rho) &= (\Phi_\tau^* a_{\Sigma_\tau})(p)(\xi, \rho) = a_{\Sigma_\tau}(\Phi_\tau(p))(\Phi_{\tau*}\xi, \Phi_{\tau*}\rho), \\ \langle \rho, A^{(0,\tau)}(p)\xi \rangle &= \langle \rho, (\Phi_\tau^* A_{\Sigma_\tau})(p)\xi \rangle = \langle \Phi_{\tau*}\rho, A_{\Sigma_\tau}(\Phi_\tau(p))(\Phi_{\tau*}\xi) \rangle, \end{aligned}$$

where g_{Σ_τ} , a_{Σ_τ} and A_{Σ_τ} are respectively the metric, the second fundamental form and the type $(1, 1)$ second fundamental tensor of $(\Sigma_\tau, g^{(0,\tau)})$.

For a continuous function $h : \Sigma_0 \rightarrow \mathbf{R}$, the graph of h over Σ_τ is defined by

$$\Sigma_{(h,\tau)} = \{X^{(0,\tau)}(p) + h(p)\nu^{(0,\tau)}(p), \quad p \in \Sigma_0\}.$$

Claim 4. For all $\eta_0 < \frac{1}{200}$, there is a constant $\delta > 0$ depending on η_0 such that for all functions $h : \Sigma_0 \rightarrow \mathbf{R}$ with $|h| < \delta$ and constants $\tau \in (-\eta, \eta)$, the surface $\Sigma_{(h,\tau)}$ is embedded. For example, if $\eta_0 = 1/200$, we can take $\delta = 1/8$.

From the proof below, note that δ can be chosen larger, as the given η_0 is smaller.

Proof. The transformations Φ_τ are continuously differentiable with respect to τ and tend to the identity as $\tau \rightarrow 0$. We want the graph of a function on the surface to be embedded; the only problematic parts are the ‘‘holes’’ in the Scherk surface. The ‘‘hole’’ in the domain $0 \leq y \leq \pi$ is at its narrowest in the plane $x = z$. If we denote by u and y the coordinates in this plane, the intersection curve of S with it is characterized by the equation $\sin y = \sinh^2(u/\sqrt{2})$. The previous equation implies that the graph of a function h with $|h| < 1/2 < \frac{1}{2} \min(\pi, 2\sqrt{2} \sinh^{-1} 1)$ over Σ_0 is embedded. The mappings Φ_τ for $\tau \in (-1/100, 1/100)$ only change the surface Σ_0 slightly and taking $\delta < 1/8$ is safe, although it is a very crude estimate. \square

For h as in the previous lemma, consider the graphs $\Sigma_{h,\tau}$ as manifolds. We can equip the surfaces $\Sigma_{h,\tau}$ with metrics $g_{\Sigma_{h,\tau}}$ induced by the embedding and the Euclidean metric in \mathbf{R}^3 . Define

$$X^{(h,\tau)}(p) = X^{(0,\tau)}(p) + h(p)\nu^{(0,\tau)}(p)$$

and the map $\Phi_{h,\tau} : \Sigma_0 \rightarrow \Sigma_{h,\tau}$ by $\Phi_{h,\tau}(p) = X^{(h,\tau)}(p)$. We can now define

$$\begin{aligned} \nu^{(h,\tau)}(p) &= \text{unit normal to } \Sigma_{h,\tau} \text{ at } \Phi_{h,\tau}(X^{(0,0)}(p)), \\ H^{(h,\tau)}(p) &= \text{mean curvature of } \Sigma_{h,\tau} \text{ at } \Phi_{h,\tau}(X^{(0,0)}(p)). \end{aligned}$$

The $(0,2)$ tensor $g^{(h,\tau)}$, the $(0,2)$ tensor $a^{(h,\tau)}$ and the $(1,1)$ tensor $A^{(h,\tau)}$ are respectively the metric, the second fundamental form and the second fundamental tensor on $\Sigma_{h,\tau}$ pulled back to Σ_0 using $\Phi_{h,\tau}$:

$$g^{(h,\tau)} = \Phi_{h,\tau}^* g_{\Sigma_{h,\tau}}, \quad a^{(h,\tau)} = \Phi_{h,\tau}^* a_{\Sigma_{h,\tau}}, \quad A^{(h,\tau)} = \Phi_{h,\tau}^* A_{\Sigma_{h,\tau}},$$

where $g_{\Sigma_{h,\tau}}$, $a_{\Sigma_{h,\tau}}$ and $A_{\Sigma_{h,\tau}}$ are respectively the metric, the second fundamental form and the second fundamental tensor on $\Sigma_{h,\tau}$.

By the definitions above, all of the vectors and tensors with superscript (\cdot, τ) are sections of Σ_0 into either \mathbf{R}^3 or a tensor bundle over Σ_0 .

3. FRÉCHET DIFFERENTIABILITY

3.1. Scaling, translating and modified equation. We wish to replace a neighborhood of the intersection of a cylinder of radius 1 about the z -axis and a plane by a surface satisfying (1); however, the surfaces $\Sigma_{h,\tau}$ are pieces of surfaces asymptotic to cylinders of radius τ^{-1} about the line $(-\tau^{-1}, 0, z)$. Working in this larger scale requires us to modify the equation above to take into account the different scale and displacement of the axis of the cylinder.

Define the map

$$(6) \quad \mathcal{H}_\tau : \mathbf{R}^3 \rightarrow \mathbf{R}^3, \quad \mathcal{H}_\tau(x, y, z) = \tau(x + \frac{1}{\tau}, y, z).$$

Lemma 5. *If $\tau \neq 0$ and $h \in W^{2,p}(\Sigma_0)$ is such that*

$$(7) \quad H^{(h,\tau)} + \tau e_1 \cdot \nu^{(h,\tau)} + \tau^2 X^{(h,\tau)} \cdot \nu^{(h,\tau)} = 0,$$

where $e_1 = (1, 0, 0)$ is the first coordinate vector in \mathbf{R}^3 , then the rescaled surface $\mathcal{H}_\tau(\Sigma_{h,\tau})$ satisfies the equation for contracting self-similar surfaces $H + X \cdot \nu = 0$, where H , X , and ν are the mean curvature, the position vector and the normal vector to $\mathcal{H}_\tau(\Sigma_{h,\tau})$ respectively.

Proof. We have $X^{(h,\tau)} = \frac{1}{\tau}X - (\frac{1}{\tau}, 0, 0) = \frac{1}{\tau}(X - (1, 0, 0))$; therefore $H^{(h,\tau)} = \tau H$, $\nu^{(h,\tau)} = \nu$, and

$$\begin{aligned} 0 &= H^{(h,\tau)} + \tau e_1 \cdot \nu^{(h,\tau)} + \tau^2 X^{(h,\tau)} \cdot \nu^{(h,\tau)} \\ &= \tau H + \tau e_1 \cdot \nu + \tau(X - (1, 0, 0)) \cdot \nu = \tau(H + X \cdot \nu). \quad \square \end{aligned}$$

3.2. Fréchet differentiability. For $\tau \in (-\eta, \eta)$ and $h \in W^{2,p}(\Sigma_0, \mathbf{R})$ so that $|h(p)| < 1/8$ for $p \in \Sigma_0$, Claim 4 guarantees that the graph of h over Σ_τ is embedded. Therefore we can define the function

$$(8) \quad F(h, \tau) = H^{(h,\tau)} + \tau e_1 \cdot \nu^{(h,\tau)} + \tau^2 X^{(h,\tau)} \cdot \nu^{(h,\tau)},$$

where $e_1 = (1, 0, 0)$ is the first coordinate vector in \mathbf{R}^3 . We can now state the main theorem of this section:

Theorem 6. *Let $p > 2$. There is a neighborhood $U \subset W^{2,p}(\Sigma_0)$ of the zero section so that the map*

$$F : U \times (-\eta, \eta) \rightarrow L^p(\Sigma_0), \\ (h, \tau) \mapsto H^{(h,\tau)} + \tau e_1 \cdot \nu^{(h,\tau)} + \tau^2 X^{(h,\tau)} \cdot \nu^{(h,\tau)}$$

is continuously Fréchet differentiable.

3.3. Preliminary lemmas. Before starting the proof of Theorem 6, we need some preliminary results. The first one shows that the metric, normal vector and second fundamental form of Σ_τ are continuously Fréchet differentiable (C^1) in τ .

Lemma 7. *The map $\tau \mapsto g^{(0,\tau)}$ is C^1 from $(-\eta, \eta) \rightarrow C^4(T_{0,2})$.*

The map $\tau \mapsto g_{(0,\tau)} = (g^{(0,\tau)})^{-1}$ is C^1 from $(-\eta, \eta) \rightarrow C^4(T_{2,0})$.

The map $\tau \mapsto \nu^{(0,\tau)}$ is C^1 from $(-\eta, \eta) \rightarrow C^3(\Sigma_0, \mathbf{R}^3)$.

The map $\tau \mapsto A^{(0,\tau)}$ is C^1 from $(-\eta, \eta) \rightarrow C^3(T_{1,1})$.

Proof. The surface Σ_0 can be covered by a finite number of coordinate charts, $\Sigma_0 = \bigcup_{l=1}^L O_l$. Using a partition of unity and the fact that sums and products of C^1 functions are C^1 , it suffices to show that $\tau \mapsto g^{(0,\tau)}|_{O_l}$ is C^1 from $(-\eta, \eta)$ into $C^4(T_{0,2}|_{O_l})$ for each l . Similarly, we only need to show differentiability of the normal vector and the second fundamental form in each O_l . These tensors can be written explicitly in coordinates; the result then follows from Lemma 3. \square

The strategy for proving Theorem 6 is to write F in terms of tensors that are continuously differentiable in τ and h . We therefore need a few results about differentiability for products and contractions, as well as for the inverse of tensors.

The Sobolev embeddings are true in our context: for $p > 2$, $W^{1,p}(T_{\alpha,\beta}) \subset L^\infty(T_{\alpha,\beta})$, and there exists a constant C independent of u so that

$$\|u\|_{L^\infty(T_{\alpha,\beta})} \leq C \|u\|_{W^{1,p}(T_{\alpha,\beta})}.$$

From this estimate, we have that the product of tensors $(u, v) \rightarrow u \otimes v$ is continuously Fréchet differentiable as a map

(i) from $W^{1,p}(T_{\alpha,\beta}) \times W^{1,p}(T_{\gamma,\delta}) \rightarrow W^{1,p}(T_{\alpha+\gamma,\beta+\delta})$,

(ii) from $L^p(T_{\alpha,\beta}) \times L^\infty(T_{\gamma,\delta}) \rightarrow L^p(T_{\alpha+\gamma,\beta+\delta})$.

In addition, a computation in coordinates shows that contraction operations, which convert a tensor of type (α, β) to one of type $(\alpha - 1, \beta - 1)$, are continuously Fréchet differentiable from $W^{1,p}(T_{\alpha,\beta}) \rightarrow W^{1,p}(T_{\alpha-1,\beta-1})$.

Define $\mathcal{O}_l = \{u \in W^{1,p}(T_{1,1}), |u(p)| < l, \forall p \in \Sigma_0\}$. The set \mathcal{O}_l is open in $W^{1,p}(T_{1,1})$. We have the following lemma for inverse of tensors.

Lemma 8. *The map*

$$\begin{aligned} f : \mathcal{O}_{1/2} &\rightarrow W^{1,p}(T_{1,1}), \\ u &\mapsto f(u) = (Id - u)^{-1} - Id \end{aligned}$$

is continuously Fréchet differentiable.

Proof. Note that $(Id - u)^{-1} = \sum_{k=0}^{\infty} u^k$ since the series converges for $u \in \mathcal{O}_{1/2}$. This implies that $\|(Id - u)^{-1}\|_{L^\infty} \leq 2$ for $u \in \mathcal{O}_{1/2}$. Moreover, we have $\|\nabla f(u)\|_{L^p} = \|\nabla (Id - u)^{-1}\|_{L^p} \leq 4\|\nabla u\|_{L^p}$. Now, using the formula

$$\begin{aligned} f(u + s) - f(u) &= (Id - u - s)^{-1} - (Id - u)^{-1} \\ &= (Id - u - s)^{-1} s (Id - u)^{-1} s (Id - u)^{-1} + (Id - u)^{-1} s (Id - u)^{-1}, \end{aligned}$$

the proof is then straightforward. □

3.4. Proof of Fréchet differentiability. We are now ready to prove Theorem 6. Recall that the function is $F(h, \tau) = H^{(h,\tau)} + \tau e_1 \cdot \nu^{(h,\tau)} + \tau^2 X^{(h,\tau)} \cdot \nu^{(h,\tau)}$. The mean curvature $H^{(h,\tau)}$ is given by $H^{(h,\tau)} = g_{(h,\tau)}^{-1} a^{(h,\tau)}$ so we have to establish differentiability for the inverse of the metric, the second fundamental form, as well as the normal vector which plays an essential role in the definition of the second fundamental form.

3.4.1. Step 1: Normal vector $\nu^{(h,\tau)}$. Let p be a point in Σ_τ and let ρ be a vector tangent to Σ_0 at p . Denote by D_ρ the derivative of \mathbf{R}^3 valued functions in the direction of ρ . A vector $\tilde{\nu}^{(h,\tau)} = \nu^{(0,\tau)} + \xi$, where ξ is a tangent vector, is normal if

$$\begin{aligned} 0 &= \langle \tilde{\nu}^{(h,\tau)}, D_\rho X^{(h,\tau)} \rangle = \langle \nu^{(0,\tau)} + \xi, D_\rho X^{(0,\tau)} + D_\rho h \nu^{(0,\tau)} + h D_\rho \nu^{(0,\tau)} \rangle \\ &= D_\rho h + \langle \xi, (Id - hA^{(0,\tau)})\rho \rangle = \langle \nabla^{(0,\tau)} h + (Id - hA^{(0,\tau)})\xi, \rho \rangle, \end{aligned}$$

for any tangent vector ρ . The normal direction is therefore given by $\tilde{\nu}^{(h,\tau)} = \nu^{(0,\tau)} - (Id - hA^{(0,\tau)})^{-1} \nabla^{(0,\tau)} h$.

Note that $\nabla^{(0,\tau)} h = \nabla^{(0,0)} h g_{(0,\tau)}^{-1}$; therefore the map $(h, \tau) \mapsto \nabla^{(0,\tau)} h$ is continuously Fréchet differentiable from $W^{2,p}(\Sigma_0, \mathbf{R}) \times (-\eta, \eta)$ to $W^{1,p}(T_{1,0})$. Using the result on the second fundamental form $A^{(0,\tau)}$ from Lemma 7, we get that the map

$$(h, \tau) \mapsto hA^{(0,\tau)} \text{ is } C^1 \text{ from } W^{2,p}(\Sigma_0) \times (-\eta, \eta) \rightarrow W^{1,p}(T_{1,1}).$$

Denote by $B_\mu(0) \subset W^{2,p}(\Sigma_0)$ the ball of radius μ about the zero section. Choose $\mu > 0$ so that $hA^{(0,\tau)} \in \mathcal{O}_{1/2}$ for $h \in B_\mu(0)$. Lemma 8 implies that the map $(h, \tau) \mapsto ((Id - hA^{(0,\tau)})^{-1} - Id)\nabla h$ is C^1 from $B_\mu(0) \times (-\eta, \eta) \rightarrow W^{1,p}(T_{1,0})$. Hence,

$$(h, \tau) \mapsto \tilde{\nu}^{(h,\tau)} - \nu^{(0,\tau)} \text{ is } C^1 \text{ from } B_\mu(0) \times (-\eta, \eta) \rightarrow W^{1,p}(T_{1,0}).$$

In order to have the normal vector $\nu^{(h,\tau)}$ instead of the normal direction $\tilde{\nu}^{(h,\tau)}$, define the function $\tilde{b} := |\tilde{\nu}^{(h,\tau)}|^{-1}$ on Σ_0 and write

$$(9) \quad \nu^{(h,\tau)} - \nu^{(0,\tau)} = \tilde{b} \tilde{\nu}^{(h,\tau)} - \nu^{(0,\tau)} = \tilde{b} (\tilde{\nu}^{(h,\tau)} - \nu^{(0,\tau)}) - (\tilde{b} - 1)\nu^{(0,\tau)}.$$

Since $\tilde{\nu}^{(h,\tau)} - \nu^{(0,\tau)}$ is a tangent vector, $\tilde{b}^{-2} = |\tilde{\nu}^{(h,\tau)} - \nu^{(0,\tau)}|^2 + 1$. Hence

$$\tilde{b} - 1 = \frac{1}{\sqrt{|\tilde{\nu}^{(h,\tau)} - \nu^{(0,\tau)}|^2 + 1}} - 1.$$

Thanks to the differentiability of $\tilde{\nu}^{(h,\tau)} - \nu^{(0,\tau)}$ and the fact that products and contractions of tensors are differentiable in $W^{1,p}$, the map $|\tilde{\nu}^{(h,\tau)} - \nu^{(0,\tau)}|^2$ is continuously Fréchet differentiable from $B_\mu(0) \times (-\eta, \eta)$ to $W^{1,p}(\Sigma_0)$ or $L^\infty(\Sigma_0)$. Since, $\tilde{\nu}^{(0,\tau)} = \nu^{(0,\tau)}$, choosing $\mu > 0$ smaller if necessary, we can assume without loss of generality that $|\tilde{\nu}^{(h,\tau)} - \nu^{(0,\tau)}|^2 < 1/2$ for $h \in B_\mu(0)$ and $\tau \in (-\eta, \eta)$. The following lemma shows that $\tilde{b} - 1$ is differentiable.

Lemma 9. *Let s be a section of $T_{0,0}$; in other words s is a function from Σ to \mathbf{R} , such that $|s(p)| \leq 1/2$ for $p \in \Sigma_0$. The map $W^{1,p}(T_{0,0}) \rightarrow W^{1,p}(T_{0,0})$, $s \mapsto f \circ s := \frac{1}{\sqrt{s+1}} - 1$ is continuously Fréchet differentiable.*

Proof. Simple estimates yield that $f \circ s \in W^{1,p}(T_{0,0})$. Noting that the function $f : (-1/2, 1/2) \rightarrow \mathbf{R}$ is smooth and has uniformly bounded derivatives, we use the Mean Value Theorem to conclude the lemma. \square

Lemma 9 implies that $\tilde{b} - 1$ is continuously Fréchet differentiable from $B_\mu(0) \times (-\eta, \eta) \rightarrow W^{1,p}(\Sigma_0)$. Since Σ_0 is bounded, \tilde{b} is also C^1 . Using equation (9) and differentiability of products of tensors from $W^{1,p} \times W^{1,p}$ to $W^{1,p}$, we have that the map

$$(10) \quad (h, \tau) \mapsto \nu^{(h,\tau)} \text{ is } C^1 \text{ from } B_\mu(0) \times (-\eta, \eta) \rightarrow W^{1,p}(\Sigma_0).$$

3.4.2. *Step 2: Inverse of metric.* Let p be a point in Σ_τ and let ξ be a vector tangent to Σ_0 at p . Denote by D_ξ the derivative in the direction of ξ . Then,

$$D_\xi X^{(h,\tau)} = D_\xi(X^{(0,\tau)} + h\nu^{(0,\tau)}) = (Id - hA^{(0,\tau)})\xi + D_\xi h \nu^{(0,\tau)}$$

and

$$\begin{aligned} g^{(h,\tau)}(\rho, \xi) &= \langle D_\rho X^{(h,\tau)}, D_\xi X^{(h,\tau)} \rangle \\ &= g^{(0,\tau)}(\rho, \xi) - 2h\langle \rho, A^{(0,\tau)}\xi \rangle + h^2\langle A^{(0,\tau)}\rho, A^{(0,\tau)}\xi \rangle + D_\rho h D_\xi h. \end{aligned}$$

Define the tensors $a(\rho, \xi) = \langle \rho, A^{(0,\tau)}\xi \rangle$, $b(\rho, \xi) = \langle A^{(0,\tau)}\rho, A^{(0,\tau)}\xi \rangle$ on Σ_0 . Since a and b do not depend on h , Lemma 7 implies that ha and h^2b are continuously Fréchet differentiable from $B_\mu \times (-\eta, \eta) \rightarrow C^1(T_{0,2})$. Therefore, the map $(h, \tau) \mapsto g^{(h,\tau)} - g^{(0,\tau)}$ is C^1 from $B_\mu(0) \times (-\eta, \eta) \rightarrow W^{1,p}(T_{0,2})$. Let $u = g_{(0,\tau)}^{-1}(g^{(0,\tau)} - g^{(h,\tau)})$. Choosing μ smaller if necessary, we can apply Lemma 8 to u and get that the map

$$(Id - u)^{-1} - Id = [Id - g_{(0,\tau)}^{-1}(g^{(0,\tau)} - g^{(h,\tau)})]^{-1} - Id = (g_{(h,\tau)}^{-1} - g_{(0,\tau)}^{-1})g^{(0,\tau)}$$

is C^1 as a map from $B_\mu(0) \times (-\eta, \eta)$ to $W^{1,p}(T_{1,1})$. Finally, we have that the map $(h, \tau) \mapsto g_{(h,\tau)}^{-1} - g_{(0,\tau)}^{-1}$ is C^1 from $B_\mu(0) \times (-\eta, \eta) \rightarrow W^{1,p}(T_{2,0})$; therefore the map

$$(11) \quad (h, \tau) \mapsto g_{(h,\tau)}^{-1} \text{ is } C^1 \text{ from } B_\mu(0) \times (-\eta, \eta) \rightarrow W^{1,p}(T_{2,0}).$$

3.4.3. *Step 3: Second fundamental form.* We start with a computation giving us an equation for the second fundamental form $a^{(h,\tau)}$ in terms of known quantities. Recall $X^{(h,\tau)} = X^{(0,\tau)} + h\nu^{(0,\tau)}$. Denote by $\nabla^{(0,\tau)}$ the covariant derivative and by $(\nabla^{(0,\tau)})^2$ the second covariant derivative on Σ_0 corresponding to the metric $g^{(0,\tau)}$. For ξ, ρ tangent vectors to Σ_0 at the point p , we have

$$\begin{aligned} (\nabla^{(0,\tau)})^2 X^{(h,\tau)}(\rho, \xi) &= (\nabla^{(0,\tau)})^2 X^{(0,\tau)}(\rho, \xi) + h(\nabla^{(0,\tau)})^2 \nu^{(0,\tau)}(\rho, \xi) \\ &\quad - A^{(0,\tau)}\rho D_\xi h - A^{(0,\tau)}\xi D_\rho h + (\nabla^{(0,\tau)})^2 h(\rho, \xi)\nu^{(0,\tau)}. \end{aligned}$$

The second derivative of $\nu^{(0,\tau)}$ is given by

$$(12) \quad (\nabla^{(0,\tau)})^2 \nu^{(0,\tau)}(\rho, \xi) = -\nabla_\rho^{(0,\tau)}(A^{(0,\tau)})\xi - [\langle A^{(0,\tau)}\rho, A^{(0,\tau)}\xi \rangle] \nu^{(0,\tau)}.$$

The second fundamental form $a^{(h,\tau)}$ is given by $a^{(h,\tau)} = |\tilde{\nu}^{(h,\tau)}|^{-1} \tilde{a}^{(h,\tau)}$, where $\tilde{\nu}^{(h,\tau)} = \nu^{(0,\tau)} - (Id - hA)^{-1} \nabla^{(0,\tau)} h$ is the normal direction on $\Sigma_{h,\tau}$ and

$$\begin{aligned} \tilde{a}^{(h,\tau)}(\rho, \xi) &= |\tilde{\nu}^{(h,\tau)}| \langle \rho, A^{(h,\tau)}\xi \rangle \\ &= \langle \tilde{\nu}^{(h,\tau)}, (\nabla^{(h,\tau)})^2 X^{(h,\tau)}(\rho, \xi) \rangle \\ &= \langle \tilde{\nu}^{(h,\tau)}, (\nabla^{(0,\tau)})^2 X^{(h,\tau)}(\rho, \xi) \rangle. \end{aligned}$$

The last equality is valid since $D_\xi X^{(h,\tau)}$ is perpendicular to $\tilde{\nu}^{(h,\tau)}$. Hence,

$$(13) \quad \begin{aligned} \tilde{a}^{(h,\tau)}(\rho, \xi) &= a^{(0,\tau)}(\rho, \xi) - h \langle A^{(0,\tau)}\rho, A^{(0,\tau)}\xi \rangle + (\nabla^{(0,\tau)})^2 h(\rho, \xi) \\ &\quad + \langle (Id - hA^{(0,\tau)})^{-1} \nabla^{(0,\tau)} h, h \nabla_\rho^{(0,\tau)} A^{(0,\tau)}\xi + A^{(0,\tau)}\rho D_\xi h + A^{(0,\tau)}\xi D_\rho h \rangle. \end{aligned}$$

We know that the map $(h, \tau) \mapsto (\nabla^{(0,0)})^2 h$ is C^1 from $B_\mu(0) \times (-\eta, \eta) \rightarrow L^p(T_{0,2})$. Let us denote by $(\Gamma^{(0,\tau)})^k_{ij}$ and $(\Gamma^{(0,0)})^k_{ij}$ the components of the Christoffel symbols corresponding to the metrics $g^{(0,\tau)}$ and $g^{(0,0)}$ respectively. By Lemma 7, the $(1, 2)$ tensor given in coordinates by $(\Gamma^{(0,\tau)})^k_{ij} - (\Gamma^{(0,0)})^k_{ij}$ is C^1 from $B_\mu(0) \times (-\eta, \eta) \rightarrow C^3(T_{1,2})$. Therefore

$$\begin{aligned} &\text{the map } (h, \tau) \mapsto (\nabla^{(0,\tau)})^2 h \text{ is } C^1 \text{ from } B_\mu(0) \times (-\eta, \eta) \rightarrow L^p(T_{0,2}), \\ &\text{the map } (h, \tau) \mapsto \nabla^{(0,\tau)} A^{(0,\tau)} \text{ is } C^1 \text{ from } B_\mu(0) \times (-\eta, \eta) \rightarrow C^2(T_{1,2}). \end{aligned}$$

The other quantities in equation (13) have been discussed previously, so it follows from the results on products and contractions of tensors that the map $(h, \tau) \mapsto \tilde{a}^{(h,\tau)}$ is C^1 from $B_\mu(0) \times (-\eta, \eta) \rightarrow L^p(T_{0,2})$. Since $|\tilde{\nu}^{(h,\tau)}|^{-1}$ is a continuously Fréchet differentiable function of $(h, \tau) \in B_\mu(0) \times (-\eta, \eta)$ into $W^{1,p}(\Sigma_0)$,

$$(14) \quad \text{the map } (h, \tau) \mapsto a^{(h,\tau)} \text{ is } C^1 \text{ from } B_\mu(0) \times (-\eta, \eta) \rightarrow L^p(T_{0,2}).$$

3.4.4. *Final step.* Combining the results for the normal vector, the metric and the second fundamental form (10) (11) and (14), we have that

$$\text{the map } (h, \tau) \mapsto F(h, \tau) \text{ is } C^1 \text{ from } B_\mu(0) \times (-\eta, \eta) \rightarrow L^p(T_{0,2}).$$

4. INVERSE FUNCTION THEOREM ARGUMENT

Our main goal in this section is to prove that, loosely speaking, for small enough boundary conditions on $\partial\Sigma_0 \cap (\{x = \pm C_0\} \cup \{z = \pm C_0\})$ and small τ , there is a function h with the boundary conditions and so that $F(h, \tau) = 0$. We will impose symmetries on the boundary conditions and the function h in order to simplify the problem and to be able to reconstruct a complete surface from the piece $\Sigma_{h,\tau}$. The symmetries do not allow us to choose the boundary values on the curves $y = \pi/2 \cap \partial\Sigma_0$ or $y = 3\pi/2 \cap \partial\Sigma_0$. The following theorem states the result more precisely.

Theorem 10. *There exists a constant C_1 so that, for any $C_0 > C_1$, there are constants $\eta_0 > 0$ and $\delta_0 > 0$ such that, if we denote by Σ_0 the surface $\Sigma_0 = \{(x, y, z) \mid \sin y = \sinh x \sinh z, -C_0 < x, z < C_0, -\pi/2 < y < 3\pi/2\}$, then for any $\tau \in (-\eta_0, \eta_0)$ and function $f \in W^{2,p}(\Sigma_0)$ satisfying $\|f\|_{W^{2,p}(\Sigma_0)} \leq \delta_0$ and*

$f(x, y, z) = -f(x, -y, -z) = f(x, \pi - y, z)$, $(x, y, z) \in \Sigma_0$, there exists a function $h \in W^{2,p}(\Sigma_0)$ having the same symmetries as f and satisfying

$$F(h, \tau) = H^{(h,\tau)} + \tau e_1 \cdot \nu^{(h,\tau)} + \tau^2 X^{(h,\tau)} \cdot \nu^{(h,\tau)} = 0,$$

$$h = f \text{ on } \partial\Sigma_0 \cap (\{x = \pm C_0\} \cup \{z = \pm C_0\}).$$

Moreover, η_0 and δ_0 can be chosen small enough so that the graph of h over Σ_τ , $\tau \in (-\eta_0, \eta_0)$, is embedded.

The proof is given in Section 5. This theorem is a direct consequence of Theorem 13 below, which uses an inverse function theorem argument. Before we can state it, we need to define the following Banach spaces.

4.1. $W^{2,p}_{Sym}(\Sigma_0)$, **Sobolev space with symmetries.** To simplify our problem, we impose symmetries on the functions h . Roughly speaking, we will only consider functions h such that the graph $\Sigma_{h,0}$ of h over Σ_0 satisfies the same symmetries as Σ_0 . More precisely, $h \in W^{2,p}_{Sym}$ if $\Sigma_{h,0}$ is invariant under

- rotation of 180° with respect to the x -axis: $h(x, -y, -z) = -h(x, y, z)$, $(x, y, z) \in \Sigma_0$,
- symmetry with respect to the plane $y = \pi/2$: $h(x, y, z) = h(x, \pi - y, z)$, $(x, y, z) \in \Sigma_0$.

Let us keep in mind that the two symmetries above, considered on the whole Scherk surface S , generate all the symmetries listed in Lemma 2. Define the Sobolev space with symmetries

$$W^{2,p}_{Sym}(\Sigma_0) = \{h \in W^{2,p}(\Sigma_0) \mid h \text{ satisfies the symmetries above}\},$$

where $W^{2,p}_{Sym}(\Sigma_0)$ ($p > 2$) is a Banach space. Since $W^{2,p}_{Sym}(\Sigma_0) \subset C^1(\Sigma_0)$ it makes sense to consider derivatives of h .

4.2. **Definition of Σ_0^* .** Let us identify pairs of points on the edges $\partial\Sigma_0 \cap \{y = -\pi/2, 3\pi/2\}$ and denote by Σ_0^* the set Σ_0 , where the points $(x, -\pi/2, z)$ and $(x, 3\pi/2, z)$ are identified. The metric on Σ_0^* is induced by the metric on Σ_0 .

The second symmetry condition in the previous section, $h(x, y, z) = h(x, \pi - y, z)$, allows us to make sense of a function $h \in W^{2,p}_{Sym}$ as a function on Σ_0^* . It is clear that

$$W^{2,p}_{Sym}(\Sigma_0) \cong W^{2,p}_{Sym}(\Sigma_0^*) =: W^{2,p}_{Sym}.$$

The next section shows that Σ^* can be conformally mapped to a part of the 2-sphere.

4.3. **Going from Σ_0^* to the sphere.** Let S be the Scherk surface defined by (2). The following results about the Gauss map of the Scherk surface S are standard (see [7] or [4]).

Lemma 11. *The Gauss map ν of S has the following properties:*

- (i) ν restricts to a diffeomorphism from $S \cap \{y \in [-\pi/2, \pi/2]\}$ onto $S^2 \cap \{y \geq 0\} \setminus \{(\pm 1, 0, 0), (0, 0, \pm 1)\}$.
- (ii) Let E_i ($i = 1, \dots, 4$) be the arcs into which the equator $S^2 \cap \{y = 0\}$ is decomposed by removing the points $(\pm 1, 0, 0)$ and $(0, 0, \pm 1)$, numbered so that E_i is in the i -th quadrant of the xz -plane. We then have

$$\nu(S \cap \{y = -\pi/2\}) = E_2 \cup E_4, \quad \nu(S \cap \{y = \pi/2\}) = E_1 \cup E_3.$$

- (iii) S has no umbilics and $\nu^*g_{S^2} = \frac{1}{2}|A|^2g$.

Σ_0 is a piece of a Scherk minimal surface. Since Σ_0 does not have any umbilics, the Gauss map $\nu : \Sigma_0 \rightarrow S^2$, $p \mapsto \nu(p)$ is conformal. Moreover, we can compute the normal vector ν explicitly using equation (2) and find

$$\nu(x, y, z) = \left(\frac{\sinh z}{\cosh z}, -\frac{-\cos y}{\cosh z \cosh x}, \frac{\sinh x}{\cosh x} \right), \text{ for } (x, y, z) \in \Sigma_0.$$

The images under the Gauss map of the curves $\{z = \pm C_0\} \cap \partial \Sigma_0$, $\{x = \pm C_0\} \cap \partial \Sigma_0$ are circles on S^2 centered at $(\pm 1, 0, 0)$ and $(0, 0, \pm 1)$ respectively.

Denote by $S_{\varphi_0}^2$ the sphere S^2 minus four balls of radius φ_0 centered at $(\pm 1, 0, 0)$ and $(0, 0, \pm 1)$:

$$S_{\varphi_0}^2 = \{(x, y, z) \in S^2 \mid x, z \in (-\cos \varphi_0, \cos \varphi_0)\}.$$

The symmetries and identification made on Σ_0 to obtain Σ_0^* allow us to define a bijective conformal continuous Gauss map ν from Σ_0^* to $S_{\varphi_0}^2$, with $\varphi_0 = \cos^{-1}(\frac{\sinh C_0}{\cosh C_0})$.

The rotation about the x -axis in Σ_0 corresponds to the reflection across the yz -plane in $S^2 \subset \mathbf{R}^3$, and the reflections across the plane $y = \pi/2$ of Σ_0 corresponds to reflection across the xz -plane in S^2 . Therefore, if we define $W_{Sym}^{2,p}(S_{\varphi_0}^2) = \{h' \in W^{2,p}(S_{\varphi_0}^2) \mid h'(x, y, z) = -h'(-x, y, z) = h'(x, -y, z)\}$, where the $W^{2,p}$ norm is taken with respect to the standard metric g_{S^2} on $S_{\varphi_0}^2$, then

$$W_{Sym}^{2,p}(\Sigma_0^*) \cong W_{Sym}^{2,p}(S_{\varphi_0}^2).$$

4.4. \mathcal{T} , the space of traces of $W_{Sym}^{2,p}(\Sigma_0)$. Denote by Σ_0^* the set Σ_0 , where we have identified the points $(x, -\pi/2, z)$ and $(x, 3\pi/2, z)$. We say that

$$f, h \in W_{Sym}^{2,p} \text{ have the same trace if } f - h \in W_0^{1,p}(\Sigma_0^*) \cap W_{Sym}^{2,p},$$

where $W_0^{1,p}(\Sigma_0^*)$ is the closure in the $W^{1,p}$ norm of smooth compactly supported functions on Σ_0^* . Define the Banach space \mathcal{T} of traces of $W_{Sym}^{2,p}$ and its norm by

$$\begin{aligned} \mathcal{T} &\cong W_{Sym}^{2,p} / (W_0^{1,p}(\Sigma_0^*) \cap W_{Sym}^{2,p}), \\ \|[f]\|_{\mathcal{T}} &= \min_{u-f \in W_0^{1,p}(\Sigma_0^*) \cap W_{Sym}^{2,p}} \|u\|_{W^{2,p}}, \quad [f] \in \mathcal{T}. \end{aligned}$$

The minimum is achieved since $W^{2,p}$ is a reflexive Banach space. Note that the trace characterizes the value of a function on $x, z = \pm C_0$. The parts of the boundary of Σ_0 on $y = -\pi/2, 3\pi/2$ are subject to our symmetry conditions; we therefore do not have the freedom to impose boundary conditions on them.

Lemma 12. For $[f] \in \mathcal{T}$, there is a unique $u \in W_{Sym}^{2,p}$ so that

$$(15) \quad \begin{aligned} \Delta u &= 0 \text{ in } \Sigma_0^*, \\ u - f &\in W_0^{1,p}(\Sigma_0^*), \end{aligned}$$

where Δ is the Laplacian computed using the metric on Σ_0^* induced by $g^{(0,0)}$ on Σ_0 .

Proof. Using the continuous conformal Gauss map $\nu : \Sigma_0^* \rightarrow S_{\varphi_0}^2$, we can consider the following equivalent problem on the sphere:

$$\begin{aligned} \Delta_{g_{S^2}} v &= 0 \text{ in } S_{\varphi_0}^2, \\ v - f \circ \nu^{-1} &\in W_0^{1,p}(S_{\varphi_0}^2), \end{aligned}$$

which is known to have a unique solution $v \in W_{Sym}^{2,p}(S_{\varphi_0}^2)$. The pull back $\nu^*g_{S^2}$ of the metric on S^2 is $\nu^*g_{S^2} = \frac{1}{2}|A|^2g$, so the function $u = \nu^*v = v \circ \nu$ is a solution to (15). It is unique since v is unique. \square

4.5. Statement of the theorem. Given C_0 a large constant, denote by Σ_0 the surface $\Sigma_0 = \{(x, y, z) \mid \sin y = \sinh x \sinh z, -C_0 < x, z < C_0, -\pi/2 < y < 3\pi/2\}$. Recall that $F(h, \tau) = H^{(h,\tau)} + \tau e_1 \cdot \nu^{(h,\tau)} + \tau^2 X^{(h,\tau)} \cdot \nu^{(h,\tau)}$. Let $W_{Sym}^{2,p}(\Sigma_0)$ and L_{Sym}^p be the Sobolev spaces with symmetries defined in Section 4.1 and let \mathcal{T} be the space of traces of $W_{Sym}^{2,p}(\Sigma_0)$ functions.

Theorem 13. *There is a constant C_1 such that for all $C_0 > C_1$, there exist a constant $\eta_0 > 0$, a neighborhood $U_{Sym} \subset W_{Sym}^{2,p}(\Sigma_0)$ of the zero section and a neighborhood $V \subset L_{Sym}^p \times \mathcal{T} \times (-\eta_0, \eta_0)$ of $(0, [0], 0)$ such that the map*

$$\begin{aligned} G : U_{Sym} \times (-\eta_0, \eta_0) &\rightarrow L_{Sym}^p \times \mathcal{T} \times (-\eta_0, \eta_0), \\ (h, \tau) &\mapsto (F(h, \tau), [h], \tau), \end{aligned}$$

is a diffeomorphism from $U_{Sym} \times (-\eta_0, \eta_0)$ to V .

Note that $G(0, 0) = (0, [0], 0)$. The proof uses the Inverse Function Theorem. The map $(h, \tau) \mapsto [h]$ is continuous and linear, and therefore continuously Fréchet differentiable. Since $F(h, \tau)$ is C^1 in its variables, G is also continuously differentiable. In order to invoke the Inverse Function Theorem, we have to show that the derivative of G at $(0, 0)$ is an isomorphism. We compute the elliptic operator characterizing $DG(0, 0)$ below. In Section 4.6, we use the conformal Gauss map to transport the operator to the 2-sphere where results on the Laplacian are readily available. The image of Σ_0^* under the Gauss map is $S_{\varphi_0}^2$, a sphere minus four small balls. Loosely speaking, since this surface is close to S^2 , the eigenvalues of the Laplacian on $S_{\varphi_0}^2$ are close to its eigenvalues on S^2 . Finally, we show that the operator on $S_{\varphi_0}^2$ which is equivalent to $DG(0, 0)$ is invertible.

Let us compute $DG(0, 0)$:

$$\begin{aligned} DG(0, 0)(u, \sigma) &= DG(0, 0)(u, 0) + DG(0, 0)(0, \sigma) \\ &= (\Delta_{g^{(0,0)}}u + |A^{(0,0)}|^2u, [u], 0) + DG(0, 0)(0, \sigma). \end{aligned}$$

Recall that $F(0, \tau) = H^{(0,\tau)} + \tau e_1 \cdot \nu^{(0,\tau)} + \tau^2 X^{(0,\tau)} \cdot \nu^{(0,\tau)}$. Therefore,

$$\begin{aligned} DF(0, 0)(0, \sigma) &= \frac{d}{dt}H(0, t\sigma)|_{t=0} + \sigma e_1 \cdot \nu^{(0,0)} \\ &= D_\tau H(0, 0)\sigma + \sigma e_1 \cdot \nu^{(0,0)} \\ &= \sigma v, \end{aligned}$$

where $v \in L_{Sym}^p(\Sigma_0^*)$ (in fact, $v \in C^1(\Sigma_0)$ by Lemma 7). Hence,

$$DG(0, 0)(u, \sigma) = (\Delta u + |A|^2u + \sigma v, [u], \sigma),$$

for some function $v \in L_{Sym}^p$ independent of (u, σ) . We discuss the properties of the operator $\Delta + |A|^2$ in the next section.

4.6. **Eigenvalues and eigenfunctions of the operator $\Delta + |A|^2$.** Recall that the Gauss map ν is conformal from $\Sigma_0^* \rightarrow S_{\varphi_0}^2$ and $\nu^*g_{S^2} = \frac{1}{2}|A|^2g$. Therefore,

$$\Delta_{\nu^*g_{S^2}} + 2 = \frac{2}{|A|^2}\Delta_g + 2 = \frac{2}{|A|^2}(\Delta_g + |A|^2).$$

To show that the kernel of $\Delta_g + |A|^2$ is trivial in $W_{Sym}^{2,p}(\Sigma_0^*) \cap W_0^{1,p}(\Sigma_0^*)$, it suffices to show that problem

$$\begin{aligned} \Delta_{g_{S^2}}\zeta + 2\zeta &= 0 \text{ in } S_{\varphi_0}^2, \\ \zeta &\in W_0^{1,p}(S_{\varphi_0}^2) \cap W_{Sym}^{2,p}(S_{\varphi_0}^2) \end{aligned}$$

only admits the zero function as a solution.

Eigenvalues and eigenfunctions of the Laplace operator on the whole sphere have been well studied and are known (see [8], for example). Denote by \mathcal{R}_{xz} the reflection with respect to the xz -plane and by \mathcal{R}_{yz} the reflection with respect to the yz -plane. We are interested in functions on the sphere that are \mathcal{R}_{xz} invariant and \mathcal{R}_{yz} antivariant (see the discussion from Section 4.3). The first eigenvalues and eigenfunctions of $M_1 = S^2$ are listed in the table below.

Eigenvalue	Eigenfunction	Properties of eigenfunction
$\lambda_{1,1} = 0$	$u_{1,1}$	\mathcal{R}_{xz} invariant
$\lambda_{1,2} = 2$	$u_{1,2} = y$	\mathcal{R}_{xz} antivariant
$\lambda_{1,3} = 2$	$u_{1,3} = x$	\mathcal{R}_{xz} invariant, \mathcal{R}_{yz} antivariant
$\lambda_{1,4} = 2$	$u_{1,4} = z$	\mathcal{R}_{xz} invariant, \mathcal{R}_{yz} invariant
6	5 eigenfunctions	

We are interested in the Laplace operator on $S_{\varphi_0}^2$ and not on S^2 . Roughly speaking, since $S_{\varphi_0}^2$ is just S^2 with very small neighborhoods removed, we expect the eigenvalues and eigenfunctions of the Laplace operator for Dirichlet problems on $S_{\varphi_0}^2$ and on S^2 not to differ much. The result is stated more precisely in Appendix B of [6]. Combined with the symmetries of the Laplace operator, this implies that, for given $\varepsilon > 0$, there is a φ_0 small enough for which there are eigenfunctions $u_{2,2}, u_{2,3}$ and $u_{2,4}$ satisfying

Eigenvalue	Eigenfunction	Properties of eigenfunction
$\lambda_{2,1} = 0$	$u_{2,1}$	
$1/2 < \lambda_{2,2} < 5/2$	$u_{2,2}$	\mathcal{R}_{xz} antivariant
$1/2 < \lambda_{2,3} < 5/2$	$u_{2,3}$	\mathcal{R}_{xz} invariant, \mathcal{R}_{yz} antivariant
$1/2 < \lambda_{2,4} < 5/2$	$u_{2,4}$	\mathcal{R}_{xz} invariant, \mathcal{R}_{yz} invariant
$13/2 < \lambda_{2,k}, k \geq 5$		

and $\|u_{1,j} - u_{2,j}\|_{L^2(S_{\varphi_0}^2)} \leq \varepsilon$, for $j = 2, 3, 4$ and $0 < \varphi < \varphi_0$.

Theorem 14. *There exists a C_1 such that for all $C_0 > C_1$, the operator $\Delta_g + |A|^2 : W_{Sym}^{2,p}(\Sigma_0^*) \cap W_0^{1,p}(\Sigma_0^*) \rightarrow L_{Sym}^p(\Sigma_0^*)$ has a trivial kernel, or equivalently, there exists a φ_1 so that, for all $\varphi_0 < \varphi_1$, the operator $\Delta_{g_{S^2}} + 2 : W_{Sym}^{2,p}(S_{\varphi_0}^2) \cap W_0^{1,p}(S_{\varphi_0}^2) \rightarrow L_{Sym}^p(S_{\varphi_0}^2)$ has a trivial kernel.*

Proof. Suppose it is not true. Then there exists a sequence of real numbers $\varphi_j \rightarrow 0$, $j \in \mathbf{N}$, and a family of functions $u_j \in W_{Sym}^{2,p}(S_{\varphi_j}^2) \cap W_0^{1,p}(S_{\varphi_j}^2)$ so that

$$\Delta_{g_{S^2}} u_j + 2u_j = 0 \text{ in } S_{\varphi_j}^2, \quad \|u_j\|_{L^2(S_{\varphi_j}^2)} = 4\pi/3.$$

Denote by $u = u_{1,3}$ the \mathcal{R}_{xz} invariant \mathcal{R}_{yz} antivariant eigenfunction corresponding to the eigenvalue 2 in S^2 :

$$\Delta_{g_{S^2}} u + 2u = 0 \text{ in } S^2, \quad \|u\|_{L^2(S^2)} = 4\pi/3.$$

Note that with this normalization, $u(x, y, z) = x$ for $(x, y, z) \in S^2$. From the discussion above and [6], we have $u_j \rightarrow u$ as $j \rightarrow \infty$ in $L^2(K)$ for any K compact, $K \subset (S^2 \setminus \{(\pm 1, 0, 0), (0, 0, \pm 1)\})$.

Claim 15. Moreover,

$$u_j \rightarrow u \text{ in } C^1(K)$$

for any K compact, $K \subset (S^2 \setminus \{(\pm 1, 0, 0), (0, 0, \pm 1)\})$.

Proof of the claim. The function $u - u_j$ satisfies $\Delta_{g_{S^2}}(u - u_j) = 0$ in $S_{\varphi_j}^2$. The claim is a direct consequence of interior estimates from standard elliptic theory, Sobolev inequalities and the convergence in L^2 on compact subsets. \square

Consider

$$\begin{aligned} 0 &= \int_{S_{\varphi_j}^2} u_j(\Delta_{g_{S^2}} u + 2u) - u(\Delta_{g_{S^2}} u_j + 2u_j) \\ (16) \quad &= \int_{S_{\varphi_j}^2} u_j \Delta_{g_{S^2}} u - u \Delta_{g_{S^2}} u_j = \int_{\partial S_{\varphi_j}^2} (-u_j \nabla_{\vec{n}} u + u \nabla_{\vec{n}} u_j) ds \\ &= 2 \int_{c_3} u \nabla_{\vec{n}} u_j ds + \int_{c_1} u \nabla_{\vec{n}} u_j ds + \int_{c_2} u \nabla_{\vec{n}} u_j ds, \end{aligned}$$

where \vec{n} is the unit vector conormal to $\partial S_{\varphi_j}^2$ pointing towards $S_{\varphi_j}^2$, and c_1, c_2 and c_3 are circles of (geodesic) radius φ_j centered at $(0, 0, 1)$, $(0, 0, -1)$ and $(1, 0, 0)$ respectively. We will show that the right hand side of (16) is positive in order to obtain a contradiction. The main contribution comes from the integral over c_3 . We bound u and ∇u from below near c_3 with the help of a maximum principle (Lemma 16) and a subsolution w_j . The integrals on the curves c_1 and c_2 are negligible because the eigenfunctions u_j are close to $u(x, y, z) = x$, which is small near $(0, 0, \pm 1)$. We use the convergence in C^1 from Claim 15 to show that the integrals on c_1 and c_2 are nonnegative.

We first examine $\int_{c_3} u \nabla_{\vec{n}} u_j ds$ and show that $\nabla_{\vec{n}} u_j > 0$ on c_3 . Note that the coefficient of the linear term is 2, which has the wrong sign to apply the maximum principle directly, so we need the following lemma.

Lemma 16. *Let Ω be a bounded domain in S^2 and suppose that $u \in C^0(\bar{\Omega})$ is a supersolution of $\Delta_{g_{S^2}} + 2$, i.e.*

$$L(u) := \Delta_{g_{S^2}} u + 2u \leq 0 \text{ in } \Omega$$

in the weak sense. If there exists a positive supersolution $\zeta \in C^0(\bar{\Omega})$, then $u \geq 0$ on $\partial\Omega$ implies $u \geq 0$ in Ω .

Proof. Consider the function $v = \frac{u}{\zeta}$, and define the operator $\tilde{L}(v) = L(\zeta v)$. On the one hand, $\tilde{L}(v) = L(\zeta v) = L(u) \leq 0$; but on the other hand, $\tilde{L}(v) = \zeta \Delta v + 2\nabla \zeta \cdot \nabla v + (\Delta \zeta + 2\zeta)v$. The coefficient in front of v is now nonpositive, so we can apply the weak maximum principle to obtain

$$\inf_{\Omega} v \geq \inf_{\partial\Omega} (\min(v, 0)).$$

Since ζ is a positive function, $v \geq 0$ on $\partial\Omega$; therefore $\inf_{\Omega} u \geq 0$. □

Denote by p the point $(1, 0, 0)$, and choose spherical coordinates $\varphi \in [0, \pi]$ and $\theta \in [0, 2\pi)$,

$$\begin{cases} x &= \cos \varphi, \\ y &= \sin \varphi \cos \theta, \\ z &= \sin \varphi \sin \theta. \end{cases}$$

The following lemma gives us the supersolution needed to apply Lemma 16 in the annulus $\delta_j < \varphi < \beta$, for β chosen below.

Lemma 17. *Let β be a small number so that $-\frac{\cos \beta}{\sin \beta} + 4\beta < 0$. The function $\zeta(\varphi, \theta) = 2\beta - \varphi$ satisfies*

$$\Delta \zeta + 2\zeta \leq 0, \quad \varphi_j < \varphi < \beta;$$

in other words, it is a supersolution of $L := \Delta + 2$ in the region $\varphi_j < \varphi < \beta$.

Proof. A computation using the Laplacian in spherical coordinates shows

$$\Delta \zeta + 2\zeta = -\frac{\cos \varphi}{\sin \varphi} + 2(2\beta - \varphi) < -\frac{\cos \beta}{\sin \beta} + 4\beta < 0, \quad \varphi_j < \varphi < \beta,$$

since $\cos \varphi > \cos \beta$ and $(\sin \varphi)^{-1} > (\sin \beta)^{-1}$ for $\varphi_j < \varphi < \beta$. □

Define $\eta_0 = 1 - \cos \beta$. The eigenfunction u is given by $u = \cos \varphi$, so $|u(\varphi, \theta) - 1| \leq \eta_0, \varphi < \beta, 0 \leq \theta < 2\pi$. From the convergence of u_j , there is a J so that $|u_j(\beta, \theta) - 1| < 2\eta_0, 0 \leq \theta < 2\pi, j > J$. For $j > J$, we have

$$\begin{aligned} L(u_j) &= \Delta u_j + 2u_j = 0, \quad \delta_j < \varphi < \beta, \\ 1 + 2\eta_0 &> u_j > 1 - 2\eta_0, \quad \varphi = \beta, \\ u_j &= 0, \quad \varphi = \varphi_j. \end{aligned}$$

We introduce an auxiliary function $w_j = \varphi - \varphi_j$. Since $L(w_j) \geq 0$, we have $L(u_j - \varepsilon w_j) \leq 0$ for $\varphi_j < \varphi < \beta$ and $u_j - \varepsilon w_j = 0$ on $\varphi = \varphi_j$. Let $\varepsilon = \frac{1-2\eta_0}{\beta-\varphi_j}$; then

$$u_j - \varepsilon w_j > 1 - 2\eta_0 - \frac{1 - 2\eta_0}{\beta - \varphi_j}(\beta - \varphi_j) = 0 \text{ on } \varphi = \beta.$$

The conditions of Lemma 16 are satisfied for $u = u_j - \varepsilon w_j$ and $\Omega = \{(\varphi, \theta) | \varphi_j < \varphi < \beta, 0 \leq \theta < 2\pi\}$. Hence, $u_j \geq \varepsilon w_j, \varphi_j < \varphi < \beta$. Since $u_j = \varepsilon w_j = 0$ on $\varphi = \varphi_j$ (which is the circle c_3) we have

$$\nabla_{\vec{n}} u_j \geq \varepsilon \nabla_{\vec{n}} w_j = \varepsilon > 0 \text{ on } \varphi = \varphi_j,$$

where \vec{n} is the inward unit vector normal to c_3 . Therefore,

$$\begin{aligned} \int_{c_3} u \nabla_{\vec{n}} u_j ds &\geq \int_{c_3} \cos \varphi_j \frac{1 - 2\eta_0}{\beta - \varphi_j} ds > 4 \cos \varphi_j \frac{1 - 2\eta_0}{\beta} \varphi_j \\ (17) \qquad \qquad &> 4 \cos \beta \left(\frac{2 \cos \beta - 1}{\beta} \right) \varphi_j. \end{aligned}$$

Let us now consider $\int_{c_1} u \nabla_{\bar{n}} u_j ds$ and $\int_{c_2} u \nabla_{\bar{n}} u_j ds$. We will show that $\nabla_{\bar{n}} u_j \geq 0$ on $c_1 \cap \{x \geq 0\}$ and $c_2 \cap \{x \geq 0\}$. In this case also, choose spherical coordinates $\varphi \in [0, \pi], \theta \in [0, 2\pi)$,

$$\begin{cases} x &= \sin \varphi \cos \theta, \\ y &= \sin \varphi \sin \theta, \\ z &= \cos \varphi, \end{cases}$$

and let ζ and β be as in Lemma 17. By definition and symmetries, we know that for all j ,

$$u_j = 0 \text{ on } \{\varphi = \varphi_j\} \cup \{\theta = \pm\pi/2\}.$$

Let $K \subset S^2 \setminus \{(\pm 1, 0, 0), (0, 0, \pm 1)\}$ be a compact set containing $\varphi = \beta$. Since $u_j \rightarrow u$ in $C^1(K)$, for $\mu > 0$, there is a J so that

$$u - \mu \cos \theta \leq u_j \leq u + \mu \cos \theta \text{ for } \theta \in (-\pi/2, \pi/2), \varphi = \beta, j > J.$$

Replacing u by its value and choosing $\mu < \sin \beta$, we get

$$0 \leq u_j \leq (\sin \beta + \mu) \cos \theta, \text{ for } \theta \in (-\pi/2, \pi/2), \varphi = \beta, j > J.$$

ζ is a supersolution of $\Delta + 2$ in this case also; hence Lemma 16 applied to u_j in the domain $\varphi_j < \varphi < \beta, -\pi/2 < \theta < \pi/2$ gives $u \geq 0$ in this domain. Since $u_j = 0$ on $\varphi = \varphi_j$,

$$\nabla_{\bar{n}} u_j \geq 0, \quad \varphi = \varphi_j, \theta \in (-\pi/2, \pi/2).$$

The function $u = \sin \varphi \cos \theta$ is also nonnegative on $c_1 \cap \{\theta \in (-\pi/2, \pi/2)\}$; therefore

$$\int_{c_1 \cap \{-\pi/2 < \theta < \pi/2\}} u \nabla_{\bar{n}} u_j ds \geq 0.$$

Using symmetries of u and u_j , we have

$$(18) \quad \int_{c_1} u \nabla_{\bar{n}} u_j ds = 2 \int_{c_1 \cap \{-\pi/2 < \theta < \pi/2\}} u \nabla_{\bar{n}} u_j ds \geq 0.$$

The integral over c_2 is treated in an analogous way, and

$$(19) \quad \int_{c_2} u \nabla_{\bar{n}} u_j ds \geq 0.$$

Equations (16), (17), (18) and (19) lead to a contradiction. Therefore, there exists a φ_1 so that, for all $\varphi_0 < \varphi_1$, the operator $\Delta_{g_{S^2}} + 2 : W_{Sym}^{2,p}(S_{\varphi_0}^2) \cap W^{1,p}(S_{\varphi_0}^2) \rightarrow L^p(S_{\varphi_0}^2)$ has a trivial kernel. \square

4.7. Proof of Theorem 13. Recall that

$$\begin{aligned} DG(0, 0) : W_{Sym}^{2,p}(\Sigma_0^*) \times \mathbf{R} &\rightarrow L_{Sym}^p(\Sigma_0^*) \times \mathcal{T} \times \mathbf{R}, \\ (u, \sigma) &\mapsto (\Delta u + |A|^2 u + \sigma v, [u], \sigma). \end{aligned}$$

In what follows, all the operators and maps are considered from $W_{Sym}^{2,p}(\Sigma_0^*) \cap W_0^{1,p}(\Sigma_0^*) \rightarrow L_{Sym}^p(\Sigma_0^*)$, unless mentioned otherwise.

The manifold Σ_0^* is bounded and smooth, so the map $u \mapsto |A|^2 u$ is compact. The operator Δ_g is Fredholm of index 0 since $\Delta_g = \frac{|A|^2}{2} \Delta_{\nu^* g_{S^2}}$ and since the Laplace operator on the sphere is invertible from $W_{Sym}^{2,p}(S_{\varphi_0}^2) \cap W_0^{1,p}(S_{\varphi_0}^2)$ to $L_{Sym}^p(S_{\varphi_0}^2)$ for $\varphi_0 \leq 1$. Thus, $\Delta_g + |A|^2$ is Fredholm of index 0.

Let C_1 be given by Theorem 14. For a choice of $C_0 > C_1$, Theorem 14 and the above discussion imply that $\Delta_g + |A|^2$ is invertible. Hence, for $v \in L_{Sym}^p(\Sigma_0^*)$ and

$\sigma \in \mathbf{R}$, $u \mapsto \Delta_g u + |A|^2 u + \sigma v$ is invertible from $W_{Sym}^{2,p}(\Sigma_0^*) \cap W_0^{1,p}(\Sigma_0^*) \rightarrow L^p_{Sym}(\Sigma_0^*)$. We derive from this that the derivative $DG(0, 0) : W_{Sym}^{2,p}(\Sigma_0^*) \times \mathbf{R} \rightarrow L^p_{Sym}(\Sigma_0^*) \times \mathcal{T} \times \mathbf{R}$ is an isomorphism.

We can apply the inverse function theorem for Banach spaces to obtain the existence of a constant $\eta_0 > 0$, a neighborhood $U_{Sym} \subset W_{Sym}^{2,p}(\Sigma_0)$ and a neighborhood $V \subset L^p_{Sym}(\Sigma_0) \times \mathcal{T} \times (-\eta_0, \eta_0)$ such that the map $G : U_{Sym} \times (-\eta_0, \eta_0) \rightarrow V$ is a diffeomorphism. □

5. COROLLARIES

Theorem 10 is an immediate corollary of Theorem 13. Let us recall it:

There exists a constant C_1 so that, for any $C_0 > C_1$, there are constants $\eta_0 > 0$ and $\delta_0 > 0$ such that for any $\tau \in (-\eta_0, \eta_0)$ and function $f \in W_{Sym}^{2,p}(\Sigma_0)$ satisfying $\|f\|_{W^{2,p}(\Sigma_0)} \leq \delta_0$, there exists a function $h \in W_{Sym}^{2,p}(\Sigma_0)$ satisfying

$$\begin{aligned} F(h, \tau) &= H^{(h,\tau)} + \tau e_1 \cdot \nu^{(h,\tau)} + \tau^2 X^{(h,\tau)} \cdot \nu^{(h,\tau)} = 0, \\ h &= f \text{ on } \partial\Sigma_0 \cap (\{x = \pm C_0\} \cup \{z = \pm C_0\}). \end{aligned}$$

Moreover, η_0 and δ_0 can be chosen small enough so that $\Sigma_{h,\tau}$, the graph of h over Σ_τ , $\tau \in (-\eta_0, \eta_0)$, is embedded.

We wish to go back to the smaller scale, so that the surfaces considered are asymptotic to cylinders of radius 1 instead of $1/\tau$. A few definitions are needed to state the next corollary.

For $N \in \mathbf{N}$ and a large constant $C > 0$, denote by Ω_C^N the domain

$$\Omega_C^N = \{(x, y, z) \in \mathbf{R}^3 \mid |x| < C, \frac{-3N\pi}{2} < y < \frac{N\pi}{2}, |z| < C\},$$

and by S_C^N the piece of Scherk surface $S_C^N = S \cap \Omega_C^N$. For $\tau \in (-\eta, \eta)$, let Φ_τ and \mathcal{H}_τ be the bending maps and the scaling maps defined by equations (4) and (6) respectively. The surface $\tilde{\Sigma}_{1/N}^C = \mathcal{H}_{1/N}(\Phi_{1/N}(S_C^N))$ is a smooth surface with N handles. For x close to $\pm C$, $\tilde{\Sigma}_{1/N}^C$ is close to a plane; for z close to $\pm C$, $\tilde{\Sigma}_{1/N}^C$ is close to a cylinder of radius 1. Moreover, $\tilde{\Sigma}_{1/N}^C$ has the following symmetries:

- it is invariant with respect to the rotation of 180 degrees about the x -axis and
- it is invariant with respect to symmetries across the planes $\theta = \pi/2N + k\pi/N$, $k = 1, \dots, 2N$, where we used cylindrical coordinates.

Theorem 1. *There is a constant C_1 so that, for any $C > C_1$, there exist an integer $N_0 > 0$ and a constant $\delta_0 > 0$ with the following properties:*

for every integer $N > N_0$ and for every function $\tilde{f} \in W^{2,p}(\tilde{\Sigma}_{1/N}^C)$ with $\|\tilde{f}\|_{W^{2,p}(\tilde{\Sigma}_{1/N}^C)} \leq \delta_0/(2N^2)$ and whose graph over $\tilde{\Sigma}_{1/N}^C$ satisfies the above symmetries, there exists a function $\tilde{h} \in W^{2,p}(\tilde{\Sigma}_{1/N}^C)$ such that

$$\begin{aligned} &\text{the graph of } \tilde{h} \text{ over } \tilde{\Sigma}_{1/N}^C \text{ satisfies } H + X \cdot \nu = 0, \\ &\text{the graph of } \tilde{h} \text{ over } \tilde{\Sigma}_{1/N}^C \text{ enjoys the above symmetries,} \\ &\tilde{h} = \tilde{f} \text{ on } \partial\tilde{\Sigma}_{1/N}^C. \end{aligned}$$

Moreover, we can choose N_0 and δ_0 so that the graph of h over $\tilde{\Sigma}_{1/N}^C$ is an embedded surface.

Proof. Lemma 5 relates the equations $F(h, \tau) = 0$ and $H + X \cdot \nu = 0$, and the invariance with respect to the planes $\theta = \pi/2N + k\pi/N$ lets us reconstruct a complete surface from the piece $\mathcal{H}_{1/N}(\Sigma_{h,1/N})$.

Fix $C > C_1$. From now on, we will not mention the dependence in C and simply write $\tilde{\Sigma}_{1/N}$ for $\tilde{\Sigma}_{1/N}^C$.

The surface $\tilde{\Sigma}_{1/N}$ contains N pieces of surface identical to $\mathcal{H}_{1/N}(\Sigma_{1/N})$. From the symmetries of f , we have $\|\tilde{f}\|_{W^{2,p}(H_{1/N}(\Sigma_{1/N}))} \leq \delta_0/(2N^3)$. Therefore $f = N\tilde{f} \circ \mathcal{H}_{1/N}$ considered as a function on $\Sigma_{1/N}$ satisfies

$$\|f\|_{W^{2,p}(\Sigma_{1/N})} \leq \delta_0/2.$$

Moreover, f enjoys the symmetries of Theorem 10. The metrics $g_\tau, \tau \in (-\eta, \eta)$, induce equivalent norms on the manifold Σ_0 since the bending map Φ from Lemma 3 is uniformly continuous from $(-\eta, \eta) \rightarrow C^5(\bar{\Omega}, \mathbf{R}^3)$. Therefore, we can choose η small enough so that

$$\|f\|_{W^{2,p}(\Sigma_0)} \leq 2\|f\|_{W^{2,p}(\Sigma_\tau)}, \quad \tau \in (-\eta, \eta).$$

For $N > N_0 > \max(\eta, \eta_0^{-1})$ with η_0 and δ_0 as in Theorem 10, there exists a function h so that $F(h, 1/N) = 0$ and $\Sigma_{h,1/N}$ is embedded. Lemma 5 implies that $\mathcal{H}_{1/N}(\Sigma_{h,1/N})$ satisfies the self-similar equation (1). We can define \tilde{h} on $\mathcal{H}_{1/N}(\Sigma_{1/N})$ by $\tilde{h} = N^{-1}h \circ (\mathcal{H}_{1/N})^{-1}$ so that the graph of \tilde{h} over $\mathcal{H}_{1/N}(\Sigma_{1/N})$ coincides with $\mathcal{H}_{1/N}(\Sigma_{h,1/N})$. We then use invariance with respect to reflections across the planes $\theta = \pi/2N + k\pi/N$ to extend the function \tilde{h} to the whole surface $\tilde{\Sigma}_{1/N}$. \square

REFERENCES

1. S. B. ANGENENT, *Shrinking doughnuts*, in Nonlinear diffusion equations and their equilibrium states, 3 (Gregynog, 1989), vol. 7 of Progr. Nonlinear Differential Equations Appl., Birkhäuser Boston, Boston, MA, 1992, pp. 21–38. MR1167827 (93d:58032)
2. S. B. ANGENENT, D. L. CHOPP, AND T. ILMANEN, *A computed example of nonuniqueness of mean curvature flow in \mathbf{R}^3* , Comm. Partial Differential Equations, 20 (1995), pp. 1937–1958. MR1361726 (96k:58045)
3. D. L. CHOPP, *Computation of self-similar solutions for mean curvature flow*, Experiment. Math., 3 (1994), pp. 1–15. MR1302814 (95j:53006)
4. U. DIERKES, S. HILDEBRANDT, A. KÜSTER, AND O. WOHLRAB, *Minimal surfaces. I*, vol. 295 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], Springer-Verlag, Berlin, 1992. Boundary value problems. MR1215267 (94c:49001a)
5. G. HUISKEN, *Asymptotic behavior for singularities of the mean curvature flow*, J. Differential Geom., 31 (1990), pp. 285–299. MR1030675 (90m:53016)
6. N. KAPOULEAS, *Complete constant mean curvature surfaces in Euclidean three-space*, Ann. of Math. (2), 131 (1990), pp. 239–330. MR1043269 (93a:53007a)
7. N. KAPOULEAS, *Complete embedded minimal surfaces of finite total curvature*, J. Differential Geom., 47 (1997), pp. 95–169. MR1601434 (99a:53008)
8. S. MONTIEL AND A. ROS, *Schrödinger operators associated to a holomorphic map*, in Global differential geometry and global analysis (Berlin, 1990), vol. 1481 of Lecture Notes in Math., Springer, Berlin, 1991, pp. 147–174. MR1178529 (93k:58053)
9. X. H. NGUYEN, *Construction of complete embedded self-similar surfaces under mean curvature flow. Part II*, preprint.
10. ———, *Construction of complete embedded self-similar surfaces under mean curvature flow. Part III*, in preparation.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WISCONSIN 53706

Current address: Department of Mathematics, Kansas State University, 138 Cardwell Hall, Manhattan, Kansas 66506