

GEOMETRY OF LIMIT SETS FOR EXPANSIVE MARKOV SYSTEMS

KATRIN GELFERT AND MICHAL RAMS

ABSTRACT. We describe the geometric and dynamical properties of expansive Markov systems.

1. INTRODUCTION

Consider a smooth map of an interval into itself. If the map is uniformly expanding, then its geometry and dynamical properties are well understood (see [12] for a contemporary view and references). The case that there exist attracting periodic points simply leads to the appearance of basins of attraction with a rather simple dynamics.

The situation which lies in between those two cases is yet of considerable interest. In the field of dynamical systems, a certain class of such interval maps has been studied by Urbański [16]. His results include, in particular, the calculation of the Hausdorff and packing dimensions and Lebesgue, Hausdorff, and packing measures of the limit set. Maps of this type appear also in the study of the so-called Markov exceptional minimal sets in foliation theory (see for example [3], [10], [1]; see [9] or [19] for extensive surveys of the field).

The goal of this paper is to describe the geometric properties of the general class of systems studied by Cantwell and Conlon [3]. It includes maps that may have marginally stable periodic orbits and hence are not (uniformly) expanding. As our main working assumption we require the map to be expansive (though not necessarily everywhere expanding). We describe its geometric properties such as the Hausdorff dimension of the limit set in terms of dynamical quantities such as entropy, topological pressure, and conformal measures (see Section 2 for the definition and notation). The following is our main result.

Received by the editors May 7, 2007.

2000 *Mathematics Subject Classification.* Primary 37E05, 28A78, 57R30.

Key words and phrases. Iterated function systems, exceptional minimal sets, thermodynamical formalism, non-uniformly hyperbolic systems, Hausdorff dimension.

The first author's research was supported by Grant EU FP6 ToK SPADE2.

The second author was supported by Grant EU FP6 ToK SPADE2 and by the Polish KBN Grant No 2P0 3A 034 25. Both authors are grateful for discussions with J. Rodriguez-Hertz that led us to Corollary 3.

Theorem 1. *Let Λ_Q be the limit set of a $C^{1+\text{Lip}}$ expansive and transitive Markov system (Λ_Q, f) . Then the Hausdorff dimension of Λ_Q satisfies*

$$\begin{aligned} \dim_{\text{H}} \Lambda_Q &= \inf\{t \geq 0: P(-t \log|f'|) = 0\} \\ &= \inf\{t \geq 0: \text{there exists a } t\text{-conformal measure supported on } \Lambda_Q\} \\ &= \sup\{\dim_{\text{H}} \mu: \mu \in \mathcal{M}_{\text{E}}, h_{\mu}(f) > 0\}. \end{aligned}$$

Moreover, Λ_Q is of zero Lebesgue measure.

Under the additional assumption that f is (piecewise) real analytic, we have $\dim_{\text{H}} \Lambda_Q < 1$ and the packing measure in this dimension is positive.

Under the assumption of Theorem 1, the zero Lebesgue measure property has been proved by Cantwell and Conlon [3]. The analogous dimension formulae of Theorem 1 have been shown in the $C^{1+\varepsilon}$ case by Urbański [16] under additional geometric assumptions. We note that we require that f be of class $C^{1+\text{Lip}}$ because our main tool for the proof of Theorem 1 is the A. Schwartz lemma. The additional analyticity assumption is necessary, as the following example shows.

Example 1. There exists a C^{∞} parabolic iterated function system in \mathbb{R} with a limit set with Hausdorff dimension equal to 1, but of zero Lebesgue measure. (Full details are given at the end of Section 5.)




FIGURE 1. The iterated function system given by (32).

Note that there exists an iterated function system in \mathbb{R} of smoothness C^1 with positive Lebesgue measure (see the example by Bowen [2]).

In Section 3 we introduce the main concept for our approach, so-called hyperbolic instants. We introduce the set H of points which have infinitely many hyperbolic instants and we prove some of its immediate properties. The conclusions in this section are quite general and, in fact, are valid for a much larger class of dynamical systems than those considered in this paper. For expansive and transitive Markov systems, most points belong to the set H . Namely we have the following theorem.

Theorem 2. *Under the assumptions of Theorem 1, a point belongs to $\Lambda_Q \setminus H$ if and only if it is either a parabolic periodic point or a parabolic preperiodic point.*

We now sketch the contents of the paper. In Section 2 we introduce the main objects of our study, namely Markov systems, topological pressure, and conformal

measures. In Section 3 we define the set H and prove some of its main properties. In Section 4 we prove Theorem 2. The proof is based on ideas from Cantwell and Conlon [3] and Matsumoto [10]. Finally, in Section 5 we prove Theorem 1. For this we use Theorem 2 together with now-standard methods from [16, 5].

2. PRELIMINARIES

We consider a general geometric construction of a Cantor-like set, modelled on a Markov geometric construction (a subshift of finite type).

2.1. Iterated function systems. Let $I \subset \mathbb{R}$ be a closed interval and I_1, \dots, I_p be a finite collection of pairwise disjoint closed subintervals of I . Let $\{g_i\}_{i=1}^p$ be a collection of bijective $C^{1+\text{Lip}}$ diffeomorphisms with range $R(g_i) \subset \text{int}(I_i)$ and domain $D(g_i) = \bigcup_{j=1}^p I_j$. Such a collection is called an *iterated function system*. We will denote

$$g_{i_1 \dots i_n} \stackrel{\text{def}}{=} g_{i_1} \circ \dots \circ g_{i_n}.$$

We consider the associated topological Markov chain $\sigma: \Sigma \rightarrow \Sigma$ defined by $\sigma(i_1 i_2 \dots) = (i_2 i_3 \dots)$ on the set

$$\Sigma \stackrel{\text{def}}{=} \{1, \dots, p\}^{\mathbb{N}}.$$

We denote $\Sigma_n = \{1, \dots, p\}^n$ and $\Sigma_* = \bigcup_{n=0}^{\infty} \Sigma_n$, where we use the convention $\Sigma_0 = \{\emptyset\}$. For each $(i_1 \dots i_n) \in \Sigma_n$ we define

$$\Delta_{i_1 \dots i_n} \stackrel{\text{def}}{=} g_{i_1 \dots i_{n-1}}(I_{i_n})$$

and $\Delta_{\emptyset} = I$.

We define

$$\Lambda \stackrel{\text{def}}{=} \bigcap_{n=1}^{\infty} \bigcup_{(i_1 \dots i_n) \in \Sigma_n} \Delta_{i_1 \dots i_n}.$$

We set

$$f(x) \stackrel{\text{def}}{=} g_i^{-1}(x) \quad \text{when } x \in R(g_i).$$

Given $x \in \Lambda$, we denote by $\Delta_n(x)$ the unique set $\Delta_{i_1 \dots i_n}$ containing x .

Given an iterated function system, for each $(i_1 i_2 \dots) \in \Sigma$ the sets $\Delta_{i_1 \dots i_n}$ form a descending sequence of non-empty compact sets, and in general the set $\bigcap_{n=1}^{\infty} \Delta_{i_1 \dots i_n}$ is either a singleton or a non-degenerate interval. We always assume that $\bigcap_{n=1}^{\infty} \Delta_{i_1 \dots i_n}$ is a singleton, that is, that

$$(1) \quad d_n \stackrel{\text{def}}{=} \max_{(i_1 \dots i_n)} |\Delta_{i_1 \dots i_n}| \rightarrow 0 \text{ as } n \rightarrow \infty,$$

and in particular that $f|_{\Lambda}$ is an expansive map. Recall that $f|_{\Lambda}$ is called *expansive* if there exists a positive constant δ such that, for any $x, y \in \Lambda$ satisfying $|f^k(x) - f^k(y)| \leq \delta$ for every $k \geq 1$, we have $x = y$. We obtain a coding map $\pi: \Sigma \rightarrow \Lambda$ given by

$$\pi(i_1 i_2 \dots) = \bigcap_{n=1}^{\infty} g_{i_1} \circ \dots \circ g_{i_{n-1}}(I_{i_n}) = \bigcap_{n=1}^{\infty} \Delta_{i_1 \dots i_n}.$$

We call the sequence $\pi^{-1}(x)$ the *symbolic expansion* of $x \in \Lambda$.

We remark that for an iterated function system for which the maps g_1, \dots, g_p are not uniformly contracting on Λ , the thermodynamic formalism of the associated symbolic dynamics does not apply since the coding map π does not in general preserve the Hölder continuity of potentials.

We further remark that it would be enough to assume only that $g_i(\bigcup I_j) \subset I_i$. For such systems one can increase all I_i and expand domains of g_i in such a way as to obtain $C^{1+\text{Lip}}$ (but not necessarily C^2 , even if the original system were smoother!) expansive iterated function system (\hat{I}_i, \hat{g}_i) with the same Λ and satisfying $\hat{g}_i(\bigcup \hat{I}_j) \subset \text{int } \hat{I}_i$.

2.2. Markov systems. We call an n -tuple $(i_1 \dots i_n) \in \Sigma_n$ a *word of length n* . Let $Q \subset \Sigma_*$ be a finite set and denote by $l(Q)$ the maximal length of words from Q . We denote by Σ_Q the set of all infinite sequences from Σ that do not contain any of the words from Q as a subword:

$$\Sigma_Q \stackrel{\text{def}}{=} \{(i_1 i_2 \dots) \in \Sigma : (i_n i_{n+1} \dots i_{n+k-1}) \notin Q \text{ for every } n, k \in \mathbb{N}\}.$$

Naturally, the same set Σ_Q may be obtained for different choices of Q . We will denote

$$l(\Sigma_Q) \stackrel{\text{def}}{=} \min l(Q),$$

where the minimum is taken over all possible choices of Q resulting in the same Σ_Q . We call the (not necessarily unique) set Q for which the minimum is achieved a *representation* of Σ_Q . The proof of the following lemma is left to the reader.

Lemma 1. *Assume Q is a representation of Σ_Q . For every $Q' \supset Q$ satisfying $l(Q') = l(Q)$ we have*

$$l(\Sigma_{Q'}) \leq l(\Sigma_Q).$$

From the definition of Σ_Q it follows that $\sigma(\Sigma_Q) \subset \Sigma_Q$, but this inclusion need not be an equality. We say that Σ_Q is *completely invariant* if and only if $\sigma(\Sigma_Q) = \Sigma_Q$, and we call $\sigma|_{\Sigma_Q}$ a *subshift of finite type* in this case. Given $(j_1 \dots j_m) \in \Sigma_*$, we denote by $j_1 \dots j_m \Sigma_Q$ the set of all infinite sequences from Σ given by

$$j_1 \dots j_m \Sigma_Q \stackrel{\text{def}}{=} \{(j_1 \dots j_m i_1 i_2 \dots) : (i_1 i_2 \dots) \in \Sigma_Q\}.$$

Proposition 1. *For any Σ_Q there exists a completely invariant set $\Sigma_{Q'}$ with $l(\Sigma_{Q'}) \leq l(\Sigma_Q)$, and there exists a finite set $\tilde{Q} \subset \Sigma_*$ such that*

$$\Sigma_{Q'} \subset \Sigma_Q \subset \bigcup_{(j_1 j_2 \dots j_m) \in \tilde{Q}} j_1 j_2 \dots j_m \Sigma_{Q'}.$$

Proof. Let Σ_Q (with representation Q) be not completely invariant, that is, let us assume that there exists such a word $(i_1 i_2 \dots) \in \Sigma_Q$ that $(i i_1 i_2 \dots) \notin \Sigma_Q$ for all $i \in \{1, \dots, p\}$. Denoting $n = l(Q)$, we see that all the words $(i i_1 i_2 \dots i_{n-1})$, $i \in \{1, \dots, p\}$, must contain a word from Q as a subword. At the same time, $(i_1 i_2 \dots i_{n-1})$ does not contain any word from Q as a subword. Let $Q' = Q \cup \{(i_1 i_2 \dots i_{n-1})\}$. Note that we have $l(Q') = l(Q)$ and

$$\Sigma_{Q'} \subset \Sigma_Q \subset \Sigma_{Q'} \cup i_1 \Sigma_{Q'}.$$

Moreover, by Lemma 1,

$$l(\Sigma_{Q'}) \leq l(\Sigma_Q).$$

If $\Sigma_{Q'}$ is not completely invariant, then we can repeat this procedure. As there exist only finitely many choices of finite sets $Q'' \subset \Sigma_*$ satisfying $l(Q'') \leq n$, we can repeat this procedure only finitely many times. This proves the proposition. \square

Denote

$$(2) \quad \Lambda_Q \stackrel{\text{def}}{=} \pi(\Sigma_Q).$$

We have $f(\Lambda_Q) \subset \Lambda_Q$ (with equality if and only if Σ_Q is completely invariant). Given Q and $x \in \Lambda_Q$, we call the sequence $(i_1 i_2 \dots) \in \Sigma$ *admissible* for x if

$$(3) \quad g_{i_n \dots i_1}(x) \in \Lambda_Q \quad \text{for all } n \in \mathbb{N}.$$

If Σ_Q is completely invariant, then for every point $x \in \Lambda_Q$ there exists an admissible sequence. The proof of the following lemma will be left to the reader.

Lemma 2. *If a sequence is admissible for x , then it is admissible for every $y \in \Delta_n(x) \cap \Lambda_Q$ with $n = l(\Sigma_Q) - 1$.*

The system $(\Lambda_Q, f, g_1, \dots, g_p)$ is called a *Markov system*. The partition $\Lambda_Q = \bigcup_{k=1}^p J_k$, where $J_k = I_k \cap \Lambda_Q$, has the property that $f(J_k)$ either contains J_ℓ or is disjoint from J_ℓ for every k, ℓ . A partition with this property is called *Markov*. We note that, in the context of dynamical systems theory, one usually considers Markov systems which are topologically conjugate to a subshift of finite type, whereas here we also want to include the more general case, which in particular will be part of our constructions in Section 4.

From now on we will always assume that Σ_Q is completely invariant.

The Markov system need not, in general, be transitive. However, if the system is not transitive, then it is a disjoint union of finitely many transitive Markov systems and we can study each of them separately. In what follows, we consider only transitive Markov systems.

2.3. Topological pressure. Let φ be a continuous function on Λ_Q . The *topological pressure* of φ (with respect to $f|_{\Lambda_Q}$) is defined by

$$(4) \quad P(\varphi) \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \log \sum_{(i_1 \dots i_n)} \exp \max_{x \in \Delta_{i_1 \dots i_n}} S_n \varphi(x),$$

where here and in the sequel the sum is taken over the cylinders with non-empty intersection with the set Λ_Q . The existence of the limit follows easily from the fact that the sum constitutes a submultiplicative sequence. Moreover, the value $P(\varphi)$ does not depend on the particular Markov partition that we use in its definition.

Denote by \mathcal{M} the family of f -invariant Borel probability measures on Λ_Q . By the variational principle we have

$$(5) \quad P(\varphi) = \max_{\mu \in \mathcal{M}} \left(h_\mu(f) + \int_{\Lambda_Q} \varphi d\mu \right),$$

where $h_\mu(f)$ denotes the entropy of f with respect to μ (see [18]).

Lemma 3. *f has tempered distortion on Λ_Q ; that is, there exists a positive sequence $(\rho_n)_n$ decreasing to 0 such that for every n we have*

$$(6) \quad \sup_{(i_1 \dots i_n)} \sup_{x, y \in \Delta_{i_1 \dots i_n}} \frac{|(f^n)'(x)|}{|(f^n)'(y)|} \leq e^{n\rho_n}.$$

Proof. By Lipschitz continuity of $f|_{\Lambda_Q}$ we have

$$\log \sup_{(i_1 \dots i_n)} \sup_{x, y \in \Delta_{i_1 \dots i_n}} \frac{|(f^n)'(x)|}{|(f^n)'(y)|} \leq c \sum_{k=1}^n d_k,$$

for some positive number c and for d_k defined in (1). By the condition (1), the sequence $(d_k)_k$ is decreasing to 0, which implies the claimed property. \square

Given $t \in \mathbb{R}$, we define the function $\varphi_t: \Lambda_Q \rightarrow \mathbb{R}$ by

$$(7) \quad \varphi_t(x) \stackrel{\text{def}}{=} -t \log|f'(x)|.$$

The tempered variation property (6) ensures in particular that in the definition of $P(\varphi_t)$ in (4) one can replace the maximum by the minimum or, in fact, by any intermediate value. Given $x \in \Lambda_Q$ we denote by $\chi(x)$ the *Lyapunov exponent* at x ,

$$\chi(x) \stackrel{\text{def}}{=} \limsup_{n \rightarrow \infty} \frac{1}{n} \log|(f^n)'(x)|.$$

Proposition 2. *The function $t \mapsto P(\varphi_t)$ is a continuous, convex, and non-increasing function of \mathbb{R} . $P(\varphi_t)$ is negative for large t if and only if there exist no f -invariant probability measures with zero Lyapunov exponent.*

Proof. The claimed properties follow immediately from general facts about the pressure together with the variational principle (5). \square

We define

$$t_0 \stackrel{\text{def}}{=} \inf\{t > 0: P(\varphi_t) \leq 0\}.$$

Proposition 3. *Suppose that $f|_{\Lambda_Q}$ is transitive. Then for every t ,*

$$(8) \quad P(\varphi_t) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \sum_{\substack{f^n(x)=x, \\ x \in \Lambda_Q}} |(f^n)'(x)|^{-t}.$$

Proof. By transitivity, for each I_j , $j = 1, \dots, p$, there exists a cylinder $\Delta(j) \stackrel{\text{def}}{=} \Delta_{i_1 \dots i_{s_j}} \subset I_j$ such that $f^{s_j}(\Delta_{i_1 \dots i_{s_j}} \cap \Lambda_Q) = \Lambda_Q$. Put $s \stackrel{\text{def}}{=} \max\{s_j: 1 \leq j \leq p\}$. For every $n \geq 1$ denote

$$\Sigma_n^0 \stackrel{\text{def}}{=} \{(i_1 \dots i_n): f^n(\Delta_{i_1 \dots i_n} \cap \Lambda_Q) = \Lambda_Q\}.$$

Recall the choice of the sequence $(\rho_n)_n$ given in (6). Notice that, by the tempered distortion property, for any $n \geq 1$ we have

$$(9) \quad \begin{aligned} \sum_{\substack{f^n(x)=x, \\ x \in \Lambda_Q}} |(f^n)'(x)|^{-t} &\geq \sum_{(i_1 \dots i_n) \in \Sigma_n^0} \min_{x \in \Delta_{i_1 \dots i_n}} |(f^n)'(x)|^{-t} \\ &\geq e^{-nt\rho_n} \sum_{(i_1 \dots i_n) \in \Sigma_n^0} |(f^n)'(x_{i_1 \dots i_n})|^{-t}, \end{aligned}$$

where $x_{i_1 \dots i_n} \in \Delta_{i_1 \dots i_n}$ is arbitrarily chosen.

Given $k > 1$, we have

$$\begin{aligned}
 & \sum_{(i_1 \dots i_k j_1 \dots j_s) \in \Sigma_{k+s}^0} |(f^{k+s})'(x_{i_1 \dots i_k j_1 \dots j_s})|^{-t} \\
 &= \sum_{(i_1 \dots i_k)} \sum_{\substack{(j_1 \dots j_s) \in \Sigma_s^0, \\ f^k(\Delta_{i_1 \dots i_k})}} \supset \Delta_{j_1 \dots j_s} |(f^{k+s})'(x_{i_1 \dots i_k j_1 \dots j_s})|^{-t} \\
 &\geq \sum_{(i_1 \dots i_k)} \sum_{\substack{(j_1 \dots j_s) \in \Sigma_s^0, \\ f^k(\Delta_{i_1 \dots i_k})}} \supset \Delta_{j_1 \dots j_s} \min_{x \in \Delta_{i_1 \dots i_k}} |(f^k)'(x)|^{-t} |(f^s)'(f^k(x_{i_1 \dots i_k j_1 \dots j_s}))|^{-t} \\
 &\geq C \sum_{(i_1 \dots i_k)} \min_{x \in \Delta_{i_1 \dots i_k}} |(f^k)'(x)|^{-t} \\
 &\geq C e^{-kt\rho_k} \sum_{(i_1 \dots i_k)} \max_{x \in \Delta_{i_1 \dots i_k}} |(f^k)'(x)|^{-t}
 \end{aligned}$$

(remember that each sum is taken over the cylinders which intersect Λ_Q), where C is some positive constant depending entirely only on f^s .

Since every $\Delta_{i_1 \dots i_n}$ contains at most one periodic point of period n , we have

$$(10) \quad \sum_{\substack{f^n(x)=x, \\ x \in \Lambda_Q}} |(f^n)'(x)|^{-t} \leq \sum_{(i_1 \dots i_n)} \max_{x \in \Delta_{i_1 \dots i_n}} |(f^n)'(x)|^{-t}.$$

From here the claimed property follows. □

We remark that the above proof used some ideas from [20, Section 5].

2.4. Conformal measures. Denote by $\mathcal{L}_\varphi \psi(x) \stackrel{\text{def}}{=} \sum_{f(y)=x, y \in \Lambda_Q} e^{\varphi(y)} \psi(y)$, $x \in \Lambda_Q$, the *Ruelle-Perron-Frobenius transfer operator* associated to φ , which is acting on the space of continuous functions $\psi: \Lambda_Q \rightarrow \mathbb{R}$. By the Schauder-Tychonov theorem, there exists a fixed point of the map $\nu \mapsto (\mathcal{L}_\varphi^* \nu(1))^{-1} \mathcal{L}_\varphi^* \nu$. This fixed point, say ν , is a conformal measure with respect to the function $\exp(P(\varphi) - \varphi)$; that is, it satisfies

$$\nu(f(A)) = \int_A e^{P(\varphi) - \varphi(x)} d\nu(x)$$

for every Borel subset A of Λ_Q such that $f|_A$ is injective (see [4] for more details on conformal measures).

In the particular case that in the definition of the transfer operator we consider the potential φ_t defined in (7) with t such that $P(\varphi_t) = 0$, such a measure ν_t satisfies

$$(11) \quad \nu_t(f(A)) = \int_A |f'(x)|^t d\nu_t(x).$$

A measure satisfying (11) is called *t-conformal* for $f|_{\Lambda_Q}$. The following lemma follows immediately from the definition. The proof will be left to the reader.

Lemma 4. *If $f|_{\Lambda_Q}$ is transitive, the t-conformal measure is strictly positive on every cylinder.*

We denote by

$$t_c \stackrel{\text{def}}{=} \inf\{t \geq 0: t\text{-conformal measure exists on } \Lambda_Q\}.$$

3. POINTS WITH INFINITELY MANY HYPERBOLIC INSTANTS

We will *a priori* not require that $f|_{\Lambda_Q}$ is expanding, neither that $f|_{\Lambda_Q}$ has bounded distortion. However, in the following, we derive that a rather large set of points already has good expanding and distortion properties.

Let us denote by

$$\text{Dist } g|_Z \stackrel{\text{def}}{=} \sup_{x,y \in Z} \frac{|g'(x)|}{|g'(y)|}$$

the maximal distortion of a map g on a set Z . We denote by H the set of points x in Λ_Q which have the property that there exist positive constants c_1, c_2 , a monotonically increasing sequence of natural numbers $(n_k)_k$ and a positive sequence $(r_k)_k$ decreasing to 0 (all, in general, depending on x) such that for every $k \geq 1$ the map f^{n_k} is defined on $B(x, r_k)$ and we have

$$\text{diam } f^{n_k}(B(x, r_k)) > c_1 \text{ and } \text{Dist } f^{n_k}|_{B(x, r_k)} < c_2.$$

We call the numbers n_k *hyperbolic instants* for x . We remark that the following statements of this section remain true if we consider Λ instead of Λ_Q ; however, we keep our restricted point of view.

We want to relate the above concept to the concept of hyperbolic times, which requires slightly stronger expansion properties of a given orbit. A number $n \in \mathbb{N}$ is called a *hyperbolic time* for a point $x \in \Lambda_Q$ with exponent α if

$$|(f^k)'(f^{n-k}(x))| \geq e^{k\alpha} \text{ for every } 1 \leq k \leq n.$$

We denote by $H_{\text{HT}}(\alpha)$ the set of points $x \in \Lambda_Q$ for which there exist infinitely many hyperbolic times with exponent α . Obviously, $H_{\text{HT}}(\alpha) \subset H_{\text{HT}}(\alpha')$ if $\alpha' < \alpha$.

The following is an immediate consequence of the generator condition (1) and Lemma 3.

Lemma 5. *We have $\chi(x) \geq 0$ for every $x \in \Lambda_Q$.*

We now show some immediate properties of the set H .

Proposition 4. *We have*

$$\{x \in \Lambda_Q : \chi(x) > 0\} = \bigcup_{\alpha > 0} H_{\text{HT}}(\alpha) \subset H.$$

A periodic point x or its preimage is in H if and only if $\chi(x) > 0$.

Remark 1. Under the prerequisites of Urbański in [16] every point in Λ except parabolic fixed points and their preimages belongs to H . Thus, H can contain points with zero Lyapunov exponent.

Remark 2. It follows from the proof of Proposition 4 that the constants c_1 and c_2 can, in fact, be chosen uniformly on any of the sets $\{x \in H : \chi(x) \geq \chi_0 > 0\}$.

Proof of Proposition 4. Since f locally extends to a $C^{1+\text{Lip}}$ diffeomorphism, for a given $\alpha > 0$ there exists $c_1 > 0$ such that

$$(12) \quad \sup_{y,z \in I_i} \max_{|f(y)-f(z)| \leq c_1} \frac{|f'(y)|}{|f'(z)|} \leq e^{\alpha/2}.$$

We choose c_1 so small that two points from Λ_Q at a distance smaller than c_1 must belong to the same I_i .

Let n be a hyperbolic time for $x \in \Lambda_Q$ with exponent $\alpha > 0$. We now prove by induction that for every y such that $f^n(y) \in B(f^n(x), c_1)$ for every $0 \leq k \leq n$ we have

$$(13) \quad |f^{n-k}(x) - f^{n-k}(y)|e^{k\alpha/2} \leq |f^n(x) - f^n(y)|.$$

For y , (13) is true for $k = 0$. Assume now that we have this property for each $k = 0, \dots, \ell$, and hence that

$$|f^{n-\ell}(x) - f^{n-\ell}(y)| \leq |f^n(x) - f^n(y)|e^{-\ell\alpha/2} < c_1 e^{-\ell\alpha/2} < c_1.$$

With (12) and the fact that $|(f^{\ell+1})'(f^{n-\ell-1}(x))| \geq e^{(\ell+1)\alpha}$ this implies that $|f^{n-\ell-1}(x) - f^{n-\ell-1}(y)| \leq |f^n(x) - f^n(y)|e^{-(\ell+1)\alpha/2}$. This shows (13).

In particular, (13) with $k = n$ implies the first claimed property since by setting $r(n) \stackrel{\text{def}}{=} c_1 e^{-n\alpha/2}$ we have

$$c_1 \leq \text{diam } f^n B(x, c_1 r(n)).$$

We now prove the bounded distortion property. By (13) and by the Lipschitz continuity of $\log|f'|$ with some Lipschitz constant $c > 0$ we have

$$\begin{aligned} \log \text{Dist } f^n|_{B(x,r(n))} &= \sup_{y,z \in B(x,r(n))} \log \frac{|f'(y)||f'(f(y))| \cdots |f'(f^{n-1}(y))|}{|f'(z)||f'(f(z))| \cdots |f'(f^{n-1}(z))|} \\ &\leq \sum_{k=0}^{n-1} c e^{-(n-k)\alpha/2} |I| \leq \sum_{k=0}^{\infty} c e^{-k\alpha/2} |I| \stackrel{\text{def}}{=} c_2 < \infty, \end{aligned}$$

which proves the inclusion $H_{\text{HT}}(\alpha) \subset H$.

To show that $\bigcup_{\alpha>0} H_{\text{HT}}(\alpha)$ coincides with the set of points with positive Lyapunov exponent, one can apply an auxiliary result due to Pliss (see, for example, [13]). However, we prefer to give a direct proof. Let $x \in \Lambda_Q$ be a point with positive Lyapunov exponent. Let us assume that $x \notin H_{\text{HT}}(\chi_0)$ for some $0 < \chi_0 < \chi(x)$, in other words, that it has at most finitely many hyperbolic times n_i with exponent χ_0 . Let $N \stackrel{\text{def}}{=} \max_i n_i$. Let $n > N$. Since n is not a hyperbolic time, there exists $k_1 < n$ such that

$$|(f^{n-k_1})'(f^{k_1}(x))| \leq e^{(n-k_1)\chi_0}.$$

If $k_1 > N$, we can continue and find numbers $k_\ell \leq N < k_{\ell-1} < \dots < k_1 < n$ such that

$$|(f^{k_{i-1}-k_i})'(f^{k_i}(x))| \leq e^{(k_{i-1}-k_i)\chi_0}, \quad i = 2, \dots, \ell.$$

Hence, we have

$$|(f^{n-k_i})'(f^{k_i}(x))| \leq e^{(n-k_i)\chi_0}.$$

But this implies that there exists some positive constant $C = C(N, \chi_0)$ such that

$$|(f^n)'(x)| \leq C e^{n\chi_0} \text{ for every } n > N$$

and hence $\chi(x) \leq \chi_0$, which is a contradiction. Thus, we can conclude that x has infinitely many hyperbolic times with any exponent smaller than $\chi(x)$.

It follows obviously from the first part of this proof that a preimage of a periodic point x is in H if $\chi(x) > 0$. We finally prove that if a periodic point x or its preimage is in H , then we must have $\chi(x) > 0$. Let $x = f^k(x)$ be a point satisfying $|(f^k)'(x)| = 1$. Let $y = g_{i_n \dots i_1}(x)$ be in H and let c_1 and c_2 be associated constants and $(n_k)_k$ and $(r_k)_k$ the associated sequences. Then

$$\text{diam } B(y, r_k) \geq |(f^{n_k})'(y)|^{-1} c_2^{-1} c_1 = |(f^n)'(y)|^{-1} |(f^{n_k-n})'(x)|^{-1} c_2^{-1} c_1$$

is uniformly bounded from below by some positive constant. In particular this means that $|\Delta_k(y)|$ stays bounded away from zero for all $k \in \mathbb{N}$ (a contradiction with (1)). This finishes the proof of Proposition 4. \square

Proposition 5. *We have $\dim_{\mathbb{H}} H \leq t_c$.*

Proof. Let $t \geq t_c$ such that there exists a t -conformal measure μ_t for $f|_{\Lambda_Q}$. Let $x \in H$ and consider the sequences $(n_k)_k$ and $(r_k)_k$ and positive constants c_1 and c_2 associated to x . Note that

$$(14) \quad 2r_k \cdot \inf_{y \in B(x, r_k)} |(f^{n_k})'(y)| \leq \text{diam } f^{n_k}(B(x, r_k)) \leq |I|.$$

By t -conformality of the measure μ_t , we have

$$\begin{aligned} \mu_t(f^{n_k}(B(x, r_k))) &= \int_{B(x, r_k)} |(f^{n_k})'(y)|^t d\mu_t(y) \\ &\leq c_2^t \inf_{y \in B(x, r_k)} |(f^{n_k})'(y)|^t \mu_t(B(x, r_k)), \end{aligned}$$

where for the second inequality we used the bounded distortion property of the map f^{n_k} at the point x . With (14) we obtain

$$\mu_t(f^{n_k}(B(x, r_k))) \leq \left(\frac{c_2}{2r_k} |I| \right)^t \mu_t(B(x, r_k)).$$

As $\text{diam}(f^{n_k}(B(x, r_k))) > c_1$, $\mu_t(f^{n_k}(B(x, r_k)))$ is bounded away from zero uniformly in k by some constant c (by Lemma 4 and a compactness argument) and hence

$$\frac{\log \mu_t(B(x, r_k))}{\log r_k} \leq \frac{-t \log(c_2 |I| 2^{-1}) + c}{\log r_k} + t.$$

This implies that the lower pointwise dimension of the measure μ_t at x can be bounded by $\underline{d}_{\mu_t}(x) \leq t$. Since $x \in H$ was arbitrary, we obtain $\dim_{\mathbb{H}} H \leq t$ (see for example [7]). \square

Note that the conformal measure μ_t is not necessarily concentrated on the set H . But we do not need this property in order to obtain the upper bound of $\dim_{\mathbb{H}} H$.

Proposition 6. *The set H has zero Lebesgue measure.*

Proof. We will show that no $x \in H$ is a Lebesgue-density point of H ; that is, we have

$$(15) \quad \liminf_{r \rightarrow 0} \frac{\text{Leb}(H \cap B(x, r))}{\text{Leb}(B(x, r))} < 1 \text{ for every } x \in H,$$

where Leb denotes the Lebesgue measure on \mathbb{R} . Since a set is nowhere dense if and only if it has zero Lebesgue measure, the assertion then follows.

Let $x \in H$ with associated sequences $(n_k)_k$ and $(r_k)_k$ and numbers $c_1 > 0$ and $c_2 \geq 1$. There exists a positive number c_3 , which depends only on c_1 but neither on n_k nor r_k , such that for every number n_k there exists an interval gap $G_k \subset f^{n_k}(B(x, r_k)) \setminus \Lambda_Q$ of length at least c_3 . Denote $\tilde{G}_k = f^{-n_k}(G_k)$. Analogously to (14), by bounded distortion of the map f^{n_k} we have

$$2r_k \cdot |(f^{n_k})'(x)| \cdot c_2^{-1} \leq \text{diam } f^{n_k}(B(x, r_k) \cap \Lambda) \leq |I|,$$

which implies

$$c_3 \leq |G_k| \leq |\tilde{G}_k| \sup_{y \in B(x, r_k)} |(f^{n_k})'(y)| \leq |\tilde{G}_k| |(f^{n_k})'(x)| c_2 \leq |\tilde{G}_k| |I| \frac{c_2^2}{2r_k}.$$

Thus we obtain

$$\frac{\text{Leb}(H \cap B(x, r_k))}{\text{Leb}(B(x, r_k))} \leq \frac{r_k(1 - c_3 c_2^{-2} |I|^{-1})}{r_k} < 1$$

for a sequence $(r_k)_k$. From here (15) follows. □

4. POINTS WITH INFINITELY MANY HYPERBOLIC INSTANTS:
GEOMETRIC DESCRIPTION

Our goal in this section is to prove Theorem 2, that is, to prove that a point belongs to $\Lambda_Q \setminus H$ if and only if it is either a parabolic periodic point or a parabolic preperiodic point.

We briefly sketch how we are going to prove this result. We first formulate the A. Schwartz lemma in our setting. We then identify the family of intervals to which this lemma is applicable. The main step in the proof is done by a bootstrapping argument: by using the A. Schwartz lemma we are able to cut out all points that visit a certain cylinder infinitely many times. By doing so, we can restrict ourselves to a new, strictly smaller, Markov system and use the A. Schwartz lemma again. This procedure terminates after finitely many steps.

Proof of Theorem 2. Since $x \in H$ if and only if $f(x) \in H$, by Proposition 1 we can restrict ourselves to the case that Σ_Q is irreducible. We assume that Q is a representation of Σ_Q .

We start with a distortion result which is based on the A. Schwartz lemma [14]. Denoting by $\text{Lip } g$ the Lipschitz constant of g , let

$$\theta \stackrel{\text{def}}{=} \max_{1 \leq i \leq p} \text{Lip } \log |g'_i|$$

and

$$(16) \quad \lambda \stackrel{\text{def}}{=} \exp(4\theta|I|).$$

Lemma 6. *Let G and K be two intersecting intervals which are contained in some I_i , $1 \leq i \leq p$, and which satisfy $\lambda|K| \leq |G|$. If $(i_1 i_2 \dots) \in \Sigma$ satisfies*

$$\sum_{k=1}^{\infty} \text{diam } g_{i_k \dots i_1}(G) \leq 2|I|,$$

then for every $n \in \mathbb{N}$ we have

$$\text{diam } g_{i_n \dots i_1}(K) \leq \text{diam } g_{i_n \dots i_1}(G) \quad \text{and} \quad \text{Dist } g_{i_n \dots i_1}|_{G \cup K} \leq \lambda.$$

Proof. We prove the statement by induction over n . The assertion is true for $n = 1$ because we have $|K| < |G|$ and

$$\log(|g'_i(x)||g'_i(y)|^{-1}) \leq \theta|x - y| < 2\theta|I|$$

for every $x, y \in G \cup K$.

Let us assume that the assertion is true for all $n < N$. We have

$$\text{Dist } g_{i_N \dots i_1}|_{G \cup K} \leq \exp(\theta \sum_{k=0}^{N-1} \text{diam } g_{i_k \dots i_1}(G \cup K)).$$

By the inductive hypothesis, for all $k < N$ we have

$$\text{diam } g_{i_k \dots i_1}(G \cup K) \leq 2 \text{diam } g_{i_k \dots i_1}(G).$$

This implies that

$$\text{Dist } g_{i_N \dots i_1}|_{G \cup K} \leq \exp(4\theta|I|) = \lambda$$

and that

$$\frac{\text{diam } g_{i_N \dots i_1}(K)}{\text{diam } g_{i_N \dots i_1}(G)} \leq \frac{|K|}{|G|} \text{Dist } g_{i_N \dots i_1}|_{G \cup K} \leq 1.$$

This proves the lemma. \square

We will call an open interval G contained in one of the intervals I_i a *gap* if $G \cap \Lambda_Q = \emptyset$ and if at least one of the endpoints of G belongs to Λ_Q .

We will apply Lemma 6 to the following geometrical situation:

Lemma 7. *Let $G \subset I_i$ be a gap, $x \in \Lambda_Q$ be an endpoint of G and let $(i_1 i_2 \dots) \in \Sigma$ be an admissible sequence for x . Assume that the points $g_{i_k \dots i_1}(x)$ are all distinct for $k \in \mathbb{N}$. Then*

$$\sum_{k=1}^{\infty} \text{diam } g_{i_k \dots i_1}(G) \leq 2|I|.$$

Proof. We will prove that $g_{i_k \dots i_1}(G)$ are pairwise disjoint for all such k for which $g_{i_k \dots i_1}$ are both orientation preserving or both orientation reversing (on I_i).

Assume that it is not true, i.e., that $g_{i_k \dots i_1}(G)$ intersects $g_{i_\ell \dots i_1}(G)$ for $g_{i_k \dots i_1}$ and $g_{i_\ell \dots i_1}$ both orientation preserving or both orientation reversing. The points $g_{i_k \dots i_1}(x)$ and $g_{i_\ell \dots i_1}(x)$ are both right endpoints (or both left endpoints) of $g_{i_k \dots i_1}(G)$ and $g_{i_\ell \dots i_1}(G)$, correspondingly. Assume that $k < \ell$. There are three possible cases.

Case 1: $y = g_{i_k \dots i_1}(x) \in g_{i_\ell \dots i_1}(G)$. In this situation $f^\ell(y) \in G$. However, $f^\ell(y) = f^{\ell-k}(x) \in \Lambda_Q$, which is a contradiction.

Case 2: $y = g_{i_\ell \dots i_1}(x) \in g_{i_k \dots i_1}(G)$. In this situation $f^k(y) \in G$. However, $f^k(y) = g_{i_{\ell-k} \dots i_1}(x) \in \Lambda_Q$ (by definition of admissible sequence), which is again a contradiction.

Case 3: $g_{i_k \dots i_1}(x) = g_{i_\ell \dots i_1}(x)$ is excluded by the assumption of the lemma. We are done. \square

Recall the definition of d_n in (1). Let G be a gap and let n be so large that

$$(17) \quad d_n \leq \frac{|G|}{3\lambda}.$$

We also assume that

$$(18) \quad n \geq l(Q).$$

Let x be an endpoint of G which is contained in Λ_Q . Denote by K the d_n -neighbourhood of $\Delta_n(x) = \Delta_{j_1 \dots j_n}$. Let $Q' \stackrel{\text{def}}{=} Q \cup \{(j_1 \dots j_n)\}$.

Remark 3. Before continuing the proof, let us recall that the range of g_i lies strictly inside I_i . Hence, Λ is strictly contained in the interior of $\bigcup g_i(\bigcup I_j)$. In particular, if $g_{i_k \dots i_1}|_K$ has distortion bounded by λ , then for any $y \in g_{i_k \dots i_1}(\Delta_n(x))$ the map f^k is well defined on $B(y, D|\Delta_{n+k}(y)|/\lambda)$ for some D not depending on k .

Lemma 8. *If the assertion of Theorem 2 is true for $\Lambda_{Q'}$, it is also true for Λ_Q .*

Proof. We denote by K_* the set of points $y \in \Lambda_Q$ for which $f^k(y) \in \Delta_n(x)$ for infinitely many $k \in \mathbb{N}$. We are going to show first that every point in K_* which is not a preimage of a parabolic periodic point is contained in H .

Note first that by Lemma 2 any sequence admissible for x is necessarily admissible for all the points in $\Delta_n(x)$. There are two cases.

Case 1: x is not a periodic point of f . Then for any sequence $(i_1 i_2 \dots)$ which is admissible for x , the points $g_{i_k \dots i_1}(x)$, $k \geq 1$, are pairwise distinct. Hence, by Lemma 7 we can apply Lemma 6 to the sets G and K and to the maps $g_{i_k \dots i_1}$. In particular we have

$$(19) \quad \text{Dist } g_{i_k \dots i_1} |_{G \cup K} \leq \lambda$$

for every $k \in \mathbb{N}$.

For any $y \in \Lambda_Q$, if $y \in g_{i_k \dots i_1}(\Delta_n(x))$ for some k -tuple $(i_k \dots i_1)$, then $f^k(\Delta_{n+k}(y)) = \Delta_n(x)$. It follows from (19) that we have

$$\text{diam } f^k \left(B \left(y, \frac{D|\Delta_{n+k}(y)|}{\lambda} \right) \right) \geq \frac{D}{\lambda^2} |\Delta_{j_1 \dots j_n}|$$

and

$$\text{Dist } f^k |_{B \left(y, \frac{D|\Delta_{n+k}(y)|}{\lambda} \right)} \leq \lambda.$$

Hence, if $f^k(y) \in \Delta_n(x)$ for infinitely many k , then $y \in H$. We conclude $K_* \subset H$ in Case 1.

Case 2: x is a periodic point of f of period m . We have $x = \pi((\ell_1 \dots \ell_m)^\mathbb{N})$ for some m -tuple $(\ell_1 \dots \ell_m)$ and

$$(20) \quad \ell_k = j_k \text{ for } k \leq \min\{m, n\}.$$

Let $y \in K_*$.

Assume for a moment that there exists $M_0 > 0$ such that $f^k(y) \in \Delta_{j_1 \dots j_n}$ implies that

$$(21) \quad f^{k-m}(y) \in \Delta_{\ell_1 \dots \ell_m}$$

for every $k > M_0$. This means that in the symbolic expansion of y from some moment on any appearance of $(j_1 \dots j_n)$ is immediately preceded by $(\ell_1 \dots \ell_m)$, that is,

$$y = \pi \left(\underset{k-m \text{ symbols}}{* \dots *} \ell_1 \dots \ell_m j_1 \dots j_n \dots \right).$$

From (20) it then follows that the word $(\ell_1 \dots \ell_m j_1 \dots j_n)$ must be preceded by $(\ell_1 \dots \ell_m)$ again, and so on. As $y \in K_*$, the word $(j_1 \dots j_n)$ appears infinitely many times. Hence the symbolic expansion of y is of the form $(i_1 \dots i_k (\ell_1 \dots \ell_m)^\mathbb{N})$; that is, y is a preimage of the periodic point x .

If y is a preimage of x , then, by Proposition 4, $y \in H$ if and only if x is an expanding periodic point.

If y is not a preimage of x , then (21) is not satisfied; that is, there exist infinitely many numbers k such that $f^k(y) \in \Delta_n(x)$ and $f^{k-m}(y) \notin \Delta_{\ell_1 \dots \ell_m}$. For any such k , the sequence $(i_k \dots i_1)$ which satisfies $y = g_{i_k \dots i_1}(f^k(y))$ is admissible for all points in $\Delta_n(x)$ and does not begin with $(\ell_m \dots \ell_1)$. Hence, the points $g_{i_\ell \dots i_1}(x)$ are pairwise distinct for every $1 \leq \ell \leq k$. By Lemma 7, we can apply Lemma 6 to the sets G and K and to the maps $g_{i_k \dots i_1}$. We obtain

$$\text{diam } f^k \left(B \left(y, \frac{D|\Delta_{n+k}(y)|}{\lambda} \right) \right) \geq \frac{D}{\lambda^2} |\Delta_{j_1 \dots j_n}|$$

and

$$\text{Dist } f^k|_{B(y, \frac{D|\Delta_{n+k}(y)|}{\lambda})} \leq \lambda$$

as in Case 1. This implies that all the points in K_* which are not preimages of parabolic periodic points belong to H if we consider the Case 2.

Consider now a point $y \in \Lambda_Q$ for which $f^{k_i}(y) \in \Delta_n(x)$ for only finitely many k_i . Then $f^k(y) \in \Lambda_{Q'}$ for every $k > \max_i k_i$. Hence,

$$\Lambda_Q \subset K_* \cup \bigcup_{(i_1 \dots i_m)} g_{i_1 \dots i_m}(\Lambda_{Q'}),$$

where the union is taken over some infinite but countable subset of Σ_* . In such a situation, to obtain the assertion of Theorem 2 for Λ_Q it is enough to prove this assertion for $\Lambda_{Q'}$. \square

Lemma 8 gives us the inductive step for the proof of Theorem 2. We start by choosing p gaps (one in each I_i) and $n \in \mathbb{N}$ such that (17) is satisfied for each gap and such that (18) is satisfied as well. To a chosen gap $G \subset I_i$ we apply Lemma 8. We see that to prove the assertion of Theorem 2 for Λ_Q it is enough to prove it for some subsystem $\Lambda_{Q'}$. Using Proposition 1 we might well assume that $\Sigma_{Q'}$ is irreducible.

Either $\Lambda_{Q'}$ is disjoint from I_i (in this case we proceed to another gap) or we can find a new gap $G' \supset G$. Its diameter is obviously not smaller than the diameter of G and hence (17) is satisfied for G' . Lemma 1 implies that (18) is satisfied for $\Lambda_{Q'}$. Hence, we can repeat the inductive step.

Since every Markov system which we consider in this chain is strictly contained in the previous one and since $l(\Sigma_Q)$ is bounded from above, the procedure must eventually terminate. In other words, at some moment we will obtain $\Lambda_{Q'} = \emptyset$. This will end the proof of Theorem 2. \square

Corollary 1. *There exist at most finitely many parabolic periodic points.*

Proof. In the proof of Theorem 2 we used an inductive procedure that produces at most one parabolic periodic point at every step. All other periodic points (not the endpoints of gaps on some step of our procedure) belong to H and thus cannot be parabolic. As the inductive procedure terminates after finitely many iterations, the number of parabolic periodic points can be at most finite. \square

The next corollary follows in the case that f is C^2 from the Mañe hyperbolicity theorem as well; see [11] for the formulation.

Corollary 2. *If there exist no parabolic periodic points, the system is uniformly hyperbolic.*

Proof. If there exist no parabolic periodic points, then, by Theorem 2, we have $H = \Lambda_Q$. Every point $x \in H$ has a cylinder neighbourhood $U(x)$ and a time $k(x)$ such that $f^{k(x)}|_{U(x)}$ is uniformly expanding. As $\Lambda_Q = H$ is compact, from sets $U(x)$ one can choose a finite covering of Λ_Q . Hence, Λ_Q is equivalent (under a change of symbolic description) to a finite uniformly expanding Markov system. \square

We say that the Markov system (Λ_Q, f) is C^1 *robustly expansive* if and only if any map g which is a sufficiently small C^1 smooth perturbation of f also satisfies (1).

Corollary 3. *The Markov system (Λ_Q, f) is C^1 robustly expansive if and only if $f|_{\Lambda_Q}$ is uniformly hyperbolic.*

Proof. Let (Λ_Q, f) be a $C^{1+\text{Lip}}$ expansive and transitive Markov system. By Corollary 1 there can exist at most a finite number of parabolic periodic points for $f|_{\Lambda_Q}$. Let us assume that there exists at least one parabolic periodic point $x \in \Lambda_Q$, and we may assume that x is a fixed point. Then we can find a perturbation g of f , which keeps x fixed, which is arbitrarily C^1 close to f and for which there is a neighborhood U of x such that $g(U) \subset \text{int}(U)$, violating (1). Hence $f|_{\Lambda_Q}$ must be uniformly hyperbolic by Corollary 2.

If $f|_{\Lambda_Q}$ is uniformly hyperbolic, then any small C^1 perturbation is also hyperbolic, and hence expansive. \square

5. PROOF OF THEOREM 1

Given $\mu \in \mathcal{M}$, we denote by

$$\chi(\mu) \stackrel{\text{def}}{=} \int \log|f'| d\mu$$

the *Lyapunov exponent* of μ .

By Theorem 2, the set $\Lambda_Q \setminus H$ is at most countable and hence has zero Lebesgue measure. Thus, it follows from Proposition 6 that the Lebesgue measure of Λ_Q is zero.

We now prove the dimension results. From Theorem 2 we know that $\dim_{\text{H}} \Lambda_Q \setminus H = 0$, and hence

$$(22) \quad \dim_{\text{H}} \Lambda_Q \leq t_c$$

follows from Proposition 5.

Note that $f|_{\Lambda_Q}$ is an open and expansive map. By [4, Theorem 3.12], for every $t \geq 0$ satisfying $P(-t \log|f'|) = 0$ there exists a t -conformal measure. This implies that

$$(23) \quad t_c \leq t_0.$$

Denote

$$D \stackrel{\text{def}}{=} \sup_{\mu} \dim_{\text{H}} \mu,$$

where the supremum is taken over all ergodic f -invariant measures supported on Λ_Q with positive entropy. D is commonly referred to as the *dynamical dimension* of $f|_{\Lambda_Q}$. Assume that $D > 0$ and consider $0 < t < D$. It follows from [8] that there exists an ergodic f -invariant measure μ with positive entropy such that $t < h_{\mu}(f)/\chi(\mu)$. This implies

$$0 < h_{\mu}(f) - t\chi(\mu) \leq P(-t \log|f'|)$$

and hence $t < t_0$. Note that $0 \leq t_0$ trivially holds true. This implies $D \leq t_0$.

Notice that for every ergodic f -invariant measure ν we have $\chi(\nu) \geq 0$, by Lemma 5. Suppose now that $P(-t \log|f'|) > 0$ for some $t \geq 0$. Then, by the variational principle, there exists an ergodic f -invariant measure ν such that $h_{\nu}(f) - t\chi(\nu) > 0$ and thus $h_{\nu}(f) > 0$. It follows then from [8] that

$$t < \frac{h_{\nu}(f)}{\chi(\nu)} = \dim_{\text{H}} \nu \leq D.$$

This implies that $P(-t \log|f'|) \leq 0$ for every $t \geq D$, and hence we have shown that

$$(24) \quad t_0 = D.$$

Notice that, by definition, we have

$$(25) \quad D \leq \dim_{\mathbb{H}} \Lambda_Q.$$

With (22)–(25) this proves the first part of the assertion.

The following construction of conformal measures omitting exceptional points is standard nowadays (see [16] or [4]). We repeat it here because it will be used for Example 1.

Proposition 7. *If f is (piecewise) real analytic, then there exists a t_0 -dimensional conformal measure ν such that $\nu(\Lambda_Q \setminus H) = 0$.*

Proof. Given $n \geq 1$, define

$$Q_n \stackrel{\text{def}}{=} Q \cup \bigcup (i_1 \dots i_n),$$

where the union is taken over all words $(i_1 \dots i_n)$ for which $\Delta_{i_1 \dots i_n} = \Delta_n(x)$ for some parabolic periodic point $x \in \Lambda_Q$. Recall the definition (2). By Proposition 1 there exists a set $\Lambda_n \subset \Lambda_{Q_n}$ which is non-empty for large enough n and which is f -invariant and, by construction and by Corollary 2, uniformly expanding with respect to f . By [17] there exists a t_n -conformal measure supported on Λ_n , denoted by

$$t_n \stackrel{\text{def}}{=} \dim_{\mathbb{H}} \Lambda_n.$$

Notice that $(t_n)_n$ is a non-decreasing sequence. Moreover, t_n is the unique zero of the equation $P_{f|_{\Lambda_n}}(\varphi_t) = 0$, where $P_{f|_{\Lambda_n}}$ denotes the topological pressure with respect to $f|_{\Lambda_n}$. It follows that $t_n \leq t_0$ for every $n \geq 1$. There exists a subsequence $(\nu_{n_k})_k$ and a number t such that ν_{n_k} converges to some probability measure ν in the weak* topology and such that t_{n_k} converges to t . Let $A \subset \Lambda_Q$ be a Borel set such that $f|_A$ is injective. Note that, by weak* convergence and by conformality, we have

$$\nu(f(A)) = \lim_{k \rightarrow \infty} \nu_{n_k}(A) = \lim_{k \rightarrow \infty} \int_A |f'(x)|^{t_{n_k}} d\nu_{n_k} = \int_A |f'(x)|^t d\nu,$$

and hence ν is t -conformal. Since there is no t' -conformal measure for $t' < t_0$, we must have $t = t_0$.

We now prove that $\nu(\Lambda_Q \setminus H) = 0$. It is sufficient to prove that there are no atoms at parabolic points. We consider a parabolic fixed point $x \in \Lambda_Q$ at which f is locally orientation preserving, and show that we have $\nu(x) = 0$. The general case then follows from the consideration of some iterate of the map f . Let us assume that there exists such a point $x \in \Lambda_Q$. Given $n \geq 1$, denote by $D_n^\pm(x)$ the right and the left interval of the set $\Delta_n(x) \setminus \Delta_{n+1}(x)$, respectively. Note that

$$(26) \quad f(D_n^\pm(x)) = D_{n-1}^\pm(x).$$

In particular, we have

$$\sum_{k=0}^{\infty} |D_k^\pm(x)| < |I|.$$

Given some number $M \geq 1$, from Lemma 6 we can conclude that for every $k \geq 1$,

$$\text{Dist } f^k |_{D_{M+k}^\pm(x)} \leq \lambda,$$

where λ is given in (16). Using this property together with the conformality of each of the measures ν_n , for every $k \geq 1$ we derive that

$$(27) \quad \begin{aligned} \nu_n(D_M^\pm(x)) &= \nu_n(f^k(D_{M+k}^\pm(x))) = \int_{D_{M+k}^\pm(x)} |(f^k)'|^{t_n} d\nu_n \\ &\geq \frac{1}{\lambda} \frac{|D_M^\pm(x)|^{t_n}}{|D_{M+k}^\pm(x)|^{t_n}} \nu_n(D_{M+k}^\pm(x)) \end{aligned}$$

if $M+k \leq n$ and $\nu_n(D_{M+k}^\pm(x)) = 0$ else, and analogously that

$$(28) \quad \nu_n(D_M^\pm(x)) \leq \lambda \frac{|D_M^\pm(x)|^{t_n}}{|D_{M+k}^\pm(x)|^{t_n}} \nu_n(D_{M+k}^\pm(x)).$$

The same estimations are valid for (ν, t_0) in place of (ν_n, t_n) . For fixed $j \geq 1$, with (26) we can conclude that

$$\nu(\Delta_j(x)) = \lim_{n \rightarrow \infty} \nu_n(\Delta_j(x)) = \lim_{n \rightarrow \infty} \left(\sum_{\ell=j}^{\infty} \nu_n(D_\ell^-(x)) + \sum_{\ell=j}^{\infty} \nu_n(D_\ell^+(x)) \right).$$

For every $j > M$ we obtain from (27) and (28),

$$\frac{\nu_n(D_M^\pm(x))}{\lambda |D_M^\pm(x)|^{t_n}} \sum_{\ell=j}^n |D_\ell^\pm(x)|^{t_n} \leq \sum_{\ell=j}^n \nu_n(D_\ell^\pm(x)) \leq \lambda \frac{\nu_n(D_M^\pm(x))}{|D_M^\pm(x)|^{t_n}} \sum_{\ell=j}^n |D_\ell^\pm(x)|^{t_n}$$

and hence

$$(29) \quad \nu(\Delta_j(x)) \leq \lambda \left(\frac{\nu(D_M^-(x))}{|D_M^-(x)|^{t_0}} \sum_{\ell=j}^{\infty} |D_\ell^-(x)|^{t_0} + \frac{\nu(D_M^+(x))}{|D_M^+(x)|^{t_0}} \sum_{\ell=j}^{\infty} |D_\ell^+(x)|^{t_0} \right),$$

together with the analogous lower bound. If we know that each of the two series in (29) converges, then we can conclude that $\lim_{j \rightarrow \infty} \nu(\Delta_j(x)) = 0$ and hence that $\nu(x) = 0$.

If f is piecewise real analytic, then we can expand it into a power series inside some one-sided neighbourhood of x :

$$f(y) = y + c(y-x)^b + O((y-x)^{b+1})$$

for $y \in (x, x+\varepsilon)$ and similarly (possibly with different c and b) in $(x-\varepsilon, x)$. This implies

$$(30) \quad C_1 \ell^{-\beta} \leq |D_\ell^+(x)| \leq C_2 \ell^{-\beta}$$

for $\beta = b/(b-1)$ and for some positive numbers C_1, C_2 , for ℓ large enough. We assume that M was large enough so that (30) is true for all $\ell \geq M$.

Existence and finiteness of the t_0 -conformal measure ν together with (27) implies that $\beta t_0 > 1$. Given $\varepsilon > 0$ such that $\beta(t_0 - \varepsilon) > 1$, we obtain

$$\sum_{\ell=M}^{\infty} |D_\ell^+(x)|^{t_0 - \varepsilon} < +\infty;$$

hence the second series in (29) converges. The first series converges for the analogous reason and we are done. \square

The real analyticity assumption is only necessary to obtain (30). We do not want to assume (30) outright, however, because we have *a priori* no control on the location of the parabolic points.

We denote by P^s the s -dimensional *packing measure*.

Corollary 4. *If f is (piecewise) real analytic, then $P^{t_0}(H) = P^{t_0}(\Lambda_Q) > 0$.*

Proof. The proof is analogous to Proposition 5. By Proposition 7, there exists a t_0 -conformal measure ν supported on H . Let $x \in H$ and consider the sequences $(n_k)_k$ and $(r_k)_k$ and positive constants c_1 and c_2 associated to x . We have

$$(31) \quad 2r_k \cdot \sup_{y \in B(x, r_k)} |(f^{n_k})'(y)| \geq \text{diam } f^{n_k}(B(x, r_k)) \geq c_1.$$

By t_0 -conformality of the measure ν , we have

$$\begin{aligned} \nu(f^{n_k}(B(x, r_k))) &= \int_{B(x, r_k)} |(f^{n_k})'(y)|^{t_0} d\nu(y) \\ &\geq c_2^{-t_0} \sup_{y \in B(x, r_k)} |(f^{n_k})'(y)|^{t_0} \nu(B(x, r_k)). \end{aligned}$$

With (31) we obtain

$$1 \geq \nu(f^{n_k}(B(x, r_k))) \geq (c_1 c_2^{-1} r_k^{-1})^{t_0} \nu(B(x, r_k))$$

and hence

$$\liminf_{r \searrow 0} \frac{\nu(B(x, r))}{r^{t_0}} \leq (c_1^{-1} c_2)^{t_0}.$$

The assertion follows from the Frostman Lemma; see [7]. □

Corollary 5. *If f is (piecewise) real analytic, then $t_0 < 1$.*

Proof. This is an immediate consequence of Proposition 6 and Corollary 4. □

We finally give the details to the example claimed in Section 1. We demonstrate that the existence of a t_0 -conformal measure which is supported on the complement of the parabolic points is not guaranteed if the map is not real analytic. Consider the map

$$(32) \quad f(x) \stackrel{\text{def}}{=} \begin{cases} x + x^2 e^{-1/x} & \text{if } x \in [0, a], \\ 10x - 9 & \text{if } x \in [\frac{9}{10}, 1], \end{cases}$$

where $a \approx 0.8095$ is the solution of the equation $a^2 e^{-1/a} + a = 1$.

Note that for $n \geq 1$ large enough we have

$$\frac{1}{\log(n-1)} \sim \frac{1}{\log n - \frac{1}{n}} = \frac{1}{\log n \left(1 - \frac{1}{n \log n}\right)} \sim \frac{1}{\log n} + \frac{1}{n(\log n)^2} = f\left(\frac{1}{\log n}\right).$$

More precisely,

$$\frac{1}{\log(n - \frac{1}{2})} < f\left(\frac{1}{\log n}\right) < \frac{1}{\log(n-2)}$$

for n large enough. Hence, from (27) we derive that $|D_k^+(0)| \sim \frac{1}{k(\log k)^2}$ for every k large enough. Obviously,

$$\sum_{k=1}^{\infty} |D_k^+(0)| < +\infty,$$

but, for any $t < 1$ we have

$$\sum_{k=1}^{\infty} |D_k^+(0)|^t = +\infty,$$

and hence there cannot exist any t -conformal measure for $t < 1$. This implies that the Hausdorff dimension of the limit set Λ_Q equals 1. However, the one-dimensional packing measure, which is equal to its Lebesgue measure, is 0.

After finishing this paper, it was pointed out to us that this example was also studied by Dobbs [6, Section 7] proving the same conclusions of full hyperbolic and Hausdorff dimension and zero Lebesgue measure. We note that a slightly different version of the above example was already introduced in [15].

REFERENCES

1. A. Biś and M. Urbański, *Geometry of Markov systems and codimension-one foliations*, preprint.
2. R. Bowen, *A horseshoe with positive measure*, *Invent. Math.* **29** (1975), 203–204. MR0380890 (52:1787)
3. J. Cantwell and L. Conlon, *Foliations and subshifts*, *Tohoku Math. J.* **40** (1988), 165–187. MR943817 (90c:57021)
4. M. Denker and M. Urbański, *On the existence of conformal measures*, *Trans. Amer. Math. Soc.* **328** (1991), 563–578. MR1014246 (92k:58155)
5. M. Denker and M. Urbański, *Hausdorff and conformal measures on Julia sets with a rationally indifferent periodic point*, *J. London Math. Soc.* **43** (1991), 107–118. MR1099090 (92k:58153)
6. N. Dobbs, *Hyperbolic dimension for interval maps*, *Nonlinearity* **19** (2006), 2877–2894. MR2273763 (2008f:37083)
7. K. Falconer, *Techniques in Fractal Geometry*, John Wiley & Sons, 1997. MR1449135 (99f:28013)
8. F. Hofbauer and P. Raith, *The Hausdorff dimension of an ergodic invariant measure for a piecewise monotonic map of the interval*, *Canad. Math. Bull.* **35** (1992), 84–98. MR1157469 (93c:58113)
9. S. Hurder, *Dynamics and the Godbillon-Vey class: A history and survey*, in *Foliations: Geometry and Dynamics* (Warsaw, 2000), World Sci. Publishing, 2002, 29–60. MR1882764 (2003b:37044)
10. S. Matsumoto, *Measure of exceptional minimal sets of codimension one foliations*, in *A Fête of Topology, Papers Dedicated to Itiro Tamura, Y. Matsumoto, T. Mizutani, and S. Morita*, eds., Academic Press, 1988, pp. 81–94. MR928398 (89g:57040)
11. W. de Melo and S. van Strien, *One-Dimensional Dynamics*, Springer, 1993. MR1239171 (95a:58035)
12. Y. Pesin, *Dimension Theory in Dynamical Systems: Contemporary Views and Applications*, Chicago Lectures in Mathematics, The University of Chicago Press, 1998. MR1489237 (99b:58003)
13. V. Pliss, *On a conjecture due to Smale*, *Differ. Uravn.* **8** (1972), 262–268. MR0299909 (45:8957)
14. A. Schwartz, *A generalization of a Poincaré-Bendixson theorem to closed two-dimensional manifolds*, *Amer. J. Math.* **85** (1963), 453–458. MR0155061 (27:5003)
15. M. Thaler, *Transformations on $[0, 1]$ with infinite invariant measures*, *Israel J. Math.* **46** (1983), 67–96. MR727023 (85g:28020)
16. M. Urbański, *Parabolic Cantor sets*, *Fund. Math.* **151** (1996), 241–277. MR1424576 (98f:58076)
17. P. Walters, *Invariant measures and equilibrium states for some mappings which expand distances*, *Trans. Amer. Math. Soc.* **236** (1978), 121–153. MR0466493 (57:6371)
18. P. Walters, *An Introduction to Ergodic Theory*, Graduate Texts in Mathematics 79, Springer, 1981. MR648108 (84e:28017)
19. P. Walczak, *Dynamics of Foliations, Groups and Pseudogroups*, IM PAN Monografie Matematyczne, Birkhäuser, 2004. MR2056374 (2005d:57042)

20. M. Yuri, *Zeta functions for certain non-hyperbolic systems and topological Markov approximations*, Ergodic Theory Dynam. Systems **18** (1998), 1589–1612. MR1658607 (2000j:37024)
21. M. Yuri, *Thermodynamic formalism for certain nonhyperbolic maps*, Ergodic Theory Dynam. Systems **19** (1999), 1365–1378. MR1721626 (2001a:37012)

INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES, UL. ŚNIADECKICH 8, 00-950
WARSZAWA, POLAND

E-mail address: `gelfert@pks.mpg.de`

URL: `http://www.pks.mpg.de/~gelfert`

Current address: Department of Mathematics, Northwestern University, 2033 Sheridan Road,
Evanston, Illinois 60208-2730

INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES, UL. ŚNIADECKICH 8, 00-950
WARSZAWA, POLAND

E-mail address: `m.rams@impan.gov.pl`

URL: `http://www.impan.gov.pl/~rams`