

A CLASS OF FINITE SIMPLE BOL LOOPS OF EXPONENT 2

GÁBOR P. NAGY

ABSTRACT. In this paper we give an infinite class of finite simple right Bol loops of exponent 2. The right multiplication group of these loops is an extension of an elementary Abelian 2-group by S_5 . The construction uses the description of the structure of such loops given by M. Aschbacher (2005). These results answer some questions of M. Aschbacher.

1. INTRODUCTION

The set Q endowed with a binary operation $x \cdot y = xy$ is a *loop* if it has a unit element 1 and the equation $xy = z$ has a unique solution whenever two of the three unknowns are given. The loop Q is a (*right*) *Bol loop* if the identity

$$((xy)z)y = x(y(zy))$$

holds for all $x, y, z \in Q$. Bol loops of exponent 2 play an important role in the theory of loops and are related to interesting group theoretical problems. Using the so-called *Baer correspondence*, this class of loops can be described by a triple (G, H, K) with group G , core-free subgroup $H \leq G$ and system of coset representatives $K \subseteq G$ such that $1 \in K$, $G = \langle K \rangle$ and $K \setminus \{1\}$ is a union of conjugacy classes of involutions.

Bol loops of exponent 2 which are not elementary Abelian groups have long been known to exist; the first construction is due to R. P. Burn [7]. Later, many infinite classes of such loops were given; see [11, 12, 13, 16]. All of these examples were solvable loops; equivalently the group G was a 2-group. The existence of nonsolvable finite Bol loops of exponent 2 was considered as one of the main open problem in the theory of loops and quasigroups. As the smallest such loop must be simple, this question was related to the existence of finite simple proper right Bol loops. Here by proper we mean right Bol loops which are not Moufang, that is, which do not satisfy the identity $x(yx) = (xy)x$. Finite simple proper Bol loops were constructed by the author [17] recently.

By [15], the solvability of a Bol loop of 2-power exponent is equivalent to having 2-power order. Later, S. Heiss [9] showed that the solvability of the loop corresponding to the triple (G, H, K) is equivalent to the solvability of the group G . The next major step was the paper [3] by M. Aschbacher. His main result gives a detailed description on the structure of the right multiplication group of minimal nonsolvable Bol loops of exponent 2. This result was achieved by using the classification of finite simple groups.

Received by the editors July 25, 2007 and, in revised form, September 19, 2007.

2000 *Mathematics Subject Classification*. Primary 20N05; Secondary 20C20, 20F29.

This paper was written during the author's Marie Curie Fellowship MEIF-CT-2006-041105.

©2009 American Mathematical Society
Reverts to public domain 28 years from publication

In this paper we apply Aschbacher's recipe to construct a class of finite simple Bol loops of exponent 2. In this way, we give a negative answer to questions 2 and 3 of [3] and [4]. The smallest member of our class has order 96. We emphasize that this example is small and the structural description of the smallest example in [3] and [4] is so precise that it was only a matter of time until somebody found it by some short computer calculation. This explains the fact that this loop was independently discovered by the author and by B. Baumeister and A. Stein [5] with a time delay of 10 days.

2. BASIC CONCEPTS

In order to make this paper self-contained, we define the basic loop theoretical concepts. Subloops, homomorphisms and normal subloops can be defined for the class of loops similarly to groups; see [6]. The loop Q is said to be *simple* if it has no proper normal subloops.

For a loop Q , the maps $L_a, R_a : Q \rightarrow Q$, $L_a(x) = ax$, $R_a(x) = xa$ are the *left* and *right multiplication maps* of Q . They are bijections of Q and generate the *left* and *right multiplication groups* of Q , respectively. The *section* $S(Q)$ of the loop Q is the set of the right multiplication maps of Q .

The right Bol identity can equivalently be expressed by $R_x R_y R_x \in S(Q)$ for all $x, y \in Q$. Bol loops are *power-associative*; that is, x^n is well defined for all $x \in Q$ and $n \in \mathbb{Z}$. The *order* of the element x is the smallest positive integer n such that $x^n = 1$. The *exponent* of Q is the smallest positive integer n such that $x^n = 1$ for each $x \in Q$.

We will formulate our results by using the *Baer correspondence* between the class of loops and the class of triples (G, H, K) where G is a group, $H \leq G$ and K is a set of coset representatives of all conjugates H^g in G , and $1 \in K$. For a given loop Q , G can be chosen to be the right multiplication group, H the stabilizer of the unit element and K the set of right multiplication maps. Conversely, for a given triple (G, H, K) , a loop can be constructed in the following way. We define the operation $x \circ y$ on K by $H(x \circ y) = Hxy$; then $Q = (K, \circ)$ is a loop. The triple satisfying the above condition will be said to be a *loop folder*.

The loop folder (G, H, K) determines a Bol loop of exponent 2 if and only if $K = \{1\} \cup \bigcup_{i \in I} C_i$, where the C_i 's are conjugacy classes of involutions in G .

3. THE FIRST CONSTRUCTION

As usual, S_5 and $PGL(2, 5)$ are the permutation groups acting on 5 and 6 points, respectively. It is well known that $S_5 \cong PGL(2, 5) \cong \text{Aut}(L_2(4)) \cong O_4^-(2)$, where $\text{Aut}(L_2(4))$ is the extension of $L_2(4) = PSL(2, 4) \cong A_5$ by a field automorphism of order 2, and $O_4^-(2)$ is the orthogonal group on a 4-dimensional orthogonal space over \mathbb{F}_2 of Witt index 1. We denote by F_{20} the affine linear group acting on \mathbb{F}_5 , $F_{20} \cong C_5 \rtimes C_4$. On the one hand, F_{20} is the Borel subgroup of $PGL(2, 5)$, that is, the stabilizer of a projective point. On the other hand, $F_{20} \leq S_5$ is a sharply 2-transitive permutation group on 5 points.

In the sequel, we define a group G which is a nonsplit extension of the elementary Abelian group of order 32 by S_5 such that the transpositions of S_5 lift to involutions in G and the even involutions of S_5 lift to elements of order 4. Despite the relatively small order of G , we found no simple description for this group; therefore our definition will be rather *ad hoc*, as well. We start with two technical lemmas.

Lemma 3.1. *We have the following presentations of groups with generators and relations.*

$$\begin{aligned} A_5 &= \langle a, b \mid a^2 = b^3 = (ab)^5 = 1 \rangle, \\ S_5 &= \langle c, d \mid c^2 = d^4 = (cd)^5 = [c, d]^3 = 1 \rangle, \\ 2.S_5 &= \langle C, D \mid C^2 = D^8 = (CD)^5 = [C, D]^3 = [C, D^4] = 1 \rangle, \end{aligned}$$

where $2.S_5$ denotes the nonsplit central extension of S_5 in which the transpositions lift to involutions. In other words, $2.S_5$ is the semidirect product of $SL(2, 5) = 2.A_5$ with a group of order 2.

Proof. The presentation for A_5 is well known. Assume $G = \langle c, d \rangle$ is the group presented by the second set of relations above. We observe first that S_5 satisfies these relations; hence no relation can collapse. Put $a = d^2$ and $b = [d, c] = [c, d]^{-1}$. Then $a^2 = b^3 = 1$ and $ab = (dc)^2 = ((cd)^2)^c$. This latter implies $(ab)^5 = 1$; hence $G_0 = \langle a, b \rangle \cong A_5$. Moreover, since $dc = (dc)^6 = (ab)^3 \in G_0$, we have $cd = dc[c, d] = dcb^{-1} \in G_0$ and

$$dbd^{-1} = cdcd^{-1} = (cd)^2d^{-2} \in G_0.$$

This means that d and cd normalize G_0 , so $G_0 \triangleleft G$. So $|G : G_0| = 2$ and $d \notin G_0$ since A_5 contains no element of order 4. This proves $G \cong S_5$. Finally, D^4 is a central involution in $H = \langle C, D \rangle$ and $H/\langle D^4 \rangle$ maps surjectively to S_5 . The order of D is 8; thus, the extension is nonsplit and the involution C covers a transposition in S_5 . □

Lemma 3.2. *The permutations*

$$\begin{aligned} c &= (1, 4)(2, 9)(3, 10)(6, 11)(7, 12)(13, 21)(14, 22)(15, 24)(16, 23)(17, 30) \\ &\quad (18, 29)(19, 31)(20, 32)(33, 35)(38, 40), \\ d &= (1, 2, 4, 6, 8, 7, 5, 3)(9, 13, 25, 18, 10, 14, 26, 17)(11, 15, 27, 20, 12, 16, \\ &\quad 28, 19)(21, 30, 38, 34, 23, 31, 40, 35)(22, 32, 39, 36, 24, 29, 37, 33) \end{aligned}$$

acting on 40 points satisfy the relations

$$(1) \quad c^2 = d^8 = (cd)^5 = [c, d]^3 = [d^4, c]^2 = [d^4, cdcd^{-2}c] = 1.$$

Moreover, with $u_1 = d^4$, $u_2 = u_1^c$, $u_3 = u_1^{cd}$, $u_4 = u_1^{cdc}$, $u_5 = u_1^{cdcd}$, $u_6 = u_1^{cdcdc}$ the identity $u_1u_2u_3u_4u_5u_6 = 1$ holds.

Proof. We leave the straightforward calculations to the reader. □

Lemma 3.3. *The group $G = \langle c, d \rangle$ given in Lemma 3.2 satisfies*

- G has an elementary Abelian normal subgroup J of order 32 such that
 - (*) $G/J \cong PGL(2, 5)$ and J is the \mathbb{F}_2 -permutation module modulo its center.
- Moreover, $[G, G]/[G, J] \cong SL(2, 5)$ and G splits over $[G, G]J$.

Proof. We claim that the conjugacy class of $u_1 = d^4$ in G is $X = \{u_1, \dots, u_6\}$. It is immediate that c induces the permutation $(u_1u_2)(u_3u_4)(u_5u_6)$ on X . Moreover, d centralizes u_1 and maps $u_2 \mapsto u_3$, $u_4 \mapsto u_5$. From the last relation in (1), it follows that $u_1^{cdc} = u_1^{cd^2}$; hence $u_3^d = u_5^c = u_4$. By $[d^4, c]^2 = 1$ we have

$$u_1^{cd^4c} = cd^4cd^4cd^4c = d^4[d^4, c]^2 = d^4 = u_1;$$

thus $u_2 = u_1^c = u_1^{cd^4} = u_4^{d^2} = u_5^d$. To see that d acts on X , we need to show that d centralizes u_6 :

$$u_6^d = u_1^{cdcdcd} = u_1^{(cd)^{-2}} = u_1^{d^{-1}cd^{-1}c} = u_1^{cd^{-1}c} = u_2^{d^{-1}c} = u_5^c = u_6.$$

The action of d on X is therefore $(u_2u_3u_4u_5)$. This not only shows that X is a conjugacy class in G , but we also have the action of G on X . Indeed, one shows by straightforward calculation that $\tilde{c} = (12)(34)(56)$ and $\tilde{d} = (2345)$ satisfy the relations of S_5 from Lemma 3.1. Since the action of S_5 on 6 points is unique, we have $G/C_G(X) \cong PGL(2, 5)$.

As $[u_1, u_2] = [d^4, cd^4c] = [d^4, c]^2 = 1$ and $PGL(2, 5)$ acts 2-transitively, $[u_i, u_j] = 1$ holds for all i, j . This means that $J = \langle X \rangle$ is an elementary Abelian 2-group and $|J| = 32$ by $u_1 \cdots u_6 = 1$. Using the presentation of $2.S_5$ from Lemma 3.1, $G/J_0 \cong 2.S_5$. This implies

$$[G, G]/[G, J] \cong [G/J_0, G/J_0] \cong 2.A_5 \cong SL(2, 5).$$

Finally, G splits over $[G, G]J$ as $c \notin [G, G]J$. \square

In the sequel, G will denote a group satisfying (*). We would like to make clear that it can be shown using the computer algebra system GAP [8] that the group given in Lemma 3.2 is the unique group with this property. However, we hope that this more general approach will help in future generalization of the constructions of this paper.

Among other properties, we show in the next lemma that for our group G , $G' = [G, G]$ is a perfect group. Actually, we found G' by using the library of perfect groups in the computer algebra system GAP [8] and constructed G as a split extension of G' by an outer automorphism of order 2.

Lemma 3.4. *Let G be a group satisfying (*) and define $J_0 = [G, J]$.*

- (i) *We have $G'' = G' = [G, G] = [G, G]J$ and $|G : G'| = 2$.*
- (ii) *$G \setminus J$ contains a unique class c^G of involutions, and $|c^G| = 80$. In particular, all involutions of $G' = [G, G] = [G, G]J$ lie in J .*
- (iii) *Let P be a Sylow 5-subgroup. Then $N_{J_0}(P) = \{1\}$ and $N_G(P) \cong C_8 \times C_5$. Moreover, if the subgroup $U \leq G$ maps onto F_{20} modulo J , then $U = N_G(P)$ or $U = N_G(P)J_0$.*

Proof. (i) Let V be the permutation \mathbb{F}_2 -module of $PGL(2, 5)$ with basis $\{u_1, \dots, u_6\}$. Due to the 2-transitivity, the orbit of the element $u_1 + u_2$ consists of the elements $u_i + u_j$, $i \neq j$, which are different modulo the center $\langle u_1 + \dots + u_6 \rangle$ of V . Hence both $PGL(2, 5)$ and $PSL(2, 5)$ act transitively on the nonidentity elements of $J_0 = [G, J]$, which implies that J_0 is a minimal normal subgroup in G and $[G, G]J$. It follows that $J_0 = [G', J_0]$ and $G''/J_0 = (G'/J_0)' = G'/J_0$ by $SL(2, 5)' = SL(2, 5)$. This means $G'' = G'$. Finally, $J \leq G'$ follows from $J/J_0 = Z(G/J_0) \leq (G/J_0)' = G'/J_0$.

(ii) Since G splits over $G' = G'J$, we can take an involution c from $G \setminus G'$; the image of c in $G/J \cong S_5$ is a transposition. As J is the permutation module modulo its center, $\dim_{\mathbb{F}_2}(C_{J_0}(c)) = 2$ and $\dim_{\mathbb{F}_2}(C_J(c)) = 3$. It is easy to check that $2.S_5 \cong G/J_0$ contains 20 noncentral involutions and they are all conjugate.

Let c' be another involution in $G \setminus J$; we want to show that c, c' are conjugate. For some $g \in G$, $(cJ_0)^{gJ_0} = c'J_0$; that is, $c^g \in c'J_0$. Hence we can assume $c \in c'J_0$, $c' = cj$ with $j \in J_0$. The element cj has order 2 if and only if $j \in C_{J_0}(c)$. On the

one hand, $c^{J_0} \subseteq cC_{J_0}(c)$. On the other hand,

$$|c^{J_0}| = |J_0 : C_{J_0}(c)| = 4 = |C_{J_0}(c)| = |cC_{J_0}(c)|.$$

This implies $c^{J_0} = cC_{J_0}(c)$ and $c' \in c^{J_0}$. No involution of G' can be conjugate to c ; hence all involutions of G' must lie in J . Finally, we show $|c^G| = 80$. As $c^g \in cJ_0$ if and only if $c^g = c^j$ for some $j \in J_0$, we have $N_G(cJ_0) = C_G(c)J_0$. Moreover, $C_{G/J_0}(cJ_0) = N_G(cJ_0)/J_0$. Thus,

$$\begin{aligned} |G : C_G(c)| &= |G : N_G(cJ_0)||C_G(c)J_0 : C_G(c)| \\ &= |G/J_0 : C_{G/J_0}(cJ_0)||J_0 : C_{J_0}(c)| \\ &= |(cJ_0)^{G/J_0}||c^{J_0}| = 20 \cdot 4 = 80. \end{aligned}$$

(iii) P acts fixed point free on the involutions of J_0 ; thus, $N_{J_0}(P) = 1$. Moreover, $5 \nmid |J| - 1$; hence P centralizes a unique element $a \in J$. Let U be a preimage of F_{20} modulo J and put $\bar{U} = U/\langle a \rangle$, $\bar{J} = J/\langle a \rangle$. Then \bar{J} is a minimal normal subgroup of \bar{U} . Since $F_{20} = \bar{U}/\bar{J}$ acts faithfully on \bar{J} , we have $C_{\bar{U}}(\bar{J}) = \bar{J}$. By [10, II.3.3. Satz], \bar{J} has a complement \bar{H} in \bar{U} , $\bar{H} \cong F_{20}$. Let H be the preimage of \bar{H} . Then H has a unique (hence normal) 5-Sylow and $H \cong C_8 \times C_5$. This shows $N_G(P) \cong C_8 \times C_5$. For the last statement, record that $U \cap J_0$ is either 1 or J_0 . \square

The following proposition will apply in all of our examples of Bol loop folders of exponent 2. We hope that it will also apply in future constructions not considered here. Recall that $O_2(G)$ is the largest normal 2-subgroup of G .

Proposition 3.5. *Assume G is a finite group, $J = O_2(G)$ and $G^+ = G/J \cong S_5$. We denote by g^+ the element of S_5 corresponding to gJ . Set $L = G'J$, K_1 the involutions in $G \setminus L$, K_0 a G -invariant subset of J containing 1 such that $K_0 \setminus \{1\}$ consists of involutions, and $H \leq G$. Set $K = K_0 \cup K_1$, $n_0 = |K_0|$, and $n_1 = |K_1 \cap aJ|$ for $a \in K_1$. Assume the following.*

- (a) $(J, H \cap J, K_0)$ is a Bol loop folder of exponent 2.
- (b) $n_0 = 2n_1$.
- (c) $|G^+ : H^+| = 6$.
- (d) For each $a \in K_1$, $C_{H \cap J}(a) = 1$.
- (e) Every involution of L is contained in J .

Then (G, H, K) is a Bol loop folder of exponent 2, and $|K| = 6n_0 = 12n_1$.

Proof. First K_1^+ is the set of transpositions of S_5 , so $|K_1^+| = 10$. This implies that n_1 is well defined. Indeed, for $a, b \in K_1$, aJ, bJ are conjugate; hence $K_1 \cap aJ, K_1 \cap bJ$ are conjugate in G . Moreover, $|K_1| = 10n_1$ and by (b),

$$|K| = |K_0| + |K_1| = 2n_1 + 10n_1 = 12n_1 = 6n_0.$$

Next by (a) and (c),

$$|G : H| = |G : HJ||HJ : H| = |G^+ : H^+||J : J \cap H| = 6|K_0| = 6n_0,$$

so $|G : H| = |K|$.

We claim $xy \notin H$ for distinct $x, y \in K$. If so, as $|G : H| = |K|$, K is a set of coset representatives for H in G . Then as K is G -invariant and $K \setminus \{1\}$ consists of involutions, (G, H, K) is a Bol loop folder of exponent 2.

If $x, y \in J$, then $x, y \in K_0$, so $xy \notin H$ by (a). Next $K_1^+ \cap H^+ = \emptyset$, so if $x \in J$ and $y \in K_1$, then $(xy)^+ = y^+ \notin H^+$, so $xy \notin H$. Thus we may take $x, y \in K_1$ and $xy \in H$. Now as K_1^+ is a set of transpositions in S_5 , the order of $(xy)^+$ is 1, 2

or 3. Since $H^+ \cong F_{20}$ has no element of order 3, we get $(xy)^2 \in J$. In particular, $D = \langle x, y \rangle$ is a 2-group. Let z be the unique involution in $\langle xy \rangle$. By $xy \in L$ and (e), $z \in H \cap J$. Moreover, x, y commute with z , which contradicts (d). \square

Remark. The fact that (e) is necessary can be seen from the counterexample $G = S_5 \times J$.

Theorem 3.6. *Assume G is a group satisfying condition (*) of Lemma 3.3. Let J_0 be the minimal normal subgroup of G and put $K = J_0 \cup c^G$. Define $H = N_G(P)$ where P is a 5-Sylow subgroup of G . Then (G, H, K) is a Bol loop folder determining a simple Bol loop of exponent 2 of order 96. Conversely, if (G, H^*, K^*) is an exponent 2 Bol loop folder, then H^* is a conjugate of H and $K^* = K$.*

Proof. With the notation of Proposition 3.5, $K_0 = J_0$ and $K_1 = c^G$. Then $n_0 = 16$, $n_1 = |c^G \cap cJ| = 80/10 = 8$ and $|G^+ : H^+| = 6$, so (b) and (c) hold. Part (e) follows from Lemma 3.4(ii). Since J is elementary Abelian, $H \cap J$ consists of 1 and the unique involution of H . This involution cannot be centralized by c ; otherwise it would be central in $G = \langle c, P, J \rangle$. Hence (d) follows. Finally, $H \cap J$ is not contained in J_0 ; therefore J_0 is a complement to $H \cap J$ in J , showing (a). By Proposition 3.5, (G, H, K) is a Bol loop folder of exponent 2.

For the converse, we observe that (G, H^*, K^*) determines a Bol loop of exponent 2 with all proper subloops solvable. Thus, by the Main Theorem of [3], H^* maps surjectively to F_{20} . By Lemma 3.4(iii), $H^* = H$ or $H^* = HJ_0$ up to conjugation. In the latter case, the loop has order 6, which is impossible. Again by Aschbacher's result, $c^G \subset K^*$. Finally, if $J_0 \not\subseteq K^*$, then K^* will contain a conjugate of the involution of H , which is not possible. This proves the theorem. \square

As the group given in Lemma 3.2 satisfies (*), we have

Corollary 3.7. *There exists a simple Bol loop of exponent 2 and order 96.* \square

Remark. The Bol loop folder (G, H, K) of Theorem 3.6 was discovered independently by B. Baumeister and A. Stein [5] (Free University of Berlin), as well.

4. S_5 -MODULES OVER \mathbb{F}_2

In this section we collect some useful facts about kS_5 -modules, where k is a field of characteristic 2.

Lemma 4.1. *The group S_5 has three absolutely irreducible representations over \mathbb{F}_2 : the trivial representation and two representations M, N of dimension 4. The two 4-dimensional modules can be distinguished by the fact that $C_M(x) = 0$ and $\dim_{\mathbb{F}_2}(C_N(x)) = 2$ for an element $x \in S_5$ of order 3. Moreover, the following hold.*

- (i) *M is the 4-dimensional irreducible component in the 6-dimensional permutation module for $S_5 \cong PGL(2, 5)$. Also, let V be the natural 2-dimensional module of $A_5 \cong SL(2, 4)$ over the field \mathbb{F}_4 and let σ be the semilinear map of V induced by the Frobenius automorphism of \mathbb{F}_4 . Then $S_5 \cong SL(2, 4) \rtimes \langle \sigma \rangle$ and V is a 4-dimensional S_5 -module over \mathbb{F}_2 . The S_5 -modules M and V are isomorphic.*
- (ii) *N is the 4-dimensional irreducible component in the 5-dimensional permutation module of S_5 . Also if N is a 4-dimensional orthogonal space of Witt index 1 over \mathbb{F}_2 , then $O(N) = O_4^-(2) \cong S_5$. Note that N has 5 singular and 10 nonsingular vectors and these are the S_5 -orbits on N .*

- (iii) N is absolutely irreducible as an A_5 -module. M is irreducible but not absolutely as an A_5 -module, the splitting field being \mathbb{F}_4 . In particular, the modules are nonisomorphic as A_5 -modules.
- (iv) N and M are isomorphic absolutely irreducible projective F_{20} -modules.

Proof. Let us first define N, M as irreducible components of the permutation modules. By [14, Table 1], they are absolutely irreducible. As S_5 has 3 classes of elements of odd order, by [1, Theorem 3.2] S_5 has no other absolutely irreducible modules over F_2 . The properties of N, M can be verified by straightforward calculations; the irreducibility as A_5 - and F_{20} -modules follows again from [14, Table 1]. We show $N_{F_{20}} \cong M_{F_{20}}$. As F_{20} has two classes of elements of odd order, F_{20} has two absolutely irreducible modules: the trivial one and $N_{F_{20}}$ coming from the 2-transitive permutation representation. So if $M_{F_{20}}$ were not isomorphic to $N_{F_{20}}$, then it could be brought to upper triangular form over $\overline{\mathbb{F}}_2$, which is clearly impossible.

It remains to show that $N_{F_{20}}$ is projective. Since N_{C_4} is isomorphic to the group algebra $\mathbb{F}_2 C_4$, it is a projective C_4 -module by [1, Theorem 4.2]. Using [1, Corollary 9.3], we obtain that N is projective as an F_{20} -module. \square

We observe that these S_5 -modules can immediately be constructed using the Steinberg Tensor Product Theorem [18, Theorem 13.1], as well.

We will now construct an S_5 -module U which will play a central role in the generalization of our first construction of a Bol loop of exponent 2.

Let $U = U_1 \oplus U_2$ be the direct sum of two copies of N as an $\mathbb{F}_2 A_5$ -module. As U_i is an orthogonal space, we can regard U as an orthogonal space which is the orthogonal direct sum of the two nondegenerate subspaces U_1 and U_2 . The stabilizer B of $\{U_1, U_2\}$ in $O(U)$ is $(G_1 \times G_2)\langle \tau \rangle$ where

$$G_i = C_{O(U)}(U_{3-i}) \cong O(U_i) \cong S_5$$

and τ is an involution interchanging U_1, U_2 . Thus B is the wreath product of S_5 with C_2 . In particular, the elements τ and $(g_1, g_2) \in G_1 G_2$ map $u_1 \oplus u_2 \in U$ to

$$(u_1 \oplus u_2)\tau = u_2 \oplus u_1 \text{ and } (u_1 \oplus u_2)(g_1, g_2) = u_1 g_1 \oplus u_2 g_2,$$

respectively. Set

$$G_0 = C_{G_1 G_2}(\tau) = \{(g, g) \mid g \in S_2\} \cong S_5,$$

and let $L = [G_0, G_0] \cong A_5$, t_0 an involution (transposition) in $G_0 \setminus L$, $t = t_0 \tau$, and $D = L\langle t \rangle$. Then $t\tau = \tau t$ and the action of t on U is

$$(u_1 \oplus u_2)t = u_2 c \oplus u_1 c,$$

where $c \in S_5$ is the transposition corresponding to t_0 . It is immediate that $D \cong S_5$.

Set $W = C_U(\tau)$. Then W is an $\mathbb{F}_2 D$ -submodule of U , and also

$$W = [U, \tau] = \{u + u\tau \mid u \in U_1\} = \{u \oplus u \mid u \in N\},$$

with the map $u \mapsto [u, \tau] = u + u\tau$ an $\mathbb{F}_2 L$ -isomorphism of U_1 with W . If Q is the quadratic form on U , then as $Q(u_1 \oplus u_2) = Q(u_1) + Q(u_2)$ and $Q(u) = Q(u\tau)$, $[u, \tau]$ is singular, so W is totally singular.

Lemma 4.2. *With the notation above, we have the following.*

- (i) U has 3 irreducible L -submodules, namely U_1, U_2 and W .
- (ii) W is the unique proper D -submodule of U .

- (iii) Let P be a Sylow 5-subgroup of D , $D_1 = N_D(P)$. Then $D_1 \cong F_{20}$ and U has precisely 3 D_1 -submodules W, T_1, T_2 .
- (iv) The orbits of D_1 on T_i have length 1, 5, 10. In particular, each member of T_i is fixed by some involution of D_1 .

Proof. By Lemma 4.1, N is projective as an F_{20} -module. Since U/W is a 4-dimensional irreducible for D_1 , U_{D_1} splits over $W_{D_1} \cong N_{D_1}$ and hence $U = W \oplus T_1$ with D_1 -submodule T_1 . Again by Lemma 4.1, N_L and N_{D_1} are completely irreducible. Schur’s lemma then implies $\text{End}_{\mathbb{F}_2 L}(N) = \text{End}_{\mathbb{F}_2 D_1}(N) = \mathbb{F}_2$. We can now apply [2, (27.14)] to obtain (i) and (iii). Part (ii) follows from (i). Finally, (iv) holds since T_i and W are D_1 -isomorphic and W is the permutation module modulo the center. □

In the next lemma, we keep using the above notation.

Lemma 4.3. (i) $\dim C_U(t) = 4$, $C_U(t) + T_1 = U$ and $C_U(t) \cap T_1 = 0$.
 (ii) Under the action of A_5 on the submodules W, U_1, U_2 of U , the lengths of the orbits are 1, 5, 10. Let S_0, S_1, S_2 be the orbits of length 5 in W, U_1, U_2 , respectively. Then $S = \{0\} \cup S_0 \cup S_1 \cup S_2$ is a (nonlinear) S_5 -invariant complement to T_1 in U ; that is, $S + T_1 = U$.

Proof. (i) We have $D = L\langle t \rangle \cong S_5$. Let us denote the element of S_5 corresponding to $a \in D$ by a^+ ; w.l.o.g. we can assume $t^+ = (12)$ and $P^+ = \langle (12345) \rangle$. Then $D_1^+ = \langle (12345), (1325) \rangle$. Let $b \in D$ such that $b^+ = (12)(35)$. Then $b \in D_1$ and b commutes with t and τ . Record that the action of b, t, τ on U is

$$\begin{aligned} b & : u_1 \oplus u_2 \mapsto u_1 b^+ \oplus u_2 b^+, \\ \tau & : u_1 \oplus u_2 \mapsto u_2 \oplus u_1, \\ t & : u_1 \oplus u_2 \mapsto u_2 t^+ \oplus u_1 t^+. \end{aligned}$$

Define the element $\tilde{c} \in B$ by

$$\tilde{c} : u_1 \oplus u_2 \mapsto u_1 t^+ \oplus u_2.$$

Then \tilde{c} commutes with b and $\tau^{\tilde{c}} = t$. Put $E_1 = \langle t, b \rangle$; clearly $E_1^{\tilde{c}} = \langle \tau, b \rangle$. On the one hand, we have

$$\begin{aligned} \dim_{\mathbb{F}_2}(C_U(E_1)) & = \dim(C_U(\tau, b)) \\ & = \dim C_{C_U(b)}(\tau) \\ & = \dim C_{C_{U_1}(b) \oplus C_{U_2}(b)}(\tau) \\ & = \dim C_{U_1}(b) \\ & = \dim C_N(b^+) = 2. \end{aligned}$$

We show on the other hand that $\dim_{\mathbb{F}_2}(C_W(E_1)) \geq 2$. Indeed, the S_5 -modules W and N are isomorphic and $\dim(C_W(t)) = \dim(C_N(t^+)) = 3$. Then $C_W(E_1) = C_{C_W(t)}(b)$ is of rank at least $\dim(C_W(t))/2 = 3/2$.

Now, from $\dim(C_W(E_1)) \geq 2$ it follows that $C_U(E_1) \leq W$. However if $C_{T_1}(t) \neq 0$, then $C_{T_1}(E_1) \neq 0$, contradicting $T_1 \cap W = 0$ and $C_U(E_1) \leq W$.

(ii) We have seen that S_i is the set of singular points in U_i for $i = 1, 2$. The action of $D_1 \cong F_{20}$ on S_i is its natural 2-transitive action on 5 points. Here $D_1 \cap L$ contains precisely 5 involutions and each member of S_i is fixed by exactly one involution of $D_1 \cap L$. Moreover, each member of T_1 is fixed by some involution of D_1 .

We have to show that for distinct $x, y \in S$, $x + y \notin T_1$; then S is a complement to T_1 by an order argument. Assume $x + y \in T_1$ and denote by a an involution of D_1 fixing $x + y$. Then x, y are the projections of $x + y$ on S_i, S_j , so a fixes the projections x and y . As $T_1 \cap U_i = 0$ for $0 \leq i \leq 2$, $x \in S_i$ and $y \in S_j$ for some $i \neq j$. If $x \in S_1$ and $y \in S_2$, then x and y are the unique fixed points of a in S_1, S_2 . By $a\tau = \tau a$, $x\tau = y$ holds, and hence $x + y = [x, \tau] \in W$, contradicting $T_1 \cap W = 0$.

Thus we may take $x \in S_0$ and $y \in S_1$. Then $x = x_1 + x_2$ with $x_i \in S_i$, and as $x \in S_0$, $x_2 = x_1\tau$ with $x_i \in S_i$. Now, x_i is the unique fixed point of a in S_i and y the unique fixed point of a in S_1 , so $y = x_1$ and $x + y = x_1 + x_2 + y = x_2 \in T_1 \cap U_2 = 0$, a contradiction. \square

5. AN INFINITE FAMILY OF SIMPLE BOL LOOPS OF EXPONENT 2

In this section, G denotes a group satisfying condition (*) of Lemma 3.3, H is the normalizer of a 5-Sylow P of G , and c is an involution from $G \setminus [G, G]J$. The S_5 -modules N and U are defined as in Section 4. Also $U = U_1 \oplus U_2 = T_1 \oplus T_2$ where U_1, U_2 are A_5 -submodules and T_1, T_2 are F_{20} -submodules. Moreover, U_1, U_2, T_1, T_2 are different from the unique S_5 -submodule W of U . All these subspaces are irreducible \mathbb{F}_2P -modules, which implies $U_i \cap T_j = 0$.

Let us fix a positive integer k and put

$$\mathcal{U} = U^k, \mathcal{U}_i = U_i^k, \mathcal{T}_i = T_i^k, \mathcal{W} = W^k.$$

Clearly, \mathcal{W} is an S_5 -submodule and $\mathcal{U} = \mathcal{U}_1 \oplus \mathcal{U}_2 = \mathcal{T}_1 \oplus \mathcal{T}_2$. We write $\mathcal{G} = G \ltimes \mathcal{U}$ where $J \triangleleft G$ acts trivially on the S_5 -module \mathcal{U} . Moreover,

$$\mathcal{J} = O_2(\mathcal{G}) = \langle J, \mathcal{U} \rangle.$$

We will consider the elementary Abelian subgroup \mathcal{J} of \mathcal{G} as an S_5 -module over the field \mathbb{F}_2 . In particular, with some abuse of notation, we will denote the group operation on \mathcal{J} additively and write $\mathcal{J} = J + \mathcal{U}$, etc. It is easy to see that

$$\mathcal{G} / \mathcal{J} \cong S_5 \text{ and } \text{soc}(\mathcal{G}) = J_0 + \mathcal{W}.$$

Moreover, since J_0 and W are nonisomorphic S_5 -modules, $J_0 \oplus W$ does not contain diagonal submodules. This implies that for any minimal submodule M of \mathcal{J} , we either have $M = J_0$ or $M \leq \mathcal{W}$.

The action of the involution $c \in G \setminus J$ on \mathcal{U} equals the action of the transposition $(12) \in S_5$; hence c interchanges $\mathcal{U}_1, \mathcal{U}_2$. This implies $|C_{\mathcal{U}}(c)| = 16^k$ and $|c^{\mathcal{U}}| = 16^k$; that is, \mathcal{U} is transitive on the involutions in $c\mathcal{U}$. As $G \cong \mathcal{G} / \mathcal{U}$ is transitive on the 80 involutions in $G \setminus J$ and as \mathcal{U} is transitive on the 16^k involutions on $c\mathcal{U}$, \mathcal{G} is transitive on the $80 \cdot 16^k$ involutions in $\mathcal{G} \setminus \mathcal{J}$.

Using Aschbacher's Main Theorem [3], we conclude that a Bol loop folder $(\mathcal{G}, \mathcal{H}, \mathcal{K})$ must have the following properties: The index of \mathcal{H} has to be $96 \cdot 16^k$; that is, \mathcal{H} must have order $40 \cdot 16^k$. The set of involutions \mathcal{K} is the union of $c^{\mathcal{G}}$ and $\mathcal{K} \cap \mathcal{J}$; thus, $|\mathcal{K} \cap \mathcal{J}| = 16 \cdot 16^k$.

There are very many possible choices for \mathcal{H} and \mathcal{K} . The most obvious choice is the following.

Proposition 5.1. *Put $\mathcal{H} = H \ltimes \mathcal{T}_1$ and $\mathcal{K} = c^{\mathcal{G}} \cup (J_0 \oplus \mathcal{W})$. Then $(\mathcal{G}, \mathcal{H}, \mathcal{K})$ is a Bol loop folder. Moreover, the homomorphism $\mathcal{G} \rightarrow G$ with kernel \mathcal{U} induces a surjective homomorphism between the loop folders $(\mathcal{G}, \mathcal{H}, \mathcal{K})$ and (G, H, K) . In other words, the Bol loop corresponding to $(\mathcal{G}, \mathcal{H}, \mathcal{K})$ is an extension of the elementary Abelian group of order 2^{2k} by the loop corresponding to (G, H, K) .*

Proof. We apply Proposition 3.5; (a), (c) and (e) are trivial. Part (b) follows from

$$n_1 = |c^{\mathcal{G}} \cap c \mathcal{J}| = |C_{\mathcal{J}}(c)| = 8|C_{\mathcal{U}}(c)| = 8 \cdot 16^k$$

and $n_0 = |J_0||\mathcal{W}| = 16 \cdot 16^k = 2n_1$. For (d), we use

$$C_{\mathcal{H} \cap \mathcal{J}}(c) = C_{H \cap J + \mathcal{T}_1}(c) = C_{H \cap J}(c) = 1,$$

as by Lemma 4.3, $\mathcal{T}_1 \cap \mathcal{T}_1^c = 0$. □

In the rest of this section, for each integer $k \geq 1$, we modify these \mathcal{H} and \mathcal{K} such that the resulting loop will be simple. Let U^* be a copy of U in \mathcal{U} such that $\mathcal{U} = U^* \oplus U^{k-1}$. We denote the subspaces corresponding to T_i, U_i, W by T_i^*, U_i^*, W^* . Let us define the set $S \subseteq U$ as in Lemma 4.3(ii) and let S^* be the corresponding subset of U^* . In order to construct the new \mathcal{K} , we simply replace W^* by S^* .

Let $\psi : J_0 \rightarrow T_1^*$ be an isomorphism of F_{20} -modules and define

$$\mathcal{T}_\psi = \{v + \psi(v) + u \mid v \in J_0, u \in T_1^{k-1}\}.$$

Then \mathcal{T}_ψ is normalized by H and we define the new subgroup \mathcal{H} of \mathcal{G} by $\mathcal{H} = H \times \mathcal{T}_\psi$.

Theorem 5.2. *Let $\mathcal{H} = H \times \mathcal{T}_\psi$, $\widetilde{\mathcal{W}} = (\mathcal{W} \setminus W^*) \cup S^*$ and $\mathcal{K} = c^{\mathcal{G}} \cup (J_0 + \widetilde{\mathcal{W}})$. Then the triple $(\mathcal{G}, \mathcal{H}, \mathcal{K})$ is a Bol loop folder such that the corresponding Bol loop is simple of exponent 2.*

Proof. Again, the first statement follows from Proposition 3.5 and Lemma 4.3; one needs to verify hypothesis (a) and (d) of the proposition only. Again (d) follows from $T_1 \cap T_1^c = 0$.

We prove (a) by showing that $\mathcal{K}_0 = \mathcal{K} \cap \mathcal{H} \mathcal{J}$ is a transversal to \mathcal{H} in $\mathcal{H} \mathcal{J}$. Since $\mathcal{K}_0 \subset \mathcal{J}$, this is equivalent to the fact that $\mathcal{K}_0 = J_0 + \widetilde{\mathcal{W}}$ is a complement of the subspace

$$\mathcal{H} \cap \mathcal{J} = H \cap J + \mathcal{T}_\psi$$

in \mathcal{J} , i.e.,

$$\mathcal{K}_0 + \mathcal{H} \cap \mathcal{J} = \mathcal{J}.$$

In order to show this, we use the identities

$$\begin{aligned} \mathcal{W} + \mathcal{T}_1 &= \mathcal{U}, \\ J_0 + \mathcal{T}_\psi &= J_0 + \mathcal{T}_1, \\ J_0 + \mathcal{H} \cap \mathcal{J} &= J_0 + H \cap J + \mathcal{T}_\psi \\ &= J_0 + H \cap J + \mathcal{T}_1. \end{aligned}$$

Then

$$\begin{aligned} J_0 + \mathcal{W} + \mathcal{H} \cap \mathcal{J} &= J_0 + \mathcal{W} + H \cap J + \mathcal{T}_\psi \\ &= J_0 + \mathcal{W} + H \cap J + \mathcal{T}_1 \\ &= J_0 + \mathcal{U} + H \cap J \\ &= J + \mathcal{U} \\ &= \mathcal{J}; \end{aligned}$$

that is, $J_0 + \mathcal{W}$ is a complement to $\mathcal{H} \cap \mathcal{J}$ in \mathcal{J} by the order argument

$$|J_0 + \mathcal{W}||\mathcal{H} \cap \mathcal{J}| = |\mathcal{J}|.$$

We constructed $\widetilde{\mathcal{W}}$ from \mathcal{W} by deleting the minimal submodule W^* and replacing it by S^* . Therefore, it is enough to show that this deformation of \mathcal{W} does not change the property of being a complement:

$$\begin{aligned} J_0 + W^* + \mathcal{H} \cap \mathcal{J} &= J_0 + W^* + H \cap J + \mathcal{T}_\psi \\ &= H \cap J + J_0 + W^* + \mathcal{T}_1 \\ &= H \cap J + J_0 + W^* + T_1^* + \mathcal{T}_1 \\ &= H \cap J + J_0 + S^* + T_1^* + \mathcal{T}_1 \\ &= J_0 + S^* + H \cap J + \mathcal{T}_\psi \\ &= J_0 + S^* + \mathcal{H} \cap \mathcal{J}. \end{aligned}$$

This proves (a); hence $(\mathcal{G}, \mathcal{H}, \mathcal{K})$ is a Bol loop folder.

It remains to show that the Bol loop \mathcal{Q} corresponding to $(\mathcal{G}, \mathcal{H}, \mathcal{K})$ is simple. Let us therefore assume that $\mathcal{Q} \rightarrow \mathcal{Q}^\sharp$ is a nontrivial surjective loop homomorphism and let $(\mathcal{G}^\sharp, \mathcal{H}^\sharp, \mathcal{K}^\sharp)$ be the loop folder of \mathcal{Q}^\sharp . Then we have a surjective homomorphism $\alpha : \mathcal{G} \rightarrow \mathcal{G}^\sharp$ with $\alpha(\mathcal{H}) = \mathcal{H}^\sharp$ and $\alpha(\mathcal{K}) = \mathcal{K}^\sharp$. Let $\mathcal{N} = \ker \alpha$ and $c^\sharp = \alpha(c) = c\mathcal{N}$. On the one hand, \mathcal{H}^\sharp is core-free; thus,

$$(2) \quad \text{core}_{\mathcal{G}}(\mathcal{H}\mathcal{N}) = \mathcal{N}.$$

On the other hand, $\mathcal{N} \leq \mathcal{J}$ since otherwise $\mathcal{H}\mathcal{N} = \mathcal{G}$ and $\mathcal{Q}^\sharp = 1$.

Let us first assume that $J_0 \leq \mathcal{N}$. Since

$$J/J_0 \leq Z(\mathcal{G}/J_0) \triangleleft \mathcal{G}/J_0 \text{ and } J/J_0 \leq \mathcal{H}\mathcal{N}/J_0,$$

we have $J \leq \mathcal{N}$ by (2). In this case the image G^\sharp of $G \leq \mathcal{G}$ is a homomorphic image of $G/J \cong S_5$. Furthermore, if $[c^\sharp, c^{\sharp g}] = 1$, then $c^\sharp c^{\sharp g}$ normalizes a Sylow 5-subgroup of G^\sharp ; thus, $c^\sharp c^{\sharp g}$ is contained in a conjugate of \mathcal{H}^\sharp , and hence $c^\sharp = c^{\sharp g}$ in this case. As the commuting graph of transpositions in S_5 is connected, $c^\sharp = c^{\sharp g}$ for all g . This means $[c, \mathcal{G}] \leq \mathcal{N}$, contradicting $\mathcal{N} \leq \mathcal{J}$.

Let us now assume $J_0 \not\leq \mathcal{N}$ and let M be a minimal normal subgroup of \mathcal{G} contained in \mathcal{N} . Then $M \leq \text{soc}(\mathcal{G}) = J_0 + \mathcal{W}$. Since J_0 and W are nonisomorphic S_5 -modules, $J_0 + \mathcal{W}$ contains no submodules isomorphic to J_0 and different from J_0 . This implies $M \leq \mathcal{W}$ and, in particular, $\mathcal{N} \cap \mathcal{W} \neq 0$.

Let us take an element $s \in S^* \setminus W^* \subseteq \mathcal{H}$. As W^* and T_0^* are complements in U^* , $s \neq 0$ has the unique decomposition $s = w + t$ with $0 \neq w \in W^*$ and $0 \neq t \in T_1^*$. Furthermore, for $0 \neq j = \psi^{-1}(t) \in J_0$, $j + t \in \mathcal{T}_\psi \leq \mathcal{H}$ holds. We claim that $j + t \in \mathcal{N}$. Indeed, we have the decomposition

$$w = (s + j) + (j + t), \quad s + j \in \mathcal{H}, \quad j + t \in \mathcal{H}.$$

If $\mathcal{N} \leq U^*$, then $M = W^*$ and $\alpha(s + j) = \alpha(j + t) \in \mathcal{H}^\sharp \cap \mathcal{H}^\sharp = 1$. In particular, $j + t \in \mathcal{N}$. If $\mathcal{N} \not\leq U^*$, then for an arbitrary element $n \in (\mathcal{N} \cap \mathcal{W}) \setminus U^*$, $w + n \in \mathcal{H}$. This means that the element $\alpha(w)$ has two $\mathcal{H}^\sharp \mathcal{H}^\sharp$ decompositions:

$$\alpha(w) = \alpha(w + n) + 0 = \alpha(s + j) + \alpha(j + t).$$

This is only possible if $j + t \in \mathcal{N}$; thus our claim is proved.

Let M' be the S_5 -submodule generated by $j + t$. Then $J_0 \leq M' \leq \mathcal{N}$ as the irreducible J_0 is not S_5 -isomorphic to a submodule of U^* . This contradiction proves the simplicity of \mathcal{Q} . \square

Remark. We have seen that there are at least two possibilities for the choice of \mathcal{H} . Also in $\widetilde{\mathcal{W}}$, we can replace any minimal submodule W^{**} by an appropriate S^{**} . This shows that there are *many* Bol loops of exponent 2 which live in the same nonsolvable group. Many of these loops are simple. Using computer calculations, we were able to construct over 30 nonisomorphic simple Bol loops of exponent 2 in \mathcal{G} in the case $k = 1$.

In fact, this phenomena is not unusual for Bol loops of exponent 2. In [12, Section 5] and [16, Theorem 5.5], the authors constructed rich classes of Bol loops of exponent 2 having the same enveloping groups, namely the wreath product $C_2^n \wr C_2$ and the extra special 2-group $E_{2^{2n+1}}^+$, respectively. In these cases, a simple parametrization of the conjugacy classes of involutions enabled a description of the associated loops. Unfortunately, the group \mathcal{G} has many conjugacy classes of involutions and these classes have no nice algebraic parametrization. Therefore, we see no way of classifying all simple Bol loops with enveloping group \mathcal{G} .

The above remark lets us make another observation. While the class of finite Bol loops of exponent 2 is very rich, the structure of the right multiplication group of a Bol loop of exponent 2 is rather restricted. In other words, while the classification of finite simple Bol loops of exponent 2 seems to be hopeless, we think that the classification of right multiplication groups of such loops could be a meaningful project.

We finish this paper with the following

Problem. Classify those almost simple groups T for which an exponent 2 Bol loop folder (G, H, K) exists such that $T \cong G/O_2(G)$.

ACKNOWLEDGEMENTS

The author would like to thank Michael Aschbacher (California Institute of Technology) for his detailed comments on the previous version of this paper. He helped to fill some gaps, to simplify the proofs and to replace computer based calculations with theoretical arguments. The author also would like to thank Peter Müller (Würzburg) for many stimulating conversations and helpful comments.

REFERENCES

- [1] J. L. ALPERIN. *Local representation theory*. Cambridge University Press, New York, 1993. MR860771 (87i:20002)
- [2] M. ASCHBACHER. *Finite group theory*. Cambridge Studies in Advanced Mathematics, 10. Cambridge University Press, Cambridge, 1986. MR895134 (89b:20001)
- [3] M. ASCHBACHER. On Bol loops of exponent 2. *J. Algebra* **288**, No. 1, 99-136 (2005). MR2138373 (2006k:20131)
- [4] M. ASCHBACHER, M. K. KINYON and J. D. PHILLIPS. Finite Bruck loops. *Trans. Am. Math. Soc.* **358**, No. 7, 3061-3075 (2005). MR2216258 (2006m:20111)
- [5] B. BAUMEISTER and A. STEIN. Private communication (2007).
- [6] R. H. BRUCK. *A survey of binary systems*. Springer-Verlag, Berlin, 1958. MR0093552 (20:76)
- [7] R. P. BURN. Finite Bol loops I. *Math. Proc. Cambridge Philos. Soc.* **84**(3) (1978), 53-66. MR0492030 (58:11194)
- [8] The GAP Group, *GAP - Groups, Algorithms, and Programming, Version 4.4.9*; 2006. <http://www.gap-system.org>.
- [9] S. HEISS. Self-invariant 1-factorizations of complete graphs, submitted for publication.
- [10] B. HUPPERT. *Endliche Gruppen I*, Springer-Verlag, Berlin, Heidelberg, 1967. MR0224703 (37:302)

- [11] H. KIECHLE. *Theory of K-loops*. Lecture Notes in Mathematics, 1778. Springer, Berlin, 2002. MR1899153 (2003d:20109)
- [12] H. KIECHLE and G. P. NAGY. On the extension of involutorial Bol loops. *Abh. Math. Semin. Univ. Hamb.* **72**, 235-250 (2002). MR1941556 (2003m:20095)
- [13] E. KOLB and A. KREUZER. Geometry of kinematic K -loops. *Abh. Math. Sem. Univ. Hamburg* **65** (1995), 189–197. MR1359127 (96k:20150)
- [14] B. MORTIMER. The modular permutation representations of the known doubly transitive groups. *Proc. London Math. Soc.* (3) **41** (1980), 1-20. MR579714 (81f:20004)
- [15] G. P. NAGY. Solvability of universal Bol 2-loops. *Comm. Algebra*, **26(2)** (1998), 549–555. MR1604103 (98k:20112)
- [16] G. P. NAGY. On the structure and number of small Frattini Bol 2-loops. *Math. Proc. Camb. Philos. Soc.* **141**, No. 3, 409-419 (2006). MR2281406 (2008a:20102)
- [17] G. P. NAGY. A class of simple proper Bol loops. *Manuscripta Math.* **127**, No. 1, 81–88 (2008). MR2429915
- [18] R. STEINBERG. Endomorphisms of linear algebraic groups. Memoirs of the American Mathematical Society, No. 80, American Mathematical Society, Providence, R.I. 1968. MR0230728 (37:6288)

BOLYAI INSTITUTE, UNIVERSITY OF SZEGED, ARADI VÉRTANÚK TERE 1, H-6720 SZEGED, HUNGARY

E-mail address: nagy@math.u-szeged.hu

Current address: Mathematisches Institut, Universität Würzburg, Am Hubland, D-97070 Würzburg, Germany