

IRREGULAR SETS, THE β -TRANSFORMATION AND THE ALMOST SPECIFICATION PROPERTY

DANIEL J. THOMPSON

ABSTRACT. Let (X, d) be a compact metric space, $f : X \mapsto X$ be a continuous map satisfying a property we call almost specification (which is slightly weaker than the g -almost product property of Pfister and Sullivan), and $\varphi : X \mapsto \mathbb{R}$ be a continuous function. We show that the set of points for which the Birkhoff average of φ does not exist (which we call the irregular set) is either empty or has full topological entropy. Every β -shift satisfies almost specification and we show that the irregular set for any β -shift or β -transformation is either empty or has full topological entropy and Hausdorff dimension.

1. INTRODUCTION

For a compact metric space (X, d) , a continuous map $f : X \mapsto X$ and a continuous function $\varphi : X \mapsto \mathbb{R}$, we define the irregular set for φ to be

$$\widehat{X}(\varphi, f) := \left\{ x \in X : \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \varphi(f^i(x)) \text{ does not exist} \right\}.$$

The irregular set arises naturally in the context of multifractal analysis. As a consequence of Birkhoff's ergodic theorem, the irregular set is not detectable from the point of view of an invariant measure. However, it is an increasingly well known phenomenon that the irregular set can be large from the point of view of dimension theory [2]. Symbolic dynamics methods have confirmed this in the uniformly hyperbolic setting [3], for certain non-uniformly hyperbolic examples [23] and for a large class of multimodal maps [31]. The irregular set has also been the focus of a great deal of work by Olsen and collaborators, for example in [1] and [16]. The papers [30, 11] studied the irregular set for maps with the specification property using a non-symbolic method inspired by the work of Takens and Verbitskiy [28].

Ruelle uses the terminology 'set of points with historic behaviour' to describe the irregular set [25]. The idea is that points for which the Birkhoff average does not exist are capturing the 'history' of the system, whereas points whose Birkhoff average converge only see average behaviour. For example, in the dynamics of the weather, the irregular points are the ones that have observed epochs of climate change. In [27], in contrast to the dimension-theoretic point of view, Takens asks for which smooth dynamical systems the irregular set has positive Lebesgue measure.

Our current work, which takes the dimension-theoretic viewpoint, is inspired by a recent innovation of Pfister and Sullivan [22], who have introduced a weak specification property called the g -almost product property, which we take the liberty of renaming as the almost specification property. They extended the variational

Received by the editors May 6, 2009 and, in revised form, November 24, 2010.
2010 *Mathematics Subject Classification*. Primary 37C45.

©2012 American Mathematical Society
Reverts to public domain 28 years from publication

principle of Takens and Verbitskiy [28] to the class of maps which satisfy this property. This variational principle is a key tool in multifractal analysis and has been extended by the author in a different direction in [29].

A striking application of the almost specification property is that it applies to every β -shift. In sharp contrast, the set of β for which the β -shift has the specification property has zero Lebesgue measure [6, 26]. The β -shift is the natural symbolic space associated to the β -transformation $f_\beta(x) = \beta x \pmod{1}$, $\beta \in (1, \infty)$, $x \in [0, 1)$. The β -transformation has been widely studied since its introduction in 1957 by Renyi [24]. The sustained interest in the study of the β -transformation arises from its connection with number theory and its special role as a model example of a one-dimensional expanding dynamical system which admits discontinuities.

Main Result 1. *When f satisfies the almost specification property, the irregular set is either empty or has full topological entropy.*

Main Result 2. *The irregular set for an arbitrary β -transformation (or β -shift) is either empty or has full entropy $\log \beta$ and Hausdorff dimension 1.*

Our second main result (stated formally as Theorem 5.1 and Theorem 5.4) is a corollary of our first main result (stated formally as Theorem 4.1). We work through the application to the β -shift carefully. We emphasise that our first main result applies to many systems other than β -transformations. For example, every ergodic toral automorphism has almost specification. This is a special case of the result that every automorphism of a compact metric abelian group that is ergodic (with respect to Haar measure) and has finite topological entropy has almost specification [15, 9, 33]. This provides another important class of examples where our first main theorem yields new results.

To undertake the proof of our first main result, we develop notions of almost spanning sets, strongly separated sets and a generalised version of the Katok formula for entropy. These should be of independent interest. We also include an alternative proof of our second main result based on approximating the β -shift from inside by shifts of finite type. We discuss the merits of the two different approaches in §5.4.

In [30], we showed that when f has the specification property, the irregular set is either empty or has full topological pressure. The method of this paper can be used to show that this more general result holds true in the almost specification setting. We restrict ourselves to the special case of entropy for clarity and brevity.

In §2, we give definitions. In §3, we establish our general version of the Katok formula for entropy. In §4, we prove our first main result. In §5, we consider arbitrary β -shifts and β -transformations and establish our second main result.

Remark. The author is grateful to a referee of this paper who informed him of the work of Durner, of which he was not aware when this work was completed. Our result on β -shifts, although not stated explicitly in Durner's work, can be obtained as a corollary of [10, Theorem 4]. Durner's work does not apply in a general setting but contains some very interesting theorems concerning β -shifts. Durner's proof implicitly uses the almost specification property of the β -shift and may be the first work in which this property was exploited.

2. DEFINITIONS AND PRELIMINARIES

Let (X, d) be a compact metric space and $f : X \mapsto X$ a continuous map. Let $C(X)$ denote the space of continuous functions from X to \mathbb{R} , and let $\varphi \in C(X)$. Let $S_n\varphi(x) := \sum_{i=0}^{n-1} \varphi(f^i(x))$. Let $\mathcal{M}_f(X)$ denote the space of f -invariant probability measures and $\mathcal{M}_f^e(X)$ denote those which are ergodic.

2.1. The irregular set. We define the irregular set for φ to be

$$(2.1) \quad \widehat{X}(\varphi, f) := \left\{ x \in X : \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \varphi(f^i(x)) \text{ does not exist} \right\}.$$

For a constant c , let $\text{Cob}(X, f, c)$ denote the space of functions cohomologous to c and $\overline{\text{Cob}(X, f, c)}$ be the closure of $\text{Cob}(X, f, c)$ in the sup norm. In [30], we proved the following lemma.

Lemma 2.1. *The following are equivalent for $\varphi \in C(X)$:*

- (a) $\frac{1}{n}S_n\varphi$ does not converge pointwise to a constant;
- (b) $\inf_{\mu \in \mathcal{M}_f(X)} \int \varphi d\mu < \sup_{\mu \in \mathcal{M}_f(X)} \int \varphi d\mu$;
- (c) $\inf_{\mu \in \mathcal{M}_f^e(X)} \int \varphi d\mu < \sup_{\mu \in \mathcal{M}_f^e(X)} \int \varphi d\mu$;
- (d) $\varphi \notin \bigcup_{c \in \mathbb{R}} \overline{\text{Cob}(X, f, c)}$;
- (e) $\frac{1}{n}S_n\varphi$ does not converge uniformly to a constant.

We also proved that if $\widehat{X}(\varphi, f)$ is non-empty, then the properties of Lemma 2.1 hold. It is a corollary of our main result (Theorem 4.1) that when f has the almost specification property, (b) implies that $\widehat{X}(\varphi, f)$ is non-empty.

2.2. The almost specification property. Pfister and Sullivan have introduced a property called the g -almost product property. We take the liberty of renaming this property as the almost specification property. The almost specification property can be verified for every β -shift (see §5.1). The version we use here is a priori weaker than that in [22] and is slightly weaker than the almost product property of [33]. First we need an auxiliary definition.

Definition 2.2. Let $\epsilon_0 > 0$. A function $g : \mathbb{N} \times (0, \epsilon_0) \mapsto \mathbb{N}$ is called a mistake function if for all $\epsilon \in (0, \epsilon_0)$ and all $n \in \mathbb{N}$, $g(n, \epsilon) \leq g(n+1, \epsilon)$ and

$$\lim_{n \rightarrow \infty} \frac{g(n, \epsilon)}{n} = 0.$$

Given a mistake function g , if $\epsilon > \epsilon_0$, we define $g(n, \epsilon) := g(n, \epsilon_0)$.

We note that for fixed $k \in \mathbb{N}$ and $\lambda > 0$, if g is a mistake function, then so is h defined by $h(n, \epsilon) = kg(n, \lambda\epsilon)$.

Definition 2.3. For $n, m \in \mathbb{N}$, $m < n$, we define the set of $(n, -m)$ index sets to be

$$I(n, -m) := \{\Lambda \subseteq \{0, \dots, n-1\}, \#\Lambda \geq n-m\}.$$

Let g be a mistake function and $\epsilon > 0$. For n sufficiently large so that $g(n, \epsilon) < n$, we define the set of (g, n, ϵ) index sets to be $I(g; n, \epsilon) := I(n, -g(n, \epsilon))$. Equivalently,

$$I(g; n, \epsilon) := \{\Lambda \subseteq \{0, \dots, n-1\}, \#\Lambda \geq n - g(n, \epsilon)\}.$$

For a finite set of indices Λ , we define

$$d_\Lambda(x, y) = \max\{d(f^j x, f^j y) : j \in \Lambda\} \text{ and } B_\Lambda(x, \epsilon) = \{y \in X : d_\Lambda(x, y) < \epsilon\}.$$

Definition 2.4. When $g(n, \epsilon) < n$, we define a ‘dynamical ball of radius ϵ and length n with $g(n, \epsilon)$ mistakes’. Let

$$\begin{aligned} B_n(g; x, \epsilon) &:= \{y \in X : y \in B_\Lambda(x, \epsilon) \text{ for some } \Lambda \in I(g; n, \epsilon)\} \\ &= \bigcup_{\Lambda \in I(g; n, \epsilon)} B_\Lambda(x, \epsilon). \end{aligned}$$

Definition 2.5. A continuous map $f : X \mapsto X$ satisfies the almost specification property if there exists a mistake function g such that for any $\epsilon_1, \dots, \epsilon_k > 0$, there exist integers $N(g, \epsilon_1), \dots, N(g, \epsilon_k)$ such that for any x_1, \dots, x_k in X and integers $n_i \geq N(g, \epsilon_i)$,

$$\bigcap_{j=1}^k f^{-\sum_{i=0}^{j-1} n_i} B_{n_j}(g; x_j, \epsilon_j) \neq \emptyset,$$

where $n_0 = 0$.

Remark 2.6. The function g can be interpreted as follows. The integer $g(n, \epsilon)$ tells us how many mistakes we are allowed to make when we use the almost specification property to ϵ shadow an orbit of length n . Henceforth, we assume for convenience and without loss of generality that $N(g, \epsilon)$ is chosen so that $g(n, \epsilon)/n < 0.1$ for all $n \geq N(g, \epsilon)$.

Remark 2.7. Pfister and Sullivan use a slightly different definition of mistake function (which they call a blow-up function). They do not allow g to depend on ϵ . An example of a function which is a mistake function under our definition but is not considered by Pfister and Sullivan is $g(n, \epsilon) = \epsilon^{-1} \log n$. Since we allow a larger class of mistake functions, the almost specification property defined here is slightly more general than the g -almost product property of Pfister and Sullivan.

Definition 2.8. A continuous map $f : X \mapsto X$ satisfies the specification property if for all $\epsilon > 0$, there exists an integer $m = m(\epsilon)$ such that for any collection $\{I_j = [a_j, b_j] \subset \mathbb{N} : j = 1, \dots, k\}$ of finite intervals with $a_{j+1} - b_j \geq m(\epsilon)$ for $j = 1, \dots, k-1$ and any x_1, \dots, x_k in X , there exists a point $x \in X$ such that

$$(2.2) \quad d(f^{p+a_j} x, f^p x_j) < \epsilon \text{ for all } p = 0, \dots, b_j - a_j \text{ and every } j = 1, \dots, k.$$

Pfister and Sullivan showed that the specification property implies the almost specification property [22] using ANY blow-up function g . To see the relation between the two concepts, we note that if f has specification and we set $g(n, \epsilon) = m(\epsilon)$ for all n larger than $m(\epsilon)$ and set $N(g, \epsilon) = m(\epsilon) + 1$, then for any x_1, \dots, x_k in X and integers $n_i \geq N(g, \epsilon)$, we have

$$\bigcap_{j=1}^k f^{-\sum_{i=0}^{j-1} n_i} B_{n_j}(g; x_j, \epsilon) \neq \emptyset.$$

The trick required to replace ϵ by $\epsilon_1, \dots, \epsilon_k$ can be found in [22].

Remark 2.9. In [33], Yamamoto studies the relationship between a number of weak forms of the specification property, including the Pfister and Sullivan definition. We note that our definition of almost specification is different from all the properties defined in [33], and the almost specification property of [33, Definition 2.1] is a special case of ours.

3. A MODIFIED KATOK ENTROPY FORMULA

The following definitions of ‘strongly separated’ and ‘almost spanning’ sets are inspired by Pfister and Sullivan and designed for use in the setting of maps with the almost specification property.

Definition 3.1. Let $Z \subseteq X$. For $m < n$, a set S is $(n, -m, \epsilon)$ separated for Z if $S \subset Z$ and for every $\Lambda \in I(n, -m)$, we have $d_\Lambda(x, y) > \epsilon$ for every $x, y \in S$. We define a set S to be $(g; n, \epsilon)$ separated if it is $(n, -g(n, \epsilon), \epsilon)$ separated. Equivalently, S is $(g; n, \epsilon)$ separated if for every $x, y \in S$

$$\#\{j \in \{0, \dots, n - 1\} : d(f^j x, f^j y) > \epsilon\} > g(n, \epsilon).$$

We think of an $(n, -m, \epsilon)$ separated set to be ‘a set which remains (n, ϵ) separated when you permit m mistakes’. In particular, a set S which is $(g; n, \epsilon)$ separated is (n, ϵ) separated in the usual sense. We define the natural dual notion of a $(g; n, \epsilon)$ spanning set.

Definition 3.2. For $m < n$, a set $S \subset Z$ is $(n, -m, \epsilon)$ spanning for Z if for all $x \in Z$, there exists $y \in S$ and $\Lambda \in I(n, -m)$ such that $d_\Lambda(x, y) \leq \epsilon$. Note that Λ depends on x and an (n, ϵ) spanning set is always $(n, -m, \epsilon)$ spanning. We define a set S to be $(g; n, \epsilon)$ spanning if it is $(n, -g(n, \epsilon), \epsilon)$ spanning.

We think of an $(n, -m, \epsilon)$ spanning set to be ‘a set which requires up to m mistakes to be (n, ϵ) spanning’. Let

$$\begin{aligned} s_n(g; Z, \epsilon) &= \sup\{\#S : S \text{ is } (g; n, \epsilon) \text{ separated for } Z\}, \\ r_n(g; Z, \epsilon) &= \inf\{\#S : S \text{ is } (g; n, \epsilon) \text{ spanning for } Z\}, \\ s_n(Z, \epsilon) &= \sup\{\#S : S \text{ is } (n, \epsilon) \text{ separated for } Z\}, \\ r_n(Z, \epsilon) &= \inf\{\#S : S \text{ is } (n, \epsilon) \text{ spanning for } Z\}. \end{aligned}$$

Lemma 3.3. Let g be any mistake function and let $h(n, \epsilon) = 2g(n, \epsilon/2)$. We have

- (1) $r_n(g; Z, \epsilon) \leq s_n(g; Z, \epsilon) \leq s_n(Z, \epsilon)$,
- (2) $s_n(h; Z, 2\epsilon) \leq r_n(g; Z, \epsilon) \leq r_n(Z, \epsilon) \leq s_n(Z, \epsilon)$.

Proof. Suppose that S is a $(g; n, \epsilon)$ separated set for Z of maximum cardinality such that S is not $(g; n, \epsilon)$ spanning. We can find

$$z \in Z \setminus \bigcup_{\Lambda \in I(g; n, \epsilon)} \bigcup_{x \in S} \overline{B_\Lambda(x, \epsilon)} = \bigcap_{\Lambda \in I(g; n, \epsilon)} \left(Z \setminus \bigcup_{x \in S} \overline{B_\Lambda(x, \epsilon)} \right).$$

Since $d_\Lambda(x, z) > \epsilon$ for all $x \in S$ and $\Lambda \in I(g; n, \epsilon)$, then $S \cup \{z\}$ is a $(g; n, \epsilon)$ separated set, which contradicts the maximality of S . Thus, every $(g; n, \epsilon)$ separated set of maximal cardinality is $(g; n, \epsilon)$ spanning.

For (2), suppose E is $(h; n, 2\epsilon)$ separated and F is $(g; n, \epsilon)$ spanning for Z . Define $\phi : E \mapsto F$ by choosing for each $x \in E$ some $\phi(x) \in F$ and some $\Lambda_x \in I(g; n, \epsilon)$ such that $d_{\Lambda_x}(x, \phi(x)) \leq \epsilon$. Suppose $x \neq y$. Let $\Lambda = \Lambda_x \cap \Lambda_y$. Since $\#\Lambda \geq n - 2g(n, \epsilon)$, then $\Lambda \in I(h; n, 2\epsilon)$. We have $d_\Lambda(\phi(x), \phi(y)) > 0$ and thus $\phi(x) \neq \phi(y)$. Thus ϕ is injective and hence $|E| \leq |F|$. □

Theorem 3.4 (Modified Katok entropy formula). Let (X, d) be a compact metric space, $f : X \mapsto X$ be a continuous map and μ be an ergodic invariant measure. For

$\gamma \in (0, 1)$ and any mistake function g , we have

$$h_\mu = \lim_{\epsilon \rightarrow 0} \liminf_{n \rightarrow \infty} \frac{1}{n} \log(\inf\{r_n(g; Z, \epsilon) : Z \subset X, \mu(Z) \geq 1 - \gamma\}).$$

The formula remains true if we replace the \liminf by \limsup and/or $r_n(g; Z, \epsilon)$ by $s_n(g; Z, \epsilon)$. The value taken by the \liminf can be bounded below independent of the choice of mistake function g .

Proof. Since $r_n(Z, \epsilon) \geq r_n(g; Z, \epsilon)$, it follows from the original Katok entropy formula that the expression on the right hand side is less than or equal to h_μ (this is the easier inequality to prove directly any way). To prove the opposite inequality, we give a method inspired by the proof of theorem A2.1 of [18].

For any $\eta > 0$, there exists $\delta, 0 < \delta \leq \eta$, a finite Borel partition $\xi = \{C_1, \dots, C_m\}$ and a finite open cover $\mathcal{U} = \{U_1, \dots, U_k\}$ of X where $k \geq m$ with the following properties:

- (1) $\text{Diam}(U_i) \leq \eta, \text{Diam}(C_j) \leq \eta$ for all $i = 1, \dots, m, j = 1, \dots, k$,
- (2) $\overline{U}_i \subset C_i$ for all $i = 1, \dots, m$,
- (3) $\mu(C_i \setminus U_i) \leq \delta$ for all $i = 1, \dots, m$ and $\mu(\bigcup_{i=m+1}^k U_i) \leq \delta$,
- (4) $2\delta \log m \leq \eta$.

This is a consequence of the regularity of the measure μ . Fix η so $1 - \gamma > \eta > 0$ and take the corresponding number δ , covering \mathcal{U} and partition ξ . Fix $Z \subset X$ with $\mu(Z) > 1 - \gamma$. Let $t_n(x)$ denote the number of $l, 0 \leq l \leq n - 1$ for which $f^l(x) \in \bigcup_{i=m+1}^k U_i$. Let $\xi_n = \bigvee_{i=0}^{n-1} f^{-i}\xi$ and $C_{\xi_n}(x)$ denote the member of the partition ξ_n to which x belongs.

Lemma 3.5. *There exists a set $A \subset Z$ and $N > 0$ with $\mu(A) \geq \mu(Z) - \delta$ such that for every $x \in A$ and $n \geq N$*

- (1) $t_n(x) \leq 2\delta n$,
- (2) $\mu(C_{\xi_n}(x)) \leq \exp(-(h_\mu(f, \xi) - \delta)n)$.

Proof. Let χ be the characteristic function of $\bigcup_{i=m+1}^k U_i$. We can write $t_n(x) = \sum_{i=0}^{n-1} \chi(f^i x)$. By Birkhoff's ergodic theorem and Egorov's theorem, we can find a set $A_1 \subset X$ with $\mu(A_1) \geq \mu(Z) - \frac{\delta}{2}$ such that for $x \in A_1$, we have uniform convergence

$$n^{-1}t_n(x) = \frac{1}{n} \sum_{i=0}^{n-1} \chi(f^i x) \rightarrow \int \chi d\mu = \mu\left(\bigcup_{i=m+1}^k U_i\right) \leq \delta.$$

Choose N_1 such that if $n \geq N_1$ and $x \in A_1$, then $t_n(x) \leq 2\delta n$.

By the Shannon-McMillan-Breiman theorem and Egorov's theorem, we can find a set $A_2 \subset X$ with $\mu(A_2) \geq \mu(Z) - \frac{\delta}{2}$ such that for $x \in A_1$, we have uniform convergence

$$-\frac{1}{n} \log \mu(C_{\xi_n}(x)) \rightarrow h_\mu(f, \xi).$$

There exists N_2 such that if $n \geq N_2$ and $x \in A_2$, then $-\frac{1}{n} \log \mu(C_{\xi_n}(x)) \leq h(f, \xi) + \delta$. Set $A = A_1 \cap A_2$ and $N = \max\{N_1, N_2\}$ and the lemma is proved. \square

Let ξ_n^* be the collection of elements C_{ξ_n} of the partition ξ_n for which $C_{\xi_n} \cap A \neq \emptyset$. Then for $n \geq N$, using property (2) of A ,

$$\begin{aligned} \#\xi_n^* &\geq \sum_{C \in \xi_n^*} \mu(C) \exp\{n(h_\mu(f, \xi) - \delta)\} \\ &\geq \mu(A) \exp\{n(h_\mu(f, \xi) - \delta)\}. \end{aligned}$$

Let 2ϵ be a Lebesgue number for \mathcal{U} and let S be $(g; n, \epsilon)$ spanning for Z . We have $Z \subseteq \bigcup_{x \in S} \bigcup_{\Lambda \in I(g; n, \epsilon)} \overline{B}_\Lambda(x, \epsilon)$. Let us fix $B = \overline{B}_\Lambda(x, \epsilon)$ for some $x \in S$ and $\Lambda \in I(g; n, \epsilon)$. Let ξ_Λ be the partition $\bigvee_{i \in \Lambda} f^{-i}\xi$. We estimate the number $p(B, \xi_\Lambda)$ of elements of the partition ξ_Λ which have non-empty intersection with $A \cap B$.

Since 2ϵ is a Lebesgue number for \mathcal{U} , then $\overline{B}(f^j x, \epsilon) \subset U_{i_j}$ for some $U_{i_j} \in \mathcal{U}$. If $i_j \in \{1, \dots, m\}$, then $f^{-j}(U_{i_j}) \subset f^{-j}(C_{i_j})$. If $i_j \in \{m+1, \dots, k\}$, then anything up to m sets of the form $f^{-j}(C_{i_j})$ may have non-empty intersection with $f^{-j}(U_{i_j})$. It follows, using property (1) of A , that

$$p(B, \xi_\Lambda) \leq m^{2\delta n} = \exp(2\delta n \log m).$$

The number $p(B, \xi_n)$ of elements of the partition ξ_n which have non-empty intersection with both A and B satisfies

$$p(B, \xi_n) \leq p(B, \xi_\Lambda) m^{g(n, \epsilon)} \leq \exp\{(2\delta n + g(n, \epsilon)) \log m\}.$$

It follows that

$$\#\xi_n^* \leq \sum_{x \in S} \sum_{\Lambda \in I(g; n, \epsilon)} p(\overline{B}_\Lambda(x, \epsilon), \xi_n) \leq G_{n, \epsilon} \#S \exp\{(2\delta n + g(n, \epsilon)) \log m\},$$

where $G_{n, \epsilon} = \sum_{i \leq g(n, \epsilon)} \binom{n}{i}$. Rearranging, we have

$$\frac{1}{n} \log \#S \geq h_\mu(f, \xi) - \delta - \left(2\delta + \frac{g(n, \epsilon)}{n}\right) \log m - \frac{1}{n} \log G_{n, \epsilon}.$$

A short combinatorial argument (see [21, Lemma 2.1]) shows that $\frac{1}{n} \log G_{n, \epsilon} \rightarrow 0$. Since $2\delta \log m < \eta$, $\text{Diam}(\xi) < \eta$, $\delta < \eta$, $\frac{g(n, \epsilon)}{n} \rightarrow 0$, and η was arbitrary, we are done. \square

As a corollary, we have a version of Theorem 3.4 for topological entropy (which we do not use).

Theorem 3.6. *Let (X, d) be a compact metric space and $f : X \mapsto X$ be a continuous map. We have*

$$h_{top}(f) = \lim_{\epsilon \rightarrow 0} \liminf_{n \rightarrow \infty} \frac{1}{n} \log r_n(g; X, \epsilon).$$

The formula remains true if we replace the \liminf by \limsup and/or $r_n(g; X, \epsilon)$ by $s_n(g; X, \epsilon)$. The value taken by the \liminf (or \limsup) is independent of the choice of mistake function g .

3.1. Topological entropy for non-compact sets. We consider the Bowen definition of topological entropy for non-compact sets. Following Pesin and Pitskel [19], we give a definition which is suitable for maps which have discontinuities. Suppose X is a compact metric space, Y is a (generally non-compact) subset of X and $f : Y \mapsto Y$ is continuous. When $f : X \mapsto X$ is continuous (which is the setting

of our main theorem), we set $Y = X$. In our final section, when we consider the β -transformation f_β , we set

$$Y = X \setminus \{\beta^{-i} : i \in \mathbb{N}\} = X \setminus \bigcup_i f_\beta^{-i}(0).$$

Let $Z \subset Y$ be an arbitrary Borel set, not necessarily compact or invariant. We consider finite and countable collections of the form $\Gamma = \{B_{n_i}(x_i, \epsilon)\}_i$. For $\alpha \in \mathbb{R}$, we define the following quantities:

$$Q(Z, \alpha, \Gamma) = \sum_{B_{n_i}(x_i, \epsilon) \in \Gamma} \exp -\alpha n_i,$$

$$M(Z, \alpha, \epsilon, N) = \inf_{\Gamma} Q(Z, \alpha, \Gamma),$$

where the infimum is taken over all finite or countable collections of the form $\Gamma = \{B_{n_i}(x_i, \epsilon)\}_i$ with $x_i \in X$ such that Γ covers Z and $n_i \geq N$ for all $i = 1, 2, \dots$. Define

$$m(Z, \alpha, \epsilon) = \lim_{N \rightarrow \infty} M(Z, \alpha, \epsilon, N).$$

The existence of the limit is guaranteed since the function $M(Z, \alpha, \epsilon, N)$ does not decrease with N . By standard techniques, we can show the existence of

$$h_{top}(Z, \epsilon) := \inf\{\alpha : m(Z, \alpha, \epsilon) = 0\} = \sup\{\alpha : m(Z, \alpha, \epsilon) = \infty\}.$$

Definition 3.7. The topological entropy of Z is given by

$$h_{top}(Z) = \lim_{\epsilon \rightarrow 0} h_{top}(Z, \epsilon).$$

See [18] for verification that the quantities $h_{top}(Z, \epsilon)$ and $h_{top}(Z)$ are well defined. When $X = Y$, we denote the topological entropy of the dynamical system (X, f) by $h_{top}(f)$ and we note that $h_{top}(X) = h_{top}(f)$. We sometimes write $h_{top}(Z, f)$ in place of $h_{top}(Z)$ when we wish to emphasise the dependence on f .

3.2. Topological entropy for shift spaces. Let Σ be a one-sided shift space (i.e., a closed, σ -invariant subset of a full one-sided shift on finitely many symbols). For shift spaces, the definition of topological entropy can be simplified and we introduce notation that reflects this. For $x = (x_i)_{i=1}^\infty$, let $C_n(x) = \{y \in \Sigma : x_i = y_i \text{ for } i = 1, \dots, n\}$. Let $Z \subset \Sigma$ be an arbitrary Borel set, not necessarily compact or invariant. We consider finite and countable collections of the form $\Gamma = \{C_{n_i}(x_i)\}_i$. For $s \in \mathbb{R}$, we define the quantities

$$Q(Z, s, \Gamma) = \sum_{C_{n_i}(x_i) \in \Gamma} \exp -s n_i,$$

$$M(Z, s, N) = \inf_{\Gamma} Q(Z, s, \Gamma),$$

where the infimum is taken over all finite or countable collections of the form $\Gamma = \{C_{n_i}(x_i)\}_i$ with $x_i \in \Sigma$ such that Γ covers Z and $n_i \geq N$ for all $i = 1, 2, \dots$. Define

$$m(Z, s) = \lim_{N \rightarrow \infty} M(Z, s, N).$$

The existence of the limit is guaranteed since the function $M(Z, s, N)$ does not decrease with N .

Lemma 3.8. *The topological entropy of $Z \subset \Sigma$ is given by*

$$h_{top}(Z) := \inf\{s : m(Z, s) = 0\} = \sup\{s : m(Z, s) = \infty\}.$$

The proof, which we omit, follows from the fact that every open ball $B(x, \epsilon)$ in Σ is a set of the form $C_n(x)$.

4. MAIN RESULT

Theorem 4.1. *Let (X, d) be a compact metric space and $f : X \mapsto X$ be a continuous map with the almost specification property. Assume that $\varphi \in C(X)$ satisfies $\inf_{\mu \in \mathcal{M}_f(X)} \int \varphi d\mu < \sup_{\mu \in \mathcal{M}_f(X)} \int \varphi d\mu$. Let $\widehat{X}(\varphi, f)$ be the irregular set for φ defined as in (2.1), then $h_{top}(\widehat{X}(\varphi, f)) = h_{top}(f)$.*

We remark that $\widehat{X}(\varphi, f) \neq \emptyset$ is a sufficient condition on φ for the theorem to apply.

Proof. Let us fix a small $\gamma > 0$, and take ergodic measures μ_1 and μ_2 such that

- (1) $h_{\mu_i} > h_{top}(f) - \gamma$ for $i = 1, 2$,
- (2) $\int \varphi d\mu_1 \neq \int \varphi d\mu_2$.

That we are able to choose μ_i to be ergodic is a slightly subtle point. Let μ_1 be ergodic and satisfy $h_{\mu_1} > h_{top}(f) - \gamma/3$. Let $\nu \in \mathcal{M}_f(X)$ satisfy $\int \varphi d\mu_1 \neq \int \varphi d\nu$. Let $\nu' = t\mu_1 + (1-t)\nu$ where $t \in (0, 1)$ is chosen sufficiently close to 1 so that $h_{\nu'} > h_{top}(f) - 2\gamma/3$. By [21], when f has the almost specification property, we can find a sequence of ergodic measures $\nu_n \in \mathcal{M}_f(X)$ such that $h_{\nu_n} \rightarrow h_{\nu'}$ and $\nu_n \rightarrow \nu'$ in the weak-* topology. Therefore, we can choose a measure belonging to this sequence which we call μ_2 which satisfies $h_{\mu_2} > h_{top}(f) - \gamma$ and $\int \varphi d\mu_1 \neq \int \varphi d\mu_2$. (We could avoid the use of the result from [21] by giving a self-contained proof along the lines of the ‘modified construction’ in [30]. We do not do so in the interest of brevity.) Choose $\delta > 0$ sufficiently small so

$$\left| \int \varphi d\mu_1 - \int \varphi d\mu_2 \right| > 4\delta.$$

Let $\rho : \mathbb{N} \mapsto \{1, 2\}$ be given by $\rho(k) = (k+1) \pmod{2} + 1$. Choose a strictly decreasing sequence $\delta_k \rightarrow 0$ with $\delta_1 < \delta$ and a strictly increasing sequence $l_k \rightarrow \infty$ so the set

$$(4.1) \quad Y_k := \left\{ x \in X' : \left| \frac{1}{n} S_n \varphi(x) - \int \varphi d\mu_{\rho(k)} \right| < \delta_k \text{ for all } n \geq l_k \right\}$$

satisfies $\mu_{\rho(k)}(Y_k) > 1 - \gamma$ for every k .

The following lemma follows readily from Theorem 3.4.

Lemma 4.2. *Define mistake functions $h_k(n, \epsilon) := 2g(n, \epsilon/2^k)$. For any sufficiently small $\epsilon > 0$, we can find a sequence $n_k \rightarrow \infty$ and a countable collection of finite sets \mathcal{S}_k so that each \mathcal{S}_k is an $(h_k; n_k, 4\epsilon)$ separated set for Y_k and satisfies*

$$\#\mathcal{S}_k \geq \exp(n_k(h_{top}(f) - 4\gamma)).$$

Furthermore, the sequence n_k can be chosen so that $n_k \geq l_k$, $n_k > N(h_k, \epsilon)$ and $h_k(n_k, \epsilon)/n_k \rightarrow 0$.

We choose ϵ sufficiently small and fix all the ingredients provided by Lemma 4.2. Our strategy is to construct a certain fractal $F \subset \widehat{X}(\varphi, f)$, on which we can define a sequence of measures suitable for an application of the following result of Takens and Verbitskiy [28].

Theorem 4.3 (Entropy Distribution Principle). *Let $f : X \mapsto X$ be a continuous transformation. Let $Z \subseteq X$ be an arbitrary Borel set. Suppose there exists $\epsilon > 0$ and $s \geq 0$ such that one can find a sequence of Borel probability measures μ_k , a constant $K > 0$ and an integer N satisfying*

$$\limsup_{k \rightarrow \infty} \mu_k(B_n(x, \epsilon)) \leq Ke^{-ns}$$

for every ball $B_n(x, \epsilon)$ such that $B_n(x, \epsilon) \cap Z \neq \emptyset$ and $n \geq N$. Furthermore, assume that at least one limit measure ν of the sequence μ_k satisfies $\nu(Z) > 0$. Then $h_{top}(Z, \epsilon) \geq s$.

4.1. Construction of the fractal F . Let us choose a sequence with $N_0 = 0$ and N_k increasing to ∞ sufficiently quickly so that

$$(4.2) \quad \lim_{k \rightarrow \infty} \frac{n_{k+1}}{N_k} = 0, \quad \lim_{k \rightarrow \infty} \frac{N_1 n_1 + \dots + N_k n_k}{N_{k+1}} = 0.$$

Let $\underline{x}_i = (x_1^i, \dots, x_{N_i}^i) \in S_i^{N_i}$. For any $(\underline{x}_1, \dots, \underline{x}_k) \in S_1^{N_1} \times \dots \times S_k^{N_k}$, by the almost specification property, we have

$$B(\underline{x}_1, \dots, \underline{x}_k) := \bigcap_{i=1}^k \bigcap_{j=1}^{N_i} f^{-\sum_{l=0}^{i-1} N_l n_l - (j-1)n_i} B_{n_i}(g; x_j^i, \frac{\epsilon}{2^i}) \neq \emptyset.$$

We define F_k by

$$F_k = \{\overline{B(\underline{x}_1, \dots, \underline{x}_k)} : (\underline{x}_1, \dots, \underline{x}_k) \in S_1^{N_1} \times \dots \times S_k^{N_k}\}.$$

Note that F_k is compact and $F_{k+1} \subseteq F_k$. Define $F = \bigcap_{k=1}^{\infty} F_k$.

Lemma 4.4. *For any $p \in F$, the sequence $\frac{1}{t_k} \sum_{i=0}^{t_k-1} \varphi(f^i(p))$ diverges, where $t_k = \sum_{i=0}^k N_i n_i$.*

Proof. Choose $p \in F$ and let $p_k := f^{t_k-1}p$. Then there exists $(x_1^k, \dots, x_{N_k}^k) \in S_k^{N_k}$ such that

$$p_k \in \bigcap_{j=1}^{N_k} f^{-(j-1)n_k} \overline{B_{n_k}(g; x_j^k, \epsilon/2^k)}.$$

For $c > 0$, let $\text{Var}(\varphi, c) := \sup\{|\varphi(x) - \varphi(y)| : d(x, y) < c\}$. We have

$$S_{n_k N_k} \varphi(p_k) \leq \sum_{j=1}^{N_k} S_{n_k} \varphi(x_j^k) + n_k N_k \text{Var}(\varphi, \epsilon/2^k) + N_k g(n_k, \epsilon/2^k) \|\varphi\|$$

and hence

$$\frac{1}{n_k N_k} S_{n_k N_k} \varphi(p_k) \leq \int \varphi d\mu_{\rho(k)} + \delta_k + \text{Var}(\varphi, \epsilon/2^k) + \frac{1}{n_k} g(n_k, \epsilon/2^k).$$

It follows that

$$\left| \frac{1}{n_k N_k} S_{n_k N_k} \varphi(p_k) - \int \varphi d\mu_{\rho(k)} \right| \rightarrow 0.$$

We can use the fact that $\frac{n_k N_k}{t_k} \rightarrow 1$ to prove that

$$\left| \frac{1}{n_k N_k} S_{n_k N_k} \varphi(p_k) - \frac{1}{t_k} S_{t_k} \varphi(p) \right| \rightarrow 0,$$

and the result follows. □

4.2. **Construction of a special sequence of measures μ_k .** We must first undertake an intermediate construction. For each $\underline{x} = (\underline{x}_1, \dots, \underline{x}_k) \in S_1^{N_1} \times \dots \times S_k^{N_k}$, we choose one point $z = z(\underline{x})$ such that

$$z \in B(\underline{x}_1, \dots, \underline{x}_k).$$

Let \mathcal{T}_k be the set of all points constructed in this way. We show that points constructed in this way are distinct and thus $\#\mathcal{T}_k = \#S_1^{N_1} \dots \#S_k^{N_k}$.

Lemma 4.5. *Let \underline{x} and \underline{y} be distinct members of $S_1^{N_1} \times \dots \times S_k^{N_k}$. Then $z_1 := z(\underline{x})$ and $z_2 := z(\underline{y})$ are distinct points. Thus $\#\mathcal{T}_k = \#S_1^{N_1} \dots \#S_k^{N_k}$.*

Proof. Since $\underline{x} \neq \underline{y}$, there exists i, j so $x_j^i \neq y_j^i$. We have $\Lambda_1, \Lambda_2 \in I(g; n_i, \epsilon/2^i)$ such that

$$d_{\Lambda_1}(x_j^i, f^a z_1) < \frac{\epsilon}{2^i} \text{ and } d_{\Lambda_2}(y_j^i, f^a z_2) < \frac{\epsilon}{2^i},$$

where $a = \sum_{l=0}^{i-1} N_l n_l + (j-1)n_i$. Let $\Lambda = \Lambda_1 \cap \Lambda_2$. Since $\Lambda \in I(2g; n_i, \epsilon/2^i)$, we have $d_\Lambda(x_j^i, y_j^i) > 4\epsilon$. Using these inequalities, we have $d_\Lambda(f^a z_1, f^a z_2) > 3\epsilon$. \square

We now define the measures on F which yield the required estimates for the Entropy Distribution Principle. We define, for each k , an atomic measure centred on \mathcal{T}_k . Precisely, let

$$\nu_k := \sum_{z \in \mathcal{T}_k} \delta_z.$$

We normalise ν_k to obtain a sequence of probability measures μ_k , i.e., we let $\mu_k := \frac{1}{\#\mathcal{T}_k} \nu_k$.

Lemma 4.6. *Suppose μ is a limit measure of the sequence of probability measures μ_k . Then $\mu(F) = 1$.*

Proof. For any fixed l and all $p \geq 0$, $\mu_{l+p}(F_l) = 1$ since $\mu_{l+p}(F_{l+p}) = 1$ and $F_{l+p} \subseteq F_l$. Suppose $\mu = \lim_{k \rightarrow \infty} \mu_{l_k}$ for some $l_k \rightarrow \infty$, then $\mu(F_l) \geq \limsup_{k \rightarrow \infty} \mu_{l_k}(F_l) = 1$. It follows that $\mu(F) = \lim_{l \rightarrow \infty} \mu(F_l) = 1$. \square

In fact, the measures μ_k converge. However, by using our version of the Entropy Distribution Principle, we do not need to use this fact and so we omit the proof (which goes like lemma 5.4 of [28]).

Let $\mathcal{B} := B_n(q, \epsilon)$ be an arbitrary ball which intersects F . Let k be the unique number which satisfies $t_k \leq n < t_{k+1}$. Let $j \in \{0, \dots, N_{k+1} - 1\}$ be the unique number so

$$t_k + n_{k+1}j \leq n < t_k + n_{k+1}(j+1).$$

We assume that $j \geq 1$ and leave the details of the simpler case $j = 0$ to the reader. The following lemma reflects the restriction on the number of points that can be in $\mathcal{B} \cap \mathcal{T}_{k+p}$.

Lemma 4.7. *For $p \geq 1$, $\mu_{k+p}(\mathcal{B}) \leq (\#\mathcal{T}_k)^{-1}(\#\mathcal{S}_{k+1})^{-j}$.*

Proof. First we show that $\mu_{k+1}(\mathcal{B}) \leq (\#\mathcal{T}_k)^{-1}(\#\mathcal{S}_{k+1})^{-j}$. We require an upper bound for the number of points in $\mathcal{T}_{k+1} \cap \mathcal{B}$. If $\mu_{k+1}(\mathcal{B}) > 0$, then $\mathcal{T}_{k+1} \cap \mathcal{B} \neq \emptyset$. Let $z = z(\underline{x}, \underline{x}_{k+1}) \in \mathcal{T}_{k+1} \cap \mathcal{B}$ where $\underline{x} \in S_1^{N_1} \times \dots \times S_k^{N_k}$ and $\underline{x}_{k+1} \in S_{k+1}^{N_{k+1}}$. Let

$$\mathcal{A}_{\underline{x}; x_1, \dots, x_j} = \{z(\underline{y}, y_{k+1}) \in \mathcal{T}_{k+1} : x_1 = y_1, \dots, x_j = y_j\}.$$

We suppose that $z' = z(\underline{y}, y_{k+1}) \in \mathcal{T}_{k+1} \cap \mathcal{B}$ and show that $z' \in \mathcal{A}_{\underline{x}; x_1, \dots, x_j}$. We have $d_n(z, z') < 2\epsilon$ and we show that this implies $x_l = y_l$ for $l \in \{1, 2, \dots, j\}$ (the

proof that $\underline{x} = \underline{y}$ is similar). Suppose that $y_l \neq x_l$ and let $a_l = t_k + (l - 1)(n_{k+1})$. There exists $\Lambda_1, \Lambda_2 \in I(g; n_{k+1}, \epsilon/2^{k+1})$ such that

$$d_{\Lambda_1}(f^{a_l} z, x_l) < \frac{\epsilon}{2^{k+1}} \text{ and } d_{\Lambda_2}(f^{a_l} z', y_l) < \frac{\epsilon}{2^{k+1}}.$$

Let $\Lambda = \Lambda_1 \cap \Lambda_2$. Since $\Lambda \in I(2g; n_{k+1}, \epsilon/2^{k+1})$, we have $d_\Lambda(x_l, y_l) > 4\epsilon$. We have

$$\begin{aligned} d_n(z, z') &\geq d_\Lambda(f^{a_l} z, f^{a_l} z') \\ &\geq d_\Lambda(x_l, y_l) - d_\Lambda(f^{a_l} z, x_l) - d_\Lambda(f^{a_l} z', y_l) \geq 3\epsilon, \end{aligned}$$

which is a contradiction. Thus, we have

$$\begin{aligned} \nu_{k+1}(\mathcal{B}) &\leq \#\mathcal{A}_{x; x_1, \dots, x_j} = (\#\mathcal{S}_{k+1})^{N_{k+1}-j}, \\ \mu_{k+1}(\mathcal{B}) &\leq (\#\mathcal{T}_{k+1})^{-1}(\#\mathcal{S}_{k+1})^{N_{k+1}-j} = (\#\mathcal{T}_k)^{-1}(\#\mathcal{S}_{k+1})^{-j}. \end{aligned}$$

Now consider $\mu_{k+p}(\mathcal{B})$. Arguing similarly to above, we have

$$\nu_{k+p}(\mathcal{B}) \leq \#\mathcal{A}_{x; x_1, \dots, x_j} (\#\mathcal{S}_{k+2})^{N_{k+1}} \dots (\#\mathcal{S}_{k+p})^{N_{k+p}}.$$

The desired result follows from this inequality by dividing by $\#\mathcal{T}_{k+p}$. □

By Lemma 4.2, we have

$$\begin{aligned} \#\mathcal{T}_k(\#\mathcal{S}_{k+1})^j &\geq \exp\{(h_{top}(f) - 4\gamma)(N_1 n_1 + N_2 n_2 + \dots + N_k n_k + j n_{k+1})\} \\ &\geq \exp\{(h_{top}(f) - 4\gamma)n\}. \end{aligned}$$

Combining this with the previous lemma gives us

$$\limsup_{l \rightarrow \infty} \mu_l(B_n(q, \epsilon)) \leq \exp\{-n(h_{top}(f) - 4\gamma)\}.$$

Applying the Entropy Distribution Principle, we have

$$h_{top}(F, \epsilon) \geq h_{top}(f) - 4\gamma.$$

Since γ and ϵ were arbitrary and $F \subset \widehat{X}(\varphi, f)$, we have $h_{top}(\widehat{X}(\varphi, f)) = h_{top}(f)$. □

5. THE β -TRANSFORMATION

In this section, let $X = [0, 1)$. For any fixed $\beta > 1$, we consider the β -transformation $f_\beta : X \mapsto X$ given by

$$f_\beta(x) = \beta x \pmod{1}.$$

As reference for the basic properties of the β -transformation, we recommend the introduction of the thesis of Maia [14]. For $\beta \notin \mathbb{N}$, let $b = [\beta]$ and for $\beta \in \mathbb{N}$, let $b = \beta - 1$. We consider the partition into $b + 1$ intervals

$$J_0 = \left[0, \frac{1}{\beta}\right), J_1 = \left[\frac{1}{\beta}, \frac{2}{\beta}\right), \dots, J_b = \left[\frac{b}{\beta}, 1\right).$$

For $x \in [0, 1)$, let $w(x) = (w_j(x))_{j=1}^\infty$ be the sequence given by $w_j(x) = i$ when $f^{j-1}x \in J_i$. We call $w(x)$ the greedy β -expansion of x and we have

$$x = \sum_{j=1}^\infty w_j(x)\beta^{-j}.$$

The β -shift $(\Sigma_\beta, \sigma_\beta)$ is the subshift defined by the closure of all such sequences in $\prod_{i=1}^\infty \{0, \dots, b\}$. Let $w(\beta) = (w_j(\beta))_{j=1}^\infty$ denote the sequence which is the lexicographic supremum of all β -expansions. The sequence $w(\beta)$ satisfies

$$\sum_{j=1}^\infty w_j(\beta)\beta^{-j} = 1,$$

so we call $w(\beta)$ the β -expansion of 1. Parry showed that the set of sequences which belong to Σ_β can be characterised as

$$w \in \Sigma_\beta \iff \sigma^k(w) \leq w(\beta) \text{ for all } k \geq 1,$$

where \leq is taken in the lexicographic ordering [17]. Parry also showed that any sequence w which satisfies $\sigma^k(w) \leq w$ is the β -expansion of 1 for some $\beta > 1$. The β -shift contains every sequence which arises as a greedy β -expansion and an additional point for every x whose β -expansion is finite (i.e., when there exists j so $w_i(x) = 0$ for all $i \geq j$). Thus the map $\pi : \Sigma_\beta \mapsto [0, 1]$ defined by

$$\pi(\underline{w}) = \sum_{j=1}^\infty w_j\beta^{-j}$$

is one-to-one except at the countably many points for which the β -expansion is finite.

Σ_β is typically not a shift of finite type (nor even a shift with specification) and the set of all β -shifts gives a natural and interesting class of subshifts. In the next section, we describe in detail the known results on specification properties for the β -shift. The key fact for our analysis is that every β -shift has the almost specification property [21]. We have (p. 179 of [32]) that $h_{top}(\sigma_\beta) = \log \beta$.

Theorem 5.1. *For $\beta > 1$, let $f_\beta : X \mapsto X$ be the β -transformation, $f_\beta(x) = \beta x \pmod{1}$. Let $\varphi \in C([0, 1])$ and assume that the irregular set for φ is non-empty (i.e., $\widehat{X}(\varphi, f_\beta) \neq \emptyset$), then $h_{top}(\widehat{X}(\varphi, f_\beta)) = \log \beta$.*


Proof. Let Σ'_β denote the set of sequences which arise as β -expansions. Recall that $\Sigma_\beta \setminus \Sigma'_\beta$ is a countable set and the restriction of π to Σ'_β is a homeomorphism satisfying $\pi \circ \sigma_\beta = f_\beta \circ \pi$. Thus, if $Z \in \Sigma_\beta$ and $Z' := Z \cap \Sigma'_\beta$, we have

$$h_{top}(Z, \sigma_\beta) = h_{top}(Z', \sigma_\beta) = h_{top}(\pi(Z'), f_\beta).$$

Suppose $\varphi : [0, 1] \mapsto \mathbb{R}$ satisfies $\widehat{X}(\varphi, f_\beta) \neq \emptyset$. Let $x \in \widehat{X}(\varphi, f_\beta)$ and let $w(x)$ be its β -expansion. We let $\overline{\varphi} \in C(\Sigma_\beta)$ be the unique continuous function which satisfies $\overline{\varphi} = \varphi \circ \pi$ on Σ'_β (this exists because we assumed φ to be continuous on $[0, 1]$). It is clear that $w(x) \in \widehat{\Sigma}_\beta(\overline{\varphi}, \sigma_\beta)$. Since the dynamical system $(\Sigma_\beta, \sigma_\beta)$ satisfies the almost specification property, it follows from Theorem 5.1 that $h_{top}(\widehat{\Sigma}_\beta(\overline{\varphi}, \sigma_\beta)) = \log \beta$. Since $\pi(\widehat{\Sigma}_\beta(\overline{\varphi}, \sigma_\beta) \cap \Sigma'_\beta) = \widehat{X}(\varphi, f_\beta)$, it follows that $h_{top}(\widehat{X}(\varphi, f_\beta)) = \log \beta$. \square

5.1. β -transformations and specification properties. There is a simple presentation of Σ_β by a labelled graph \mathcal{G}_β due to Blanchard and Hansel [5]. We describe the construction of \mathcal{G}_β when the β -expansion of 1 is not eventually periodic. We refer the reader to [21] for the slightly different construction required when the β -expansion of 1 is eventually periodic (in this case, Σ_β is a sofic shift [4] and therefore has specification).

Let v_1, v_2, \dots be a countable set of vertices. We draw a directed edge from v_i to v_{i+1} and label it with the value $w_i(\beta)$ for all $i \geq 1$. If $w_i(\beta) \geq 1$, we draw a directed edge from v_i to v_1 labelled with the value 0. If $b = 1$, the construction is complete. If $b > 1$, then for all $j \in \{2, \dots, b\}$ and all $w_i(\beta) \geq j$, we draw a directed edge from v_i to v_1 labelled with the value $j - 1$. Note that if $w_i(\beta) = 0$, the only edge which starts at v_i is the edge from v_i to v_{i+1} labelled by 0, and if $w_i(\beta) \neq 0$ there is always an edge from v_i to v_1 . We have $w \in \Sigma_\beta$ iff w labels an infinite path of directed edges of \mathcal{G}_β which starts at the vertex v_1 . The following figure depicts part of the graph \mathcal{G}_β for a value of β satisfying $(w_j(\beta))_{j=1}^6 = (2, 0, 1, 0, 0, 1)$.



An arbitrary subshift Σ on $b + 1$ symbols is a closed shift-invariant subset of $\prod_{i=1}^{\infty} \{0, \dots, b\}$. We define v to be admissible word of length $n \geq 1$ for Σ if there exists $x \in \Sigma$ such that $v = (x_1, \dots, x_n)$. The specification property of Definition 2.8 can easily be seen to be equivalent to the following property in the case of an arbitrary subshift.

Definition 5.2. A subshift Σ has the specification property if there exists $M > 0$ such that for any two admissible words w_1 and w_2 , there exists a word w of length less than M such that $w_1 w w_2$ is an admissible word.

We now return to the β -shift Σ_β . Define

$$z_n(\beta) = \min\{i \geq 0 : w_{n+i}(\beta) \neq 0\}.$$

Equivalently, $z_n(\beta) + 1$ is the minimum number of edges required to travel from v_n to v_1 . The β -shift fails to have the specification property iff ‘blocks of consecutive zeroes in the β expansion of 1 have unbounded length’, i.e., if $z_n(\beta)$ is unbounded [4]. Consider concatenations of the word $c_n := (w_1(\beta), \dots, w_n(\beta))$ with some other admissible word v . We can see from the graph \mathcal{G}_β that the length of the shortest word w such that $c_n w v$ is an admissible word is $z_n(\beta)$ (the word w is a block of zeroes of length $z_n(\beta)$). Now for $x \in \Sigma_\beta$, we define $z_n(x) \geq 0$ to be the length of the shortest word w required so that for any admissible word v , $(x_1, \dots, x_n) w v$ is an admissible word. Note that for all $x \in \Sigma_\beta$, $z_n(x) \leq \max\{z_i(\beta) : 1 \leq i \leq n\}$. Thus Σ_β has specification iff $z_n(\beta)$ is bounded. Buzzi shows that the set of β for which this situation occurs has Lebesgue measure 0.

Pfister and Sullivan [21] used the graph \mathcal{G}_β to observe that every β -shift has the almost specification property. Their strategy is to ‘jump ship’ on the last non-zero entry of an admissible word. More precisely, every β -shift has the following property. Given any admissible word w , there is a word w' which differs from w

only by one symbol, such that $w'v$ is admissible for every admissible word v . The modified word w' is given by replacing the last non-zero entry of the word w by a 0. This property is best seen from inspection of the graph \mathcal{G}_β and is the content of proposition 5.1 of [21]. It can easily be seen that this property implies the almost specification property.

5.2. Hausdorff dimension of the irregular set for the β -shift. We equip Σ_β with the metric $d_\beta(x, y) = \frac{1}{\beta^n}$ where n is the smallest integer such that $x_{n+1} \neq y_{n+1}$. Let $C_n(x) = \{y \in \Sigma_\beta : x_i = y_i \text{ for } i = 0, \dots, n-1\}$. For $Z \subset \Sigma_\beta$, we define the diameter of the set to be $\text{Diam}(Z) = |Z| = \sup\{d_\beta(x, y) : x, y \in Z\}$. We have

$$\begin{aligned} \text{Diam}(C_n(w(\beta))) &= \frac{1}{\beta^{n+z_{n+1}(\beta)}}, \\ \frac{1}{\beta^n} &\geq \text{Diam}(C_n(x)) \geq \frac{1}{\beta^{n+z_{n+1}(x)}}. \end{aligned}$$

Because there is no uniform lower bound on the diameter of a cylinder when $z_n(\beta)$ is unbounded, it is plausible that the relation between Hausdorff dimension and topological entropy could not be straightforward. Indeed, because of this potential difficulty, in [21, 22], it seems that Pfister and Sullivan establish a relationship between topological entropy and Hausdorff dimension only for β -shifts which satisfy an additional hypothesis on $w(\beta)$ (which holds for Lebesgue almost every β). In fact, the following lemma tells us that the relationship between Hausdorff dimension and topological entropy is straightforward for every β -shift. While this lemma is surely known to some experts as a folklore theorem and is a corollary of more general results of Climenhaga [8, Theorem 2.4], we include an elementary direct proof to clarify the situation. For Hausdorff dimension, we fix the notation

$$H(Z, \alpha, \delta) = \inf\left\{\sum_i |B(x_i, \delta_i)|^\alpha : Z \subseteq \bigcup_i B(x_i, \delta_i), \delta_i \leq \delta\right\}.$$

$H(Z, \alpha) = \lim_{\delta \rightarrow 0} H(Z, \alpha, \delta)$ and $\text{Dim}_H(Z) = \sup\{\alpha : H(Z, \alpha) = \infty\}$. We sometimes write $\text{Dim}_H(Z, d)$ in place of $\text{Dim}_H(Z)$ when we wish to emphasise the dependence on the metric d . We note that the map $x \mapsto w(x)$ is bi-Lipschitz with respect to the metric d_β and thus for $Z \subset [0, 1)$, $\text{Dim}_H(Z) = \text{Dim}_H(\pi^{-1}(Z), d_\beta)$.

Lemma 5.3. *For arbitrary $Z \subset \Sigma_\beta$, we have $\log \beta \text{Dim}_H(Z) = h_{\text{top}}(Z)$.*

Proof. That $\log \beta \text{Dim}_H(Z) \leq h_{\text{top}}(Z)$ is a standard argument which follows from the fact that $\text{Diam}(C_n(x)) \leq \frac{1}{\beta^n}$. For the other inequality, let $\gamma_N = \beta^{-N}$, and consider a cover of Z by metric balls $B(x_i, \delta_i)$ with $\delta_i < \gamma_N$. We modify the radii of the balls to create a ‘better’ cover. For each δ_i , let n_i be the unique natural number so that $\beta^{-(n_i+1)} < \delta_i \leq \beta^{n_i}$. Then $n_i \geq N$ for all i and $|B(x_i, \delta_i)| \leq \beta^{-(n_i+1)}$. If $|B(x_i, \delta_i)| = \beta^{-(n_i+1)}$, let $\rho_i = \delta_i$ and $m_i = 1$. If not, $|B(x_i, \delta_i)| = \beta^{-(n_i+m_i)}$ for some $m_i \geq 2$. In this case, let $\rho_i = \beta^{-(n_i+m_i-1)}$. We have $\rho_i \leq \delta_i$ and $B(x_i, \delta_i) = B(x_i, \rho_i) = C_{n_i+m_i}(x_i)$. The collection $\Gamma = \{C_{n_i+m_i}(x_i)\}$ covers Z , and thus

$$\begin{aligned} \sum |B(x_i, \delta_i)|^\alpha &= \sum |B(x_i, \rho_i)|^\alpha = \sum \beta^{-(n_i+m_i)\alpha} = \sum \exp(-\alpha(n_i + m_i) \log \beta) \\ &= Q(Z, \alpha \log \beta, \Gamma). \end{aligned}$$

Taking infimums, we have $H(Z, \alpha, \gamma_N) \geq M(Z, \alpha \log \beta, N)$. Taking the limit as $N \rightarrow \infty$, it follows that if $m(Z, \alpha \log \beta) = \infty$, then $H(Z, \alpha) = \infty$ and the inequality follows. \square

We obtain the following theorem as an immediate corollary of Lemma 5.3 and Theorems 4.1 and 5.1.

Theorem 5.4. *For every $\beta > 1$,*

- (1) *If $\varphi \in C(\Sigma_\beta)$ and $\widehat{\Sigma}_\beta(\varphi, \sigma_\beta) \neq \emptyset$, then $\text{Dim}_H(\widehat{\Sigma}_\beta(\varphi, \sigma_\beta)) = 1$,*
- (2) *If $\varphi \in C([0, 1])$ and $\widehat{X}(\varphi, f_\beta) \neq \emptyset$, then $\text{Dim}_H(\widehat{X}(\varphi, f_\beta)) = 1$.*

Remark 5.5. We may also consider the Billingsley dimension Dim_ν of the irregular set (with respect to a reference measure ν) [21, 7]. We remark that when ν is equivalent to Lebesgue measure, then $\text{Dim}_\nu = \text{Dim}_H$. Every β -transformation has an invariant measure ν_β which is equivalent to Lebesgue (and is the measure of maximal entropy). Therefore, it is a corollary of Theorem 5.4 that if $\varphi \in C(\overline{X})$ and $\widehat{X}(\varphi, f_\beta) \neq \emptyset$, then $\text{Dim}_{\nu_\beta}(\widehat{X}(\varphi, f_\beta)) = 1$.

5.3. An alternative approach for β -shifts. We describe an alternative proof of Theorem 5.4 based on approximating β -shifts from inside by n -step shifts of finite type. We discuss the relative merits of the two approaches in §5.4. The key quoted result in this method is a version of Theorem 5.1 in the special case of n -step Markov shifts. We note that the ‘almost specification’ method of proof applies in far greater generality and a self-contained version of the proof described below would be comparable in length.

Proof. Recall that any sequence (a_n) on a finite number of symbols which satisfies $\sigma^k(a_n) \leq (a_n)$ for all $k \geq 0$ arises as $w(\beta)$ for some $\beta > 1$. Fix $\beta > 1$ and write $w_i := w_i(\beta)$. Let $\beta(n)$ be the simple β -number corresponding to the sequence $(w_1, w_2, \dots, w_n, 0, 0, 0, \dots)$. An elementary argument [17] shows that $\beta(n) \rightarrow \beta$. It is clear that $\Sigma_{\beta(n)}$ can be considered to be a subsystem of Σ_β (the subshift $\Sigma_{\beta(n)}$ corresponds to the set of labels of edges of infinite paths that only visit the first n vertices of \mathcal{G}_β).

Now suppose $\varphi \in C(\Sigma_\beta)$ is a function for which the irregular set is non-empty. Then there exists $x, y \in \Sigma_\beta$ such that

$$\lim_{n \rightarrow \infty} \frac{1}{n} S_n \varphi(x) \neq \lim_{n \rightarrow \infty} \frac{1}{n} S_n \varphi(y).$$

Let $\delta > 0$ and $N_1 \in \mathbb{N}$ be such that for $n \geq N_1$,

$$\left| \frac{1}{n} S_n \varphi(x) - \frac{1}{n} S_n \varphi(y) \right| > 4\delta.$$

Pick N_2 sufficiently large that

$$\sup\{|\varphi(w) - \varphi(v)| : w, v \in \Sigma_\beta, w_i = v_i \text{ for } i = 1, \dots, N_2\} < \delta.$$

For any $n \geq N = \max\{N_1, N_2\}$, let us choose $x' \in C_N(x) \cap \Sigma_{\beta(n)}$ and $y' \in C_N(y) \cap \Sigma_{\beta(n)}$. We have for all $m \geq N$,

$$\left| \frac{1}{m} S_m \varphi(x') - \frac{1}{m} S_m \varphi(y') \right| > 2\delta.$$

Thus the restriction of φ to $\Sigma_{\beta(n)}$ does not have trivial spectrum of Birkhoff averages and by Lemma 2.1, our main theorem gives us

$$(5.1) \quad h_{top}(\widehat{\Sigma}_{\beta(n)}(\varphi, \sigma_{\beta(n)})) = h_{top}(\sigma_{\beta(n)}) = \log \beta(n).$$

We remark that $\Sigma_{\beta(n)}$ is an n -step Markov shift (and thus has specification), so formula (5.1) also follows for Hölder continuous φ from theorem 9.3.2 of [2]. Note that $\widehat{\Sigma}_{\beta(n)}(\varphi, \sigma_{\beta(n)}) \subset \widehat{\Sigma}_{\beta}(\varphi, \sigma_{\beta})$.

By Lemma 5.3, any subset $Z \subset \Sigma_{\beta(n)}$ satisfies $\text{Dim}_H(Z, d_{\beta}) = h_{top}(Z)/\log \beta$. In particular, $\text{Dim}_H(\Sigma_{\beta(n)}, d_{\beta}) = \log(\beta(n))/\log \beta$. Thus

$$\text{Dim}_H(\widehat{\Sigma}_{\beta}(\varphi, \sigma_{\beta}), d_{\beta}) \geq \sup \left\{ \text{Dim}_H(\widehat{\Sigma}_{\beta(n)}(\varphi, \sigma_{\beta(n)}), d_{\beta}) \right\} = \sup \left\{ \frac{\log \beta(n)}{\log \beta} \right\} = 1.$$

□

5.4. Some final comments. We give some closing remarks, comparing the ‘almost specification’ method of Theorem 4.1 with the ‘approximation from inside’ method of §5.3.

Remark 5.6. An almost sofic shift [13] is defined to be a shift space Σ for which one can find a sequence of subshifts of finite type Σ_n such that $\Sigma_n \subset \Sigma$ and $\lim_{n \rightarrow \infty} h_{top}(\Sigma_n, \sigma) = h_{top}(\Sigma, \sigma)$. By our previous reasoning, every β -shift is almost sofic. We remark that the proof of Theorem 5.3 shows that if (Σ, σ) is an almost sofic shift, $\varphi \in C(\Sigma)$ and $\widehat{\Sigma}(\varphi, \sigma) \neq \emptyset$, then $h_{top}(\widehat{\Sigma}(\varphi, \sigma)) = h_{top}(\sigma)$.

Remark 5.7. We point out that Theorem 4.1 applies to systems where the approach of §5.3 cannot apply, even in the setting of shift spaces. It is not too difficult to construct a shift space (Σ, σ) such that

- (1) (Σ, σ) has positive topological entropy but no periodic points;
- (2) (Σ, σ) satisfies almost specification but not specification.

We give an idea of the construction. We define $\Sigma := \bigcap_{n \geq 1} \Sigma_n$, where $\Sigma_n \subset \prod_{i=1}^{\infty} \{0, 1\}$ are suitably chosen topologically mixing shifts of finite type satisfying $\Sigma_{n+1} \subset \Sigma_n$ for all $n \geq 1$. For $i \geq 1$, let $\epsilon_i = \frac{1}{2^{i+1}} \log 2$ and N_i be an increasing sequence of integers to be chosen later.

Recall that a finite collection of forbidden words defines a shift of finite type. For a word u and an integer N , we use the notation u^N to denote the concatenation of N copies of u . Let \mathcal{F}_1 denote the set of words

$$\mathcal{F}_1 = \{(1)^{N_1}, (0)^{N_1}\}$$

and define Σ_1 as the shift defined by the forbidden words \mathcal{F}_1 . It follows easily from theorem 7.13 of [32] that if we choose N_1 to be large enough, then $h_{top}(\Sigma_1, \sigma) > \log 2 - \epsilon_1$. It is clear that Σ_1 contains no fixed points.

We define Σ_k inductively as the shift defined by the collection of forbidden words $\bigcup_{i=1}^k \mathcal{F}_i$, where

$$\mathcal{F}_k = \{v^{N_k} : v \text{ is an admissible word of length } k \text{ in } \Sigma_{k-1}\},$$

with N_k chosen large enough so that $h_{top}(\Sigma_k, \sigma) > h_{top}(\Sigma_{k-1}, \sigma) - \epsilon_k$. It is clear that Σ_k contains no periodic points of period less than or equal to k .

It is easily verified that $h_{top}(\Sigma, \sigma) = \lim_{k \rightarrow \infty} h_{top}(\Sigma_k, \sigma) > 0$. Since Σ contains no periodic points, it contains no shifts of finite type. We claim that (Σ, σ) satisfies almost specification but not specification and briefly explain why. The key fact is that if a word w belongs to \mathcal{F}_k , we can change it to a word not in \mathcal{F}_k by changing

any one of its entries (from a 0 to a 1 or vice versa). Furthermore, one can modify w by changing a single entry in at least $|w|(1 - 2/N_1)$ positions so that the new word is not in $\bigcup_{i=1}^k \mathcal{F}_i$. This abundance of choice when we wish to adjust an inadmissible word to make it admissible gives rise to the almost specification property.

It is not hard to find Σ -admissible words u and w_k such that uvw_k is Σ -inadmissible for any word v of length less than or equal to k . Hence, (Σ, σ) does not have specification. However, by changing u and w_k at a small number of entries (relative to the length of u and w_k), we can easily find new words u^* and w_k^* such that $u^*w_k^*$ is Σ -admissible. This leads to the almost specification property.

Remark 5.8. We could study the class of systems (X, f) such that for every $\epsilon > 0$, there exists $n \in \mathbb{N}$ and a compact f^n -invariant subset $Y \subset X$ satisfying

- (1) (Y, f^n) is a topological factor of a shift of finite type;
- (2) $h_{\text{top}}(Y, f^n) > n(h_{\text{top}}(X, f) - \epsilon)$.

We call (Y, f^n) a horseshoe for (X, f) . To study irregular sets for (X, f) , it should suffice to study the intersection of the irregular set with the horseshoe. As demonstrated by our previous remark, systems with almost specification do not necessarily contain any horseshoes, so we could not prove Theorem 4.1 using this approach. Also, theorems on the existence of horseshoes typically require smoothness of the system (see, for example, theorem S.5.9 of [12]), whereas our ‘almost specification’ approach is a topological approach to a topological question. However, we do note that examples exist that do not have specification but where an ‘approximation from inside’ approach could yield results. For example, a continuous interval map which is not mixing contains horseshoes but does not have specification (see corollary 15.2.10 of [12]). Thus, the ‘almost specification’ approach and the ‘approximation from inside’ approach both have their own merits.

Remark 5.9. In [20], Petersen studies some interesting families of shift spaces which first arose in an applied setting. He shows that some of these are almost sofic and some are not. It would be interesting to investigate the almost specification property in this setting.

ACKNOWLEDGEMENTS

This work constitutes part of the author’s Ph.D., which was supported by the EPSRC. The author would like to thank his supervisors, Mark Pollicott and Peter Walters, for many useful discussions and reading draft versions of this work, for which the author is most grateful. The author would also like to thank Thomas Jordan for comments on a draft version of this work and Charles-Edouard Pfister for a useful discussion.

REFERENCES

1. I.S. Baek, L. Olsen, and N. Snigireva, *Divergence points of self-similar measures and packing dimension*, Adv. Math. **214** (2007), no. 1, 267–287. MR2348031 (2009a:28020)
2. L. Barreira, *Dimension and recurrence in hyperbolic dynamics*, Progress in Mathematics, vol. 272, Birkhäuser, 2008. MR2434246 (2009i:37056)
3. L. Barreira and J. Schmeling, *Sets of “non-typical” points have full topological entropy and full Hausdorff dimension*, Israel J. Math. **116** (2000), 29–70. MR1759398 (2002d:37040)
4. A. Bertrand-Mathis, *Développement en base θ* , Bull. Soc. Math. France **114** (1986), 271–323. MR878240 (88e:11067)
5. F. Blanchard and G. Hansel, *Systèmes codés*, Theor. Comput. Sci. **44** (1986), 17–49. MR858689 (88m:68029)

6. J. Buzzi, *Specification on the interval*, Trans. Amer. Math. Soc. **349** (1997), no. 7, 2737–2754. MR1407484 (97i:58043)
7. H. Cajar, *Billingsley dimension in probability spaces*, Lecture Notes in Mathematics, vol. 892, Springer-Verlag, 1981. MR654147 (84a:10055)
8. V. Climenhaga, *Bowen's equation in the non-uniform setting*, To appear in Ergodic Theory Dynam. Systems, 2010.
9. M. Dateyama, *The almost weak specification property for ergodic group automorphisms of abelian groups*, J. Math. Soc. Japan **42** (1990), 341–351. MR1041229 (91b:28014)
10. A. Durner, *Distribution measures and Hausdorff dimensions*, Forum Math. **9** (1997), 687–705. MR1480551 (98i:11060)
11. C. Ercai, T. Küpper, and S. Lin, *Topological entropy for divergence points*, Ergodic Theory Dynam. Systems **25** (2005), no. 4, 1173–1208. MR2158401 (2006c:37014)
12. A. Katok and B. Hasselblatt, *Introduction to the modern theory of dynamical systems*, Encyclopedia of Mathematics and Its Applications, vol. 54, Cambridge University Press, 1995. MR1326374 (96c:58055)
13. D. Lind and B. Marcus, *An introduction to symbolic dynamics and coding*, Cambridge University Press, 1995. MR1369092 (97a:58050)
14. B. Maia, *An equivalent system for studying periodic points of the β -transformation for a Pisot or a Salem number*, Ph.D. thesis, University of Warwick, 2007, http://www.warwick.ac.uk/~marcq/bmaia_thesis.pdf.
15. B. Marcus, *A note on periodic points for toral automorphisms*, Manotsh. Math. **89** (1980), 121–129. MR572888 (81f:28016)
16. L. Olsen and S. Winter, *Multifractal analysis of divergence points of deformed measure theoretical Birkhoff averages. II.*, Bull. Sci. Math. **6** (2007), 518–558. MR2351308 (2010b:28023)
17. W. Parry, *On the β -expansions of real numbers*, Acta Math. Hung. **11** (1960), 401–416. MR0142719 (26:288)
18. Y.B. Pesin, *Dimension theory in dimensional systems: Contemporary views and applications*, University of Chicago Press, Chicago, 1997. MR1489237 (99b:58003)
19. Y.B. Pesin and B.S. Pitskel, *Topological pressure and the variational principle for non-compact sets (English translation)*, Funct. Anal. Appl. **18** (1984), 307–318. MR775933 (86i:28031)
20. K. Petersen, *Chains, entropy, coding*, Ergodic Theory Dynam. Systems **6** (1986), 415–448. MR863204 (88i:28040)
21. C.-E. Pfister and W.G. Sullivan, *Large deviations estimates for dynamical systems without the specification property. Applications to the β -shifts*, Nonlinearity **18** (2005), 237–261. MR2109476 (2005h:37015)
22. ———, *On the topological entropy of saturated sets*, Ergodic Theory Dynam. Systems **27** (2007), 929–956. MR2322186 (2008f:37036)
23. M. Pollicott and H. Weiss, *Multifractal analysis of Lyapunov exponent for continued fraction and Manneville-Pomeau transformations and applications to Diophantine approximation*, Comm. Math. Phys. **207** (1999), 145–171. MR1724859 (2000j:37027)
24. A. Renyi, *Representations for real numbers and their ergodic properties*, Acta Math. Acad. Sci. Hungar. **8** (1957), 477–493. MR0097374 (20:3843)
25. D. Ruelle, *Historic behaviour in smooth dynamical systems*, Global Analysis of Dynamical Systems (H. W. Broer, B. Krauskopf, and G. Vegter, eds.), Bristol: Institute of Physics Publishing, 2001. MR1858471
26. J. Schmeling, *Symbolic dynamics for β -shifts and self-normal numbers*, Ergodic Theory Dynam. Systems **17** (1997), 675–694. MR1452189 (98c:11080)
27. F. Takens, *Orbits with historic behaviour, or non-existence of averages*, Nonlinearity **21** (2008), T33–T36. MR2396607 (2009a:37043)
28. F. Takens and E. Verbitskiy, *On the variational principle for the topological entropy of certain non-compact sets*, Ergodic Theory Dynam. Systems **23** (2003), no. 1, 317–348. MR1971209 (2004b:37041)
29. D.J. Thompson, *A variational principle for topological pressure for certain non-compact sets*, J. London Math. Soc. **80** (2009), no. 3, 585–602. MR2559118 (2010k:37040)
30. ———, *The irregular set for maps with the specification property has full topological pressure*, Dyn. Syst. **25** (2010), no. 1, 25–51.
31. M. Todd, *Multifractal analysis for multimodal maps*, Preprint, arXiv:0809.1074v2, 2008.

32. P. Walters, *An introduction to ergodic theory*, Graduate Texts in Mathematics, vol. 79, Springer, New York, 1982. MR648108 (84e:28017)
33. K. Yamamoto, *On the weaker forms of the specification property and their applications*, Proc. Amer. Math. Soc. **137** (2009), no. 11, 3807–3814. MR2529890 (2010f:37027)

DEPARTMENT OF MATHEMATICS, THE PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK,
PENNSYLVANIA 16802

E-mail address: thompson@math.psu.edu

Current address: Department of Mathematics, The Ohio State University, 100 Math Tower,
231 West 18th Avenue, Columbus, Ohio 43210