

THE NONLOW COMPUTABLY ENUMERABLE DEGREES ARE NOT INVARIANT IN \mathcal{E}

RACHEL EPSTEIN

ABSTRACT. We study the structure of the computably enumerable (c.e.) sets, which form a lattice \mathcal{E} under set inclusion. The upward closed jump classes \mathbf{L}_n and \mathbf{H}_n have all been shown to be definable by a lattice-theoretic formula, except for $\overline{\mathbf{L}}_1$, the nonlow degrees. We say a class of c.e. degrees is *invariant* if it is the set of degrees of a class of c.e. sets that is invariant under automorphisms of \mathcal{E} . All definable classes of degrees are invariant. We show that $\overline{\mathbf{L}}_1$ is not invariant, thus proving a 1996 conjecture of Harrington and Soare that the nonlow degrees are not definable, and completing the problem of determining the definability of each jump class. We prove this by constructing a nonlow c.e. set D such that for all c.e. $A \leq_T D$, there is a low set B such that A can be taken by an automorphism of \mathcal{E} to B .

1. INTRODUCTION

Definability is one of the fundamental themes in computability theory. The problem of determining which classes of Turing degrees are definable in the languages of Turing reduction and set inclusion has long been a major topic of study among computability theorists. In particular, the question of which jump classes of computably enumerable (c.e.) degrees are definable in the language of set inclusion has been studied for over 40 years. We complete the answer to this question by showing that the class of nonlow c.e. degrees is not definable.

1.1. The main result. The computably enumerable (c.e.) degrees are a particularly important class of Turing degrees. They have been studied extensively since Post first asked in 1944 [10] whether there is a c.e. degree strictly between $\mathbf{0}$ and $\mathbf{0}'$. Such degrees were found independently by Friedberg and Muchnik in the 1950s using the priority method, which we will use here. The c.e. sets can be equivalently defined as the domains of partial computable functions and as Σ_1^0 sets. The definition we use here is that a set C is *computably enumerable* (c.e.) if there is a uniformly computable sequence of computable sets $\{C_s\}_{s \in \omega}$ such that $C = \bigcup_s C_s$. A degree is said to be c.e. if it contains a c.e. set.

Another important class of degrees below $\mathbf{0}'$ are the low degrees. A degree $\mathbf{d} < \mathbf{0}'$ is *low* (or \mathbf{L}_1) if $\mathbf{d}' = \mathbf{0}'$. The concept of lowness has been generalized to reflect the behavior of the n^{th} jump of \mathbf{d} . In particular, a degree \mathbf{d} is *low_n* (or \mathbf{L}_n) if $\mathbf{d}^{(n)} = \mathbf{0}^{(n)}$. Similarly, a degree \mathbf{d} is *high* (or \mathbf{H}_1) if $\mathbf{d}' = \mathbf{0}''$ and is *high_n* (or \mathbf{H}_n) if

Received by the editors January 28, 2011 and, in revised form, April 8, 2011.

2010 *Mathematics Subject Classification.* Primary 03D25.

Key words and phrases. Computably enumerable, recursively enumerable, definability, automorphisms, invariance.

The author would like to thank Bob Soare for many helpful comments and conversations.

©2012 American Mathematical Society
Reverts to public domain 28 years from publication

$\mathbf{d}^{(n)} = \mathbf{0}^{(n+1)}$. We call these classes and their complements *jump classes of degrees*. We will also sometimes call a set low_n or high_n if it is of low_n or high_n degree.

Among the jump classes, the low degrees are perhaps the most studied and most significant class. For instance, Gödel's incompleteness theorem tells us that there is no computable completion of Peano arithmetic. However, Peano arithmetic has a low completion. As we will see in §1.3, all upward-closed jump classes except for the nonlow degrees have been shown to be definable in the lattice of c.e. sets \mathcal{E} under set inclusion. We will show that the nonlow degrees are in fact not definable, setting them apart from all other jump classes.

Theorem 1.1 (Main theorem). *The nonlow degrees are not definable in \mathcal{E} .*

1.2. Methods of the proof. If we were to try to show that the nonlow degrees were definable in \mathcal{E} , we would look for some lattice-theoretic property P such that $P(A)$ holds for some A in every nonlow degree, but $P(A)$ does not hold for any low set A . We instead will show that the nonlow degrees are not definable, so we must show that there is no such property P . We can use automorphisms to achieve this.

Definition 1.2. A class of sets $\mathcal{S} \subseteq \mathcal{E}$ is *invariant* if it is closed under $\text{Aut}(\mathcal{E})$, the set of all automorphisms of \mathcal{E} . A class of degrees \mathcal{C} is *invariant* if

$$\mathcal{C} = \{\text{deg}(W) \mid W \in \mathcal{S}\},$$

where \mathcal{S} is invariant.

All classes of degrees definable in \mathcal{E} are invariant because automorphisms preserve all lattice-theoretic properties. Thus, in order to show that a class is not definable, it suffices to show that it is not invariant.

To prove Theorem 1.1, we will show that the nonlow degrees are not invariant under automorphisms of \mathcal{E} . We actually prove something stronger. It suffices to show that all sets in a certain nonlow degree can be taken by automorphisms to low sets. However, we will show that all sets Turing reducible to a particular nonlow degree can be taken by automorphisms to low sets.

Theorem 1.3. *There is a nonlow c.e. set D such that for all c.e. $A \leq_T D$, there exists a low c.e. set B and an automorphism of \mathcal{E} taking A to B .*

Theorem 1.1 follows because Theorem 1.3 shows that the nonlow degrees are not invariant.

One technique used in the proof is the Harrington-Soare automorphism method from [4]. However, we will not be able to simply build an automorphism from A to B using only this method, as it does not allow for any restraint to be put on either set A or B . Instead, we must break up the proof into two parts, performed simultaneously. The first is to build a partial automorphism on the complements of A and B , and then to extend it to a full automorphism. The Extension Theorem from [13] does not work here, as we cannot get an effective automorphism. Instead, we modify the Δ_3^0 automorphism method from [4] so that it can act as an extension theorem.

As we will see in §1.3, our set D must be low_2 . In §3, we construct a nonlow c.e. set that is low_2 . While this theorem is already known, it illustrates the technique we will use to prove our main theorem. While we will not specifically make our set D low_2 , the fact that the construction succeeds guarantees that D is low_2 .

Both the Harrington-Soare Δ_3^0 automorphism construction and the construction of a low₂ set that is nonlow are performed on trees of strategies. We will combine these trees into one tree that will provide the framework for building an automorphism from A to B while guaranteeing that D is nonlow and B is low. In order to combine these two methods, we must make significant changes to both. In §4, we will discuss the methods of the proof in more detail before we begin the formal proof.

1.3. Historical background. We can examine the c.e. sets and degrees by looking at the structure \mathcal{R} of the c.e. degrees under Turing reducibility or at the structure \mathcal{E} of the c.e. sets under set inclusion. Notice that \mathcal{E} forms a lattice with least element \emptyset and greatest element ω .

Definition 1.4. A class of sets \mathcal{S} is *definable in \mathcal{E}* if we can describe \mathcal{S} in the language of set inclusion, and a class of sets \mathcal{S} is *definable in \mathcal{R}* if we can describe \mathcal{S} in the language of Turing reducibility. A class of degrees \mathbf{D} is *definable in \mathcal{E} or \mathcal{R}* if there is a class of sets \mathcal{S} definable in \mathcal{E} or \mathcal{R} such that $\mathbf{D} = \{\deg(W) \mid W \in \mathcal{S}\}$.

It is natural to ask which jump classes of degrees are definable in the structures \mathcal{E} and \mathcal{R} . We will discuss the problem for \mathcal{E} and refer to \mathcal{R} only for comparison. For the structure \mathcal{E} , the first progress on this problem came in the form of Martin's Theorem 1.5. We say a c.e. set M is *maximal* if for every c.e. set W_e , either $W_e \subseteq^* M$ or $W_e \cup M =^* \omega$. The property of being a maximal set is lattice-theoretic, or definable in \mathcal{E} .

Theorem 1.5 (Martin [8]). *The high degrees \mathbf{H}_1 are exactly the degrees of the maximal sets.*

Thus, the high degrees are definable in \mathcal{E} . This theorem marked the beginning of investigations as to which jump classes of degrees are definable in \mathcal{E} .

It is not difficult to see that the degree $\mathbf{0}$ is definable in \mathcal{E} , as it is the degree of the least element \emptyset of the lattice \mathcal{E} . In addition, the set of all nonzero degrees is definable in \mathcal{E} as the degrees of $\{A \in \mathcal{E} \mid \bar{A} \notin \mathcal{E}\}$. Harrington showed in 1986 (see Soare [15]) that the degree $\mathbf{0}'$ is definable by finding a definition for the class of creative sets, which is a class of sets known to be complete.

A c.e. set A is called *atomless* if A is not contained in any maximal set. The atomless sets are clearly a class definable in \mathcal{E} .

Theorem 1.6 (Lachlan [6]). *All atomless sets are nonlow₂.*

Theorem 1.7 (Shoenfield [11]). *Every nonlow₂ c.e. degree contains an atomless c.e. set.*

This shows that the degrees of the atomless c.e. sets are the nonlow₂ degrees. Thus, the nonlow₂ c.e. degrees are definable in \mathcal{E} .

The first nondefinability results came in 1995, many years after the first definability results.

Theorem 1.8 (Cholak [1]; Harrington-Soare [4]). *Every noncomputable c.e. set can be taken by an automorphism of \mathcal{E} to a high set.*

This theorem showed that the downward closed jump classes \mathbf{L}_n , $n > 0$ and $\bar{\mathbf{H}}_n$, $n > 0$ are not invariant and thus not definable. Harrington and Soare [4] used prompt sets to show that $\bar{\mathbf{H}}_0$, the class of incomplete c.e. degrees, is not definable.

This completed the problem of showing the noninvariance of the noncomputable downward closed jump classes.

Definition 1.9. A c.e. set A is *prompt* if there is an enumeration $\{A_s\}$ of A and a computable function p such that for all s , $p(s) \geq s$, and for all e ,

$$W_e \text{ infinite} \implies (\exists x) (\exists s) [x \in W_{e, \text{at } s} \ \& \ A_s \upharpoonright x \neq A_{p(s)} \upharpoonright x],$$

where $C \upharpoonright x$ denotes the first x bits of C .

Promptness is a property of degrees, as every set Turing equivalent to a prompt set is also prompt.

Theorem 1.10 (Harrington-Soare [4]). *For all prompt sets A , there exists $B \in \mathbf{0}'$ such that A is automorphic to B .*

It is well known that there is a low prompt degree \mathbf{d} , so every set $A \in \mathbf{d}$ can be taken by an automorphism of \mathcal{E} to a complete set B . This reconfirms that the low degrees are not invariant, and neither is any other class of degrees that contains the low degrees and does not contain $\mathbf{0}'$. Thus, every downward closed jump class \mathbf{L}_n for $n \geq 1$ and $\overline{\mathbf{H}}_n$ for $n \geq 0$ is neither invariant nor definable in \mathcal{E} .

The only remaining jump classes in 1996 were the upward closed jump classes $\overline{\mathbf{L}}_n$ for $n = 1$ and $n \geq 3$ and \mathbf{H}_n for $n \geq 2$.

Theorem 1.11 (Cholak-Harrington [2]). *For $n \geq 2$, \mathbf{H}_n and $\overline{\mathbf{L}}_n$ are definable.*

This is analogous to the result of Nies, Shore, and Slaman [9] for the c.e. degrees $(\mathcal{R}, <_T)$ showing that \mathbf{L}_n and \mathbf{H}_n for $n \geq 2$ (and, equivalently, their complements) are definable in \mathcal{R} .

The only remaining case for the structure \mathcal{E} is the class of nonlow degrees, $\overline{\mathbf{L}}_1$. We will show that this class differs from all the other upward closed jump classes, as it is the only one that is not definable. For the structure \mathcal{R} of c.e. degrees under Turing reducibility, all jump classes are known to be definable except for the class of low degrees, which is still unknown. In both structures \mathcal{E} and \mathcal{R} , the single jump is more difficult to define than the double jump.

2. DEFINITIONS AND NOTATION

We give some basic definitions that will be used throughout this paper. Definitions specific to the automorphism construction will appear in §5.

Let $C \subset \omega$. Then C can be viewed as an element of 2^ω by means of its characteristic function. For any set C , node α , or path f , we write $C \upharpoonright n$ to denote the initial segment of C of length n and similarly for $\alpha \upharpoonright n$ and $f \upharpoonright n$.

Let Φ_e^C be the e^{th} Turing functional with oracle C . We define the use $\varphi_e^C(n)$ of the computation $\Phi_e^C(n)$ by

$$\varphi_e^C(n) = \begin{cases} m + 1, & \text{if } \Phi_e^C(n) \downarrow, \text{ where } m \text{ is the largest bit of } C \\ & \text{queried in the computation } \Phi_e^C(n) \\ \text{undefined,} & \text{otherwise.} \end{cases}$$

Let C be a c.e. set with computable enumeration $C = \bigcup_{s \in \omega} C_s$. We write $\Phi_e^C(n)[s]$ to mean $\Phi_{e,s}^{C_s}(n)$, the result of the Turing functional Φ_e with oracle C_s on input n , after s steps of computation. Similarly, we write $\varphi_e^C(n)[s]$ for the use of that computation.

Let $\langle x, y \rangle$ be the standard pairing function from ω^2 to ω . We will also use $\langle a, b, c \rangle = \langle \langle a, b \rangle, c \rangle$, allowing us to view sets as three-dimensional matrices, where each “row” of the matrix is itself a two-dimensional matrix.

A tree T is a subset of $\omega^{<\omega}$ such that if $\alpha \in T$, then every initial segment of α is in T . Let $[T]$ be the set of infinite paths through T , where h is an infinite path through T if $h \upharpoonright n \in T$ for all n . Let $\alpha, \beta, \gamma, \delta, \dots$ range over T . Let $|\alpha|$ denote the length of α . Let $\alpha \preceq \beta$ ($\alpha \prec \beta$) denote that string β extends (properly extends) α . Let λ denote the empty string, and α^- denote the predecessor of α if $\alpha \neq \lambda$. Let $\alpha \hat{\ } \beta$ denote the concatenation of string α followed by string β . In this paper, we work with trees in $\omega^{<\omega}$.

3. A LOW_2 NONLOW C.E. SET D

The existence of a low_2 nonlow $_1$ c.e. set D follows from the Sacks Jump Theorem [12] but that construction involves a two-stage process, one stage computable in \mathbf{O}' , and is not useful in a computable construction for building automorphisms which we need here. Now we present another construction which will be key for our main theorem. It expands and corrects a sketch by Harrington and Soare [5], §5.2.

Theorem 3.1. *There is a low_2 nonlow c.e. set D .*

3.1. Listing all Δ_2^0 partial functions. Let $\{\widehat{\varphi}_e\}_{e \in \omega}$ be a listing of all Δ_2^0 functions. That is, define $\widehat{\varphi}_e(x) = \lim_z \varphi_e(x, z)$ if the limit exists and if φ_e is total, and $\widehat{\varphi}_e(x) \uparrow$ otherwise, where $\{\varphi_e\}_{e \in \omega}$ is the usual listing of all partial computable functions. It is useful to have a computable total function $\widetilde{\varphi}_{e,s}(x)$ which approximates the Δ_2^0 function $\widehat{\varphi}_e(x)$ as follows.

Define $\widetilde{\varphi}_{e,s}(x) = \varphi_{e,s}(x, z)$, where z is the maximum element such that $z \leq s$ and $\varphi_{e,s}(x, v) \downarrow$ for all $v \leq z$, with a default output of 0 if no such z exists. Now the function $\widetilde{\varphi}$ is primitive recursive in the variables e, s , and x , and clearly

$$(1) \quad \widehat{\varphi}_e(x) = y \implies \lim_s \widetilde{\varphi}_{e,s}(x) = y.$$

3.2. Making D nonlow. To make D nonlow we must diagonalize over all Δ_2^0 functions which could make D low.

We will meet for every j the requirement,

$$(2) \quad P_j : \widehat{\varphi}_j \text{ is not the characteristic function of } \{x : W_x \cap \overline{D} \neq \emptyset\}.$$

This actually ensures the stronger property that \overline{D} is not semi-low [15, p. 72], and hence that D is not low.

The strategy for P_j . We define a set Z_j , whose index $g(j)$ is known a priori by the Recursion Theorem, so that $Z_j = W_{g(j)}$, where g is a computable function. We describe the strategy in terms of a Lachlan game as in Lachlan [7] in which the Blue player (us) and the Red player (the opponent) simultaneously construct c.e. sets.

Step 1a. Initially, $Z_j = \emptyset$, and $Z_j \cap \overline{D} = \emptyset$. Wait until (if ever) the opponent recognizes this fact by producing a stage t_0 such that $\widetilde{\varphi}_{j,t_0}(g(j)) \downarrow = 0$. Then choose a fresh element $\Gamma_{t_0}^j = x_1 > x_0$ not yet in D , and put it in Z_j so that (at least temporarily) $\overline{D} \cap Z_j \neq \emptyset$.

Step 1b. Wait until (if ever) the opponent produces a stage t_1 such that $\widetilde{\varphi}_{j,t_1}(g(j)) \downarrow = 1$. Then put $\Gamma_{t_1}^j = x_1$ into D and repeat the procedure. (That is, go to Step 2a as above with a fresh element $x_2 > x_1$ not yet in D .)

3.2.1. *The Σ_2 -outcome for the P_j strategy.* The Σ_2 -outcome is that this process is repeated finitely often. Then $\tilde{\varphi}_{j,t}(g(j))$ gives the wrong answer for almost every t . Therefore, requirement P_j is satisfied. Furthermore, P_j appoints at most finitely many witnesses $\{x_1 < x_2 < \dots < x_n\}$. This will cause only a finite set of injuries to the lower priority negative requirements described below.

3.2.2. *The Π_2 -outcome for the P_j strategy.* The Π_2 -outcome is that this process is repeated infinitely often. Then $\lim_t \tilde{\varphi}_{j,t}(g(j)) [t]$ diverges. Therefore, requirement P_j is satisfied because $\hat{\varphi}_j(g(j))$ fails to give the correct value according to (1). In this case, the P_j strategy enumerates an infinite set $\{x_1 < x_2 < \dots\}$ of elements into D . However, this set is not only computable but is enumerated in strictly increasing order. Therefore, its effect on opposing negative requirements is easy to analyze.

Let Γ_s^j or $\Gamma^j[s]$ denote the value of the witness x_n for P_j at the end of stage s if it exists, and the maximum of s and all previous witnesses otherwise. Choose the witnesses in $\omega^{[j]}$ in increasing order and such that $\Gamma_s^j \leq \Gamma_{s+1}^j$. We think of Γ^j as a movable marker which is monotonically nondecreasing on a strictly increasing sequence of elements $\{x_1 < x_2 < \dots\}$ and goes to infinity in the Π_2 case. Therefore, Γ^j provides a lower bound for any future elements which will be contributed to D by P_j .

3.3. **Making D low₂.** Define $\text{Tot}^A := \{e : \Phi_e^A \text{ is total}\}$ and $\text{Tot} := \text{Tot}^\emptyset$. Recall that $\text{Tot}^A \equiv_{\text{T}} A''$ and indeed Tot^A is Π_2^A -complete. The requirements must guarantee that

$$(3) \quad \text{Tot}^X \leq_{\text{T}} \text{Tot},$$

which implies that X is low₂ by [15, Theorem IV.3.2 and Exer. IV.4.5]. Define

$$(4) \quad \Psi_{e,s}^X(y) = \begin{cases} \Phi_{e,s}^X(y) & \text{if } (\forall z \leq y) [\Phi_{e,s}^X(z) \downarrow] \\ \text{undefined} & \text{otherwise.} \end{cases}$$

Define the use function, $\psi_{e,s}(y) := \varphi_e^D(y)[s]$, also written $\psi_e(y)[s]$, if $\Psi_e^D(y)[s]$ is defined, and let $\psi_{e,s}(y)$ be undefined otherwise. Note that Ψ_e^X is either total or has finite domain, and Ψ_e^X is total if and only if Φ_e^X is total. The first outcome with $X = D$ is the Π_2 -outcome for Ψ_e^D and the second is the Σ_2 -outcome. To achieve condition (3) we define the following *pseudo-requirements*:

$$(5) \quad N_{e,y} : (\exists^\infty s) [\Psi_e^D(y)[s] \downarrow] \iff \Psi_e^D(y) \downarrow .$$

Let N_e be the conjunction, $N_e = \& \{N_{e,y} : y \in \omega\}$. Of course, we cannot literally satisfy these negative pseudo-requirements $N_{e,y}$ for every e and y or else D would be low because they are exactly the lowness requirements of the low simple set construction in [15, p. 111]. Instead these $N_{e,y}$ are like the pseudo-requirements Q_e in the proof of the Sacks Jump Theorem in [15, p. 138]. *Attempting* to satisfy them causes us to put on a restraint to preserve a computation $\psi(y)[s]$ whenever possible. This restraint may be injured infinitely often (but only computably) by a higher priority positive requirement and $\mathbf{0}''$ can discover that injury and can calculate when computations in (5) will no longer be injured and whether $\Psi_e^D(y)$ converges or not. Therefore, we use (5) to achieve the following:

$$(6) \quad (\forall y)(\exists^\infty s) [\Psi_e^D(y)[s] \downarrow] \iff \Psi_e^D \text{ is total.}$$

The left hand side is Π_2 and therefore computable in $\mathbf{0}''$. The right hand side is Π_2^D -complete. Therefore, $D'' \leq_T \mathbf{0}''$ and D is low₂.

3.4. The interference between requirements. We now consider the interference between opposing positive and negative requirements.

3.4.1. *One P_j versus one N_e with $j < e$.* Consider one positive requirement P_j and one negative requirement N_e where $j < e$ so that P_j has higher priority. Our strategy for N_e will be split into two parts, one in case P_j has the Π_2 -outcome and one in case P_j has the Σ_2 -outcome. We say that N_e is *injured* by P_j whenever the strategy for P_j enumerates a witness $x_n < \psi_e(y)[s]$ into D .

The Σ_2 -outcome for P_j . If P_j has the Σ_2 outcome and N_e is guessing correctly at the number of witnesses it will enumerate in D , then N_e can wait until those witnesses have all been enumerated in D before starting its strategy.

The Π_2 -outcome for P_j . Suppose P_j has the Π_2 -outcome and N_e is guessing this correctly. Then N_e knows that the marker Γ_j for P_j is moving monotonically to infinity. N_e can replace the raw use function $\psi_e(y)$ by a *modified* use function $\widehat{\psi}_e(y)$ which is $\psi_e(y)$ whenever $\psi_e(y)[s] < \Gamma^j[s]$ and undefined otherwise. Any real computation with use $\psi_e(y)$ will eventually have this use below Γ_j and therefore it will also equal $\widehat{\psi}_e(y)$. However, P_j will never cause a witness $x < \widehat{\psi}_e(y)$ to enter D , so there will never be any injury to the function $\widehat{\psi}_e(y)$.

If there are two positive requirements of higher priority, P_j and P_k for $j, k < e$, then as the upper boundary for N_e we take the minimum of $\Gamma^j[s]$ and $\Gamma^k[s]$. Again there will be no injury to $\widehat{\psi}_e(y)$. This is formalized for the general case in (7) and (8).

3.4.2. *One N_e versus one P_j with $e < j$.*

The Σ_2 -outcome for N_e . In the Σ_2 -outcome for N_e there is some least y such that $\Psi^D(y)$ is undefined. Therefore, by the definition of Ψ_e and (5) there will be at most finitely many stages when a new computation $\psi_e(z)$ appears for any z and only finitely many times N_e wants to impose a new restraint. Whenever this happens, we restart the P_j strategy which is therefore restarted (and injured) at most finitely often.

The Π_2 -outcome for N_e . Even though N_e has a higher priority than P_j it must guarantee (5) for *all* y not just for $y = e$. Therefore, N_e cannot simply preserve $\psi_e(y)$ with priority N_e against all followers x_n for P_j as in the construction of a low simple set.

The solution is to use the natural ω ranking of all negative pairs $\langle e, y \rangle$ against the ω ranking $\langle j, n \rangle$ of all positive pairs. P_j guesses that $\widehat{\psi}_e(y)$ will eventually be defined for all y . When P_j is ready to define a new x_n it can wait until $\widehat{\psi}_e(y)[s]$ is defined for all y with $\langle e, y \rangle < \langle j, n \rangle$ and can then appoint $x_n > \psi_e(y)[s]$ for such a y .

Note that the computation $\psi_e(y)[s]$ cannot be destroyed at any stage $t > s$ because the witnesses $x_k, k < n$, have already entered D by stage s and x_n the first active witness after s exceeds the use. Of course, a computation $\widehat{\psi}_e(z)$ may be injured at some stage $t > s$ because it may have become defined after x_n was appointed. However, $\widehat{\psi}_e(z)$ can be injured by only finitely many witnesses x_k and at most once by each. Therefore, the subrequirement $N_{e,y}$ of (5) is met for every y .

But when P_j is ready to appoint x_n it need only wait for $\widehat{\psi}_e(y)$ to be defined for all $\langle e, y \rangle < \langle j, n \rangle$ which must eventually occur. The fact that x_n is ready to be appointed means that all witnesses $x_k, k < n$, have already entered D . Therefore, these computations cannot later change and x_n can carry out its strategy as before.

Notice that this argument succeeds only because when we define $x_n > \widehat{\psi}_e(y)[s]$ we know that $\widehat{\psi}_e(y)[s]$ cannot later change and hence that x_n can never injure this computation. If we have two or more positive requirements we have to worry that *another* positive requirement may contribute a follower $z < \widehat{\psi}_e(y)[s]$ to D at some later stage $t > s$ which allows $\widehat{\psi}_e(y)$ to be redefined so that $x_n < \widehat{\psi}_e(y)[t]$. This threatens to allow x_n to later enter D and injure the $\widehat{\psi}_e(y)$ computation. We address this obstacle in §3.7.

3.5. A tree argument. To complete the proof we convert these strategies into a tree construction as described in [15, Chapter XIV]. Put the empty node λ on T . If $\alpha \in T$ and $|\alpha| = 2e$, then we call α “even” and we associate with α a version of the strategy N_e , and put the nodes $\alpha \widehat{0}$ and $\alpha \widehat{1}$ on T which represent the Π_2 and Σ_2 -outcomes, respectively, of the α strategy. If $\beta \in T$ and $|\beta| = 2i + 1$, then we call β “odd” and we associate with β a version of the strategy for P_i , and put the outcomes $\beta \widehat{0}$ and $\beta \widehat{p}$ for all $p \geq 1$ on T , representing the Π_2 and all possible Σ_2 -outcomes of the β strategy ($\beta \widehat{p}$ means a guess of exactly $p - 1$ β -witnesses enumerated into D). The order of the successor nodes to β will be $\beta \widehat{0}$ on the left, and the others on the right in the order of ω^* . The true path f on T and its computable approximation $\{f_s\}_{s \in \omega}$ such that $f = \liminf_s f_s$ will be defined below.

Suppose $\beta \in T$ and $|\beta| = 2j + 1$. Then β has an associated witness function Γ_s^β for carrying out its strategy for P_j as above. At each stage s there is at most one such witness $\Gamma_s^\beta = x_n^\beta$, the n^{th} witness appointed by β , and it can contribute at most that witness to D at stage $s + 1$. We arrange that $\Gamma_s^\beta \leq \Gamma_{s+1}^\beta$. (The following is the general case of the explanation in §3.4.1 in the paragraph on the Π_2 -outcome for P_j .)

Suppose $\alpha \in T$ and $|\alpha| = 2e$. Define

$$(7) \quad \Lambda_s^\alpha = \min\{\Gamma_s^\beta : \beta \widehat{0} \preceq \alpha \ \& \ |\beta| \text{ is odd }\}, \text{ and}$$

$$(8) \quad \widehat{\psi}_{\alpha,s}(y) = \begin{cases} \psi_{e,s}(y) & \text{if } \psi_{e,s}(y) \downarrow < \Lambda_s^\alpha \text{ and (for all odd } \beta) \\ & [\beta \widehat{p} \preceq \alpha \implies \text{all } p - 1 \text{ } \beta\text{-witnesses have entered } D] \\ \text{undefined} & \text{otherwise.} \end{cases}$$

Note that $\lim_s \Lambda_s^\alpha = \infty$ if $\alpha \prec f$. For each $\beta \prec \alpha, |\beta| = 2i + 1$, β may put elements into D without respecting the computations $\psi_{\alpha,s}(y)$ which α is trying to protect. Assume $\alpha \prec f$. Then no such β will ever contribute to D an element $x \leq \widehat{\psi}_{\alpha,s}(y)$. Choose a stage s_0 such that for each $\beta \prec \alpha, |\beta| = 2i + 1$, and if $\beta \widehat{p} \preceq \alpha$ for $p \neq 0$, or $\beta <_L \alpha$, then β does not contribute an element to D at any stage $s \geq s_0$. Now no $\beta < \alpha$ will contribute an element $x \leq \widehat{\psi}_{\alpha,s}(y)$ at any stage $s > s_0$.

But for odd length β of *lower* priority $\beta \succeq \alpha \widehat{0}$, the β witnesses are arranged as for N_e in §3.3. That is, such a β can appoint the n^{th} witness, x_n^β , only after $\widehat{\psi}_{\alpha,s}(y) \downarrow$ for all $y, \langle \alpha, y \rangle \leq \langle \beta, n \rangle$, and must exceed their values. (Here, we identify the nodes of T with their Gödel numbering under some fixed effective numbering of T .) As discussed in §3.7, this is not quite as simple as it sounds. However, as we will see, for each y the computation $\widehat{\psi}_{\alpha,s}(y)$ can be injured at most finitely often,

giving

$$(9) \quad N_\alpha : (\forall y)[(\exists^\infty s)[\widehat{\psi}_\alpha^D(y)[s]\downarrow] \iff \widehat{\psi}_\alpha^D(y)\downarrow].$$

Finally, $\text{Tot}^A \leq_T \mathbf{0}''$ because to determine whether Φ_e^D is total, we use $\mathbf{0}''$ to compute the node on the true path $\alpha = f \upharpoonright 2e$. Now by N_α we know that Ψ_e is total if and only if $(\forall y)(\exists^\infty s)[\widehat{\psi}_\alpha^D(y)[s]\downarrow]$, which is a Π_2 statement.

The ordering of the nodes on the tree is important. Note that the positive requirement nodes follow a different ordering, in that the 0 node is on the left, and the rest of the nodes are in the order of ω^* .

Definition 3.2. Let $\alpha, \beta \in T$.

(i) Unless otherwise specified, α is to the left of β ($\alpha <_L \beta$) if

$$(\exists a, b \in \omega) (\exists \gamma \in T) [\gamma \hat{<} a \preceq \alpha \ \& \ \gamma \hat{<} b \preceq \beta \ \& \ a < b].$$

(ii) If $h \in [T]$ we say $\alpha <_L h$ if there exists $\beta \prec h$ such that $\alpha <_L \beta$ and similarly for $h <_L \alpha$.

3.6. The construction. Fix $T = 2^{<\omega}$. Simultaneously with the construction of D at each stage we define a function $f_s \in T$, $|f_s| = 2s$, such that $f = \liminf_s f_s$. We say that stage s is an α -stage if $\alpha \prec f_s$. For each even $\alpha \in T$ define the function

$$(10) \quad l(\alpha, s) = \max \{x : (\forall y < x)[\widehat{\psi}_\alpha^D(y)[s]\downarrow]\}.$$

Stage $s = 0$. Define $f_0 = \lambda$ to be the empty node on T .

Stage $s + 1$. Define f_{s+1} by a series of substages t , $0 \leq t \leq 2s$, where we define δ^t at substage t , as follows, and then define $f_{s+1} = \delta^{2s}$.

Substage t , $0 \leq t \leq 2s$. Define $\delta^0 = \lambda$. Given δ^t , perform the following action.

Case 1. $|\delta^t|$ is even. Let $\alpha = \delta^t$. Let v be the last α -stage less than s . If $l(\alpha, v) < l(\alpha, s)$, then define $\delta^{t+1} = \delta^t \hat{<} 0$, and otherwise let $\delta^{t+1} = \delta^t \hat{<} 1$.

Case 2. $|\delta^t| = 2j + 1$. Let $\beta = \delta^t$. Here β is associated with the strategy for P_j described in §3.2. (In particular, we assume a set Z_β is being defined whose index $g(\beta)$ is known a priori, such that $W_{g(\beta)} = Z_\beta$, g is the computable function as in §3.2 Step 1a but adjusted with β in place of j , and that our objective is to arrange that either $\lim_s \tilde{\varphi}_{j,s}(g(\beta))$ diverges, or gives the wrong answer about whether $W_{g(\beta)} \cap \overline{D} \neq \emptyset$.)

Subcase 1. (Like §3.2 Step 1a) Suppose $Z_{\beta,s} \cap \overline{D}_s = \emptyset$, $\tilde{\varphi}_{j,s}(g(\beta)) \downarrow = 0$, and there is no β -witness in \overline{D}_s . Let the β -witnesses previously appointed be $x_1^\beta \dots x_{n-1}^\beta$. Suppose

$$(11) \quad (\forall \text{ even } \alpha) \left[\alpha \hat{<} 0 \prec \beta \implies (\forall p)[\langle \alpha, p \rangle \leq \langle \beta, n \rangle \implies \widehat{\psi}_{\alpha,s}(p)\downarrow] \right], \text{ and}$$

$$(12) \quad (\forall \text{ odd } \xi) \left[\begin{array}{l} \xi \hat{<} p \prec \beta \implies x_{n'}^\xi \in D_s \text{ for all } n' < p, \\ \text{and } (\xi \hat{<} 0 \prec \beta \ \& \ \langle \xi, n' \rangle < \langle \beta, n \rangle) \implies x_{n'}^\xi \in D_s \end{array} \right].$$

Define y to be the smallest $z \in \overline{D}_s \cap \omega^{[s]}$, where β is identified with its Gödel number, such that $\Gamma_s^\beta < z < \Lambda_s^\beta$, and the following holds:

$$(13) \quad (\forall \text{ even } \alpha) \left[\alpha \hat{<} 0 \prec \beta \implies (\forall \langle \alpha, p \rangle < \langle \beta, n \rangle) [\widehat{\psi}_{\alpha,s}(p)\downarrow < z] \right] \\ \text{and } (\forall \text{ odd } \gamma <_L \beta) [z > \Gamma_s^\gamma].$$

The above equation says that we choose a new β -witness y to be bigger than the last β -witness and all γ -witnesses to the left of β , smaller than the higher priority witnesses, and that preserves higher priority restraint functions for the negative requirements. If y exists, then appoint y as the next β -witness, call it x_n^β , define $\Gamma_{s+1}^\beta = x_n^\beta$, and enumerate x_n^β in $Z_{\beta,s+1}$.

Subcase 2. (Like §3.2 Step 1b) Suppose that $\tilde{\varphi}_{j,s}(g(\beta)) \downarrow = 1$, Γ_s^β is defined and equal to some β -witness x_n^β , and that

$$(14) \quad (\forall \text{ odd } \xi)[\xi \frown 0 \prec \beta \implies \Gamma_s^\xi = x_p^\xi \in D_{s+1}].$$

Then enumerate x_n^β in D_{s+1} , and define

$$(15) \quad \Gamma_{s+1}^\beta = \max \{\Gamma_s^\beta, s + 1\}.$$

(Equation (14) says that for every odd ξ with $\xi \frown 0 \prec \beta$, i.e., for which β is guessing that ξ has the Π_2 outcome of infinitely many witnesses, β must wait until ξ actually puts its witness $\Gamma_s^\xi = x_p^\xi$ into D_{s+1} during an earlier substage of stage $s + 1$ before β is allowed to simultaneously contribute its own witness at a later substage of stage $s + 1$.)

Subcase 3. Otherwise, if $\Gamma_s^\beta = x_n^\beta$ for some β -witness x_n^β , then define $\Gamma_{s+1}^\beta = \Gamma_s^\beta$, and if not, then define $\Gamma_{s+1}^\beta = \max \{\Gamma_s^\beta, s + 1\}$.

To complete Case 3 define $\delta^{t+1} = \delta^t \frown 0$ if Subcase 2 holds, and otherwise $\delta^{t+1} = \delta^t \frown p + 1$, where p is the number of elements β has enumerated into D . Complete the construction by defining $f_{s+1} = \delta^{2s}$ and by initializing every odd node β such that $f_{s+1} <_L \beta$, that is, by canceling any β -witness, and defining $\Gamma_{s+1}^\beta = \max \{\Gamma_s^\beta, s + 1\}$.

3.7. The obstacle. Before we begin the formal verification, we will discuss the primary obstacle in the proof, which was missing from the proof given in [5]. The obstacle appears in proving that along the true path, requirement N_α in (9) is met.

The tricky part is showing that for a node $\beta \succ \alpha$ with witness x_n^β , where $\langle \beta, n \rangle > \langle \alpha, y \rangle$, that x_n^β cannot injure $\hat{\psi}_\alpha^D(y)[t]$ for any t . Now, when x_n^β was appointed as a witness at some stage s , it must have been chosen to respect the $\langle \alpha, y \rangle$ computation $\hat{\psi}_\alpha^D(y)[s]$. The question is whether x_n^β still respects the $\langle \alpha, y \rangle$ computation $\hat{\psi}_\alpha^D(y)[t]$ at any stage $t > s$. If it does not, then x_n^β could injure the $\langle \alpha, y \rangle$ computation when x_n^β enters D .

This raises a new question: is it possible for x_n^β to be defined at stage s , when the $\langle \alpha, y \rangle$ computation was less than x_n^β , and then enter D at stage $t > s$, when the $\langle \alpha, y \rangle$ computation is greater than x_n^β ? Note that the only way for this to happen is if something injures the $\langle \alpha, y \rangle$ computation between stages s and t . In the verification, we will show by examining each type of node that it is not possible for this injury to occur. Thus, x_n^β cannot injure $\hat{\psi}_\alpha^D(y)[t]$ for any $t > s$.

3.8. The verification. We now verify that this construction establishes Theorem 3.1. First define the *true path* f of the construction by induction on e . Let $f \upharpoonright 0 = \lambda$, the empty node. Given $\alpha = f \upharpoonright 2e$ define $f(2e) = 0$ if $(\forall y)[(\exists^\infty s)[\hat{\psi}_\alpha^D(y)[s] \downarrow]]$, and $f(2e) = 1$ otherwise. Given $\beta = f \upharpoonright (2e + 1)$ define $f(2e + 1) = 0$ if β appoints infinitely many β -witnesses and $f(2e + 1) = p + 1$ if β appoints p β -witnesses that eventually enter D .

Lemma 3.3. *Fix $e \in \omega$. Let $\alpha = f \upharpoonright 2e$. Then the construction has satisfied the requirement,*

$$(16) \quad N_\alpha : \quad (\forall y) \left[(\exists^\infty s) [\widehat{\psi}_\alpha^D(y)[s] \downarrow] \iff \widehat{\psi}_\alpha^D(y) \downarrow \right],$$

so that D is low₂.

Proof. Fix $\alpha = f \upharpoonright 2e$ and fix $y \in \omega$. The right to left direction is obvious. For the other direction, assume $(\exists^\infty s) [\widehat{\psi}_\alpha^D(y)[s] \downarrow]$. Assume by induction that

$$(17) \quad (\exists t)(\forall s > t)(\forall z < y) [\widehat{\psi}_\alpha^D(z)[s] \downarrow = \widehat{\psi}_\alpha^D(z)].$$

Consider all the finitely many pairs $\langle \beta, n \rangle < \langle \alpha, y \rangle$. Let p be the maximum of $|\beta|$ for all such β . Choose some odd ξ , such that $|\xi| > p$, and $\alpha \widehat{\ } 0 \prec \xi \widehat{\ } 0 \prec f$. Let t be as in (17). Choose $s_0 > t$ such that for all $s \geq s_0$: (i) if $\langle \beta, n \rangle < \langle \alpha, y \rangle$, then the β -witness x_n^β does not become newly defined or enter D at stage s ; (ii) $f_s \not\prec_L \xi$ (so every odd $\beta \prec_L \xi$ has ceased to act by stage s); and (iii) every β with $\beta \widehat{\ } p \prec \xi$, $p \neq 0$, has ceased to act by stage s . Thus, by choice of s_0 no witness x_n^β , for $\beta \prec_L \xi$, $\beta \widehat{\ } p \prec \xi$, or $\langle \beta, n \rangle < \langle \alpha, y \rangle$ will ever contribute a β -witness to D after stage s_0 . In other words, these three types of witnesses cannot injure the $\langle \alpha, y \rangle$ computation after stage s_0 .

Choose $s_1 \geq s_0$ such that ξ contributes a ξ -witness x_m^ξ to D at s_1 . This action of ξ causes initialization of all β , $\xi \prec_L \beta$, and cancellation of all such β -witnesses at stage s_1 . By (14), all β with $\beta \widehat{\ } 0 \prec \xi$ must have contributed their witnesses to D by stage s_1 , and so have no β -witnesses in existence at the end of stage s_1 . Thus, for all β with $\xi \prec_L \beta$ or $\beta \widehat{\ } 0 \prec \xi$, no β -witnesses currently in existence at stage s_1 will ever injure the $\langle \alpha, y \rangle$ computation.

The above arguments show that for any $\beta \prec \xi$ or to the right or left of ξ , at stage s_1 , β has no witness that could possibly injure the computation $\widehat{\psi}_{\alpha, s_1}(y)$. We have also shown that if $\langle \beta, n \rangle < \langle \alpha, y \rangle$, then x_n^β will not injure the $\langle \alpha, y \rangle$ computation. The only remaining case is when $\xi \prec \beta$ and $\langle \beta, n \rangle > \langle \alpha, y \rangle$. This is the case discussed in §3.7. We describe it here in detail.

Consider $\xi \prec \beta$. Suppose $\langle \beta, n \rangle > \langle \alpha, y \rangle$ and x_n^β is a witness that exists at stage s_1 . If x_n^β was appointed when $\widehat{\psi}_{\alpha, t}(y)$ was the same as at stage s_1 , then x_n^β must be larger than $\widehat{\psi}_{\alpha, s_1}(y)$, so it will not injure it. However, if x_n^β was appointed when $\widehat{\psi}_{\alpha, t}(y)$ was smaller than it is at stage s_1 , then at some stage after s_1 , x_n^β could injure the current computation. Now, the only way for this to happen is for something else to injure the computation before stage s_1 and after x_n^β has been appointed. The injury could not come from a node $\gamma \prec_L \beta$ because that would reset the β -witness. It could not come from a node $\gamma \succ \beta$ because by (14), γ must wait for β to enumerate its witness before enumerating its own. Suppose the injury came from a node γ to the right of β . Then to avoid being reset, x_n^β must be defined before the γ -witness. Thus, by (13), γ must have its witness larger than x_n^β , which means that γ could not have injured the $\langle \alpha, y \rangle$ computation.

Now, if $\gamma \widehat{\ } p \prec \beta$ for any $p \neq 0$, then by (12), β must wait for all γ -witnesses to enter D before defining its own witness. If $\gamma \widehat{\ } 0 \prec \beta$, then β must choose its witness smaller than the γ -witness, so γ cannot injure the $\langle \alpha, y \rangle$ computation. Thus, no β -witness for $\xi \prec \beta$ appointed before stage s_1 can possibly injure $\widehat{\psi}_{\alpha, s_1}(y)$.

Hence, at the end of stage s_1 , the only β -witnesses in existence which have the priority to injure a computation $\widehat{\psi}_{\alpha,t}(y)$ for some $t \geq s_1$ are those which are permanently dormant and will never enter D .

Consider any new β -witness x_n^β appointed after stage s_1 . We cannot have $\beta <_L \xi$ by (ii), or $\beta \widehat{<} p \prec \xi$ by (iii). If $\alpha \widehat{<} 0 \prec \beta$, then we must have $\widehat{\psi}_{\alpha,s}(y) \downarrow < x_n^\beta$ by (72). If $f <_L \beta$, then x_n^β must exceed $\widehat{\psi}_{\alpha,s}(y)$ if the latter converges when x_n^β is appointed, and if not, x_n^β must be canceled whenever it later converges. (Note that any x_n^β newly appointed at s must exceed $\widehat{\psi}_\alpha^D(y)[s]$ because the former must exceed s by the definition of Γ_s^β , and the latter must be less than s by the usual convention on the use function.) Choose $k > s_1$ such that $\widehat{\psi}_{\alpha,k}(y) \downarrow$. Then by induction on v we have for all $v \geq k$ that $\widehat{\psi}_{\alpha,v}(y) \downarrow = \widehat{\psi}_{\alpha,k}(y) \downarrow = \widehat{\psi}_\alpha(y)$, and no new witness is appointed less than this value. Hence, $(\exists k)(\forall s > k)[\widehat{\psi}_\alpha^D(y)[s] \downarrow]$. By the discussion in §3.3, D is low₂. □

Lemma 3.4. *For every $j \in \omega$ the strategy of node $\beta = f \upharpoonright (2j + 1)$ has satisfied the requirement,*

$$(18) \quad P_j : \widehat{\varphi}_j \neq \text{characteristic function of } \{x : W_x \cap \overline{D} \neq \emptyset\},$$

so that D is not low₁.

Proof. Fix $\beta = f \upharpoonright (2j + 1)$. Suppose that $\widehat{\varphi}_j$ is the Δ_2^0 characteristic function of $\{i : W_i \cap \overline{D}\}$. Hence, for all i , $\lim_s \widetilde{\varphi}_{j,s}(i)$ exists and equals $\widehat{\varphi}_j(i)$. (Here $\widehat{\varphi}_j$ and $\widetilde{\varphi}_{j,s}(i)$ were defined in §3.1.)

Suppose $\widetilde{\varphi}_{j,s}(g(\beta)) \downarrow = 0$ for all $s \geq s_0$ for some s_0 . Since $\beta \prec f$ there is some stage $s_1 > s_0$ after which β is never initialized. Also for all α , $\alpha \widehat{<} 0 \prec \beta$, $\widehat{\psi}_\alpha$ is total by Lemma 3.3. Hence, β can wait for a witness $y = x_n^\beta$ in (72) to appear at some stage $x > s_1$. Then according to Case 3, Subcase 1, β will enumerate x_n^β into $Z_\beta = W_{g(\beta)}$ causing $Z_\beta \cap \overline{D} \neq \emptyset$. Since $\widetilde{\varphi}_{j,t}(g(\beta)) = \widetilde{\varphi}_{j,s}(g(\beta))$ for all $t > s$, β will never enumerate x_n^β into D , so $\lim_s \widetilde{\varphi}_{j,s}(g(\beta)) = 0$, but $W_{g(\beta)} \cap \overline{D} \neq \emptyset$, a contradiction.

Next suppose $\lim_s \widetilde{\varphi}_{j,s}(g(\beta)) \downarrow = 1$ for all $s \geq s_0$ for some s_0 . For every s there can be at most one element in $Z_s^\beta - D_s$ and that must be $\Gamma_s = x_n^\beta$ for some n . Since $\beta \prec f$ we know that every ξ with $\xi \widehat{<} 0 \prec \beta$ will have the Π_2 -outcome. Hence, for each such x_n^β , there will come a stage by (73) at which x_n^β is enumerated in D under Subcase 2. But no new β -witness can be appointed after stage s_0 under Subcase 1. Hence, $\lim_s \widetilde{\varphi}_{j,s}(g(\beta)) \downarrow = 1$, but $W_{g(\beta)} \cap \overline{D} = \emptyset$, a contradiction. □

While we do not use this theorem itself in our proof of Theorem 1.3, we will have the same positive requirements in our theorem, as well as a nearly identical interaction between the positive and negative requirements. In the proof of Theorem 1.3, notice the similarity between Lemma 3.3 above and Lemma 8.8. More directly, Lemma 3.4 appears below with some minor changes as Lemma 8.16.

4. PROOF STRATEGY

As in Theorem 3.1, we will build our c.e. set D on a tree. The positive requirements that make D nonlow are exactly the same as in the theorem above. However, the negative requirements will now be part of the automorphism construction. Our tree will be much more intricate than the tree used in Theorem 3.1 because it will include the entire automorphism construction.

Recall that we are trying to show the following theorem:

Theorem 1.3. *There is a nonlow c.e. set D such that for all c.e. $A \leq_T D$, there exists a low c.e. set B and an automorphism of \mathcal{E} taking A to B .*

We will consider the case for a *single* set $A = \Psi^D = W_i$. The theorem follows as explained in §4.7.

4.1. The negative requirements. Let us first examine the special case where $D = A$. The set B is low, which means that it satisfies a stability property saying that if infinitely many elements enter W_e before entering B , then infinitely many elements remain in $W_e \cap \overline{B}$. In order to guarantee an automorphism taking A to B , we will need to match the flow of elements into A with a flow of elements into B , so we will need some sort of stability on A to meet the stability of B . However, if we guarantee this stability property on the whole tree, we would in fact have that \overline{A} was semi-low, which is exactly what the positive requirements are guaranteeing that it is not. What we will actually achieve is stability along the true path only. Because the stability is only along the true path, A will not be semi-low, but will have enough stability to build the automorphism. In Theorem 3.1, this is analogous to the pseudo-requirement of making D low in order to actually achieve that D is just low₂.

For the general case when $A \neq D$, the pseudo-requirement of stability will be the same. However, we cannot restrain elements from A directly, so we will instead restrain D , which will in turn keep the desired elements out of A , via the reduction $A = \Psi^D$.

4.2. The Harrington-Soare automorphism method. We must build a nonlow set D , a low set B , and an automorphism from $A = \Psi^D$ to B . We modify the Harrington-Soare automorphism method as explained below, while simultaneously adding elements to D to ensure that D is nonlow and restraining D to ensure that A is automorphic to B .

Harrington and Soare developed their Δ_3^0 automorphism construction in [4]. The construction is performed on a computable tree. Take the lattice \mathcal{E} and quotient out by the ideal of finite sets to get the lattice \mathcal{E}^* of c.e. sets up to finite difference. Soare [15, page 343] showed that if there is an automorphism of \mathcal{E}^* taking A to B , then there is one of \mathcal{E} as well, so we can work in \mathcal{E}^* . We first make two lists $\{U_i\}_{i \in \omega}$ and $\{V_i\}_{i \in \omega}$ of all the c.e. sets up to finite difference. We let $U_0 = A$. We imagine the U_i sets as being on the A -side of the construction, and the V_i sets as being on the B -side. We build c.e. sets $\{\widehat{U}_i\}_{i \in \omega}$ and $\{\widehat{V}_i\}_{i \in \omega}$ such that there is an automorphism taking U_i to \widehat{U}_i . The inverse of the automorphism will take V_i to \widehat{V}_i . We define B to be \widehat{U}_0 .

The e -state of an element x tells us which c.e. sets U_i and \widehat{V}_i , or \widehat{U}_i and V_i , contain the element x , for all $i < e$. In order to build an automorphism taking a set A to a set B , it suffices to ensure that the same e -states contain infinitely many elements on each side of the construction.

We build two identical trees, T and \widehat{T} , according to rules that will be explained later. Elements move through the nodes on the trees, starting from the root. If x is at node α or some node δ that extends α on T , then x can be enumerated into the sets U_α and \widehat{V}_α , where $U_\alpha =^* U_{e_\alpha}$ for some $e_\alpha \in \omega$ such that the coding is invariant

over nodes of the same length and similarly for elements on \widehat{T} . The automorphism we build takes U_α to \widehat{U}_α and V_α to \widehat{V}_α for all α on the true path, which is Δ_3^0 .

We can think of the construction as a game between the Red and Blue players. The Red player (thought of as the opponent) controls the sets U_i and V_i , and the Blue player (thought of as us) controls \widehat{U}_i and \widehat{V}_i . The Blue player wins if the sets define an automorphism. In the Harrington-Soare construction of [4], if Red enumerates an element x at node α into the set $A = U_0$, then Blue matches it by enumerating the first available \widehat{x} into B such that x and \widehat{x} are in the same e_α -state.

However, in our theorem, we want B to be low, so Blue cannot be allowed to choose any element to put into B . It is problematic if Red enumerates so many elements into A that Blue must either ignore the lowness restraint on B or fail to put enough elements into B to build an automorphism. We must therefore ensure that Red always leaves infinitely many elements from every e -state in \overline{A} . In other words, we must satisfy the stability property from §4.1, at least along the true path.

4.3. The modified automorphism method. We use a method similar to the Harrington-Soare Δ_3^0 automorphism construction. However, we must make B low, which means that we must have a restraint on B to guarantee stability. In the Harrington-Soare construction of [4], elements moved among the nodes on two copies of our tree of strategies, corresponding to the A/\overline{A} and B/\overline{B} sides of the construction. Now, if we let A have free rein, elements may enter A that will not be matched on the B side due to the restraint on B . We must therefore keep elements in \overline{A} by somehow restraining A . As we don't control A itself, the only way to do this is to put a restraint on the set D .

In our new construction, we build four identical trees, corresponding to \overline{A} , A , \overline{B} , and B . Elements move through the \overline{A} and \overline{B} trees until they are enumerated into A or B , when they switch to the corresponding tree. We now must take into account the fact that elements are disappearing from two of the trees and appearing in the others. These mechanics have not appeared before now.

In addition to the nodes on the tree that deal with the automorphism, we must put nodes on the tree corresponding to the positive requirement of making D non-low. The automorphism nodes themselves act as negative requirement nodes. They tell us which elements we want to restrain to keep out of A .

As mentioned previously, the mechanics of the positive and negative requirements are very similar to those involved in constructing a nonlow c.e. set that is low_2 as in §3.

4.4. A tree argument for automorphisms. We perform our construction on a tree as in §3.5, but with the addition of automorphism nodes that will also act as negative requirement nodes. We shall define in §6.2 a computable tree T with true path f . If $\alpha \in T$ and $\alpha \widehat{a} \prec f$, then a is the correct guess at the outcome of α .

For each $n \in \omega$ there is some $m_n \in \omega$ such that for every $\alpha \in T$ of length m_n , \widehat{U}_α will be a potential candidate for \widehat{U}_n and if $\alpha \prec f$, then $U_\alpha =^* U_n$ and \widehat{U}_α will be the correct candidate for \widehat{U}_n . Thus, f will specify the sequence $\{\widehat{U}_{f \upharpoonright m_n}\}_{n \in \omega}$ which will be the desired sequence $\{\widehat{U}_n\}_{n \in \omega}$ and similarly for V_n . In a tree construction f is not in general computable but only $\mathbf{0}''$ -computable, so the sequences $\{\widehat{U}_n\}_{n \in \omega}$ and $\{\widehat{V}_n\}_{n \in \omega}$ will only have a $\mathbf{0}''$ -computable (i.e., Δ_3^0) presentation.

We say an e -state is *well-visited* if infinitely many elements enter that state during the construction. An e -state is *well-resided* if infinitely many elements remain in

that state, and *emptied* if it is well-visited but not well-resided. If $\alpha \in T$ and $|\alpha| = 5e + j$, for $0 < j \leq 4$, then associate with α a version of the automorphism strategy. Each α will have associated sets \mathcal{M}_α , \mathcal{B}_α , and \mathcal{R}_α , where \mathcal{M}_α is a set of e -states, $\mathcal{R}_\alpha \subseteq \mathcal{M}_\alpha$, and $\mathcal{B}_\alpha \subseteq \mathcal{M}_\alpha - \mathcal{R}_\alpha$. These represent α 's guess at which states are well-visited, states whose emptying is attributed to the Red player, and states whose emptying is attributed to the Blue player, respectively. In addition each α has an associated $k_\alpha \in \omega$, which represents α 's guess at the least element such that no larger element will ever enter a non-well-visited e -state. Further restrictions for which nodes appear on the tree are given in Definition 6.2.

Each of the sets \mathcal{M}_α , etc., will be divided into two subsets \mathcal{M}_α^A and $\mathcal{M}_\alpha^{\bar{A}}$, depending on whether or not $A = U_0$ is in each state. We can think of these as being on two different trees, for A and \bar{A} . In addition, we will have duals $\widehat{\mathcal{M}}_\alpha$, etc., corresponding to B and \bar{B} . As mentioned in §4.3, we will refer to these as four separate, but identical, trees, one corresponding to each of A, \bar{A}, B , and \bar{B} .

If $\beta \in T$ and $|\beta| = 5e + 4$, then associate with β a version of the strategy for P_e , and put the outcomes $\beta \hat{\ } 0$ and $\beta \hat{\ } p$ for all $p \geq 1$ on T , representing the Π_2 outcome and all possible Σ_2 -outcomes of the β -strategy. The nodes below β have order type $1 + \omega^*$ with $\beta \hat{\ } 0$ on the left, and the others to the right arranged in the order of ω^* . Note that the nodes guessing at the positive strategy are of length $5e + 5$, so these do not conflict with β 's own guesses for the automorphism requirements. We also associate $\beta \hat{\ } p$ with $\mathcal{M}_{\beta \hat{\ } p} = \mathcal{M}_\beta$, etc. The true path f on T and its computable approximation $\{f_s\}_{s \in \omega}$ such that $f = \liminf_s f_s$ will be defined below.

Suppose $\beta \in T$ and $|\beta| = 5j + 4$. Then β has an associated witness function Γ_s^β for carrying out its strategy for P_j as in the construction of a low₂ set that is not low in §3.2.

Suppose $\alpha \in T$. Let $A = \Psi^D$. The stage s guess at whether y is in A is $\Psi^D(y)[s] = \Psi_s^{D_s}(y)$. The use of this computation is denoted by $\psi_s(y)$. The following functions are modifications of those in (7) and (8). Define

$$(19) \quad \Lambda_s^\alpha = \min\{\Gamma_s^\beta : \beta \hat{\ } 0 \preceq \alpha \ \& \ |\beta| \equiv 4 \pmod{5}\}, \text{ and}$$

$$(20) \quad \widehat{\psi}_{\alpha,s}(y) = \begin{cases} \psi_{e,s}(y) & \text{if } \psi_{e,s}(y) \downarrow < \Lambda_s^\alpha \text{ and (for all } |\beta| \equiv 4 \pmod{5}) \\ & [\beta \hat{\ } k \preceq \alpha \implies \text{all } k - 1 \text{ } \beta\text{-witnesses have entered } D] \\ \text{undefined} & \text{otherwise.} \end{cases}$$

For notational simplicity, we will write $\widehat{\Psi}_\alpha(x)[s] = 0$ to denote $\Psi^D(x)[s] = 0$ and $\widehat{\psi}_{\alpha,s}(x) \downarrow$. This means that at stage s , D_s believes that $x \notin A$, and all nodes $\beta \prec \alpha$ will no longer injure the computation.

4.5. Making $\mathcal{L}^*(A) \cong \mathcal{L}^*(B)$. We know by Soare [14] that since B is low, $\mathcal{L}^*(B) \cong \mathcal{E}^*$, where $\mathcal{L}^*(C)$ is the lattice of c.e. sets modulo finite difference intersected with the complement of C (this is isomorphic to the lattice of c.e. sets modulo finite difference containing C). The negative requirements stem from the automorphism requirements, which must guarantee that $\mathcal{L}^*(A) \cong \mathcal{L}^*(B)$. We use a modified version of the automorphism construction in Harrington and Soare [4]. Elements flow through the tree as in that paper, but when they enter $A = U_0$, they disappear from the tree (as they switch from the \bar{A} -tree to the A -tree).

We would like to match the well-resided e -states that do not contain A with the well-resided e -states that do not contain B . However, elements may be falling into

A that we cannot match by putting elements into B because B is low. Thus, we need to restrain elements in \bar{A} so that no well-visited state in $\mathcal{M}_\alpha^{\bar{A}}$ is emptied by A alone. In other words, we need stability.

For each $k \in \omega$ and $\nu_e \in \mathcal{M}_\alpha$, we will choose an element $y_{\langle e,k \rangle} \in \nu_e$ and restrain D to keep $y_{\langle e,k \rangle}$ out of A . This restraint will be the modified use function $\widehat{\psi}_\alpha(y_{\langle e,k \rangle})[s]$. In fact, we will not put x into any set \widehat{V}_α or U_α unless $\widehat{\Psi}_\alpha^D(x)[s] = 0$.

For a given e with $\nu_e \in \mathcal{M}_\alpha^{\bar{A}}$ and stage s , define $y_{\langle e,k \rangle}^s$ by recursion on k . This is our current guess at $y_{\langle e,k \rangle}$, where $y_{\langle e,k \rangle}$ is an element permanently in \bar{A} . Suppose $y_{\langle e,i \rangle}^s$ has been defined for all $i < k$. Let $y_{\langle e,k \rangle}^s = y_{\langle e,k \rangle}^{s-1}$ if $\widehat{\Psi}_\alpha(y_{\langle e,k \rangle}^{s-1})[s-1] = 0$. Otherwise, check if there is a y that entered ν_e at stage s under Step 1 of the construction in §7.3 (or Step 6C if $\alpha = \lambda$) such that

$$(21) \quad y > \max\{y_{\langle e,k-1 \rangle}^s, y_{\langle e,k \rangle}^t \mid t < s\}, \quad y \in \nu_{e,s}, \quad \text{and} \quad \widehat{\Psi}_\alpha(y)[s-1] = 0.$$

If such a y exists, set it as $y_{\langle e,k \rangle}^s$. Otherwise, $y_{\langle e,k \rangle}^s$ is undefined. To summarize, $y_{\langle e,k \rangle}^s$ is a value that enters ν_e under Step 1 and appears to not be in A . It stays as $y_{\langle e,k \rangle}^s$ until it no longer appears to be in \bar{A} .

For a positive requirement β of lower priority, β can appoint the n^{th} witness, x_n^β , only after $\widehat{\psi}_{\alpha,s}(y_{\langle e,k \rangle}) \downarrow$ for all $\langle \alpha, e, k \rangle \leq \langle \beta, n \rangle$, and must exceed their values. (Here, we identify the nodes of T with their Gödel numbering under some fixed effective numbering of T .) We will show that for each e and k the computation $\widehat{\psi}_{\alpha,s}(y_{\langle e,k \rangle}^s)$ can be injured at most finitely often. The proof is similar to the proof that the negative requirements are satisfied in the construction of a low₂ set that is nonlow.

4.6. Building B . We must also build B low such that $\mathcal{M}_\alpha^A = \widehat{\mathcal{M}}_\alpha^B$. That is, we want to match the well-visited states that contain A with the well-visited states that contain B . So far we have only discussed keeping elements out of A , but not putting elements into B . We will ensure $\mathcal{G}_\alpha = \widehat{\mathcal{G}}_\alpha$, where

$$(22) \quad \mathcal{G}_\alpha = \{\nu \mid (\exists^\infty x)(\exists s)[x \in A_{\text{at } s} \text{ and } \nu(\alpha, x, s) = \nu]\}$$

and $\widehat{\mathcal{G}}_\alpha$ is defined similarly for B in place of A . We can then force $\mathcal{M}_\alpha^A = \widehat{\mathcal{M}}_\alpha^B$ by performing the automorphism steps on elements that have already entered A or B .

To make B low, we will meet the negative requirement:

$$(23) \quad N_j^B : \quad (\exists^\infty s) \Phi_j^B(j)[s] \downarrow \implies \Phi_j^B(j) \downarrow .$$

In addition, we must ensure that no well-visited state is emptied by B . To satisfy this, we will wait for two new elements to come into the state before allowing one of them to enter B .

For the positive requirement, we need to ensure that $\mathcal{G}_\alpha = \widehat{\mathcal{G}}_\alpha$. To do this, we meet:

$$(24) \quad P_{e,x}^B : \quad x \text{ enters } A \text{ from state } \nu_e \implies (\exists y)[y \text{ enters } B \text{ from } \widehat{\nu}_e].$$

When x enters A from α -state ν_e , we add $\langle \alpha, x, e \rangle$ to a list Λ , and later find a witness y to enumerate into B in the state $\widehat{\nu}_e$.

4.7. The hidden master tree. In order to show that $\overline{\mathbf{L}}_1$ is noninvariant, we must show that for every c.e. $A \leq_T D$, there is a low B such that A is automorphic to B . However, so far we have only discussed this for a single $A = W_i = \Psi^D = \Phi_e^D$. To show this for all $A \leq_T D$, we will use a much bigger master tree with infinitely many levels on the tree corresponding to each pair $\langle e, i \rangle$, representing the possibility that $A = W_i = \Phi_e^D$.

Let $n_{\langle e, i \rangle}$ be the least of any of the lengths corresponding to $\langle e, i \rangle$. Then nodes of length $n_{\langle e, i \rangle}$ will be extended by guesses at whether we believe $\Phi_e^D = W_i$. Let α have length $n_{\langle e, i \rangle}$. Then $\alpha \hat{=} 0$ represents the guess that $W_i = \Phi_e^D$, and $\alpha \hat{=} 1$ represents the guess that $W_i \neq \Phi_e^D$. We say s is an α -state if the approximation to the true path at stage s contains α . If s is an α -state and the length of agreement between $W_{i,s}$ and $\Phi_e^D[s]$ has increased since the previous α -state, we let our approximation to the true path at stage s go through $\alpha \hat{=} 0$, and otherwise, through $\alpha \hat{=} 1$. For each α of length $n_{\langle e, i \rangle}$, we create a tree T_α made up of the nodes affiliated with $\langle e, i \rangle$ that extend $\alpha \hat{=} 0$ as well as all positive requirement nodes extending $\alpha \hat{=} 0$. Note that T_α will already be guessing at all positive requirement nodes that α extends. If these guesses are incorrect, then only finitely much activity will occur on T_α . The automorphism machinery for $\langle e, i \rangle$ will only act at stages when the approximation to the true path goes through $\alpha \hat{=} 0$. This machinery acts upon the tree T_α . This is the tree mentioned in §4.4 and throughout the rest of the proof, for a fixed e, i , and α .

The ordering of the nodes on the master tree is as follows: height 1 nodes guess at whether $W_0 = \Phi_0^D$, and the next four levels correspond to the first four automorphism requirement nodes on the tree $T_0 = T_{\langle 0, 0 \rangle}$, followed by height 6 nodes, which are guesses at the number of witnesses for P_0 . These are followed by the guess for $\langle e, i \rangle = 1$, then the second four automorphism levels of T_0 and the first four automorphism levels of T_1 , and then P_1 . Thus, the space between the P_i 's increases as more and more automorphism nodes are added.

In the end, the trees T_α will look and act exactly as the tree T in the other sections of this proof with one small difference. Each positive node will need to respect finitely many restraints put on by the finitely many additional $T_{\alpha'}$, for $\alpha' <_L \alpha$. As the restraint is finite, this is a negligible change.

To summarize, the addition of multiple A 's does not change the theorem except that it adds more nodes to the tree that must obey the same rules. It will cause D to be restrained more. The case for a single e and i is the heart of the problem. We need not refer back to this master tree, as it simply tells us that we may perform the construction for a specific $A = W_i = \Phi_e^D$ and that we may weave these constructions together to get the general proof.

5. AN OVERVIEW OF THE AUTOMORPHISM MACHINERY

This section, which contains important definitions, is largely taken from the Harrington-Soare Δ_3^0 automorphism paper [4]. The few changes are primarily in differentiating between the A and \overline{A} parts of the proof.

5.1. Background. As mentioned previously, by [15, page 343], building an automorphism of \mathcal{E} is equivalent to building one of \mathcal{E}^* , the quotient lattice of \mathcal{E} modulo the ideal \mathcal{F} of finite sets. To do this we fix two copies of the natural numbers ω and $\widehat{\omega}$. We let variables $x, y, \dots (\hat{x}, \hat{y}, \dots)$ range over ω ($\widehat{\omega}$). Occasionally, we shall

specify the definitions and action for only the ω -side, meaning that the $\widehat{\omega}$ -side will be entirely dual or simplified in an obvious way. However, there are quite a few differences between the ω and $\widehat{\omega}$ sides that were not needed in the Harrington-Soare paper [4], and we will make these differences explicit.

As mentioned in §4.2, we view the construction of the automorphism Φ as a game between two players in the sense of Lachlan [7]. The Red player produces two standard indexings $\{U_n\}_{n \in \omega}$ and $\{V_n\}_{n \in \omega}$ of the c.e. sets (up to finite difference), where we view U_n as being on the ω -side and V_n on the $\widehat{\omega}$ -side. The Blue player responds by building c.e. sets $\{\widehat{U}_n\}_{n \in \omega}$ on the $\widehat{\omega}$ -side and $\{\widehat{V}_n\}_{n \in \omega}$ on the ω -side. The condition necessary to show that this correspondence $\Phi(U_n) = \widehat{U}_n$ and $\widehat{V}_n = \Phi^{-1}(V_n)$ is an automorphism is best stated in terms of the following notion of full e -state. We restrict our indexings $\{U_n\}_{n \in \omega}$ to those where $U_0 = W_i = A$.

Definition 5.1. Given two sequences of c.e. sets $\{X_n\}_{n \in \omega}$ and $\{Y_n\}_{n \in \omega}$, define $\nu(e, x)$, the *full e -state* of x with respect to (w.r.t.) $\{X_n\}_{n \in \omega}$ and $\{Y_n\}_{n \in \omega}$ to be the triple $\langle e, \sigma(e, x), \tau(e, x) \rangle$, where

$$\begin{aligned} \sigma(e, x) &= \{i : i \leq e \ \& \ x \in X_i\}, \text{ and} \\ \tau(e, x) &= \{i : i \leq e \ \& \ x \in Y_i\}. \end{aligned}$$

To see that Φ is an automorphism it suffices to satisfy the requirement,

$$(25) \quad \begin{aligned} (\forall \nu)(\exists^\infty x \in \omega)[\nu(e, x) = \nu \text{ w.r.t. } \{U_n\}_{n \in \omega} \text{ and } \{\widehat{V}_n\}_{n \in \omega}] \\ \iff (\exists^\infty \hat{y} \in \hat{\omega})[\nu(e, \hat{y}) = \nu \text{ w.r.t. } \{\widehat{U}_n\}_{n \in \omega} \text{ and } \{V_n\}_{n \in \omega}]. \end{aligned}$$

Definition 5.2. Given computable enumerations $\{X_s\}_{s \in \omega}$ and $\{Y_s\}_{s \in \omega}$ of c.e. sets X and Y , define $X \setminus Y = \{z : (\exists s)[z \in X_s - Y_s]\}$.

5.2. The α -section S_α , α -region R_α , and c.e. set Y_α . We divide up the ω -side into disjoint α -sections, S_α , for $\alpha \in T$. We shall define during the construction in §7 a function $\alpha(x, s)$ with range T which indicates that x is in section $S_{\alpha(x, s)}$ at the end of stage s , and we shall guarantee that $\alpha(x) = \lim_s \alpha(x, s)$ exists. The α -region R_α consists of all S_γ such that $\alpha \preceq \gamma$. For each stage s we define

$$\begin{aligned} S_{\alpha, s} &= \{x : \alpha(x, s) = \alpha\}, \\ R_{\alpha, s} &= \{x : \alpha \preceq \alpha(x, s)\}, \text{ and} \\ Y_{\alpha, s} &= \bigcup \{R_{\alpha, t} : t \leq s\}. \end{aligned}$$

Define $S_{\alpha, \infty} = \{x : \alpha(x) = \alpha\}$ and $R_{\alpha, \infty} = \{x : \alpha(x) \preceq \alpha\}$. Note that these sets contain all x in either the A -tree or the \overline{A} -tree. When we need to distinguish between them, we will write them as $S_{\alpha, s}^A, S_{\alpha, s}^{\overline{A}}$, etc.

We shall guarantee that for all $\alpha \in T, \alpha \neq \lambda$,

$$(26) \quad Y_\alpha \setminus Y_{\alpha^-} = \emptyset, \text{ and}$$

$$(27) \quad \begin{aligned} \alpha \prec f \implies \quad R_{\alpha, \infty} =^* Y_\alpha =^* \omega, \\ R_{\alpha, \infty}^A =^* Y_\alpha^A =^* A, \text{ and} \\ R_{\alpha, \infty}^{\overline{A}} =^* Y_\alpha^{\overline{A}} =^* \overline{A}. \end{aligned}$$

We shall ensure (26) by making x enter S_{α^-} before x enters R_α . Also x will enter R_α at most once (although x may later leave R_α). During the construction in §7 we shall define a computable sequence $\{f_s\}_{s \in \omega}$ such that $f = \liminf_s f_s$.

If $f_s <_L \alpha$ for some $s \geq x$ we say x is α -ineligible at all stages $t \geq s$, and we insist that $x \notin S_{\alpha,t}$. Hence, $R_{\alpha,\infty} = \emptyset$ for all α with $f <_L \alpha$. Secondly, Y_α will be finite for all $\alpha <_L f$. Finally, $S_{\alpha,\infty}$ will be finite for all α . These three facts imply (27).

5.3. The α -states $\nu(\alpha, x, s)$, and lists $\mathcal{E}_\alpha, \mathcal{F}_\alpha, \mathcal{M}_\alpha$. For conceptual simplicity we do as little action as possible at each node $\alpha \in T$. If $|\alpha| \equiv 1 \pmod 5$ ($|\alpha| \equiv 2 \pmod 5$), we consider one new U set (V set). If $|\alpha| \equiv 3 \pmod 5$ ($|\alpha| \equiv 4 \pmod 5$), we consider new α -states $\nu(\hat{\nu})$ which may be non-well-resided on Y_α (\hat{Y}_α). If $|\alpha| \equiv 0 \pmod 5$ we make no new commitments for the automorphism machinery but we instead act for the positive requirements. We shall arrange for all $n \in \omega$ that for $\alpha < f$,

$$(28) \quad |\alpha| = 5n + 1 \implies U_\alpha =^* U_n, \text{ and}$$

$$(29) \quad |\alpha| = 5n + 2 \implies V_\alpha =^* V_n.$$

We let U_α and \hat{U}_α (V_α and \hat{V}_α) be undefined if $|\alpha| \not\equiv 1 \pmod 5$ ($|\alpha| \not\equiv 2 \pmod 5$). We let $e_\alpha(\hat{e}_\alpha)$ correspond to n in (28) (respectively (29)). Specifically, define $e_\lambda = \hat{e}_\lambda = -1$ and if $|\alpha| \equiv 1 \pmod 5$, then let $e_\alpha = e_{\alpha^-} + 1$, and otherwise let $e_\alpha = e_{\alpha^-}$. Define \hat{e}_α similarly with $|\alpha| \equiv 2 \pmod 5$ in place of $|\alpha| \equiv 1 \pmod 5$. Hence, $e_\alpha > e_{\alpha^-}$ ($\hat{e}_\alpha > \hat{e}_{\alpha^-}$) if and only if $|\alpha| \equiv 1 \pmod 5$ ($|\alpha| \equiv 2 \pmod 5$).

Definition 5.3. An α -state is a triple $\langle \alpha, \sigma, \tau \rangle$ where $\sigma \subseteq \{0, \dots, e_\alpha\}$ and $\tau \subseteq \{0, \dots, \hat{e}_\alpha\}$. The only λ -state is $\nu_{-1} = \langle \lambda, \emptyset, \emptyset \rangle$.

The construction in §7 will produce a simultaneous computable enumeration $U_{\alpha,s}, V_{\alpha,s}, \hat{U}_{\alpha,s}, \hat{V}_{\alpha,s}$, for $\alpha \in T$ and $s \in \omega$, of these c.e. sets which we use in the following definition.

Definition 5.4. (i) The α -state of x at stage s , $\nu(\alpha, x, s)$, is the triple $\langle \alpha, \sigma(\alpha, x, s), \tau(\alpha, x, s) \rangle$, where

$$\sigma(\alpha, x, s) = \{e_\beta : \beta \preceq \alpha \ \& \ e_\beta > e_{\beta^-} \ \& \ x \in U_{\beta,s}\},$$

$$\tau(\alpha, x, s) = \{\hat{e}_\beta : \beta \preceq \alpha \ \& \ \hat{e}_\beta > \hat{e}_{\beta^-} \ \& \ x \in \hat{V}_{\beta,s}\}.$$

(ii) The final α -state of x is $\nu(\alpha, x) = \langle \alpha, \sigma(\alpha, x), \tau(\alpha, x) \rangle$, where $\sigma(\alpha, x) = \lim_s \sigma(\alpha, x, s)$ and $\tau(\alpha, x) = \lim_s \tau(\alpha, x, s)$.

For each $\alpha \in T$ we define the following sets of α -states called lists:

$$\mathcal{E}_\alpha^C = \{\nu : (\exists^\infty x)(\exists s)[x \in S_{\alpha,s}^C - \bigcup \{S_{\alpha,t} : t < s\} \ \& \ \nu(\alpha, x, s) = \nu]\},$$

where C is either A or \bar{A} . We define $\mathcal{E}_\alpha = \mathcal{E}_\alpha^A \cup \mathcal{E}_\alpha^{\bar{A}}$. Let

$$\mathcal{F}_\alpha = \{\nu : (\exists^\infty x)(\exists s)[x \in R_{\alpha,s} \ \& \ \nu(\alpha, x, s) = \nu]\}.$$

We may also define \mathcal{F}_α^A and $\mathcal{F}_\alpha^{\bar{A}}$ by replacing $R_{\alpha,s}$ with $R_{\alpha,s}^A$ or $R_{\alpha,s}^{\bar{A}}$. Then $\mathcal{F}_\alpha = \mathcal{F}_\alpha^A \cup \mathcal{F}_\alpha^{\bar{A}}$.

Note that \mathcal{E}_α consists of states well-visited by elements x when they first enter R_α , and \mathcal{F}_α those states well-visited while they remain in Y_α , so $\mathcal{E}_\alpha \subseteq \mathcal{F}_\alpha$. Note that \mathcal{E}_α^A is the set of all states that are well-visited as entry states by elements that have never been in S_α . This excludes entry states that are well-visited only by elements switching from $S_\alpha^{\bar{A}}$ to S_α^A . However, we will show that in fact these entry states are included, as they will also be well-visited by elements that have not yet been in S_α . Each $\alpha \in T$ will have an associated list \mathcal{M}_α which is roughly

α 's "guess" at the true \mathcal{F}_α such that if $\alpha \prec f$, then $\mathcal{M}_\alpha = \mathcal{F}_\alpha$. For $\alpha \prec f$ we shall achieve $\mathcal{M}_\alpha = \mathcal{F}_\alpha$ by ensuring the following properties of \mathcal{M}_α :

$$(30) \quad \mathcal{E}_\alpha \subseteq \mathcal{M}_\alpha,$$

$$(31) \quad (\forall^\infty x) \left[\begin{array}{l} \text{if } x \in Y_{\alpha,s}, \nu_0 = \nu(\alpha, x, s) \in \mathcal{M}_\alpha, \\ \text{and Red causes enumeration of } x \text{ so that} \\ \nu_1 = \nu(\alpha, x, s + 1), \text{ then } \nu_1 \in \mathcal{M}_\alpha \end{array} \right],$$

$$(32) \quad (\forall^\infty x) \left[\begin{array}{l} \text{if } x \in Y_{\alpha,s}, \nu_0 = \nu(\alpha, x, s) \in \mathcal{M}_\alpha, \\ \text{and Blue causes enumeration of } x \text{ so that} \\ \nu_1 = \nu(\alpha, x, s + 1), \text{ then } \nu_1 \in \mathcal{M}_\alpha \end{array} \right].$$

(Here $(\forall^\infty x)$ denotes "for almost every x ".) Blue enumeration which satisfies (32) is called α -legal. Two main constraints on Blue's moves will be (30) and (32). Clearly, (30), (31), and (32) guarantee

$$(33) \quad \mathcal{F}_\alpha \subseteq \mathcal{M}_\alpha.$$

During Step 1 of the construction in §7 we shall promptly pull elements $x \in Y_{\alpha-,s}$ into $S_{\alpha,s+1}$ in order to ensure

$$(34) \quad \mathcal{M}_\alpha \subseteq \mathcal{E}_\alpha.$$

Hence, by (33), (34), and $\mathcal{E}_\alpha \subseteq \mathcal{F}_\alpha$ we have

$$(35) \quad \mathcal{M}_\alpha = \mathcal{F}_\alpha = \mathcal{E}_\alpha.$$

On the $\widehat{\omega}$ -side we have dual definitions for the above items by replacing $\omega, x, U_\alpha, \widehat{V}_\alpha, A, \overline{A}$ by $\widehat{\omega}, \widehat{x}, \widehat{U}_\alpha, V_\alpha, B, \overline{B}$, respectively. These dual items will be denoted by $\widehat{\nu}(\alpha, \widehat{x}, s), \widehat{S}_\alpha, \widehat{R}_\alpha, \widehat{Y}_\alpha, \widehat{\mathcal{E}}_\alpha, \widehat{\mathcal{F}}_\alpha$, and $\widehat{\mathcal{M}}_\alpha$. We write hats over the α -states, e.g. $\widehat{\nu}_1 = \nu(\alpha, \widehat{x}, s)$, to indicate α -states for elements $\widehat{x} \in \widehat{\omega}$. We shall ensure

$$(36) \quad \widehat{\mathcal{M}}_\alpha = \{\widehat{\nu} : \nu \in \mathcal{M}_\alpha\},$$

which implies by (35) that the well-visited α -states on both sides coincide. We also define \mathcal{M}_α^A , etc. when we restrict to the A -tree or the \overline{A} -tree.

Definition 5.5. Given α -states $\nu_0 = \langle \alpha, \sigma_0, \tau_0 \rangle$ and $\nu_1 = \langle \alpha, \sigma_1, \tau_1 \rangle$, we have:

- (i) $\nu_0 \leq_R \nu_1$ if $\sigma_0 \subseteq \sigma_1$ and $\tau_0 = \tau_1$.
- (ii) $\nu_0 \leq_B \nu_1$ if $\tau_0 \subseteq \tau_1$ and $\sigma_0 = \sigma_1$.
- (iii) $\widehat{\nu}_0 \leq_R \widehat{\nu}_1$ if $\widehat{\sigma}_0 = \widehat{\sigma}_1$ and $\widehat{\tau}_0 \subseteq \widehat{\tau}_1$.
- (iv) $\widehat{\nu}_0 \leq_B \widehat{\nu}_1$ if $\widehat{\sigma}_0 \subseteq \widehat{\sigma}_1$ and $\widehat{\tau}_0 = \widehat{\tau}_1$.
- (v) $\nu_0 <_R \nu_1$ ($\nu_0 <_B \nu_1$) if $\nu_0 \leq_R \nu_1$ ($\nu_0 \leq_B \nu_1$) and $\nu_0 \neq \nu_1$, and similarly for $\widehat{\nu}_0 <_R \widehat{\nu}_1$ and $\widehat{\nu}_0 <_B \widehat{\nu}_1$.

The intuition is that if $\nu_0 = \nu(\alpha, x, s)$ and $\nu_0 <_R \nu_1$ ($\nu_0 <_B \nu_1$), then Red (Blue) can enumerate x in the necessary U sets (\widehat{V} sets) causing $\nu_1 = \nu(\alpha, x, s + 1)$. For $\widehat{\nu}_0$ and $\widehat{\nu}_1$ the role of σ and τ is reversed because on the $\widehat{\omega}$ -side Blue (Red) plays the \widehat{U} sets (V sets), and hence

$$(37) \quad [\nu_0 <_R \nu_1 \iff \widehat{\nu}_0 <_B \widehat{\nu}_1] \ \& \ [\nu_0 <_B \nu_1 \iff \widehat{\nu}_0 <_R \widehat{\nu}_1].$$

Definition 5.6. Let $\beta \preceq \alpha \in T$, ν_0 be an α -state $\nu_0 = \langle \alpha, \sigma_0, \tau_0 \rangle$, and \mathcal{C}_α be a set of α -states. Define

- (i) $\nu_0 \upharpoonright \beta = \langle \beta, \sigma_1, \tau_1 \rangle$, where we define $\sigma_1 = \sigma_0 \cap \{0, \dots, e_\beta\}$ and we define $\tau_1 = \tau_0 \cap \{0, \dots, \widehat{e}_\beta\}$,
- (ii) $\nu_1 \preceq \nu_0$ (read " ν_0 extends ν_1 ") if $\nu_0 \upharpoonright \beta = \nu_1$, and

(iii) $\mathcal{C}_\alpha \upharpoonright \beta = \{\nu \upharpoonright \beta : \nu \in \mathcal{C}_\alpha\}$.

(iv) Given a finite set of α -states, $\{\nu(\alpha, \sigma_i, \tau_i)\}_{i \in I}$, we then define $\bigcup_{i \in I} \{\nu(\alpha, \sigma_i, \tau_i)\} := \langle \alpha, \sigma, \tau \rangle$, where $\sigma = \bigcup_{i \in I} \sigma_i$, and $\tau = \bigcup_{i \in I} \tau_i$.

Definition 5.7. A node $\alpha \in T$ is \mathcal{M} -inconsistent if $e_\alpha > e_\beta$, where $\beta = \alpha^-$, and there are α -states $\nu_0 <_B \nu_1$ such that $\nu_0 \in \mathcal{M}_\alpha$ and $\nu_1 \upharpoonright \beta \in \mathcal{M}_\beta$ but $\nu_1 \notin \mathcal{M}_\alpha$. Otherwise α is \mathcal{M} -consistent.

We shall take action in Step 3 of the construction in §7 to ensure that α is \mathcal{M} -consistent if $\alpha < f$.

5.4. Well-resided α -states and the lists \mathcal{R}_α and \mathcal{B}_α . Define the set of well-resided α -states,

$$(38) \quad \mathcal{K}_\alpha = \{\nu_1 : (\exists^\infty x)[x \in Y_\alpha \ \& \ \nu(\alpha, x) = \nu_1]\}.$$

Likewise define $\widehat{\mathcal{K}}_\alpha$ for the $\widehat{\omega}$ -side. To satisfy the automorphism requirement (25) we must show for $\alpha < f$ that

$$(39) \quad \widehat{\mathcal{K}}_\alpha = \{\widehat{\nu} : \nu \in \mathcal{K}_\alpha\}.$$

To achieve (39) note that unlike \mathcal{E}_α and \mathcal{F}_α of §5.3, \mathcal{K}_α is Π_3^0 not Π_2^0 , so α cannot guess at \mathcal{K}_α directly. Instead we guess at disjoint sets \mathcal{R}_α and \mathcal{B}_α such that for $\alpha < f$, $\mathcal{M}_\alpha - (\mathcal{R}_\alpha \cup \mathcal{B}_\alpha) = \mathcal{K}_\alpha$. These sets, \mathcal{R}_α and \mathcal{B}_α , correspond to those $\nu \in \mathcal{M}_\alpha$ that α believes are being emptied by Red and Blue, respectively.

To define \mathcal{R}_α and \mathcal{B}_α fix $\alpha \in T$, let $\beta = \alpha^-$, and assume that $\mathcal{R}_\gamma, \mathcal{B}_\gamma$ and their duals $\widehat{\mathcal{R}}_\gamma, \widehat{\mathcal{B}}_\gamma$ have been defined for all $\gamma < \alpha$. We decompose \mathcal{R}_α into the disjoint union,

$$(40) \quad \mathcal{R}_\alpha = \mathcal{R}_\alpha^\alpha \sqcup \mathcal{R}_\alpha^{<\alpha}, \text{ where}$$

$$(41) \quad \mathcal{R}_\alpha^{<\alpha} := \{\nu : \nu \in \mathcal{M}_\alpha \ \& \ \nu \upharpoonright \beta \in \mathcal{R}_\beta\}, \text{ and}$$

$$(42) \quad \mathcal{R}_\alpha^\alpha := \mathcal{R}_\alpha - \mathcal{R}_\alpha^{<\alpha}.$$

Note that $\mathcal{R}_\alpha^{<\alpha}$ is determined by $\mathcal{R}_\beta, \beta < \alpha$, but $\mathcal{R}_\alpha^\alpha$ may contain new elements and for $\alpha < f$ it has the meaning described below in (44). Likewise, let $\mathcal{B}_\alpha = \mathcal{B}_\alpha^\alpha \sqcup \mathcal{B}_\alpha^{<\alpha}$, where $\mathcal{B}_\alpha^{<\alpha}$ is defined as in (41) but with \mathcal{B}_β in place of \mathcal{R}_β .

If $|\alpha| \not\equiv 3 \pmod 5$ define $\mathcal{R}_\alpha^\alpha = \widehat{\mathcal{B}}_\alpha^\alpha = \emptyset$. If $|\alpha| \equiv 3 \pmod 5$ we let $\mathcal{M}_\alpha = \mathcal{M}_\beta$ (since α -states are β -states because $e_\alpha = e_\beta$ and $\widehat{e}_\alpha = \widehat{e}_\beta$). We define the Π_2^0 predicate,

$$(43) \quad F(\beta, \nu) \equiv (\forall x)[[x > |\beta| \ \& \ x \in Y_\beta] \implies \nu(\alpha, x) \neq \nu],$$

and we allow $\mathcal{R}_\alpha^\alpha \neq \emptyset$ with the intention that for $\alpha < f$,

$$(44) \quad \mathcal{R}_\alpha^\alpha = \{\nu : \nu \in \mathcal{M}_\alpha - (\mathcal{R}_\alpha^{<\alpha} \cup \mathcal{B}_\alpha^{<\alpha}) \ \& \ F(\beta, \nu)\}.$$

We define

$$(45) \quad \widehat{\mathcal{B}}_\alpha^\alpha := \{\widehat{\nu} : \nu \in \mathcal{R}_\alpha^\alpha\}.$$

If $|\alpha| \not\equiv 4 \pmod 5$ define $\widehat{\mathcal{R}}_\alpha^\alpha = \mathcal{B}_\alpha^\alpha = \emptyset$. If $|\alpha| \equiv 4 \pmod 5$ we allow $\widehat{\mathcal{R}}_\alpha^\alpha \neq \emptyset$ (using the duals of (40)–(44) where e.g. in the dual of (43) we use \widehat{Y}_β in place of Y_β), and we define

$$(46) \quad \mathcal{B}_\alpha^\alpha := \{\nu : \widehat{\nu} \in \widehat{\mathcal{R}}_\alpha^\alpha\}.$$

At most one of $\mathcal{R}_\alpha^\alpha$ and $\widehat{\mathcal{R}}_\alpha^\alpha$ is nonempty, so by (46), (45), and (44),

$$(47) \quad \mathcal{R}_\alpha^\alpha \cap \mathcal{B}_\alpha^\alpha = \emptyset \ \& \ ((\mathcal{R}_\alpha^\alpha \cup \mathcal{B}_\alpha^\alpha) \cap (\mathcal{R}_\alpha^{<\alpha} \cup \mathcal{B}_\alpha^{<\alpha}) = \emptyset),$$

and hence

$$(48) \quad \mathcal{R}_\alpha \cap \mathcal{B}_\alpha = \emptyset.$$

If $\alpha \prec f$, then $\nu \in \mathcal{R}_\alpha$ implies $F(\alpha^-, \nu)$ and hence

$$(49) \quad (\forall \nu \in \mathcal{R}_\alpha)(\forall x \in Y_\alpha)(\forall s) [\nu(\alpha, x, s) = \nu \implies (\exists t > s)[\nu(\alpha, x, t) \neq \nu]].$$

It will be Blue's responsibility to change the α -state of x if $\nu(\alpha, x, s) \in \mathcal{B}_\alpha$ and $x \in R_\alpha$. However, $\mathcal{B}_\alpha \cap \mathcal{R}_\alpha = \emptyset$ so if $\nu(\alpha, x, s) = \nu \in \mathcal{R}_\alpha$, then Blue can wait for Red to change the α -state of x to meet (49). That is,

$$(50) \quad (\forall \nu \in \mathcal{R}_\alpha)(\forall x \in R_\alpha)(\forall s) \left[\begin{array}{l} \text{if } \nu(\alpha, x, s) = \nu, \text{ then it is an} \\ \alpha\text{-admissible move for Blue to restrain} \\ x \text{ from further Blue enumeration until} \\ (\exists t > s)[\nu(\alpha, x, s) <_R \nu(\alpha, x, t)] \end{array} \right].$$

Definition 5.8. A node $\alpha \in T$ is \mathcal{R} -consistent if

$$(51) \quad (\forall \nu_0 \in \mathcal{R}_\alpha)(\exists \nu_1)[\nu_0 <_R \nu_1 \ \& \ \nu_1 \in \mathcal{M}_\alpha],$$

and \mathcal{R} -inconsistent otherwise.

By applying (50), Blue will ensure that α is \mathcal{R} -consistent for $\alpha \prec f$. Now (51), (45), and (37) imply for $\alpha \prec f$ that

$$(52) \quad (\forall \hat{\nu}_0 \in \widehat{\mathcal{B}}_\alpha)(\exists \hat{\nu}_1)[\hat{\nu}_0 <_B \hat{\nu}_1 \ \& \ \hat{\nu}_1 \in \widehat{\mathcal{M}}_\alpha].$$

By repeatedly applying (52), Blue can achieve $\hat{\nu}_1 \in \widehat{\mathcal{M}}_\alpha - \widehat{\mathcal{B}}_\alpha$. That is,

$$(53) \quad (\exists \text{ function } \hat{h}_\alpha) [\hat{h}_\alpha : \widehat{\mathcal{B}}_\alpha \rightarrow (\widehat{\mathcal{M}}_\alpha - \widehat{\mathcal{B}}_\alpha) \ \& \ (\forall \hat{\nu} \in \widehat{\mathcal{B}}_\alpha)[\hat{\nu} <_B \hat{h}_\alpha(\hat{\nu})]].$$

It is Blue's responsibility to move any $\hat{x} \in \widehat{R}_\alpha$ for which $\nu(\alpha, \hat{x}, s) = \hat{\nu}_0 \in \widehat{\mathcal{B}}_\alpha$ to the target state $\hat{\nu}_1 = \hat{h}_\alpha(\hat{\nu}_0)$ (and where \hat{h} is called the target function) so that Blue can achieve

$$(54) \quad (\forall \hat{x} \in \widehat{R}_\alpha)(\forall s) [\nu(\alpha, \hat{x}, s) \in \widehat{\mathcal{B}}_\alpha \implies (\exists t > s)[\nu(\alpha, \hat{x}, t) \in \widehat{\mathcal{M}}_\alpha - \widehat{\mathcal{B}}_\alpha]],$$

and hence Blue will cause every state $\hat{\nu}_0 \in \widehat{\mathcal{B}}_\alpha$ to be emptied. To achieve (54) on \widehat{R}_α it suffices to achieve the following on \widehat{S}_γ for each $\gamma \succeq \alpha$,

$$(55) \quad (\forall \hat{x} \in \widehat{S}_\gamma)(\forall s) [\nu(\gamma, \hat{x}, s) \in \widehat{\mathcal{B}}_\gamma \implies (\exists t > s)[\nu(\gamma, \hat{x}, t) \in \widehat{\mathcal{M}}_\gamma - \widehat{\mathcal{B}}_\gamma]].$$

We often refer to the dual of (53), which asserts

$$(56) \quad (\exists \text{ function } h_\alpha) [h_\alpha : \mathcal{B}_\alpha \rightarrow (\mathcal{M}_\alpha - \mathcal{B}_\alpha) \ \& \ (\forall \nu \in \mathcal{B}_\alpha)[\nu <_B h_\alpha(\nu)]],$$

and which enables us to achieve the dual of (55),

$$(57) \quad (\forall x \in S_\gamma)(\forall s) [\nu(\gamma, x, s) \in \mathcal{B}_\gamma \implies (\exists t > s)[\nu(\gamma, x, t) \in \mathcal{M}_\gamma - \mathcal{B}_\gamma]].$$

Finally, we have ensured, for all $\gamma \prec f$ and all $\nu_0 \in \mathcal{M}_\gamma$,

$$(58) \quad (\exists <^\infty x)[x \in Y_\gamma \ \& \ \nu(\gamma, x) = \nu_0] \implies (\exists \alpha)_{\gamma \prec \alpha \prec f} [\{\nu_1 \in \mathcal{M}_\alpha : \nu_1 \upharpoonright \gamma = \nu_0\} \subseteq \mathcal{R}_\alpha \cup \mathcal{B}_\alpha].$$

To check (58) fix $\gamma \prec f$ and $\nu_0 \in \mathcal{M}_\gamma$. By (27) $Y_\gamma =^* \omega$, so if the hypothesis of (58) holds, then we can choose b such that

$$(59) \quad (\forall x \in \omega)[x > b \implies \nu(\gamma, x) \neq \nu_0].$$

Choose $\alpha \prec f$ such that $\alpha \succ \gamma$, $|\alpha| > b$ and $|\alpha| \equiv 3 \pmod 5$. Consider any $\nu_1 \in \mathcal{M}_\alpha$ such that $\nu_1 \upharpoonright \gamma = \nu_0$. If $\nu_1 \notin \mathcal{R}_\alpha^{<\alpha} \cup \mathcal{B}_\alpha^{<\alpha}$, then $F(\alpha^-, \nu_1)$ holds, so $\nu_1 \in \mathcal{R}_\alpha^\alpha$ by (44), and hence $\nu_1 \in \mathcal{R}_\alpha$ by (40).

Equations (45), (49), (54), (58) and their duals guarantee (39), as long as the well-visited α -states are equal as well.

5.5. Verifying the automorphism requirement. We shall arrange that $\lim_{\alpha \prec f} e_\alpha = \infty$. By (28) and (29) the sets $\{U_\alpha\}_{\alpha \prec f}$ and $\{V_\alpha\}_{\alpha \prec f}$ constitute skeletons for $\{W_n\}_{n \in \omega}$. By (35), its dual, and (36) we know that the well-visited α -states on the ω -side and $\widehat{\omega}$ -side coincide. By (39) the well-resided α -states also coincide, so (25) is satisfied. The construction in §7 and verification in §8 will demonstrate that the equations of §5.2, §5.3, and §5.4 are satisfied. First we need a few more definitions.

6. THE TREE AND THE TRUE PATH

In this section, we define the tree of strategies and the true path f through the tree.

6.1. The well-visited states and \mathcal{F}_β^+ . In §4.4 we said that every $\alpha \in T$ would have an associated set \mathcal{M}_α such that $\mathcal{M}_\alpha = \mathcal{F}_\alpha$ if $\alpha \prec f$. However, although this is the *property* we want \mathcal{M}_α to have, we cannot simply *define* \mathcal{M}_α to be α 's guess at \mathcal{F}_α because that definition would be circular. Rather we must define here a certain set \mathcal{F}_β^+ which depends only on β , and then let \mathcal{M}_α be α 's guess at \mathcal{F}_β^+ so that $\mathcal{M}_\alpha = \mathcal{F}_\beta^+ (= \mathcal{F}_\alpha)$ for $\alpha \prec f$.

Fix $\alpha \in T$ such that $e_\alpha > e_\beta$ for $\beta = \alpha^-$. Define the c.e. set $Z_{e_\alpha} = \bigcup_s Z_{e_\alpha, s}$, where

$$(60) \quad Z_{e_\alpha, s+1} := \{x : x \in U_{e_\alpha, s+1} \ \& \ x \in Y_{\beta, s}\}.$$

Define the α -state function $\nu^+(\alpha, x, s)$ exactly as for $\nu(\alpha, x, s)$ in Definition 5.4 but with $Z_{e_\alpha, s}$ in place of $U_{\alpha, s}$.

Define

$$(61) \quad \mathcal{F}_\beta^{\bar{A}^+} = \{\nu : (\exists^\infty x)(\exists s)[x \in Y_{\beta, s}^{\bar{A}}, \nu^+(\alpha, x, s) = \nu, 0 \notin \nu, \text{ and } \widehat{\Psi}_\alpha(x)[s] = 0]\}$$

and

$$(62) \quad \mathcal{F}_\beta^{A^+} = \{\nu : (\exists^\infty x)(\exists s)[x \in Y_{\beta, s}^A \ \& \ \nu^+(\alpha, x, s) = \nu]\}.$$

Let $\mathcal{F}_\beta^+ = \mathcal{F}_\beta^{\bar{A}^+} \cup \mathcal{F}_\beta^{A^+}$. Define

$$(63) \quad k_\beta^+ = \min\{y : (\forall x > y)(\forall s) \left[[x \in Y_{\beta, s} \ \& \ \nu^+(\alpha, x, s) = \nu_1] \implies \nu_1 \in \mathcal{F}_\beta^+ \right]\}.$$

If $e_\alpha > e_\beta$ we also define $\widehat{\mathcal{F}}_\beta^+ = \{\widehat{\nu} : \nu \in \mathcal{F}_\beta^+\}$. (Note that Z_{e_α} and hence \mathcal{F}_β^+ and k_β^+ depend only upon β not α and thus α can make guesses \mathcal{M}_α and k_α for \mathcal{F}_β^+ and k_β^+ .)

If $\widehat{e}_\alpha > \widehat{e}_\beta$ we first define

$$(64) \quad \widehat{\mathcal{F}}_\beta^+ = \{\nu : (\exists^\infty x)(\exists s)[x \in Y_{\beta, s} \ \& \ \nu^+(\alpha, x, s) = \nu]\}.$$

Define k_β^+ using the dual of (63). Now we can define $\mathcal{F}_\beta^+ = \{\nu : \widehat{\nu} \in \widehat{\mathcal{F}}_\beta^+\}$. (Note that there is no \widehat{k}_β^+ only k_β^+ .)

Every $\alpha \in T$ will have associated items \mathcal{M}_α and k_α such that $\mathcal{M}_\alpha = \mathcal{F}_\beta^+$ and $k_\alpha = k_\beta^+$ for $\alpha \prec f$. We allow x to enter Y_α only if $x > k_\alpha$. If $e_\alpha = e_\beta$ and $\widehat{e}_\alpha = \widehat{e}_\beta$ we define $\mathcal{F}_\beta^+ = \mathcal{F}_\beta$, $\widehat{\mathcal{F}}_\beta^+ = \widehat{\mathcal{F}}_\beta$, and $k_\beta^+ = k_\beta$. If

$$(65) \quad (\exists x)(\exists s)[x \in Y_{\alpha,s} \ \& \ \nu(\alpha, x, s) \notin \mathcal{M}_\alpha],$$

then we say that α is *provably incorrect* at all stages $t \geq s$ and we ensure that $\alpha \not\prec f$.

6.2. The definition of the tree T .

Definition 6.1. We say that $\alpha \in T$ is *consistent* if α is \mathcal{M} -consistent (Definition 5.7) and \mathcal{R} -consistent (Definition 5.8).

Note that by clause (i) in the following Definition 6.2 of T ,

$$(66) \quad \beta \in T \implies [\beta \text{ inconsistent} \iff \beta \text{ is a terminal node on } T].$$

We shall show that if $\alpha \prec f$, then α is consistent, and therefore $\lim_{\alpha \prec f} e_\alpha = \infty$, so the argument of §5.5 applies.

Definition 6.2. Put $\lambda \in T$ and define $\mathcal{M}_\lambda = \mathcal{R}_\lambda = \mathcal{B}_\lambda = \langle \lambda, \emptyset, \emptyset \rangle$, and $k_\lambda = e_\lambda = \widehat{e}_\lambda = -1$. If $\beta \in T$ we put $\alpha = \beta \widehat{\langle \mathcal{M}_\alpha, \mathcal{R}_\alpha, \mathcal{B}_\alpha, k_\alpha \rangle}$ in T providing the following conditions hold:

- (i) β is consistent (as defined in Definition 6.1),
- (ii) \mathcal{M}_α is a set of α -states, $\mathcal{R}_\alpha \subseteq \mathcal{M}_\alpha$, $\mathcal{B}_\alpha \subseteq \mathcal{M}_\alpha$, and $\mathcal{R}_\alpha \cap \mathcal{B}_\alpha = \emptyset$,
- (iii) $\mathcal{M}_\alpha \upharpoonright \beta \subseteq \mathcal{M}_\beta$,
- (iv) $[e_\alpha = e_\beta \ \& \ \widehat{e}_\alpha = \widehat{e}_\beta] \implies \mathcal{M}_\alpha = \mathcal{M}_\beta \ \& \ k_\alpha = k_\beta$,
- (v) $\mathcal{R}_\alpha^{<\alpha} := \{\nu \in \mathcal{M}_\alpha : \nu \upharpoonright \beta \in \mathcal{R}_\beta\} \subseteq \mathcal{R}_\alpha$,
- (vi) $\mathcal{B}_\alpha^{<\alpha} := \{\nu \in \mathcal{M}_\alpha : \nu \upharpoonright \beta \in \mathcal{B}_\beta\} \subseteq \mathcal{B}_\alpha$,
- (vii) $\mathcal{R}_\alpha^\alpha := \mathcal{R}_\alpha - \mathcal{R}_\alpha^{<\alpha} \neq \emptyset \implies |\alpha| \equiv 3 \pmod 5$,
- (viii) $\mathcal{B}_\alpha^\alpha := \mathcal{B}_\alpha - \mathcal{B}_\alpha^{<\alpha} \neq \emptyset \implies |\alpha| \equiv 4 \pmod 5$,
- (ix) $|\alpha| \equiv 0 \pmod 5 \implies \langle \mathcal{M}_\alpha, \mathcal{R}_\alpha, \mathcal{B}_\alpha, k_\alpha \rangle = \langle \mathcal{M}_\beta, \mathcal{R}_\beta, \mathcal{B}_\beta, k_\beta \rangle$. These nodes correspond to the positive requirement, so they are also associated with either a 0 representing the Π_2 -outcome, or any $p \in \omega$, $p > 0$, representing the Σ_2 -outcomes. These latter have the ω^* -ordering, with 0 the leftmost node and 1 the rightmost.

In addition, each $\alpha \in T$ has associated dual sets $\widehat{\mathcal{M}}_\alpha$, $\widehat{\mathcal{R}}_\alpha$, and $\widehat{\mathcal{B}}_\alpha$, which are determined from \mathcal{M}_α , \mathcal{B}_α , and \mathcal{R}_α , by (36), (46), and (45), respectively. Also α has associated integers e_α and \widehat{e}_α (depending only on $|\alpha|$) as defined at the beginning of §5.3. (We identify the finite object $\langle \mathcal{M}_\alpha, \mathcal{R}_\alpha, \mathcal{B}_\alpha, k_\alpha \rangle$ with an integer under some effective coding, so we may regard $T \subseteq \omega^{<\omega}$.) For $\mathcal{M}_\alpha \supset \mathcal{M}_{\alpha'}$, where $|\alpha| = |\alpha'|$, we order the nodes so that $\alpha <_L \alpha'$ and similarly for \mathcal{R}_α and \mathcal{B}_α . We also ensure that if $k_\alpha < k_{\alpha'}$, then $\alpha <_L \alpha'$.

Definition 6.3. The *true path* $f \in [T]$ is defined by induction on n . We define $f \upharpoonright 1$ to be the node ρ of length 1 such that \mathcal{M}_ρ contains all possible ρ -states and $k_\rho = 0$. Note there are exactly two states in \mathcal{M}_ρ , the state containing $U_\rho =^* A$ and the state that does not contain U_ρ . For $\widehat{\mathcal{M}}_\alpha$, the two states correspond to the one containing $\widehat{U}_\rho = B$ and the one that does not. When we refer to B_s , we are actually referring to $\widehat{U}_{\rho,s}$. The approximation of the true path f_s will go through ρ for all s .

Let $\beta = f \upharpoonright n$ be consistent. Then $f \upharpoonright (n + 1)$ is the $<_L$ -least $\alpha \in T$, $\alpha \succ \beta$, of length $m = n + 1$ such that:

- (i) For $m = 5e$, $f(n) = 0$ if β appoints infinitely many witnesses, and $f(n) = p + 1$ if β enumerates p witnesses into D ,
- (ii) $m \equiv 1 \pmod 5 \implies \mathcal{M}_\alpha = \mathcal{F}_\beta^+ \quad \& \quad k_\alpha = k_\beta^+$,
- (iii) $m \equiv 2 \pmod 5 \implies \widehat{\mathcal{M}}_\alpha = \widehat{\mathcal{F}}_\beta^+ \quad \& \quad k_\alpha = k_\beta^+$,
- (iv)

$$m \equiv 3 \pmod 5 \implies \begin{aligned} \mathcal{R}_\alpha &= \{ \nu : \nu \in \mathcal{M}_\alpha - (\mathcal{R}_\alpha^{<\alpha} \cap \mathcal{B}_\alpha^{<\alpha}) \quad \& \quad F(\beta, \nu) \} \\ &\quad \& \quad \widehat{\mathcal{B}}_\alpha^\alpha = \{ \hat{\nu} : \nu \in \mathcal{R}_\alpha^\alpha \}, \end{aligned}$$

(v)

$$m \equiv 4 \pmod 5 \implies \begin{aligned} \widehat{\mathcal{R}}_\alpha^\alpha &= \{ \hat{\nu} : \hat{\nu} \in \widehat{\mathcal{M}}_\alpha - (\widehat{\mathcal{R}}_\alpha^{<\alpha} \cup \widehat{\mathcal{B}}_\alpha^{<\alpha}) \quad \& \quad \widehat{F}(\beta, \nu) \} \\ &\quad \& \quad \mathcal{B}_\alpha^\alpha = \{ \nu : \hat{\nu} \in \widehat{\mathcal{R}}_\alpha^\alpha \}, \end{aligned}$$

- (vi) unless otherwise specified in (i)–(iv), \mathcal{M}_α , \mathcal{R}_α , \mathcal{B}_α , and k_α take the values \mathcal{M}_β , \mathcal{R}_β , \mathcal{B}_β , and k_β , respectively.

For a consistent $\beta = f \upharpoonright n$, note that \mathcal{F}_β^+ is just a finite set of states and k_β^+ is an integer, so clearly α exists. We shall prove that if $\alpha \prec f$, then α is consistent, so the true path f exists and is infinite. Note that each of the conditions in Definition 6.3 is Π_2^0 . A computable approximation to the true path, the limit inferior of which will be the true path itself, is given along with the D -Module in the next section.

7. THE CONSTRUCTION

Let T be the tree defined in §6.2. We simultaneously construct D and the sets $U_\alpha, \widehat{U}_\alpha, V_\alpha, \widehat{V}_\alpha$. At each stage we also define a function $f_s \in T$, $|f_s| = 2s$, such that $f = \liminf_s f_s$. We say that stage s is an α -stage if $\alpha \prec f_s$.

7.1. Order of the construction. We can think of our construction happening on four identical trees, corresponding to \overline{A} , \overline{B} , A , and B . Every $x \in \omega$ begins on the \overline{A} tree and moves through the nodes. If $x \in A$, it will eventually move from the \overline{A} tree to the A tree. This happens for each $\hat{x} \in \hat{\omega}$ for the B side as well.

In the following construction, Steps 1-5 are for the \overline{A} tree. For the A tree, the Step 1-5 construction is exactly the construction in the Harrington-Soare paper, except without distinguishing between S^0 and S^1 . Similarly, Steps $\widehat{1} - \widehat{5}$, which apply to the \overline{B} and B trees are also as in the Harrington-Soare construction in [4], except with the addition of switching elements from the \overline{B} tree to the B tree. These steps are identical (or dual) to the Steps 1-5 mentioned here, except that any mention of keeping elements out of A , such as in (1.8), is omitted.

At the beginning of each stage s , we perform one of Steps 1-5 of the A or \overline{A} tree construction if any applies. If no step can act, we perform one of Steps $\widehat{1} - \widehat{5}$ for the B and \overline{B} trees. Regardless of whether we performed any of these steps, we continue to Step B and the D -Module. If none of the Steps 1-5 or $\widehat{1} - \widehat{5}$ has acted on any of the four trees, we proceed to Step 6 *after* performing Step B and the D -Module. Step 6 applies to all four trees.

7.2. The automorphism construction. We perform the automorphism construction as in the Δ_3^0 paper, with some important changes. We switch elements x from the \bar{A} tree to the A tree under Step 4 when they enter $A = W_i$. Let $U_0 = A = W_i$.

To *initialize* node α means: to remove every $x \in S_{\alpha,s}$ ($\hat{x} \in \hat{S}_{\alpha,s}$), and put x in S_β (\hat{x} in \hat{S}_β) for $\beta = \alpha \cap f_{s+1}$ (where $\alpha \cap \delta$ denotes the longest γ such that $\gamma \preceq \alpha$ and $\gamma \preceq \delta$).

We present in the next section Steps 1–5 for the construction and a final Step 6 at which we define f_{s+1} . These properties will produce the automorphism.

Besides adding the D and B constructions, we must also make some changes to the old automorphism construction. The primary difference between this construction and the Harrington-Soare construction of [4] is that we now must only allow elements to move in the \bar{A} tree if they appear to be in \bar{A} . Each of Steps 1-5 now requires that the element x must satisfy the condition that $\hat{\Psi}^D(x)[s] = 0$. This condition says that D believes x is in \bar{A} . This condition is actually only important for Step 1, but we must impose it on all steps because otherwise, another step, such as Step 3, could empty out a state before Step 1 has the ability to pull elements from that state down to the next node. By requiring that all steps wait for elements to appear to be in \bar{A} , Step 1 will have priority over the other steps.

Note that in order to ensure that Step 3 can act, we allow it to go immediately after Step 1, so that elements pulled into S_α by Step 1 will still appear to be in \bar{A} when Step 3 acts. Otherwise, Step 3 may never have the ability to act. We only allow Step 3 to act in this way for half of the elements pulled into a state by Step 1, ensuring that the state is not prematurely emptied.

7.3. The steps of the construction. Stage $s = 0$. For all $\alpha \in T$ define $U_{\alpha,0} = V_{\alpha,0} = \hat{U}_{\alpha,0} = \hat{V}_{\alpha,0} = \emptyset$, and define $m(\alpha, 0) = 0$. Define $Y_{\lambda,0} = \hat{Y}_{\lambda,0} = \emptyset$, and $f_0 = \lambda$.

Let $B_0 = \emptyset$, and define $\delta_0 = \lambda$, the empty node on T .

Stage $s + 1$. Find the least $n < 6$ such that Step n applies to some $x \in Y_{\alpha,s}$, and perform the indicated action. This can apply to either the A or \bar{A} trees. If there is no such n , then likewise find the least $n < 6$ such that Step \hat{n} applies to some $\hat{x} \in \hat{Y}_{\alpha,s}$, and perform the indicated action. This can apply to either the B or \bar{B} trees.

Next, check if Step B applies, and perform the indicated action if it does.

Then perform the D -Module. Note that both Step B and the D -Module should be performed after Steps 1–5 and Steps $\hat{1}$ – $\hat{5}$, regardless of whether these steps actually acted.

If none of Steps 1–5 or Steps $\hat{1}$ – $\hat{5}$ acted, then perform Step 6 after Step B and the D -Module. (It is important that these steps be performed in the indicated order.)

7.4. Steps 1–5. Step 1. (Prompt pulling of x from R_β to S_α to ensure $\mathcal{M}_\alpha \subseteq \mathcal{E}_\alpha$) Suppose $\langle \alpha, \nu_i \rangle$ is the first unmarked entry on the list \mathcal{L}_s defined in Step 6 such that the following conditions hold for some x , where $\nu_i = \langle \alpha, \sigma_i, \tau_i \rangle$, and $A \notin \sigma_i$ unless $e_\alpha = 0$.

(1.1) $x \in R_{\beta,s} - Y_{\alpha,s}$,

(1.2) $x > k_\alpha$, $x > |\alpha|$, and $x > p$ for $\alpha = \beta \hat{\ } p$ and $|\alpha| = 5e$,

- (1.3) x is α -eligible (i.e., $\neg(\exists t)[x \leq t \leq s \ \& \ f_t < \alpha]$),
- (1.4) $\neg[\alpha(x, s) <_L \alpha]$,
- (1.5) $x > m(\alpha, s)$,
- (1.6) $\nu(\beta, x, s) = \nu_i \upharpoonright \beta$,
- (1.7) $e_\alpha > e_\beta \implies \nu^+(\alpha, x, s) = \nu_i$,
- (1.8) if $A \notin \sigma_i$, $\widehat{\Psi}_\alpha(x)[s] \downarrow = 0$.

Action. Choose the least x corresponding to $\langle \alpha, \nu_i \rangle$, and do the following.

- (1.9) Mark the α -entry $\langle \alpha, \nu_i \rangle$ on \mathcal{L}_s , and suppose this is the k^{th} occurrence of $\langle \alpha, \nu_i \rangle$ on \mathcal{L}_s .
- (1.10) Move x to S_α .
- (1.11) If $e_\alpha > e_\beta$ and $e_\alpha \in \sigma_i$, then enumerate x in $U_{\alpha, s+1}$.
- (1.12) If $\widehat{e}_\alpha > \widehat{e}_\beta$ and $\widehat{e}_\alpha \in \tau_i$, then enumerate x in $\widehat{V}_{\alpha, s+1}$. (Hence, $\nu(\alpha, x, s+1) = \nu_i$. Also $\nu_i \in \mathcal{M}_\alpha$ because $\langle \alpha, \nu_i \rangle \in \mathcal{L}$ implies $\nu_i \in \mathcal{M}_\alpha$.)
- (1.13) If k is odd and $A \notin \sigma_i$, immediately perform Step 3, if it applies.
- (1.14) If $A \in \sigma_i$, switch x to the A tree, and enumerate $\langle \alpha, x, i \rangle$ in Λ .

Step 2. (Move x from S_β to S_α , so $Y_\alpha =^* \omega$) Suppose there is an x such that

- (2.1) $x \in S_{\beta, s}$,
- (2.2) $x > |\alpha|$ and $x > k_\alpha$, and $x > p$ for $\alpha = \beta \widehat{\ } p$ and $|\alpha| = 5e$,
- (2.3) x is α -eligible,
- (2.4) $x < m(\alpha, s)$,
- (2.5) $\widehat{\Psi}_\alpha(x)[s] \downarrow = 0$,
- (2.6) α is the $<_L$ -least $\gamma \in T$ with $\gamma^- = \beta$ satisfying (2.1)–(2.5).

Action. Choose the least pair $\langle \alpha, x \rangle$ and

- (2.7) move x from S_β to S_α .

(In Step 2 we need (2.4), so Y_α will not grow while α is waiting for another prompt pulling under Step 1.)

Step 3. (For α \mathcal{M} -inconsistent to ensure $\alpha \not\prec f$) Suppose for $\alpha \in T$ there exists x such that,

- (3.1) $e_\alpha > e_\beta$,
- (3.2) $x \in S_{\alpha, s}$,
- (3.3) $\nu(\alpha, x, s) = \nu_0 \in \mathcal{M}_\alpha$,
- (3.4) $(\exists \nu_1)[\nu_0 <_B \nu_1 \ \& \ \nu_1 \upharpoonright \beta \in \mathcal{M}_\beta \ \& \ \nu_1 \notin \mathcal{M}_\alpha]$,
- (3.5) $\widehat{\Psi}_\alpha(x)[s] \downarrow = 0$.

Action. Choose the least such pair $\langle \alpha, x \rangle$ and

- (3.6) enumerate x in $\widehat{V}_{\delta, s+1}$ for all $\delta \prec \alpha$ such that $e_\delta \in \tau_1$. (This action causes $\nu(\alpha, x, s+1) = \nu_1$. Hence, α is provably incorrect at all stages $t \geq s+1$ so $\alpha \not\prec f$.)

Step 4. (Delayed Red enumeration into U_α) Suppose $x \in R_{\alpha, s}$, or $x \in S_{\lambda, s}$ and let $\alpha = \rho$, and

- (4.1) $e_\alpha > e_\beta$,
- (4.2) $x \notin U_{\alpha, s}$,
- (4.3) $x \in Z_{e_\alpha, s} := U_{e_\alpha, s} \cap Y_{\beta, s-1}$, and
- (4.3) if $\alpha \neq \rho$, then $\widehat{\Psi}_\alpha(x)[s] \downarrow = 0$.

Action. Choose the least such pair $\langle \alpha, x \rangle$ and

(4.5) enumerate x in $U_{\alpha, s+1}$.

(4.6) If $e_\alpha = 0$, switch x to the A tree, and enumerate $\langle \gamma, x, i \rangle$ in the list Λ for $x \in S_\gamma, s$ for $\nu(\gamma, x, s) = \nu_i$.

Step 5. (Blue emptying of state $\nu \in \mathcal{B}_\alpha$) Suppose for $\alpha \in T$ there exists x such that either Case 1 or Case 2 holds.

Case 1. Suppose

(5.1) $\nu(\alpha, x, s) = \nu_0 \in \mathcal{B}_\alpha$, say $\nu_0 = \langle \alpha, \sigma_0, \tau_0 \rangle$,

(5.2) $x \in S_{\alpha, s}$,

(5.3) α is \mathcal{M} -consistent and \mathcal{R} -consistent,

(5.4) $\widehat{\Psi}_\alpha(x)[s] \downarrow = 0$.

Action. Choose the least such pair $\langle \alpha, x \rangle$. Let $\nu_1 = h_\alpha(\nu_0) >_B \nu_0$, where h_α is a target function satisfying (56). Let $\nu_1 = \langle \alpha, \sigma_1, \tau_1 \rangle$.

(5.5) Enumerate $x \in \widehat{V}_\delta$ for all $\delta \preceq \alpha$ such that $\widehat{e}_\delta > \widehat{e}_{\delta^-}$ and also $e_\delta \in \tau_1 - \tau_0$. (Hence, $\nu(\alpha, x, s + 1) = \nu_1$.)

Case 2. Suppose that (5.1) holds and

(5.6) $x \in S_{\gamma, s}$ where $\gamma^- = \alpha$, and

(5.6) γ is either \mathcal{M} -inconsistent or \mathcal{R} -inconsistent,

(5.7) $\widehat{\Psi}_\alpha(x)[s] \downarrow = 0$.

Action. Perform the same action as in Case 1 to achieve $\nu(\alpha, x, s + 1) = \nu_1$.

(In (5.6) note that by (66) $\gamma \in T$ implies (5.3) for $\alpha = \gamma^-$, so h_α exists in Case 2. Note in Step 5, Case 2 that the enumeration may not be γ -legal, i.e., perhaps $\nu(\gamma, x, s + 1) \notin \mathcal{M}_\gamma$, but this will not matter because we shall prove that $\gamma \not\prec f$ if γ is inconsistent. Hence, it only matters that the enumeration is α -legal, i.e., $\nu(\alpha, x, s) \in \mathcal{M}_\alpha$.)

7.5. Step B and the D -Module. Step B. Let $\langle \alpha, x, e \rangle$ be the least entry on the list Λ such that there is a y that has never before caused action on this step, satisfying

(B.1) $y \in \widehat{S}_\alpha^A$,

(B.2) $\widehat{\nu}(\alpha, y, s) = \widehat{\nu}_e$,

(B.3) for all $i < \langle \alpha, x, e \rangle$, $\varphi_i^B(i)[s] \downarrow \implies \varphi_i^B(i)[s] < y$.

Action. (B.4) If $\langle \alpha, x, e \rangle$ is not checked, check $\langle \alpha, x, e \rangle$, and do not enumerate y into B . If $\langle \alpha, x, e \rangle$ has been checked already, enumerate y into B , switching y to the B tree, and remove $\langle \alpha, x, e \rangle$ from the list Λ . This will leave infinitely many elements in \overline{B} , while still matching the flow into A .

The D -Module (and approximation to the true path). Define f_{s+1} by a series of substages t , $0 \leq t \leq 2s$, where we define δ^t at substage t , as follows, and later define $f_{s+1} = \delta^{2s}$. (Note that in the full theorem as described in §4.7, we must perform this for multiple different trees T_α at the same time to get the approximation to the true path.)

Substage t , $0 \leq t \leq 2s$. Define $\delta^0 = \lambda$ and $\delta^1 = \rho$ as in Definition 6.3. Given δ^t , perform the following action.

Case 1. $|\delta^t| \equiv 0$ (or 1) mod 5. Let $\beta = \delta^t$. Given δ^t let $v \leq s$ be maximal such that $\delta^t \preceq f_v$ if v exists and let $v = 0$ otherwise. For $|\delta^t| \equiv 0$ mod 5, let α be the leftmost node (i.e. the largest possible \mathcal{M}_α) with $\alpha^- = \beta$ such that for all $\nu \in \mathcal{M}_\alpha^{\bar{A}}$,

$$(67) \quad \begin{aligned} & |\{x : (\exists t \leq s)[x \in Y_{\beta,t}^{\bar{A}}, \nu^+(\alpha, x, t) = \nu, 0 \notin \nu, \ \& \ \widehat{\Psi}_\alpha(x)[t] = 0]\}| \\ & > |\{x : (\exists t \leq v)[x \in Y_{\beta,t}^{\bar{A}}, \nu^+(\alpha, x, t) = \nu, 0 \notin \nu, \ \& \ \widehat{\Psi}_\alpha(x)[t] = 0]\}|, \end{aligned}$$

for all $\nu \in \mathcal{M}_\alpha^A$,

$$(68) \quad \begin{aligned} & |\{x : (\exists t \leq s)[x \in Y_{\beta,t}^{\bar{A}} \& \nu^+(\alpha, x, t) = \nu]\}| \\ & > |\{x : (\exists t \leq v)[x \in Y_{\beta,t}^{\bar{A}} \& \nu^+(\alpha, x, t) = \nu]\}|, \end{aligned}$$

and

$$(69) \quad \begin{aligned} k_\alpha = & \quad (\mu u < s)(\forall s', u < s' < s) \\ & \quad [[x \in Y_{\beta,s'} \ \& \ \nu^+(\alpha, x, s') = \nu_1] \implies \nu_1 \in \mathcal{M}_\alpha]. \end{aligned}$$

Set $\delta^{t+1} = \alpha$. If no such α exists, let δ^{t+1} be the rightmost extension α of δ^t such that $k_\alpha = s$. The dual is similar, for $|\delta^t| \equiv 1$ mod 5, except that it uses only equations (68) and (69).

Case 2. $|\delta^t| \equiv 2$ (or 3) mod 5. Let $\beta = \delta^t$. Given δ^t let $v \leq s$ be maximal such that $\delta^t \preceq f_v$ if v exists and let $v = 0$ otherwise. For each α with $\alpha^- = \beta$, let $g(\alpha, s)$ be the greatest $h < s$ such that for all $x \leq h$, $\nu(\beta, x, s) \notin \mathcal{R}_\alpha^\alpha$. Let δ^{t+1} be the leftmost α such that $g(\alpha, s) > g(\alpha, v)$. If no such α exists, let δ^{t+1} be the rightmost extension of β of length $|\delta^t| + 1$. The dual for $|\delta^t| \equiv 3$ mod 5 is similar.

Case 3. $|\delta^t| \equiv 4$ mod 5. Let $\beta = \delta^t$. Here β is associated with the strategy for P_j described in §3.2. These correspond to what we called odd nodes in the construction of a nonlow low₂ set. (In particular, we assume a set Z_β is being defined whose index $g(\beta)$ is known a priori, such that $W_{g(\beta)} = Z_\beta$, g is the computable function as in §3.2 Step 1a but adjusted with β in place of j , and that our objective is to arrange that either $\lim_s \tilde{\varphi}_{j,s}(g(\beta)) \uparrow$, or gives the wrong answer about whether $W_{g(\beta)} \cap \bar{D} \neq \emptyset$.)

Subcase 1. (Like §3.2 Step 1a) Suppose $Z_{\beta,s} \cap \bar{D}_s = \emptyset$, $\tilde{\varphi}_{j,s}(g(\beta)) \downarrow = 0$, and there is no β -witness in \bar{D}_s . Let the β -witnesses previously appointed be $x_1^\beta \dots x_{n-1}^\beta$. Suppose

$$(70) \quad \left[(\forall \alpha) \left[\alpha \preceq \beta \implies \begin{aligned} & (\forall e, k \text{ where } \nu_e \in \mathcal{M}_\alpha) \\ & \langle \alpha, e, k \rangle \leq \langle \beta, n \rangle \implies y_{\langle e, k \rangle}^{s+1} \text{ exists } \ \& \ \widehat{\psi}_{\alpha,s}(y_{\langle e, k \rangle}^{s+1}) \downarrow \end{aligned} \right] \right]$$

and for all ξ with $|\xi| \equiv 4$ mod 5,

$$(71) \quad \begin{aligned} \xi \widehat{\prec} p \prec \beta & \implies x_n^\xi \in D_s \text{ for all } n' < p, \\ \text{and } \xi \widehat{\prec} 0 \prec \beta \ \& \ \langle \xi, n' \rangle < \langle \beta, n \rangle & \implies x_n^\xi \in D_s. \end{aligned}$$

Define y to be the smallest $z \in \bar{D}_s \cap \omega^{[\beta]}$, where β is identified with its Gödel number, such that $\Gamma_s^\beta < z < \Lambda_s^\beta$, and

$$(72) \quad \begin{aligned} & (\forall \alpha \preceq \beta) \left[(\forall e, k \text{ with } \nu_e \in \mathcal{M}_\alpha \ \& \ \langle \alpha, e, k \rangle < \langle \beta, n \rangle) \left[\widehat{\psi}_{\alpha,s}(y_{\langle e, k \rangle}) \downarrow < z \right] \right] \\ & \text{and } (\forall \gamma <_L \beta) [z \text{ is greater than any current witness at } \gamma]. \end{aligned}$$

If y exists, then appoint y as the next β -witness, call it x_n^β , define $\Gamma_{s+1}^\beta = x_n^\beta$, and enumerate x_n^β in $Z_{\beta,s+1}$.

Subcase 2. (Like §3.2 Step 1b) Suppose that $\tilde{\varphi}_{j,s}(g(\beta)) \downarrow = 1$, Γ_s^β is defined and equal to some β -witness x_n^β , and that for all ξ with $|\xi| \equiv 4 \pmod 5$,

$$(73) \quad \xi \hat{<} 0 \prec \beta \implies \Gamma_s^\xi = x_p^\xi \in D_{s+1}.$$

Then enumerate x_n^β in D_{s+1} , and define

$$(74) \quad \Gamma_{s+1}^\beta = \max \{ \Gamma_s^\beta, s + 1 \}.$$

(Equation (73) says that for every ξ with $|\xi| \equiv \pmod 5$ and $\xi \hat{<} 0 \prec \beta$, i.e., for which β is guessing that ξ has the Π_2 -outcome of infinitely many witnesses, β must wait until ξ actually puts its witness $\Gamma_s^\xi = x_p^\xi$ into D_{s+1} during an earlier substage of stage $s + 1$ before β is allowed to simultaneously contribute its own witness at a later substage of stage $s + 1$.)

Subcase 3. Otherwise. If $\Gamma_s^\beta = x_n^\beta$ for some β -witness x_n^β , then define $\Gamma_{s+1}^\beta = \Gamma_s^\beta$, and if not, then define $\Gamma_{s+1}^\beta = \max \{ \Gamma_s^\beta, s + 1 \}$.

To complete Case 3 define $\delta^{t+1} = \delta^t \hat{<} 0$ if Subcase 2 holds, and $\delta^{t+1} = \delta^t \hat{<} p + 1$, otherwise, where p is the number of β -witnesses that have been enumerated into D . Complete the construction by defining $f_{s+1} = \delta^{2s}$ and by P-initializing every node β , $|\beta| \equiv \pmod 5$, such that $f_{s+1} <_L \beta$ or there exists $\alpha <_L \beta$ and some x such that $\hat{\psi}_{\alpha,s+1}(x) \downarrow$ and $\hat{\psi}_{\alpha,s}(x) \uparrow$. P-initializing β means canceling any β -witness, and defining $\Gamma_{s+1}^\beta = \max \{ \Gamma_s^\beta, s + 1 \}$.

7.6. Step 6. (Defining $m(\alpha, s + 1)$, \mathcal{L}_{s+1} and $Y_{\lambda,s+1}$) This step applies to all four copies of the tree. It should be performed only if none of Steps 1-5 ($\hat{1}$ - $\hat{5}$) applied at this stage.

Substep 6A. (Defining $m(\alpha, s + 1)$, \mathcal{L}_{s+1} , and their duals) For every $\alpha \preceq f_{s+1}$ if every α -entry $\langle \alpha, \nu \rangle$ on \mathcal{L}_s and every α -entry $\langle \alpha, \hat{\nu} \rangle$ on $\hat{\mathcal{L}}_s$ is marked we say that the lists are α -marked and we

$$(6.1) \text{ define } m(\alpha, s + 1) = m(\alpha, s) + 1, \text{ and}$$

(6.2) add to the bottom of list \mathcal{L}_s ($\hat{\mathcal{L}}_s$) a new (unmarked) α -entry $\langle \alpha, \nu \rangle$ ($\langle \alpha, \hat{\nu} \rangle$) for every such $\alpha \neq \lambda$ and every $\nu \in \mathcal{M}_\alpha$. Let the resulting list be \mathcal{L}_{s+1} ($\hat{\mathcal{L}}_{s+1}$).

If the lists are not both α -marked, then let $m(\alpha, s + 1) = m(\alpha, s)$, $\mathcal{L}_{s+1} = \mathcal{L}_s$ and $\hat{\mathcal{L}}_{s+1} = \hat{\mathcal{L}}_s$.

Substep 6B. (Emptying R_α to the right of f_{s+1}) For every α such that $f_{s+1} <_L \alpha$, initialize α .

Substep 6C. (Filling Y_λ and \hat{Y}_λ) Choose the least $x \notin Y_{\lambda,s}$ ($\hat{x} \notin \hat{Y}_{\lambda,s}$) and $x < s$. Put x in S_λ^A (\hat{x} in \hat{S}_λ^B).

For each $x \in Y_{\lambda,s+1}$ ($\hat{x} \in \hat{Y}_{\lambda,s+1}$) let $\alpha(x, s + 1)$ ($\alpha(\hat{x}, s + 1)$) denote the unique γ such that $x \in S_{\gamma,s+1}$ ($\hat{x} \in \hat{S}_{\gamma,s+1}$).

(Note that after each application of Step 6, the other Steps 1-5 and Steps $\hat{1}$ - $\hat{5}$ can apply only finitely often until the next application of Step 6 as we prove in Lemma 8.7.)

8. THE VERIFICATION

We now verify that this construction establishes the theorem. Many of the lemmas in this section are similar to lemmas in [4]. The most significant and important new or altered lemmas are Lemma 8.8 and Lemma 8.9, which contain the key elements of combining the automorphism construction with the construction of a nonlow set. Lemma 8.12 also has important changes.

First define the *true path* f of the construction by induction on e . Let $f \upharpoonright 0 = \lambda$, the empty node. Given $\beta = f \upharpoonright 5e$ define $f(5e) = 0$ if β appoints infinitely many β -witnesses and $f(5e) = p + 1$ if p β -witnesses are ever enumerated into D . For $\alpha = f \upharpoonright n$, where $n \neq 5e$, define $f(n)$ as in the automorphism construction in Harrington-Soare [4].

Lemma 8.1. *At stage $s + 1$,*

(i) *if x enters R_α , $\alpha \neq \lambda$, then Step 1 or Step 2 applies to α and x . (This does not hold for R_α^A .)*

(ii) *if x moves from S_α to S_δ , then one of the following steps must apply to x : Step 1_δ for $\delta <_L \alpha$ or $\delta^- = \alpha$; Step 2_δ for δ such that $\delta^- = \alpha$; or Step 6_α Substep B applying to α , so $f_{s+1} <_L \alpha$;*

(iii) *if $x \in S_{\alpha,s}$ is enumerated in a red set U_α at stage $s + 1$, then Step 1 or Step 4 must apply to x ;*

(iv) *if $x \in S_{\alpha,s}$ is enumerated in a blue set \widehat{V}_α , then Step 1, Step 3, or Step 5 must apply to x .*

Proof. This follows directly from the construction. □

Lemma 8.2 (True Path Lemma). $f = \liminf_s f_s$.

Proof. Since f is Π_2^0 , we know that such a computable approximation is possible. The definition of f_s in the D -Module has been formulated precisely so that $f = \liminf_s f_s$, where f is as in Definition 6.3. □

Lemma 8.3. $\mathcal{M}_\rho = \mathcal{F}_\lambda^+$ and $k_\rho = k_\lambda^+$.

Proof. We know that A is infinite and co-infinite, so there are infinitely many elements x such that $\nu^+(\lambda, x, s)$ will contain U_0 and infinitely many that will not, and will appear to D not to contain U_0 . Thus, \mathcal{F}_λ^+ contains both possible states. Because of this, it is easy to see that k_λ^+ must be 0, as it is not possible for any element to be in a state that is not well-visited, since both states are well-visited. □

8.1. The lemmas of motion, Y_α , and $\alpha(x, s)$. We now verify the properties we stated in the three subsections §5.2, §5.3, and §5.4. For each lemma there are obvious dual lemmas with similar proofs unless we state and prove the dual explicitly.

Lemma 8.4. *For all $\alpha \in T$,*

(i) $f <_L \alpha \implies R_{\alpha,\infty} = \emptyset$,

(ii) $\alpha <_L f \implies Y_\alpha =^* \emptyset$,

(iii) $\alpha \prec f \implies Y_{<\alpha} := \bigcup \{Y_\delta : \delta <_L \alpha\} =^* \emptyset$.

Proof. (i) Given x choose $s > x$ such that $f_s <_L \alpha$. By Step 6B, $R_{\alpha,s} = \emptyset$. Now x is γ -ineligible for all $t \geq s$ and all $\gamma \succeq \alpha$, so $x \notin S_{\gamma,t}$ and hence $x \notin R_{\alpha,t}$ by (1.3) and (2.3).

(ii) Assume $\alpha <_L f$. Then $\alpha < f_s$ for finitely many s and there are only finitely many α -entries $\langle \alpha, \nu \rangle$ on the list \mathcal{L} under (6.2). Hence, finitely many x enter S_α under Step 1 because every such x must mark some unmarked α -entry on \mathcal{L} . Thus, $m(\alpha) := \lim_s m(\alpha, s) < \infty$ since \mathcal{L} will be α -marked at most finitely often. Hence, by (2.4), Step 2 moves only finitely many x into R_α . But each x enters R_α only under Step 1 or Step 2, so $Y_\alpha =^* \emptyset$.

(iii) Immediate by (ii) since $f_s <_L \alpha$ finitely often, so there are only finitely many $\beta <_L \alpha$ such that $Y_\beta \neq \emptyset$. □

Lemma 8.5. *For every $\alpha \in T$ if $\alpha \neq \lambda$ and $\beta = \alpha^-$, then*

- (i) $Y_\alpha \setminus Y_\beta = \emptyset$ and $Y_\alpha \subseteq Y_\beta$ (however, elements may enter Y_α^A at the same time as they enter Y_β^A),
- (ii) $(\forall x)(\exists^{\leq 1} s)[x \in R_{\alpha, s+1} - R_{\alpha, s}]$,
- (iii) $U_\alpha \setminus Y_\alpha = \widehat{V}_\alpha \setminus Y_\alpha = \emptyset$, except that elements may enter $A = U_\rho$ ($B = \widehat{U}_\rho$) before entering Y_ρ (\widehat{Y}_ρ).
- (iv) If $\alpha < f$, then

$$(75) \quad (\exists v_\alpha)(\forall x)(\forall s \geq v_\alpha) \left[\begin{array}{l} x \in R_{\alpha, s}^{\bar{A}} \implies (\forall t \geq s)[x \in R_{\alpha, t}^{\bar{A}} \text{ or } x \in R_{\alpha, t}^A] \\ \text{and } x \in R_{\alpha, s}^A \implies (\forall t \geq s)[x \in R_{\alpha, t}^A] \end{array} \right].$$

Proof. (i) Suppose $x \in Y_{\alpha, s+1} - Y_{\alpha, s}$. Then at stage $s + 1$ either Step 1, Step 2, or switching applies to x and α , so $x \in Y_{\beta, s}$ by (1.1) and (2.1), unless switching occurred, in which case $x \in Y_{\beta, s+1}^A$.

(ii) Suppose $x \in R_{\alpha, s+1} - R_{\alpha, s}$ and $x \in R_{\alpha, t} - R_{\alpha, t+1}$ for some $t > s$. Then $x < s$ by Step 6C. Hence, by Lemma 8.1(ii) at stage $t + 1$ either: (1) Step 6B applies to α and x , or (2) Step 1 applies to δ and x for some $\delta <_L \alpha$, $\delta = \alpha(x, t + 1)$. If (1), then $f_{s+1} <_L \alpha$, so x is γ -ineligible at all stages $v \geq t + 1$ and all $\gamma \succeq \alpha$, and x can never reenter R_α because of (1.3) and (2.3). If (2), then by Lemma 8.1(ii), (1.4), and induction on $v \geq t$, either for all $v \geq t$, $\alpha(x, v) <_L \alpha$ so $x \notin R_{\alpha, v}$, or else Step 6B applies at stage $v + 1$ to x and some $\eta <_L \alpha$, $\eta = \alpha(x, v)$, in which case the argument for (1) shows that $x \notin R_{\alpha, w}$, for all $w \geq v$.

(iii) Enumeration of x in $U_{\alpha, s+1}$ ($\widehat{V}_{\alpha, s+1}$) takes place only under Step 1, in which case $x \in Y_{\alpha, s+1}$, or under Step 4 (respectively, Step 3 or Step 5), in which case $x \in Y_{\alpha, s}$ already. However, x may enter $U_{\rho, s+1}$ directly from S_λ by Step 1.

(iv) Assume $\alpha < f$. Choose v_α such that for $s \geq v_\alpha$, $f_s \not<_L \alpha$, and no $\beta <_L \alpha$ acts at stage s , and hence $Y_{<\alpha, s} = Y_{<\alpha}$. Thus, if $x \in R_{\alpha, s}$ for $s \geq v_\alpha$, then x cannot be pulled to S_γ for $\gamma <_L \alpha$ by Step 1 $_\gamma$ and x cannot be removed from R_α by Step 6B, so x must remain in $R_{\gamma, t}$ for all $t \geq s$ or else be enumerated into A if it has not already been. □

Lemma 8.6. *For all $x \in \omega$,*

- (i) $\alpha(x) := \lim_s \alpha(x, s)$ exists, and
- (ii) x is enumerated in at most finitely many c.e. sets U_γ , \widehat{V}_γ , and hence for $\alpha = \alpha(x)$,

$$\nu(\alpha, x) := \lim_s \nu(\alpha, x, s) \text{ exists.}$$

Proof. (i) By (1.2), (2.2), and Lemma 8.1(i), $x \in S_{\alpha, s}$ implies $x > |\alpha|$ and $x > p$ if $\alpha = \beta \widehat{p}$ and $|\alpha| = 5e$. Fix x , let $\gamma = f \upharpoonright x$ and choose $s > v_\gamma$ (as defined in Lemma 8.5(iv)) such that $\gamma < f_s$. Let $\delta_0 = \alpha(x, s)$. Clearly, $\delta_0 <_L \gamma$ or $\delta_0 \preceq \gamma$ by

Step 6B. Also by induction on $t \geq s$, if $\delta_k = \alpha(x, t)$ and $\delta_{k+1} = \alpha(x, t + 1)$, then $\delta_{k+1} <_L \delta_k$ or $\delta_{k+1} \succ \delta_k$ because Step 1 or Step 2 must have applied to δ_k and x at stage $t + 1$ since Step 6B cannot apply to x after stage v_γ . But there is no infinite sequence $\{\delta_0, \delta_1, \dots\}$ such that for all k , $\delta_{k+1} <_L \delta_k$ or $\delta_{k+1} \succ \delta_k$, where x still satisfies (1.2) and (2.2).

(ii) By (i) choose $t_x \geq v_\gamma$ such that $\alpha(x, s) = \alpha$ for all $s \geq t_x$. Then $\nu(\alpha, x, s) \subseteq \nu(\alpha, x, s + 1)$ for all $s \geq t_x$. Hence,

$$\nu(\alpha, x) = \bigcup \{ \nu(\alpha, x, s + 1) : s \geq t_x \},$$

where this union is defined as in Definition 5.6(iv). □

Lemma 8.7. (i) *Step 6 applies infinitely often.*

(ii) *If the hypotheses of some Steps 1–5 (Steps $\widehat{1}$ – $\widehat{5}$) remain satisfied, then that step eventually applies.*

Proof. (i) If Step 6 applies at stage s , then the finitely many $x \in Y_{\lambda, s}$ ($\widehat{x} \in \widehat{Y}_{\lambda, s}$) remain the same until the next application of Step 6. Each later application of Steps 1–5 (Steps $\widehat{1}$ – $\widehat{5}$) chooses some $x(\widehat{x})$ to change position or to be enumerated in some set U_γ or \widehat{V}_γ (\widehat{U}_γ or V_γ). By Lemma 8.6, this can happen at most finitely often for each $x \in Y_{\lambda, s}$ ($\widehat{x} \in \widehat{Y}_{\lambda, s}$). Hence, Step 6 applies at some stage $t > s$.

(ii) Step 6 cannot apply at stage t if the hypotheses for some Steps 1–5 (Steps $\widehat{1}$ – $\widehat{5}$) are satisfied because the latter steps are performed before Step 6 by the basic construction in §7. □

8.2. Showing that $\mathcal{M}_\alpha \subseteq \mathcal{E}_\alpha$. The following lemma is based on Lemma 3.3. While the statement may seem entirely different, both lemmas show the satisfaction of negative requirements, which are interacting with positive requirements in the same way.

Lemma 8.8. *For all $\alpha \prec f$ and all $\nu_e \in \mathcal{M}_\alpha^{\overline{A}}$,*

$$(\exists^\infty s)[y_{\langle e, k \rangle}^s \text{ exists}] \implies \lim_s y_{\langle e, k \rangle}^s \text{ exists and is in } \overline{A}.$$

Proof. Induct on the length of α and on k . Let $\nu_e \in \mathcal{M}_\alpha^{\overline{A}}$. Assume true for all $\beta \prec \alpha$ and all $\langle e, i \rangle$ with $i < k$.

Let s_0 be such that $y_{\langle e, i \rangle}^s \in \overline{A}$ for all $s \geq s_0$ and all $i < k$.

Consider all the finitely many pairs $\langle \beta, n \rangle \prec \langle \alpha, e, k \rangle$. Let p be the maximum of $|\beta|$ for such a β . Choose some ξ associated with the positive requirement of making D nonlow (i.e. $|\xi| \equiv 4 \pmod{5}$), such that $|\xi| > p$, and $\alpha \prec \xi \widehat{0} \prec f$. Choose $s_1 \geq s_0$ such that for all $s \geq s_1$: (a) if $\langle \beta, n \rangle \prec \langle \alpha, e, k \rangle$, then the β -witness x_n^β does not become newly defined or enter D at stage s ; (b) $f_s \not\prec_L \xi$ (so every $\beta \prec_L \xi$ with $|\beta| \equiv 4 \pmod{5}$ has ceased to act by stage s); and (c) every β with $\beta \widehat{p} \prec \xi$ for $p \neq 0$ has ceased to act by stage s . By choice of s_1 no witness x_n^β , for $\beta \prec_L \xi$ or $\beta \widehat{p} \prec \xi$ for $p \neq 0$, will ever contribute a β -witness to D after stage s_1 .

Choose $s_2 \geq s_1$ such that ξ contributes a ξ -witness x_m^ξ to D at s_2 . This action of ξ causes P-initialization of all β with $\xi \prec_L \beta$, and cancellation of all such β -witnesses at stage s_2 . By (73), all β , $\beta \widehat{0} \prec \xi$, must have contributed their witnesses to D at stage s_2 , and so have no β -witnesses in existence at the end of stage s_2 .

Choose $s_3 \geq s_2$ such that $y_{\langle e, k \rangle}^{s_3}$ is defined.

Consider $\xi \prec \beta$. Suppose $\langle \beta, n \rangle > \langle \alpha, e, k \rangle$ and x_n^β is a witness that exists at stage s_3 . If x_n^β was appointed when $\widehat{\psi}_{\alpha,t}(y_{\langle e,k \rangle}^t) \geq \widehat{\psi}_{\alpha,s_3}(y_{\langle e,k \rangle}^{s_3})$, then x_n^β must be larger than $\widehat{\psi}_{\alpha,s_3}(y_{\langle e,k \rangle}^{s_3})$, so it will not injure it. However, if x_n^β was appointed when $\widehat{\psi}_{\alpha,t}(y_{\langle e,k \rangle}^t) < \widehat{\psi}_{\alpha,s_3}(y_{\langle e,k \rangle}^{s_3})$, then at some stage after s_3 , x_n^β could injure the current computation. Now, the only way for this to happen is for something else to injure the computation before stage s_3 and after x_n^β has been appointed. The injury could not come from a node $\gamma <_L \beta$ because that would reset the β -witness. It could not come from a node $\gamma \succ \beta$ because by (73), γ must wait for β to enumerate its witness before enumerating its own. Suppose the injury came from a node γ to the right of β . Then to avoid the γ -witness being reset, x_n^β must be defined before the γ -witness. Thus, by (72), γ must have its witness larger than x_n^β , which means that γ could not have injured the $\langle \alpha, y \rangle$ computation. Now, if $\gamma \widehat{p} \prec \beta$ for any $p \neq 0$, then by (71), β must wait for all γ -witnesses to enter D before defining its own witness. If $\gamma \widehat{0} \prec \beta$, then by (72), β must choose its witness smaller than the γ -witness, so γ cannot injure the $\langle \alpha, y \rangle$ computation. Thus, no β -witness for $\xi \prec \beta$ appointed before stage s_1 can possibly injure $\widehat{\psi}_{\alpha,s_3}(y_{\langle e,k \rangle}^{s_3})$.

By the time $y_{\langle e,k \rangle}^{s_3}$ is defined, the only β -witnesses in existence which have the priority to injure a computation $\widehat{\psi}_{\alpha,t}(x)$ for some $t \geq s_3$ are those which are permanently dormant and will never enter D .

Consider any new β -witness x_n^β appointed at or after stage s_3 . We cannot have $\beta <_L \xi$ by (b), or $\beta \widehat{p} \prec \xi$ by (c). If $\alpha \preceq \beta$, then we must have $\widehat{\psi}_{\alpha,s}(y_{\langle e,k \rangle}^s) \downarrow < x_n^\beta$ by (72). If $f <_L \beta$, then x_n^β must exceed $\widehat{\psi}_{\alpha,s}(y_{\langle e,k \rangle}^s)$ if the latter converges when x_n^β is appointed, and if not, x_n^β must be canceled whenever it later converges. (Note that any x_n^β newly appointed at s must exceed $\widehat{\psi}_\alpha^D(y_{\langle e,k \rangle}^s)[s]$ because the former must exceed s by the definition of Γ_s^β , and the latter must be less than s by the usual convention on the use function.)

Hence, $\widehat{\psi}_{\alpha,s_3}(y_{\langle e,k \rangle}^{s_3})$ is never injured, so $y_{\langle e,k \rangle}^{s_3} = \lim_s y_{\langle e,k \rangle}^s$ and does not enter A . □

Lemma 8.9. *Let $\alpha \prec f$. Then*

- (i) $(\forall \gamma <_L f)[m(\gamma) := \lim_s m(\gamma, s) < \infty]$,
- (ii) $m(\alpha) := \lim_s m(\alpha, s) = \infty$,
- (iii) $(\forall \nu_e \in \mathcal{M}_\alpha^A)(\forall k)(\exists^\infty s)[y_{\langle e,k \rangle}^s \text{ exists}]$,
- (iv) $(\forall \nu_e \in \mathcal{M}_\alpha^A)(\forall k)[\lim_s y_{\langle e,k \rangle}^s \text{ exists and is not in } A]$,
- (v) $\mathcal{E}_\alpha \supseteq \mathcal{M}_\alpha$, and
- (vi) $\widehat{\mathcal{E}}_\alpha \supseteq \widehat{\mathcal{M}}_\alpha$.

Proof. (i) If $\gamma <_L f$, then $\gamma \prec f_s$ for finitely many s , so finitely many γ -entries are ever added to \mathcal{L} and hence \mathcal{L} is γ -marked finitely often and $m(\gamma) < \infty$.

For (ii)-(vi), induct on α .

For the base case, $\alpha = \lambda$, we have (ii) because we never put any entries on the λ -list, so every time Step 6 acts, $m(\lambda, s)$ increases. Part (iii) follows by the fact that \bar{A} is infinite, and Lemma 8.8 gives (iv). For (v) and (vi), we have $\mathcal{M}_\lambda \subset \mathcal{E}_\lambda$ and its dual simply because Step 6 acts infinitely often.

Assume the theorem is true for all $\gamma \prec \alpha$.

Suppose for a contradiction that $m(\alpha) < \infty$, say $m(\alpha, s) = m_0$ for all $s \geq s_0$.

Claim 1. Every α -entry $\langle \alpha, \nu_1 \rangle$ on \mathcal{L} ($\langle \alpha, \widehat{\nu}_1 \rangle$ on $\widehat{\mathcal{L}}$) is eventually marked.

Proof.

Case 1. The first unmarked entry is in \mathcal{M}_α^A .

Suppose that some α -entry $\langle \alpha, \nu_1 \rangle$ on \mathcal{L} is never marked, where $\nu_1 \in \mathcal{M}_\alpha^A$. Hence, by Step 6A there are only finitely many α -entries on \mathcal{L} . Choose $s_1 \geq s_0$ such that every α -entry on \mathcal{L} and every entry on \mathcal{L} preceding $\langle \alpha, \nu_1 \rangle$ which will ever be marked is marked by stage s_1 , $Y_{<\alpha, s_1} = Y_{<\alpha}$, and for all $x \leq m_0$, $x \in Y_{\alpha, s_1}$ iff $x \in Y_\alpha$.

Suppose $e_\alpha > e_\beta$. Then $\nu_1 \in \mathcal{M}_\alpha = \mathcal{F}_\beta^+$ since $\alpha \prec f$. Thus,

$$(\exists^\infty x)(\exists s > s_1)[x \in Y_{\beta, s}^A \ \& \ \nu^+(\alpha, x, s) = \nu_1].$$

By the choice of s_1 almost every such x also satisfies (1.1)–(1.8). Thus, some such x is moved to S_α under Step 1 at some stage $s + 1 > s_1$ and the entry $\langle \alpha, \nu_1 \rangle$ is then marked, contrary to hypothesis.

Suppose $e_\alpha = e_\beta$. Now $\nu_1 \in \mathcal{M}_\alpha$, so $\nu_1 \upharpoonright \beta \in \mathcal{M}_\beta$. By the induction hypothesis, $\nu_1 \upharpoonright \beta \in \mathcal{E}_\beta$. Thus

$$(\exists^\infty x)(\exists s > s_1)[x \in Y_{\beta, s}^A \ \& \ \nu(\beta, x, s) = \nu_1 \upharpoonright \beta].$$

By the choice of s_1 almost every such x also satisfies (1.1)–(1.8). Thus, some such x is moved to S_α under Step 1 at some stage $s + 1 > s_1$ and the entry $\langle \alpha, \nu_1 \rangle$ is then marked, contrary to hypothesis.

Case 2. The first unmarked entry is in $\mathcal{M}_\alpha^{\bar{A}}$.

Suppose that some α -entry $\langle \alpha, \nu_1 \rangle$ on \mathcal{L} is never marked, where $\nu_1 \in \mathcal{M}_\alpha^{\bar{A}}$. Hence, by Step 6A there are only finitely many α -entries on \mathcal{L} . Choose $s_1 \geq s_0$ such that every α -entry on \mathcal{L} and every entry on \mathcal{L} preceding $\langle \alpha, \nu_1 \rangle$ which will ever be marked is marked by stage s_1 , $Y_{<\alpha, s_1} = Y_{<\alpha}$, and for all $x \leq m_0$, $x \in Y_{\alpha, s_1}$ iff $x \in Y_\alpha$.

Suppose $e_\alpha > e_\beta$. Then $\mathcal{M}_\alpha = \mathcal{F}_\beta^+$ since $\alpha \prec f$. So

$$(\exists^\infty x)(\exists s > s_1)[x \in Y_{\beta, s}, \nu^+(\alpha, x, s) = \nu_1, \widehat{\Psi}_\alpha^D(x)[s] \downarrow = 0].$$

By the choice of s_1 almost every such x also satisfies (1.1)–(1.7) of Step 1. Thus, some such x is moved to S_α under Step 1 at some stage $s + 1 > s_1$ and the entry $\langle \alpha, \nu_1 \rangle$ is then marked, contrary to hypothesis.

Suppose $e_\alpha = e_\beta$. Now $\nu_1 \in \mathcal{M}_\alpha$, so $\nu_1 \upharpoonright \beta \in \mathcal{M}_\beta$. By the induction hypothesis,

$$(\forall k)(\forall^\infty s)[\exists x = y_{\langle e', k \rangle}^s \text{ where } x \notin A],$$

where $\nu_{e'} = \nu_1 \upharpoonright \beta$. By the choice of s_1 , almost all of these must stay in S_β . Since x is not allowed to change states without appearing to not be in A , these x values must at some stage be in S_β , appear to be in \bar{A} , and be in state $\nu_1 \upharpoonright \beta$. By the choice of s_1 almost every such x also satisfies (1.1)–(1.7). Thus, some x is moved to S_α under Step 1 at some stage $s + 1 > s_1$ and the entry $\langle \alpha, \nu_1 \rangle$ is then marked, contrary to hypothesis. This proves the claim.

To complete the proof of (ii) use the claim to find $s > s_0$ such that $\alpha \prec f_{s+1}$ and every α -entry on \mathcal{L}_s and $\widehat{\mathcal{L}}_s$ is marked. Now by Step 6A, $m(\alpha, s+1) > m(\alpha, s) = m_0$, contrary to the choice of s_0 .

For (iii), induct on k . Suppose it is true for all $i < k$. Then by Theorem 8.8, if $k > 0$, then $\lim_s y_{\langle e, k-1 \rangle}^s = y_{\langle e, k-1 \rangle}$ exists and is not in A . Since $m(\alpha) = \infty$ by part (ii), we know that infinitely many elements are pulled up into ν_e by Step 1. Thus,

$y_{(e,k)}^s$ will be infinitely often defined. (Note that $y_{(e,k)}^s$ may be defined during stage s before any elements enter D in stage s .) Part (iv) follows by Theorem 8.8.

Parts (v) and (vi) follow by (ii), since every element of \mathcal{M}_α is marked by Step 1. □

8.3. Completing the automorphism verification.

Lemma 8.10. $\alpha \prec f \implies R_{\alpha,\infty} =^* Y_\alpha =^* Y_\lambda = \omega$.

Proof. By Lemma 8.7(i), Step 6C must eventually put every element $x \in \omega$ into Y_λ . By induction we may assume $R_{\beta,\infty} =^* Y_\beta =^* \omega$, for $\beta^- = \alpha$. By Lemma 8.9, $m(\alpha) = \infty$ and $m(\gamma) < \infty$ for all $\gamma <_L \alpha$ with $\gamma^- = \beta$.

By Lemma 8.4, $Y_{<\alpha} =^* \emptyset$ and almost every $x \in R_\beta$ not yet in R_α must eventually lie in S_β . Hence, almost every $x \in R_\beta^{\bar{A}}$ not yet in $R_\alpha^{\bar{A}}$ must eventually satisfy the conditions of Step $2^{\bar{A}}$ and must eventually move to S_α by Step 2, or else be enumerated into A . Once in A , almost every $x \in R_\beta^A$ not yet in R_α^A must eventually satisfy the conditions of Step 2^A and be moved to S_α by Step 2. By Lemma 8.5(iv) almost every such $x \in \bar{A}$ will remain in R_α forever. □

Lemma 8.11. $\alpha \prec f \implies \alpha$ is \mathcal{M} -consistent.

Proof. Let $\alpha \prec f$ and $\beta = \alpha^-$. Assume for a contradiction that α is not \mathcal{M} -consistent. Then $e_\alpha > e_\beta$ and there exist $\nu_0 \in \mathcal{M}_\alpha$, $\nu_1 \notin \mathcal{M}_\alpha$, $\nu_0 <_B \nu_1$ and $\nu_1 \upharpoonright \beta \in \mathcal{M}_\beta$.

By (66), α is a terminal node on T , so $S_\alpha = R_\alpha$. By Lemmas 8.10 and 8.5(iv), $S_{\alpha,\infty} =^* \omega$ and no $x \in S_{\alpha,s}$, $s > v_\alpha$, later leaves S_α . By Lemma 8.9, $\mathcal{E}_\alpha \supseteq \mathcal{M}_\alpha$, and

$$(\exists^\infty x)(\exists s)[x \in S_{\alpha,s+1} - S_{\alpha,s} \text{ by Step 1, and after Step 1, } x \text{ is in state } \nu_0].$$

Choose any such x and $s > v_\alpha$. At the end of Step 1, Step 3 will act and enumerate x into such sets that $\nu(\alpha, x, s + 1) = \nu_1$. Thus, α is provably incorrect at all stages $v > s + 1$, so $\alpha \not\prec f$. □

Lemma 8.12. *If $\alpha \prec f$, then*

- (i) $\widehat{\mathcal{M}}_\alpha = \{\widehat{\nu} : \nu \in \mathcal{M}_\alpha\}$,
- (ii) $\mathcal{M}_\alpha = \mathcal{F}_\alpha = \mathcal{E}_\alpha$, and
- (iii) $\widehat{\mathcal{M}}_\alpha = \widehat{\mathcal{F}}_\alpha = \widehat{\mathcal{E}}_\alpha$.

Proof. Fix $\alpha \prec f$, and let $\beta = \alpha^-$. Now (i) holds by the definitions of \mathcal{M}_α and $\widehat{\mathcal{M}}_\alpha$. Assume (ii) and (iii) for β . We know $\mathcal{E}_\alpha \subseteq \mathcal{F}_\alpha$ by their definitions, and $\mathcal{M}_\alpha \subseteq \mathcal{E}_\alpha$ by Lemma 8.9. Thus, to prove (ii) (and (iii)) it suffices to prove $\mathcal{F}_\alpha \subseteq \mathcal{M}_\alpha$ (and $\widehat{\mathcal{F}}_\alpha \subseteq \widehat{\mathcal{M}}_\alpha$). Induct on α . Suppose the theorem holds for all $\beta \prec \alpha$.

Case 1. $e_\alpha = e_\beta$ and $\widehat{e}_\alpha = \widehat{e}_\beta$.

Then $\mathcal{M}_\alpha = \mathcal{M}_\beta$. Also $\mathcal{F}_\alpha \subseteq \mathcal{F}_\beta$ since $Y_\alpha \subseteq Y_\beta$. Finally, $\mathcal{M}_\beta = \mathcal{F}_\beta$ by the inductive hypothesis (ii) for β . Hence,

$$\mathcal{F}_\alpha \subseteq \mathcal{F}_\beta = \mathcal{M}_\beta = \mathcal{M}_\alpha,$$

so (ii) holds for α . Likewise, $\widehat{\mathcal{F}}_\alpha \subseteq \widehat{\mathcal{M}}_\alpha$, so (iii) holds for α .

Before considering Cases 2 and 3 we need a technical sublemma.

Sublemma. *If $e_\alpha > e_\beta$, $\nu_2 = \langle \alpha, \sigma_2, \tau_2 \rangle \in \mathcal{F}_\beta^+$, and $\nu_1 = \langle \alpha, \sigma_1, \tau_2 \rangle$, where $\sigma_1 = \sigma_2 - \{e_\alpha\}$, then $\nu_1 \in \mathcal{F}_\beta^+$ also.*

Proof. Suppose $\nu_2 \in \mathcal{F}_\beta^+$. Then $\nu_3 = \nu_2 \upharpoonright \beta \in \mathcal{M}_\beta$.

Suppose $\nu_3 \in \mathcal{M}_\beta^A$. By Lemma 8.9, for every k , there is an x that enters ν_3 by Step 1, and immediately becomes the final $y_{\langle e, k \rangle}^s$ for the appropriate e . Hence,

$$(\exists^\infty x)(\exists s)[x \in Y_{\beta, s} - Y_{\beta, s-1}, \nu(\beta, x, s) = \nu_3, \widehat{\Psi}_\beta^D(x)[s] \downarrow = 0].$$

However, for each such x and s , $x \notin Z_{e_\alpha, s}$ (by the definition of $Z_{e_\alpha, s}$ in §6.1) so $\nu^+(\alpha, x, s) = \nu_1$. Hence, $\nu_1 \in \mathcal{F}_\beta^+$ by the definition of \mathcal{F}_β^+ because $\widehat{\psi}_{\beta, s}(x) = \widehat{\psi}_{\alpha, s}(x)$ since β is not a positive requirement node.

Suppose $\nu_3 \in \mathcal{M}_\beta^A$. Then $\nu_3 \in \mathcal{F}_\beta^A = \mathcal{E}_\beta^A$ by the inductive hypothesis for β . Hence, by the definition of \mathcal{E}_β^A ,

$$(\exists^\infty x)(\exists s)[x \in Y_{\beta, s}^A - Y_{\beta, s-1} \ \& \ \nu(\beta, x, s) = \nu_3].$$

But for each such x and s , $x \notin Z_{e_\alpha, s}$ (by the definition of $Z_{e_\alpha, s}$ in §6.1) so $\nu^+(\alpha, x, s) = \nu_1$. Hence, $\nu_1 \in \mathcal{F}_\beta^+$ by the definition of \mathcal{F}_β^+ . \square

The dual of the Sublemma is proved as in the \mathcal{M}_β^A case above, with B or \overline{B} in place of A .

Case 2. $e_\alpha > e_\beta$.

We prove $\mathcal{F}_\alpha \subseteq \mathcal{M}_\alpha$ and its dual $\widehat{\mathcal{F}}_\alpha \subseteq \widehat{\mathcal{M}}_\alpha$ in the following claims. (The proof of Case 3, $\widehat{e}_\alpha > \widehat{e}_\beta$, is similar, but slightly simpler. In particular, the proof of $\widehat{\mathcal{F}}_\alpha \subseteq \widehat{\mathcal{M}}_\alpha$ is dual to Claim 1 for the A -side of the tree, and it is not necessary to differentiate the B and \overline{B} sides in the proof. The proof of $\mathcal{F}_\alpha \subseteq \mathcal{M}_\alpha$ is dual to Claims 2a, 2b, and 2c.)

Claim 1. $\mathcal{F}_\alpha \subseteq \mathcal{M}_\alpha$.

Proof. Suppose $\nu_1 \in \mathcal{F}_\alpha$. Let $\nu_1 = \langle \alpha, \sigma_1, \tau_1 \rangle$. Then

$$(76) \quad (\exists^\infty x)(\exists s)[x \in Y_{\alpha, s} \ \& \ \nu(\alpha, x, s) = \nu_1].$$

Note that $Y_{\alpha, s} \subseteq Y_{\beta, s}$ and $\nu(\alpha, x, s) \leq_R \nu^+(\alpha, x, s)$ because $U_{\alpha, s} \subseteq Z_{e_\alpha, s}$.

Subcase 1. $\nu_1 \in \mathcal{F}_\alpha^A$.

The $Y_{\alpha, s}$ in (76) can be taken to be $Y_{\alpha, s}^A$. Now, we have

$$(77) \quad (\exists^\infty x)(\exists s)[x \in Y_{\beta, s}^A \ \& \ \nu(\beta, x, s) = \nu_1 \upharpoonright \beta].$$

By the inductive hypothesis, $\nu_1 \upharpoonright \beta \in \mathcal{F}_\beta \subset \mathcal{M}_\beta$. As in Lemma 8.9 (iv), infinitely many of these x values stay in \overline{A} forever, and appear to be in \overline{A} from the moment they enter S_β^A in state $\nu_1 \upharpoonright \beta$. For each such x that enters S_β^A , at s , either $\nu^+(\alpha, x, s) = \nu_1$, in which case $\nu_1 \in \mathcal{F}_\beta^+$ since x appears to be in \overline{A} , or $\nu^+(\alpha, x, s) = \nu_2$, where $\nu_2 = \langle \alpha, \sigma_2, \tau_1 \rangle$ and where $e_\alpha \notin \sigma_1$ and $\sigma_2 = \sigma_1 \cup \{e_\alpha\}$. Now $\nu_2 \in \mathcal{F}_\beta^+$ since $Y_{\alpha, s} \subseteq Y_{\beta, s}$, so $\nu_1 \in \mathcal{F}_\beta^+ = \mathcal{M}_\alpha$ by the Sublemma.

Subcase 2. $\nu_1 \in \mathcal{F}_\alpha^A$.

The $Y_{\alpha,s}$ in (76) can be taken to be $Y_{\alpha,s}^A$. First suppose

$$(78) \quad (\exists^\infty x)(\exists s)[x \in Y_{\alpha,s}^A \ \& \ \nu^+(\alpha, x, s) = \nu_1].$$

Then $\nu_1 \in \mathcal{F}_\beta^+$ by definition of \mathcal{F}_β^+ because $Y_{\alpha,s}^A \subseteq Y_{\beta,s}^A$, and $\mathcal{F}_\beta^+ = \mathcal{M}_\alpha$ since $\alpha \prec f$.

If (78) fails, then for almost every x in (76), $\nu^+(\alpha, x, s) = \nu_2 >_R \nu_1$, so $\nu_2 = \langle \alpha, \sigma_2, \tau_1 \rangle$, where $e_\alpha \notin \sigma_1$ and $\sigma_2 = \sigma_1 \cup \{e_\alpha\}$. Now $\nu_2 \in \mathcal{F}_\beta^+$ since $Y_{\alpha,s}^A \subseteq Y_{\beta,s}^A$, so $\nu_1 \in \mathcal{F}_\beta^+ = \mathcal{M}_\alpha$ by the Sublemma. □

Claim 2. $\widehat{\mathcal{F}}_\alpha \subseteq \widehat{\mathcal{M}}_\alpha$.

Proof. We establish Claim 2 by the next three claims. □

Claim 2a. $\widehat{\mathcal{E}}_\alpha \subseteq \widehat{\mathcal{M}}_\alpha$.

Proof. Assume $\widehat{\nu}_1 \in \widehat{\mathcal{E}}_\alpha$. Hence,

$$(\exists^\infty \widehat{x})(\exists s)[\widehat{x} \in \widehat{S}_{\alpha,s+1} - \widehat{Y}_{\alpha,s} \ \& \ \nu(\alpha, \widehat{x}, s + 1) = \widehat{\nu}_1].$$

For every such \widehat{x} and s , \widehat{x} must have entered $\widehat{S}_{\alpha,s+1}$ under Step $\widehat{1}$ or Step $\widehat{2}$. If Step $\widehat{1}$ applied, then we marked an entry $\langle \alpha, \widehat{\nu}_1 \rangle$ on $\widehat{\mathcal{L}}_s$, so $\widehat{\nu}_1 \in \widehat{\mathcal{M}}_\alpha$ by the definition of $\widehat{\mathcal{L}}$ in Step 6. If Step $\widehat{2}$ applied, then $\widehat{x} \notin \widehat{U}_{\alpha,s+1}$ because $\widehat{x} \notin \widehat{U}_{\alpha,s}$ by Lemma 8.5(iii) and no enumeration takes place at stage $s + 1$ under Step $\widehat{2}$. Hence, $e_\alpha \notin \sigma_1$, where $\nu_1 = \langle \alpha, \sigma_1, \tau_1 \rangle$.

Let $\nu_3 = \nu_1 \upharpoonright \beta$. Now $\widehat{\nu}_3 \in \widehat{\mathcal{F}}_\beta = \widehat{\mathcal{M}}_\beta$, so $\nu_3 \in \mathcal{M}_\beta = \mathcal{F}_\beta$ and thus either $\nu_1 \in \mathcal{F}_\beta^+$ or $\nu_2 \in \mathcal{F}_\beta^+$, where $\nu_2 = \langle \alpha, \sigma_1 \cup \{e_\alpha\}, \tau_1 \rangle$, by the same reasoning in Claim 1a. But if $\nu_2 \in \mathcal{F}_\beta^+$, then $\nu_1 \in \mathcal{F}_\beta^+$ by the Sublemma. In either case $\nu_1 \in \mathcal{F}_\beta^+ = \mathcal{M}_\alpha$, so $\widehat{\nu}_1 \in \widehat{\mathcal{M}}_\alpha$. □

Claim 2b. If $\widehat{x} \in \widehat{Y}_{\alpha,s}$, $\widehat{\nu}_1 = \widehat{\nu}(\alpha, \widehat{x}, s) \in \mathcal{M}_\alpha$, $s > v_\alpha$ of Lemma 8.5(iv), and Red causes enumeration of \widehat{x} so that $\widehat{\nu}_2 = \widehat{\nu}(\alpha, \widehat{x}, s + 1)$, then $\widehat{\nu}_2 \in \widehat{\mathcal{M}}_\alpha$.

Proof. Suppose this enumeration occurs. Then $\widehat{\nu}_1 <_R \widehat{\nu}_2$, so $\nu_1 <_B \nu_2$ by (37). Now $\widehat{\nu}_1 \in \widehat{\mathcal{M}}_\alpha$, so $\nu_1 \in \mathcal{M}_\alpha$. But α is \mathcal{M} -consistent by Lemma 8.11, so $\nu_2 \in \mathcal{M}_\alpha$, and hence $\widehat{\nu}_2 \in \widehat{\mathcal{M}}_\alpha$. □

Claim 2c. If $\widehat{x} \in \widehat{Y}_{\alpha,s}$, $\widehat{\nu}_1 = (\alpha, \widehat{x}, s) \in \widehat{\mathcal{M}}_\alpha$, $s > v_\alpha$ of Lemma 8.5(iv), and Blue causes enumeration of \widehat{x} so that $\widehat{\nu}_2 = \widehat{\nu}(\alpha, \widehat{x}, s + 1)$, then $\widehat{\nu}_2 \in \widehat{\mathcal{M}}_\alpha$.

Proof. Suppose $\widehat{x} \in \widehat{Y}_{\alpha,s}$ and Blue causes this enumeration at stage $s + 1$, so $\widehat{\nu}_1 <_B \widehat{\nu}_2$. Since $s > v_\alpha$, $\widehat{x} \in \widehat{R}_{\alpha,s} \cap \widehat{R}_{\alpha,s+1}$. Hence, Step $\widehat{1}$, Step $\widehat{3}$, or Step $\widehat{5}$ applies to \widehat{x} at stage $s + 1$ for some $\gamma \succeq \alpha$. If Step $\widehat{1}_\gamma$ or Step $\widehat{5}_\gamma$ applies, then $\widehat{\nu}_3 = \widehat{\nu}(\gamma, \widehat{x}, s + 1) \in \widehat{\mathcal{M}}_\gamma$, so $\widehat{\nu}_2 = \widehat{\nu}_3 \upharpoonright \alpha \in \widehat{\mathcal{M}}_\alpha$. (Here Step $\widehat{5}_\gamma$ means Step $\widehat{5}$, Case 1 for $\widehat{x} \in \widehat{Y}_{\gamma,s}$ or Step $\widehat{5}$, Case 2 for $\widehat{x} \in \widehat{Y}_{\delta,s}$ where $\gamma = \delta^-$.) If Step $\widehat{3}_\gamma$ applies, then $\gamma \not\succeq \alpha$ (since α is \mathcal{M} -consistent and γ is not) and $\widehat{\nu}_3 = \widehat{\nu}(\gamma^-, \widehat{x}, s + 1) \in \widehat{\mathcal{M}}_{\gamma^-}$ by (3.4), so $\widehat{\nu}_2 = \widehat{\nu}_3 \upharpoonright \alpha \in \widehat{\mathcal{M}}_\alpha$. This completes the proof of Claim 2c. □

Claims 2a, 2b, and 2c establish Claim 2 because Claim 2a shows that any entry α -state is in $\widehat{\mathcal{M}}_\alpha$, and any α -state that is well-visited by elements coming from states in $\widehat{\mathcal{M}}_\alpha$ must also be in $\widehat{\mathcal{M}}_\alpha$. So $\widehat{\mathcal{F}}_\alpha \subseteq \widehat{\mathcal{M}}_\alpha$.

Lemma 8.13. $\alpha \prec f \implies \alpha$ is \mathcal{R} -consistent.

Proof. Assume for a contradiction that $\alpha \prec f$ and α is not \mathcal{R} -consistent. Now α is either of length 3 mod 5 or length 4 mod 5. Assume the former, so the \mathcal{R} -inconsistency occurs on the A -side of the construction. Choose $\nu_e \in \mathcal{R}_\alpha$ such that for all $\nu \in \mathcal{M}_\alpha$, $\nu_e \not\prec_R \nu$. By (66) α is a terminal node on T , so $S_\alpha = R_\alpha$. By Lemmas 8.10 and 8.5(iv), $S_{\alpha,\infty} =^* \omega$ and no $x \in S_{\alpha,s}$, $s > v_\alpha$, later leaves S_α . Now $\nu_e \in \mathcal{R}_\alpha \subseteq \mathcal{M}_\alpha = \mathcal{E}_\alpha$ by Lemma 8.12, so

$$(\exists^\infty x)(\exists s > v_\alpha)[x \in S_{\alpha,s+1} - Y_{\alpha,s} \ \& \ \nu(\alpha, x, s) = \nu_e].$$

Note that for states associated with \overline{A} , infinitely many of these x values are in fact $y_{\langle e,k \rangle}$ values for some $k \in \omega$, and thus stay in \overline{A} . For each such x and s , since α is a terminal node, neither Step 1 nor Step 2 can apply to x at any stage $t > s + 1$. Now Step 3 cannot apply to $x \in S_{\alpha,t}$ because α is \mathcal{M} -consistent by Lemma 8.11. Furthermore, Step 5 cannot apply to $x \in S_{\alpha,t}$ while $\nu(\alpha, x, t) = \nu_e$ because $\nu_e \in \mathcal{R}_\alpha$ and $\mathcal{R}_\alpha \cap \mathcal{B}_\alpha = \emptyset$. But if $\nu(\alpha, x, t) = \nu_e$ for all $t \geq s$, then x witnesses that $F(\alpha^-, \nu_e)$ fails, so $\nu_e \in \mathcal{R}_\alpha$ contradicts $\alpha \prec f$. Hence, either x is enumerated in A or Step 4 applies to x . There must be infinitely many x such that Step 4 applies since infinitely many elements in ν_e never enter A . For the dual case, we get that B cannot empty ν_e because for every element that we enumerate into B from ν_e , we leave one element in \overline{B} . Hence, for infinitely many x , Step 4 applies to $x \in S_{\alpha,t}$ at some stage $t + 1 > s + 1$ such that $\nu_e = \nu(\alpha, x, s) = \nu(\alpha, x, t)$, $\nu = \nu(\alpha, x, t + 1)$, and $\nu_e \prec_R \nu$. Choose ν such that this happens for infinitely many $x \in S_\alpha$. Now $\nu \in \mathcal{F}_\alpha$, so $\nu \in \mathcal{M}_\alpha$ by Lemma 8.12. \square

Lemma 8.14. If $\alpha \prec f$ and $\nu_1 \in \mathcal{B}_\alpha$, then $\{x : x \in Y_\alpha \ \& \ \nu(\alpha, x) = \nu_1\} =^* \emptyset$.

Proof. Fix $\alpha \prec f$ and $\nu_1 \in \mathcal{B}_\alpha$. Let v_α be as in Lemma 8.5(iv). Assume for a contradiction that $x \in R_{\alpha,s}$ for some $s > v_\alpha$ and that for all $t \geq s$, $\gamma = \alpha(x, t)$, and $\nu_1 = \nu(\alpha, x, t)$. Now $\gamma \succeq \alpha$ and $\alpha \in T$, so by the condition on T in Definition 6.2 (vi) we have $\nu'_1 \in \mathcal{B}_\gamma$ for all $\nu'_1 \in \mathcal{M}_\gamma$ such that $\nu'_1 \upharpoonright \alpha = \nu_1$. \square

Case 1. γ is consistent. Then Step 5, Case 1 applies to x and γ at some stage $t + 1 > s$, so $\nu'_1 = \nu(\gamma, x, t)$, $\nu'_2 = \nu(\gamma, x, t + 1)$, $\nu'_1 \prec_B \nu'_2$, and $\nu'_2 \in \mathcal{M}_\gamma - \mathcal{B}_\gamma$. Hence, $\nu_2 = \nu'_2 \upharpoonright \alpha \in \mathcal{M}_\alpha - \mathcal{B}_\alpha$, and $\nu(\alpha, x, t + 1) = \nu_2 >_B \nu_1$.

Case 2. Otherwise. Then at some stage $t + 1 > s$, Step 5, Case 2 applies to x and $\delta = \gamma^- \succeq \alpha$, so $\nu(\alpha, x, t + 1) = \nu_2 >_B \nu_1$ as in Case 1 but with δ in place of γ . \square

Lemma 8.15. The correspondence $U_\alpha \leftrightarrow \widehat{U}_\alpha$ and $\widehat{V}_\alpha \leftrightarrow V_\alpha$, $\alpha \prec f$, defines an automorphism of \mathcal{E}^* .

Proof. Choose $\alpha \prec f$. By Lemmas 8.11 and 8.13, α is \mathcal{M} -consistent and α is also \mathcal{R} -consistent. Hence, α is consistent by Definition 6.1. Thus, by the Definition 6.3, f is infinite, and hence $\lim_{\alpha \prec f} e_\alpha = \infty$.

By Lemma 8.10, $Y_\alpha =^* \omega$; by Lemma 8.12, we have (35), its dual, and (36) (so the well-visited α -states on ω coincide with those on $\widehat{\omega}$); and by Lemma 8.14 and its dual, we have (39). It immediately follows that the automorphism requirement (25) is satisfied as remarked in §5.5. \square

8.4. **Showing that D is not low and that B is low.** The next lemma is a slight modification of Lemma 3.4.

Lemma 8.16. *For every $j \in \omega$ the strategy of every positive requirement node $\beta \prec f$ has satisfied the requirement,*

$$(79) \quad P_j : \quad \widehat{\varphi}_j \neq \text{characteristic function of } \{x : W_x \cap \overline{D} \neq \emptyset\},$$

so that D is not low_1 .

Proof. Fix $\beta = f \upharpoonright (5j + 4)$. Suppose that $\widehat{\varphi}_j$ is the Δ_2^0 characteristic function of $\{i : W_i \cap \overline{D} \neq \emptyset\}$. Hence, for all i , $\lim_s \widetilde{\varphi}_{j,s}(i)$ exists and equals $\widehat{\varphi}_j(i)$. (Here $\widehat{\varphi}_j$ and $\widetilde{\varphi}_{j,s}(i)$ were defined in §3.1.)

Suppose $\widetilde{\varphi}_{j,s}(g(\beta)) \downarrow = 0$ for all $s \geq s_0$ for some s_0 . Since $\beta \prec f$ there is some stage $s_1 > s_0$ after which β is never P-initialized as in Subcase 3 of the D -Module because if $\widehat{\psi}_\gamma(x)$ is ever defined for $\gamma <_L \beta$ after the true path permanently leaves γ , all witnesses to the right of γ get reset to prevent injury, so $\widehat{\psi}_\gamma(x)$ will never be redefined. Also for all α , $\alpha \preceq \beta$, α guesses correctly at the well-visited states, and by Lemma 8.9, for each $e \in \mathcal{M}_\alpha$ and each k , $y_{\langle e,k \rangle}^s$ is defined at some stage. Hence, β waits for a witness $y = x_n^\beta$ in (72) to appear at some stage $x > s_1$. Then according to Case 3, Subcase 1 of the D -Module, β will enumerate x_n^β into $Z_\beta = W_{g(\beta)}$ causing $Z_\beta \cap \overline{D} \neq \emptyset$. Since $\widetilde{\varphi}_{j,t}(g(\beta)) = \widetilde{\varphi}_{j,s}(g(\beta))$ for all $t > s$, β will never enumerate x_n^β into D , so $\lim_s \widetilde{\varphi}_{j,s}(g(\beta)) = 0$, but $W_{g(\beta)} \cap \overline{D} \neq \emptyset$, a contradiction.

Next suppose $\lim_s \widetilde{\varphi}_{j,s}(g(\beta)) \downarrow = 1$ for all $s \geq s_0$ for some s_0 . For every s there can be at most one element in $Z_s^\beta - D_s$ and that must be $\Gamma_s = x_n^\beta$ for some n . Since $\beta \prec f$ we know every ξ , $\xi \widehat{\sim} 0 \prec \beta$, will have the Π_2 -outcome. Hence, for each such x_n^β , there will come a stage by (73) at which x_n^β is enumerated in D under Subcase 2. But no new β -witness can be appointed after stage s_0 under Subcase 1. Hence, $\lim_s \widetilde{\varphi}_{j,s}(g(\beta)) \downarrow = 1$, but $W_{g(\beta)} \cap \overline{D} = \emptyset$, a contradiction. \square

Lemma 8.17. *The set B is low_1 .*

Proof. Suppose there exist infinitely many s such that $\Phi_i^B(i)[s] \downarrow$.

Let s_0 be the least s such that for all $\langle \alpha, x, e \rangle \leq i$, $\langle \alpha, x, e \rangle$ has either been removed from the list Λ already or will never be removed from the list (this can happen if it is never added to the list, or if it is added but never matched). Since each $\langle \alpha, x, e \rangle$ can enter Λ only once, then after stage s_0 , no y will enter B in order to match $\langle \alpha, x, e \rangle$. Let $s > s_0$ be some stage with $\Phi_i^B(i)[s] \downarrow$. Then by (B.3), nothing can enter B below the use of this computation. So $(\forall t > s) [\Phi_i^B(i)[t] \downarrow]$. So either $(\forall^\infty s) [\Phi_i^B(i)[s] \downarrow]$ or $(\forall^\infty s) [\Phi_i^B(i)[s] \uparrow]$. Thus, B is low. \square

We have constructed a set D that is nonlow (by Lemma 8.16) such that for $A = W_i = \Psi^D$, there is a B that is low (by Lemma 8.17) such that an automorphism of \mathcal{E}^* takes A to B (by Lemma 8.15). By [15, page 343], building an automorphism of \mathcal{E}^* is equivalent to building one of \mathcal{E} . By §4.7, every c.e. $A \leq_{\text{T}} D$ can be taken to a low set B by an automorphism of \mathcal{E} . In particular, every c.e. set A of the same degree as D is automorphic to a low set B . Thus, the nonlow degrees cannot be invariant, and so cannot be definable. This completes the proof of Theorem 1.3. The nonlow degrees are the only upward closed jump class that is not definable.

REFERENCES

- [1] P. A. Cholak, Automorphisms of the Lattice of Recursively Enumerable Sets, *Mem. Amer. Math. Soc.* **113** (1995). MR1227497 (95f:03064)
- [2] P. A. Cholak and L. A. Harrington, On the Definability of the Double Jump in the Computationally Enumerable Sets, *J. Math. Log.* **2** (2002), 261–296. MR1938925 (2003h:03063)
- [3] L. Harrington and R. I. Soare, Definability, automorphisms, and dynamic properties of computably enumerable sets, *Bull. Symbolic Logic*, **2** (1996), 199–213. MR1396855 (97d:03057)
- [4] L. Harrington and R. I. Soare, The Δ_3^0 automorphism method and noninvariant classes of degrees, *J. Amer. Math. Soc.*, **9** (1996), 617–666. MR1311821 (96j:03060)
- [5] L. Harrington and R. I. Soare, Definable properties of the computably enumerable sets, Proceedings of the Oberwolfach Conference on Computability Theory, *Ann. Pure Appl. Logic*, **94** (1998), 97–125. MR1640265 (99f:03055)
- [6] A. H. Lachlan, Degrees of recursively enumerable sets which have no maximal superset, *J. Symbolic Logic* **33** (1968), 431–443. MR0236016 (38:4314)
- [7] A. H. Lachlan, On some games which are relevant to the theory of recursively enumerable sets, *Ann. of Math. (2)* **91** (1970), 291–310. MR0284333 (44:1562)
- [8] D. A. Martin, Classes of recursively enumerable sets and degrees of unsolvability, *Z. Math. Logik Grundlagen Math.*, **12** (1966), 295–310. MR0224469 (37:68)
- [9] A. Nies, R. A. Shore, and T. A. Slaman. Definability in the recursively enumerable degrees. *Bull. Symbolic Logic* **2** (1996), 392–404. MR1460314 (98g:03105)
- [10] S. L. Post, Recursively enumerable sets of positive integers and their decision problems, *Bull. Amer. Math. Soc.*, **50** (1944), 284–316. Reprinted in Davis (1965), 304–337. MR0010514 (6:29f)
- [11] J. R. Shoenfield, Degrees of classes of r.e. sets, *J. Symbolic Logic* **41** (1976), 695–696. MR0485293 (58:5140)
- [12] G. E. Sacks, Recursive enumerability and the jump operator, *Trans. Amer. Math. Soc.* **108** (1963), 223–239. MR0155747 (27:5681)
- [13] R. I. Soare, Automorphisms of the lattice of recursively enumerable sets, Part I: Maximal sets, *Ann. of Math. (2)* **100** (1974), 80–120. MR0360235 (50:12685)
- [14] R. I. Soare, Automorphisms of the lattice of recursively enumerable sets, Part II: Low sets, *Ann. Math. Logic* **22** (1982), 69–107. MR661478 (83k:03048)
- [15] R. I. Soare, *Recursively Enumerable Sets and Degrees: A Study of Computable Functions and Computationally Generated Sets*, Springer-Verlag, Heidelberg, 1987. MR882921 (88m:03003)

DEPARTMENT OF MATHEMATICS, FACULTY OF ARTS AND SCIENCES, HARVARD UNIVERSITY,
1 OXFORD STREET, CAMBRIDGE, MASSACHUSETTS 02138

E-mail address: repstein@math.harvard.edu