

THE METAPLECTIC CASSELMAN-SHALIKA FORMULA

PETER J. MCNAMARA

ABSTRACT. This paper studies spherical Whittaker functions for central extensions of reductive groups over local fields. We follow the development of Chinta and Offen to produce a metaplectic Casselman-Shalika formula for tame covers of all unramified groups.

1. INTRODUCTION

Suppose that G is an unramified reductive group over a nonarchimedean local field F . By definition, this means that G is the generic fibre of a smooth reductive group scheme over the valuation ring O_F , or equivalently that G is quasi-split and splits over an unramified extension of F . The Casselman-Shalika formula is an explicit formula for the spherical Whittaker function that is associated to an unramified principal series representation of $G(F)$.

In this paper, we replace $G(F)$ by a central extension of it by a finite cyclic group, and develop a Casselman-Shalika formula for this so-called metaplectic group. Our main result is the union of Theorems 8.1, 13.1 and 14.1. Theorem 8.1 is the metaplectic Casselman-Shalika formula, computing the metaplectic Whittaker function in terms of a certain Weyl group action. Theorems 13.1 and 14.1 describe how a simple reflection acts in this Weyl group action. Along the way, in Section 12, we prove a metaplectic Gindikin-Karpelevic formula.

Our approach is to follow the technique of Chinta and Offen [CO] who have shown how to generalise the approach of Casselman and Shalika [CS] to provide a formula for a Whittaker function on the metaplectic cover of GL_r . The purpose of this paper is to show how their technique generalises to the more general case of covers of unramified groups.

This paper naturally splits into two parts. In the first part of this paper, we work in the generality of considering any finite cyclic cover of any reductive p -adic group modulo Assumptions 2.5 and 2.8. These assumptions are valid in the most important case when G is simply connected and unramified. For more general G these are more subtle issues and there is some discussion of this in the body of the paper. This part of the paper culminates in the aforementioned Theorem 8.1, and closely follows the approach of Chinta and Offen. The second part begins with Section 9 and develops the necessary extra results to enable one to compute this Whittaker function when the underlying reductive group is unramified.

To conclude, we compare our computation of the metaplectic Whittaker function with the objects appearing in the local part of a Weyl group multiple Dirichlet series constructed by Chinta and Gunnells [CG].

Received by the editors April 23, 2014 and, in revised form, October 15, 2014 and October 19, 2014.

2010 *Mathematics Subject Classification*. Primary 22E50.

Throughout, we work with an assumption that $q \equiv 1 \pmod{2n}$ (where q is the cardinality of the residue field and n is the degree of the central extension) which we feel is worth remarking on. In writing this paper, we have strived to work with the greatest possible range of groups G . In this generality, we are not sure how to proceed at times under what might be considered the more natural assumption that $q \equiv 1 \pmod{n}$. As an indication of the simplification this assumption entails, one may look at the setup of Weissman [W], where the assumption $q \equiv 1 \pmod{2n}$ corresponds to trivialising his second twist χ . However we expect that working only with $q \equiv 1 \pmod{n}$ will ultimately only introduce signs into various formulae we encounter.

We have chosen not to discuss the spherical function as we do not believe we can offer anything novel to say about it. The techniques of Casselman [C] are sufficient to compute the metaplectic spherical function; for example see [CO, §10] for a treatment of the case when $G = GL_n$.

The author would like to thank D. Bump, G. Chinta, G. Hainke, O. Offen, A. Puskas and some anonymous help for useful discussions.

2. THE METAPLECTIC GROUP

Fix a positive integer n . Let F be a non-archimedean local field with valuation ring O_F , uniformiser ϖ and residue field of order q . We assume $q \equiv 1 \pmod{2n}$. Let μ_n denote the group of n -th roots of unity, this is a cyclic group of order n . Fix, once and for all, an embedding $\epsilon : \mu_n \rightarrow \mathbb{C}^\times$.

Let G be a connected reductive algebraic group over F . Let S be a maximal split torus of G and let T be a maximal torus of G containing S . We use the following theorem to find a split reductive subgroup G' of G that will be of use to us.

Theorem 2.1 ([BT, Théorème 7.2]). *Let $\Phi = \Phi(S, G)$ be the root system of G relative to S , and let Φ' be the set of roots a for which $2a$ is not a root. Let Δ be a choice of simple roots in Φ' . For each $a \in \Delta$, let E_a be a one-dimensional subgroup of the root subgroup of a , and let V be the group generated by the E_a . Then G possesses a unique split reductive subgroup G' containing S and V . The torus S is a maximal torus of G' and the root system is $\Phi' = \Phi(G', S)$. In particular, the Weyl groups of G and G' are isomorphic.*

Let \overline{F} be a separable closure of F . We write W for the Weyl group of G and $W_{\overline{F}}$ for the Weyl group of $G_{\overline{F}}$, the base change of G to \overline{F} . Consider the geometric cocharacter lattice $X_*(T_{\overline{F}}) = \text{Hom}_{\overline{F}}(\mathbb{G}_m, T_{\overline{F}})$. This is equipped with actions of both $W_{\overline{F}}$ and the Galois group $\text{Gal}(\overline{F}/F)$.

Brylinski and Deligne [BD, Theorem 7.2] provide a classification of central extensions of G by K_2 as sheaves on the big Zariski site over $\text{Spec}(F)$. Since $H^1(F, K_2) = 0$, at the level of F -points, this gives a central extension of G by $K_2(F)$. We push forward such a central extension by the Hilbert symbol $K_2(F) \rightarrow \mu_n$ to obtain a central extension \tilde{G} of G by μ_n . Explicitly, such a central extension is a short exact sequence of topological groups

$$1 \rightarrow \mu_n \rightarrow \tilde{G} \rightarrow G \rightarrow 1$$

with μ_n lying in the centre of \tilde{G} . We write $p : \tilde{G} \rightarrow G$ for the projection map. On occasion, we shall find it necessary to express \tilde{G} in terms of a 2-cocycle on G . When this is the case, we typically denote the section $G \mapsto \tilde{G}$ by \mathfrak{s} and the chosen

2-cocycle by σ . Note that \mathbf{s} is not a homomorphism, the multiplication in \tilde{G} is given by $\mathbf{s}(g_1g_2) = \mathbf{s}(g_1)\mathbf{s}(g_2)\sigma(g_1, g_2)$.

All representations of \tilde{G} which we consider will be representations where the central μ_n acts by scalars by the embedding ϵ . Such representations are called *genuine*. We often invoke the convention of omitting ϵ from the notation.

The major piece of data classifying central extensions of G by K_2 is a $\text{Gal}(\overline{F}/F)$ and $W_{\overline{F}}$ -invariant quadratic form $Q : \text{Hom}_{\overline{F}}(\mathbb{G}_m, T_{\overline{F}}) \rightarrow \mathbb{Z}$. Let us fix once and for all such a quadratic form Q .

When the group G does not split over F , the metaplectic cover can be constructed via a descent process starting from the split case. We will now give a brief description of the most relevant part of this construction when the group G is unramified. This is the case which is of greatest interest to us. This process is described with more detail in [BD, §12.11]. Let E be a degree d unramified extension of F over which G splits.

We think about the construction of \tilde{G} as a three-step process. First one constructs the central extension of $G(F)$ by $K_2(F)$. Then one pushes forward by the tame symbol $K_2(F) \rightarrow k^\times$ to arrive at an extension of G by k^\times . One finally pushes forward this extension via the operation of raising to the $(q - 1)/n$ -th power to obtain the metaplectic group \tilde{G} .

The Brylinski-Deligne construction shows in this case that \tilde{G} can be realised as a subgroup of a group $\widehat{G}(\overline{E})$ which is a central extension of $G(E)$ by the group of $n \frac{q^d - 1}{q - 1}$ -th roots of unity.

For any subgroup H of G , we denote by \tilde{H} the inverse image of H in \tilde{G} .

Let $B(x, y) = Q(x + y) - Q(x) - Q(y)$ be the bilinear form on $X_*(T_{\overline{F}})$ associated to Q . The commutator map $[\cdot, \cdot] : \tilde{T} \times \tilde{T} \rightarrow \mu_n$ factors through $p \times p : \tilde{T} \times \tilde{T} \rightarrow T \times T$ to give a well-defined map from $T \times T$ to μ_n , also denoted $[\cdot, \cdot]$. It takes the following form [BD, Corollary 3.14]:

$$(2.1) \quad [x^\lambda, y^\mu] = (x, y)^{B(\lambda, \mu)},$$

where $(\cdot, \cdot) : F^\times \times F^\times \rightarrow \mu_n$ is the Hilbert symbol.

There is an explicit formula for the Hilbert symbol in the tame case which we are in, namely

$$(2.2) \quad (x, y) = \left(\overline{(-1)^{v(x)v(y)} \frac{x^{v(y)}}{y^{v(x)}}} \right)^{\frac{q-1}{n}},$$

where the bar means to take the reduction modulo ϖ of an element of O_F . A choice of an embedding $\mathbb{F}_q^\times \rightarrow \mathbb{C}^\times$ is needed to identify the image of (\cdot, \cdot) as lying in μ_n .

Let us now restrict our attention to the central extension \tilde{G}' of G' . There is a natural inclusion of cocharacter groups $X_*(S) \subset X_*(T_{\overline{F}})$. The restriction of Q to $X_*(S)$ is a W -invariant quadratic form and \tilde{G}' is the central extension of G' associated to Q by the Brylinski-Deligne theory.

Since G' is split, we can, and will, make use of the theory developed in [Mc2] where only covers of split groups were considered.

We may identify the Bruhat-Tits building of G' with a subset of the Bruhat-Tits building of G . Pick a hyperspecial point in the apartment corresponding to S , and let \mathbf{G} be the corresponding group scheme over O_F with special fibre G via

Bruhat-Tits theory. We let $K = \mathbf{G}(O_F)$; this is a maximal compact subgroup of G .

We will need to define a lift for any element of W into G . To achieve this, it suffices to work inside the group G' . By a theorem of Tits [T], for each simple reflection $s_\alpha \in W$, we can choose $w_\alpha \in \mathbf{G}'(O_F)$ such that the collection of w_α 's so obtained satisfies the braid relations. Furthermore, if we consider the natural projection from the group generated by the w_α to W , the kernel is an elementary abelian 2-group contained in S . Thus for any $w \in W$, we define a lift by writing $w = s_{\alpha_1} \cdots s_{\alpha_N}$ as a reduced product of simple reflections, and letting the lift be the product $w_{\alpha_1} \cdots w_{\alpha_N}$. Let us denote by W_0 the subgroup of G generated by the w_α .

Let M be the centraliser of S in G . This is a minimal Levi subgroup of G .

Lemma 2.2. *The kernel of the projection from W_0 to W lies in the centre of \widetilde{M} .*

Proof. For a general element $m \in \widetilde{M}$, $p(m)$ is semisimple, so lies in a maximal torus T' of G . The central extension \widetilde{G} is obtainable via a descent process from a central extension of $G(E)$ where E is some Galois extension of F . To check whether m and z commute, it suffices to perform this computation in the central extension of $G(E)$. We choose E so that the torus T' splits. This reduces the problem to the split case.

Now suppose G is split and z is in the aforementioned kernel. Note that $z^2 = 1$. If $m \in \widetilde{S}$, then by the explicit formula (2.1) for the commutator, we compute $[m, z] = 1$, since $(q-1)/n$ is even and hence the Hilbert symbol $(-1, x)$ is always 1. \square

At all stages, unless explicitly mentioned otherwise, we choose normalisations of Haar measures such that the intersection with the maximal compact subgroup K has volume 1.

A subgroup J of G is said to be split in the extension if there is a section $J \mapsto \widetilde{J}$ that is a group homomorphism. When this occurs, we also denote the image of J in \widetilde{G} by J .

Theorem 2.3 ([Mc2, Proposition 4.1]). *Any unipotent subgroup of G has a unique splitting.*

Remark 2.4. In the reference quoted the splitting is said to be canonical but an examination of the proof shows it is unique.

We will make the following assumption, necessary to make sense of the notion of a spherical Whittaker function.

Assumption 2.5. *The subgroup K has a splitting.*

When G is not simply connected, the splitting or otherwise of a Brylinski-Deligne extension over K is subtle. For example [GG, §4.6] gives an example of a Brylinski-Deligne extension of PGL_2 by μ_2 which does not split over $PGL_2(O_F)$. In the same paper a criterion for splitting the subgroup K is given [GG, Corollary 4.2] for all split G . The family of central extensions \widetilde{G} which satisfy the condition of Assumption 2.5 via this criterion contains a large family of interesting central extensions.

We now explain why, when G is simply connected and unramified, making this assumption on the splitting of K does not entail any loss of generality.

If G is split, then Assumption 2.5 is true by [GG, Corollary 4.2]. Now consider any unramified simply connected G . It is possible to push forward the central extension \widetilde{G} by the inclusion $\mu_n \hookrightarrow S^1$ and get a central extension of G by the group of complex numbers of norm 1. Doing so does not in any way change the representation theory of \widetilde{G} . The group \widetilde{K} which is now a central extension of K by S^1 is a subgroup of a split group \widetilde{L} which is the inverse image in the central extension of a maximal compact subgroup after making an unramified base change where G splits. Therefore the group \widetilde{K} splits under the central extension by S^1 , which is sufficient for our representation theoretic purposes.

The splitting of K is not unique in general. We will choose once and for all such a splitting. Since the group W_0 lies in K , this allows us to define a lift of any element $w \in W$ to an element of \widetilde{G} , which we will by an abuse of notation also call w .

We will need to choose a lift of $X_*(S)$ into \widetilde{G} . To achieve this, we first use the uniformiser ϖ to determine a homomorphism from $X_*(S)$ to G , namely $\lambda \mapsto \varpi^\lambda$. Then we have a central extension

$$(2.3) \quad 1 \rightarrow \mu_n \rightarrow \widetilde{X}_*(S) \rightarrow X_*(S) \rightarrow 1$$

which is abelian by the commutator formula (2.1). At this point we are using the assumption that $2n$ divides $q - 1$ which has the consequence that $(\varpi, \varpi) = 1$.

Since $X_*(S)$ is a free abelian group, the short exact sequence (2.3) of abelian groups splits. We choose a splitting and also denote it by $\lambda \mapsto \varpi^\lambda \in \widetilde{G}$.

Now that we have a splitting of K , we define a subgroup H of \widetilde{M} as follows:

$$H = \{h \in \widetilde{M} \mid [h, \eta] \in K \text{ for all } \eta \in \widetilde{M} \cap K\}.$$

If G happens to be quasi-split, then this is simply the centraliser in \widetilde{M} of $M \cap K = \widetilde{M} \cap K$.

Lemma 2.6. *Let $s \in S$. Then $s(s^n) \in H$.*

Proof. Fix $m \in \widetilde{M}$. As S is central in M , the map $\psi_m : S \rightarrow \mu_n$ defined by $\psi_m(s) = [m, s]$ is a group homomorphism, hence trivial on all n -th powers. \square

For any coroot α , define $n_\alpha = n / \gcd(n, Q(\alpha))$.

Lemma 2.7. *Suppose G is unramified and let α be a coroot. Then $\varpi^{n_\alpha \alpha} \in H$.*

Proof. Under these assumptions, M is a maximal torus of G . We have to show that $\varpi^{n_\alpha \alpha}$ commutes with all of \widetilde{M} . To do this, it suffices to pass to an extension field where G is split; then we can use the commutator formula (2.1) to deduce the result. \square

Write Λ' for the lattice $\widetilde{M}/(\mu_n(M \cap K))$ and Λ for the sublattice $H/(\mu_n(M \cap K))$. Lemma 2.6 shows that Λ is a finite index in Λ' . If G is unramified, Λ' is canonically isomorphic to the cocharacter lattice $X_*(S)$.

We will make one more assumption, which will be used in the construction of the principal series representations.

Assumption 2.8. *The quotient group $H/(M \cap K)$ is abelian.*

In the unramified case, the following lemma proves that this assumption is always satisfied.

Lemma 2.9. *If G is unramified, then H is abelian.*

Proof. Since G is unramified, M is abelian and $M = S(M \cap K)$. Now suppose that $h_1, h_2 \in H$. Write $h_i = s_i k_i$ with $s_i \in \tilde{S}$ and $k_i \in M \cap K$. The only nontrivial part is to show that s_1 and s_2 commute. But s_i commutes with $\tilde{S} \cap K$, so by [Mc2, Lemma 5.3], we're done. \square

The following elementary observation will often be used.

Lemma 2.10. *Let A be a compact topological group and $\chi : A \rightarrow \mathbb{C}^\times$ a nontrivial homomorphism. Then $\int_A \chi(a) da = 0$.*

3. UNRAMIFIED REPRESENTATIONS

Let P be a minimal parabolic F -subgroup of G containing S and let U be its unipotent radical. The quotient P/U is canonically isomorphic to the Levi subgroup $M = Z_G(S)$. Thinking of M as a quotient of P makes the constructions we make more canonical, for example $I(\chi)$ will not depend on the choice of a minimal parabolic subgroup up to canonical isomorphism.

Consider the complex algebraic variety

$$Y = \{ \chi \in \text{Hom}(H/(\tilde{M} \cap K), \mathbb{C}^\times) \mid \chi(\zeta) = \epsilon(\zeta) \text{ for all } \zeta \in \mu_n \}.$$

There is a short exact sequence of groups

$$(3.1) \quad 0 \rightarrow \mu_n \rightarrow H/(\tilde{M} \cap K) \rightarrow \Lambda \rightarrow 0,$$

where the middle term is abelian by Assumption 2.8. As Λ is free, this sequence splits, so we obtain a noncanonical isomorphism $Y \cong \text{Spec}(\mathbb{C}[\Lambda])$.

Consider a (complex) point $\chi \in Y$. We define the corresponding unramified representation $(\pi_\chi, i(\chi))$ of \tilde{M} as follows: Given χ , we inflate it to a one-dimensional representation of the subgroup H . We define $i(\chi)$ to be the induction of this representation from H to \tilde{M} . Note that $i(\chi)$ is finite dimensional.

We also construct an unramified principal series representation $I(\chi)$ of \tilde{G} . First we use the canonical surjection $\tilde{P} \rightarrow \tilde{M}$ to consider $i(\chi)$ as a representation of \tilde{P} . The representation $I(\chi)$ is defined to be the induction of $i(\chi)$ from \tilde{P} to \tilde{G} .

Concretely, $I(\chi)$ is the space of all locally constant functions $f : \tilde{G} \rightarrow i(\chi)$ such that

$$f(pg) = \delta^{1/2}(p)\pi_\chi(p)f(g)$$

for all $p \in \tilde{P}$ and $g \in \tilde{G}$ where δ is the modular quasi-character of \tilde{P} . The action of \tilde{G} on $I(\chi)$ is given by right translation.

The group W_0 acts on \tilde{M} by conjugation, and Lemma 2.2 shows that this descends to an action of W on \tilde{M} . As W_0 is a subgroup of K , this induces an action of W on the subgroup H , under which the subgroup $M \cap K$ is stable. Hence we obtain an action of W on Y . If G is unramified, then Λ is a sublattice of $X_*(S)$ which allows us to choose the splitting $\lambda \mapsto \varpi^\lambda$ of (3.1). This choice makes the noncanonical isomorphism $Y \cong \text{Spec}(\mathbb{C}[\Lambda])$ Weyl group invariant.

We now define isomorphisms between the underlying vector spaces of $i(\chi)$ and $i(w\chi)$. As an induced representation, the underlying vector space of $i(\chi)$ is

$$\{ f : \tilde{M} \rightarrow \mathbb{C} \mid f(mh) = \chi(h)f(m) \text{ for all } h \in H, m \in \tilde{M} \}.$$

Our choice of isomorphism $i(\chi) \xrightarrow{\sim} i(w\chi)$ is $f \mapsto {}^w f$, where

$${}^w f(m) = f(w^{-1}mw).$$

These form a transitive system of isomorphisms.

Theorem 3.1. *The map $f \mapsto f(1)$ is an isomorphism between $I(\chi)^K$ and $i(\chi)^{\widetilde{M} \cap K}$. These are both one-dimensional vector spaces.*

Proof. The argument of [Mc2, Lemma 6.3] applies in this case without change. \square

The identification of the spaces $i(\chi)$ and $i(w\chi)$ constructed above can be construed as an action of W on $i(\chi)$. Under this action, the subspace $i(\chi)^{\widetilde{M} \cap K}$ is invariant. Let us pick a nonzero vector v_0 in this subspace. By Theorem 3.1, we choose a spherical vector $\phi_K^{(\chi)} \in I(\chi)^K$ for each χ in a W -orbit in a compatible manner such that $\phi_K^{(w\chi)}(1) = v_0$.

4. INTERTWINING OPERATORS

For each positive root α , there is an associated root subgroup U_α of U with Lie algebra equal to $\text{Lie}(G)_\alpha + \text{Lie}(G)_{2\alpha}$. For any $w \in W$, define U_w to be the group equal to the product

$$\prod_{\alpha > 0, w\alpha < 0} U_\alpha.$$

This is also canonically isomorphic to the left quotient $(U \cap wUw^{-1}) \backslash U$.

The (unnormalised) intertwining operators $T_w : I(\chi) \rightarrow I(w\chi)$ are defined by

$$(T_w f)(g) = \int_{U_w} f(w^{-1}ug) du$$

whenever this is absolutely convergent, and by a standard process of meromorphic continuation in general, for example following [Mc2, §7]. It is a routine calculation to show that T_w does indeed map $I(\chi)$ into $I(w\chi)$ as claimed. We denote by X the open subset of Y on which all the intertwining operators T_w have no poles.

When $w = s$ is a simple reflection, then we freely identify U_s with the intersection of U and the corresponding standard Levi subgroup M_s .

Proposition 4.1. *Suppose that w_1 and w_2 are two elements of W such that $\ell(w_1w_2) = \ell(w_1) + \ell(w_2)$. Then the intertwining operators satisfy the identity $T_{w_1w_2} = T_{w_1}T_{w_2}$.*

Proof. There is a measure preserving bijection from $U_{w_1} \times U_{w_2}$ to $U_{w_1w_2}$ given by $(u_1, u_2) \mapsto w_1u_2w_1^{-1}u_1$. The remainder of the proof is a standard manipulation involving Fubini's theorem. \square

Let B denote the fraction field of the coordinate ring $\mathcal{O}(Y)$.

Theorem 4.2. *There exists a nonzero element $c_w(\chi) \in B$ such that for all $\chi \in Y$,*

$$T_w \phi_K^{(\chi)} = c_w(\chi) \phi_K^{(w\chi)}.$$

Remark 4.3. In Theorem 12.1 we will provide a more precise statement for unramified groups.

Proof. By Proposition 4.1, we may assume without loss of generality that w is a simple reflection s . Also by meromorphic continuation, we may assume without loss of generality that the defining integral for T_s converges.

The function $T_s\phi_K^{(\chi)}$ is guaranteed to be K -invariant, and hence by Theorem 3.1 is a scalar multiple of $\phi_K^{(s\chi)}$. Thus $c_s(\chi)$ exists as a function on the open subset of Y on which the intertwining operator T_s has no poles. To compute $c_s(\chi)$, it suffices to evaluate $(T_s\phi_K^{(\chi)})(1)$.

There is a filtration on U_s induced by a valuation of root datum. This consists of the data of a compact subgroup $U_{s,r}$ of U_s for each $r \in \mathbb{R}$ with the property that $U_{s,r'} \subset U_{s,r}$ if $r' \leq r$. We let $C_r = U_{s,r} \setminus \bigcup_{r' < r} U_{s,r'}$. First we collate some facts about these sets.

If $u \in C_r$ and $r > 0$, then $s^{-1}u \in \mu_n \varpi^{r\alpha} U_s K$. Conjugation by ϖ^α sends C_r to C_{r+2} , and there are only finitely many orbits of nonempty C_r under the action of conjugation by ϖ^α .

There is a decomposition of U_s into the disjoint union

$$U_s = (U_s \cap K) \cup \left(\bigcup_{r>0} C_r \right).$$

We apply this decomposition to the domain of integration in the equation

$$T_s\phi_K^{(\chi)}(1) = \int_{U_s} \phi_K^{(\chi)}(s^{-1}u)du.$$

The integral over $U_s \cap K$ is equal to v_0 . For the integrals over the C_r , we use the fact that $T_s\phi_K^{(\chi)}(1) \in i(s\chi)^{\widetilde{M} \cap K}$ to see that only those r for which $\varpi^{r\alpha} \in H$ can give a nonzero contribution. This provides an expression for $c_s(\chi)$ of the form

$$c_s(\chi) = 1 + \sum_{r>0, \varpi^{r\alpha} \in H} d_r \chi(\varpi^{r\alpha})$$

for some constants d_r .

Note that by Lemma 2.6, $\varpi^{n\alpha} \in H$. In passing from the integral over C_r to that over C_{r+2n} via conjugation by $\varpi^{n\alpha}$, the integrand has been multiplied by $(\delta^{1/2}\chi)(\varpi^{2n\alpha})$ while the change of coordinates contributes a factor of $\delta(\varpi^{n\alpha})^{-1}$. Hence one integral is $\chi(\varpi^{2n\alpha})$ times the other. Thus our expression for $c_s(\chi)$ is actually a sum of geometric series, so is an element of B as required. \square

In light of the above result, we now define a renormalised version of the intertwining operators. Let $\overline{T}_w = c_w(\chi)^{-1}T_w$. The upshot is that we have the equation

$$(4.1) \quad \overline{T}_w\phi_K^{(\chi)} = \phi_K^{(w\chi)}$$

as well as the following proposition.

Proposition 4.4. *For any $w_1, w_2 \in W$, we have*

$$(4.2) \quad \overline{T}_{w_1 w_2} = \overline{T}_{w_1} \overline{T}_{w_2}.$$

Proof. The lattices Λ and $X_*(S)$ are commensurable, so the set of characters χ with trivial stabiliser under the W -action is dense in Y . Hence it suffices to prove the proposition for such χ . As in [Mc2, Theorem 7.1], we use [BZ, Theorem 5.2] to conclude that $\dim \text{Hom}_{\overline{G}}(I(\chi), I(w\chi)) = 1$. The proposition now follows from Theorem 4.2. \square

5. IWAHORI INVARIANTS

The structure and results of this section closely follow [CO, §3.3]. Let I be the Iwahori subgroup of G , maximal with respect to intersection with P among all Iwahori subgroups contained in K .

Proposition 5.1. *The dimension of the space of vectors in an unramified principal series representation $I(\chi)$ that are invariant under the Iwahori subgroup I is equal to $|W|$.*

Proof. One has $\tilde{P}\backslash\tilde{G}/I \cong W$. Thus $\dim(I(\chi)^I) \leq |W|$.

For any $w \in W$, we can define $\phi_w \in I(\chi)^I$ by $\phi_w(g) = \phi_K(g)$ if $g \in \tilde{P}wI$ and $\phi_w(g) = 0$ otherwise. These functions are obviously linearly independent so the proposition is proved. \square

Let w_0 denote the longest element in W . Let $\phi_{w_0}^{(\chi)}$ be the function in $I(\chi)$ supported on $\tilde{P}w_0I$ and taking the value v_0 at w_0 .

Proposition 5.2 ([CO, Lemma 5]). *A basis for the space $I(\chi)^I$ can be given by the elements $T_w\phi_{w_0}^{(w^{-1}\chi)}$ as w ranges over W .*

Proof. Let us write f_w for the function $T_w\phi_{w_0}^{(w^{-1}\chi)}$. Suppose that $f_w(v) \neq 0$ for some $v \in W$. By definition,

$$f_w(v) = \int_{U_w} \phi_{w_0}^{(w^{-1}\chi)}(w^{-1}uv)du.$$

For this to be nonzero, it is necessary that $w^{-1}Uv \cap \tilde{P}w_0I \neq \emptyset$. Since the group I admits an Iwahori factorisation, we have the containment $\tilde{P}w_0I \subset \tilde{P}w_0\tilde{P}$. Now $w^{-1}Uv \subset Pw^{-1}PvP$ and since these are double cosets in a Tits system, this intersects Pw_0P nontrivially only if $\ell(w^{-1}) + \ell(v) \geq \ell(w_0)$. Furthermore, if there is equality $\ell(w^{-1}) + \ell(v) = \ell(w_0)$, the intersection is nonempty if and only if $w^{-1}v = w_0$.

From the above considerations, it suffices to show that $f_w(ww_0) \neq 0$. Since the support of ϕ_{w_0} is $\tilde{P}w_0I$, we need to understand when $w^{-1}uww_0 \in Pw_0I$ with $u \in U_w$. We rewrite this as $w^{-1}uw \in P \cdot w_0Iw_0^{-1}$. As $w^{-1}uw \in U$ and the group $w_0Iw_0^{-1}$ admits an Iwahori factorisation, this only happens when $w^{-1}uw \in (w_0Iw_0^{-1}) \cap U^-$. For such u , $w^{-1}uww_0 \in w_0I$ and hence $\phi_{w_0}^{(w^{-1}\chi)}(w^{-1}uww_0) = v_0$.

Therefore the above integral for $f_w(ww_0)$ reduces to an integral of v_0 over a compact subset of U_w with positive measure, hence the integral is nonzero, as required. \square

Proposition 5.3. *The spherical function ϕ_K can be expanded as*

$$\phi_K = \sum_{w \in W} c_{w_0}(w^{-1}\chi)\overline{T}_w\phi_{w_0}^{(w^{-1}\chi)}.$$

Proof. By the previous proposition, there exist $d_w(\chi)$ such that

$$(5.1) \quad \phi_K^{(\chi)} = \sum_{w \in W} d_w(\chi)\overline{T}_w\phi_{w_0}^{(w^{-1}\chi)}.$$

Let us apply \overline{T}_u to this equation. Via (4.1) and (4.2), we arrive at

$$\sum_{w \in W} d_w(u\chi)\overline{T}_w\phi_{w_0}^{(w^{-1}u\chi)} = \sum_{w \in W} d_w(\chi)\overline{T}_{uw}\phi_{w_0}^{(w^{-1}u\chi)}.$$

Again using the fact that we have a basis of $I(u\chi)^I$, we compare coefficients to obtain $d_w(\chi) = d_{uw}(u\chi)$. Thus it suffices to establish the value of $d_{w_0}(\chi)$. We now evaluate (5.1) at the identity:

$$\phi_K^{(\chi)}(1) = \sum_{w \in W} d_w(\chi) c_{w_0}(w\chi)^{-1} \int_{U_w} \phi_{w_0}^{(w^{-1}\chi)}(w^{-1}u) du.$$

As in the proof of Proposition 5.2, $w^{-1}u \in Pw_0I$ if and only if $w = w_0$ and $u \in U \cap I$. Thus only one term survives in this sum, and the surviving integral is the integral of v_0 over a set of measure one. We end up with $d_{w_0}(\chi) = c_{w_0}(w_0^{-1}\chi)$ which implies the result. □

6. WHITTAKER FUNCTIONALS

A group homomorphism $\psi : U \rightarrow \mathbb{C}^\times$ is said to be an unramified character if for each simple reflection $s \in W$, the restriction of ψ to the intersection $U_s = U \cap M_s$ with the corresponding Levi subgroup M_s is trivial on $U_s \cap K$ and nontrivial on all open subgroups of U_s with a larger abelianisation than $U_s \cap K$. We fix one such unramified character ψ . By abuse of notation we also use ψ to denote the corresponding group homomorphism from U^- to \mathbb{C}^\times obtained by precomposing ψ with conjugation by w_0 .

Definition 6.1. A Whittaker functional on a representation (π, V) of \tilde{G} is a linear functional W on V such that $W(\pi(u)v) = \psi(u)v$ for all $u \in U$ and $v \in V$.

Theorem 6.2. *There is an isomorphism between $i(\chi)^*$ and the space of Whittaker functionals on $I(\chi)$ given by $\lambda \mapsto W_\lambda$ with*

$$W_\lambda(\phi) = \lambda \left(\int_{U^-} \phi(uw_0)\psi(u) du \right).$$

Proof. This follows from [BZ, Theorem 5.2] and the Bruhat decomposition. □

Define $W^{(\chi)} : I(\chi) \rightarrow i(\chi)$ by

$$W^{(\chi)}(\phi) = \int_{U^-} \phi(uw_0)\psi(u) du.$$

We use $R(g)$ to denote the action of $g \in \tilde{G}$ on a function by right translation. A group element $t \in \tilde{M}$ is said to be dominant if $t(U \cap K)t^{-1} \subset K$.

Lemma 6.3 ([CO, Lemma 7]). *Let $t \in \tilde{M}$. Then*

$$W^{(\chi)}(R(t)\phi_{w_0}^{(\chi)}) = \begin{cases} \delta^{1/2}(t)\pi_\chi(w_0tw_0^{-1})v_0 & \text{if } t \text{ is dominant,} \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Let $t^* = w_0tw_0^{-1}$ and $u' = (t^*)^{-1}ut^* \in U^-$. Since the group I admits an Iwahori factorisation, we have $u'w_0 \in \tilde{P}w_0I$ if and only if $u' \in U^- \cap w_0Iw_0^{-1}$. Therefore

$$\phi_{w_0}^{(\chi)}(uw_0t) = \phi_{w_0}^{(\chi)}(t^*u'w_0) = \delta^{1/2}(t^*)\pi_\chi(t^*)\phi_{w_0}^{(\chi)}(u'w_0).$$

Since the support of $\phi_{w_0}^{(\chi)}$ is $\tilde{P}w_0I$, our above remark shows that this expression vanishes unless $u' \in U^- \cap w_0Iw_0^{-1}$, which implies $u'w_0 \in w_0I$. Since $\phi_{w_0}^{(\chi)}$ is I -invariant on the right, $\phi_{w_0}^{(\chi)}(u'w_0) = \phi_{w_0}^{(\chi)}(w_0)$ under these circumstances. Therefore

$$\phi_{w_0}^{(\chi)}(uw_0t) = \begin{cases} \delta^{1/2}(t^*)\pi_\chi(t^*)v_0 & \text{if } u' \in U^- \cap w_0Iw_0^{-1}, \\ 0 & \text{otherwise.} \end{cases}$$

Now we can compute

$$\begin{aligned} W^{(\chi)}(R(t)\phi_{w_0}^{(\chi)}) &= \int_{U^-} \phi_{w_0}^{(\chi)}(uw_0t)\psi(u)du \\ &= \delta^{1/2}(t^*)\pi_\chi(t^*)v_0 \int_{t^*(U^- \cap K)(t^*)^{-1}} \psi(u)du. \end{aligned}$$

This final integral vanishes unless ψ is identically 1 on the subgroup being integrated over, which happens exactly when t is dominant, by our assumption that ψ is unramified. In this case, we get a factor of the volume of the group appearing, which is $\delta(t) = \delta^{-1}(t^*)$. This completes the proof. \square

7. CONSTRUCTING THE CHINTA-GUNNELLS ACTION

Let \mathcal{E} be the holomorphic vector bundle on Y whose fibre over a point $\chi \in Y$ is canonically isomorphic to $i(\chi)$. Its dual \mathcal{E}^* is another holomorphic vector bundle on Y whose fibre over a point χ is canonically isomorphic to $i(\chi)^*$. Let \mathcal{F} be the holomorphic vector bundle on Y whose fibre over a point χ is canonically isomorphic to the space of Whittaker functionals on $I(\chi)$.

Theorem 6.2 provides us with an isomorphism between \mathcal{E}^* and \mathcal{F} .

Let V be the open dense subset of Y where the rational functions $c_w(\chi)$ have neither zeroes nor poles.

For any $\chi \in V$, the space $\text{End}(I(\chi))$ is one dimensional by [Mc2, Theorem 7.1(2)], which is valid in this more general context with the same proof. Hence this endomorphism space consists only of scalars. The map \overline{T}_s^2 is an endomorphism of $I(\chi)$ sending ϕ_K to itself, hence must be the identity.

This fact allows us to define an action of the Weyl group W on the total space of $\mathcal{F}|_V$. For $w \in W$ and \mathcal{W} a point of the total space of $\mathcal{F}|_V$, let

$$w \cdot \mathcal{W} = \mathcal{W} \circ \overline{T}_w.$$

If \mathcal{W} is a Whittaker functional on $I(\chi)$, then the composite $\mathcal{W} \circ \overline{T}_w$ is easily seen to be a Whittaker functional on $I(w^{-1}\chi)$.

Via the isomorphism from \mathcal{F} to \mathcal{E}^* , we transport the W -action on the total space of $\mathcal{F}|_V$ to a W -action on the total space of $\mathcal{E}^*|_V$. This is not related to any more naive action on \mathcal{E}^* . From our point of view, the Chinta-Gunnells action will be the induced W -action on the space of algebraic global sections $\Gamma(V, \mathcal{E}^*)$. We will explain in the final section of this paper why this agrees with the action constructed in [CG].

Note that the fact that the W -action is algebraic has not yet been justified. We postpone the verification of this fact until the end of Section 9.

8. FORMAL COMPUTATION OF THE WHITTAKER FUNCTION

Our aim is to compute the general spherical Whittaker function

$$\mathcal{W}_\chi(g) = W^{(\chi)}(R(g)\phi_K).$$

This is an $i(\chi)$ -valued function on \widetilde{G} satisfying

$$\mathcal{W}_\chi(\zeta u g k) = \zeta \psi(u) \mathcal{W}_\chi(g)$$

for all $\zeta \in \mu_n$, $u \in U$, $g \in \widetilde{G}$ and $k \in K$. The Iwasawa decomposition takes the form $G = UMK$, so in order to compute \mathcal{W}_χ , it suffices to know the values taken by \mathcal{W}_χ on \widetilde{M} . This is what we shall concentrate our efforts on.

Theorem 8.1. *Suppose $\lambda \in i(\chi)^*$ and $t \in \widetilde{M}$. The Whittaker function $\mathcal{W}_\chi(t)$ vanishes unless t is dominant. If t is dominant, then we have*

$$\lambda(\mathcal{W}_\chi(t)) = \delta^{1/2}(t) \sum_{w \in W} c_{w_0}(w^{-1}\chi)(w \cdot \lambda)(\pi_{w^{-1}\chi}(w_0 t w_0^{-1})v_0).$$

Proof. Using Lemma 5.3, the fact that T_w is an intertwining operator and the definition of the W -action, we compute

$$\begin{aligned} \lambda(\mathcal{W}_\chi(t)) &= \lambda(W^{(\chi)}R(t)\phi_K) \\ &= \sum_{w \in W} c_{w_0}(w^{-1}\chi)\lambda(W^{(\chi)}(R(t)\overline{T}_w(\phi_{w_0}^{(w^{-1}\chi)}))) \\ &= \sum_{w \in W} c_{w_0}(w^{-1}\chi)\lambda(W^{(\chi)}(\overline{T}_w(R(t)\phi_{w_0}^{(w^{-1}\chi)}))) \\ &= \sum_{w \in W} c_{w_0}(w^{-1}\chi)(w \cdot \lambda)(W^{(w^{-1}\chi)}(R(t)\phi_{w_0}^{(w^{-1}\chi)})) \end{aligned}$$

and the result now follows from Lemma 6.3. □

9. SETTING UP THE EXPLICIT COMPUTATION

Let Γ be a set of left coset representatives for H in \widetilde{M} . A particularly nice, but still noncanonical, choice can be made whenever G is unramified and this is carried out when G is split in Section 15. The vectors $\{\pi_\chi(a)v_0\}_{a \in \Gamma}$ form a basis of $i(\chi)$. Incidentally, this yields a trivialisation of the vector bundle \mathcal{E} .

Let $\lambda_a^{(\chi)}$ denote the corresponding dual basis of $i(\chi)^*$, and let $W_a^{(\chi)}$ be the Whittaker functional corresponding to $\lambda_a^{(\chi)}$ under the bijection of Theorem 6.2.

We now introduce the change of basis coefficients as in [KP, §I.3].

As has already been mentioned, for any $a \in \Gamma$ and any $w \in W$, the composition $W_a^{(w\chi)} \circ \overline{T}_w$ is a Whittaker functional on $I(\chi)$. We expand it in the basis $\{W_b^{(\chi)}\}_{b \in \Gamma}$, defining coefficients $\tau_{a,b}^{(w,\chi)}$ by

$$(9.1) \quad W_a^{(w\chi)} \circ \overline{T}_w = \sum_{b \in \Gamma} \tau_{a,b}^{(w,\chi)} W_b^{(\chi)}.$$

Any explicit computation of metaplectic Whittaker functions reduces via Theorem 8.1 to computing the Weyl group action, and hence these coefficients.

Let K_1 be an open compact subgroup of G normalised by W and admitting an Iwahori factorisation with respect to P . For each $b \in \widetilde{M}$, define $f_b \in I(\chi)^{K_1}$ to be a function in this space supported on $\widetilde{P}w_0K_1$ and taking the value $\pi_\chi(b)v_0$ at w_0 .

Lemma 9.1. *We have $W_a^{(\chi)}(f_b) = 0$ unless $aH = bH$, in which case $W_a^{(\chi)}(f_a) = |U^- \cap K_1|$.*

Proof. Suppose that $u \in U^-$ and $f_b(uw_0) \neq 0$. The support of f_b is $\widetilde{P}w_0K_1$ and since K_1 is invariant under conjugation by w_0 this implies that $u \in \widetilde{P}K_1$. As K_1 has the Iwahori factorisation $K_1 = (\widetilde{P} \cap K_1)(U^- \cap K_1)$, we have $u \in \widetilde{P}(U^- \cap K_1)$. As $U^- \cap \widetilde{P} = \{1\}$, this forces $u \in U^- \cap K_1$.

Therefore

$$W_a^{(\chi)}(f_b) = \lambda_a^{(\chi)} \left(\int_{U^-} f_b(uw_0)\psi(u)du \right) = \lambda_a^{(\chi)} \left(\int_{U^- \cap K_1} \pi_\chi(b)v_0\psi(u)du \right).$$

By the definition of $\lambda_a^{(\chi)}$, this is zero unless $aH = bH$ and when $a = b$, we are integrating the constant 1 over the group $U \cap K_1$ so we get the measure $|U \cap K_1|$ as the answer. □

The following corollary is immediate.

Corollary 9.2.

$$\tau_{a,b}^{(w,\chi)} = \frac{(W_a^{(w\chi)} \circ \overline{T}_w)(f_b)}{|U^- \cap K_1|}.$$

Now let us restrict our attention to the case where $w = s = s_\alpha$, a simple reflection corresponding to the simple coroot α of G' . From our definition, the above corollary is equivalent to

$$(9.2) \quad \tau_{a,b}^{(s,\chi)} = \frac{1}{c_s(\chi)|U^- \cap K_1|} \lambda_a^{(s\chi)} \left(\int_{U^-} \int_{U_s} f_b(s^{-1}nuw_0)dn\psi(u)du \right).$$

Lemma 9.3. *Suppose that $n \in U_s$ is not equal to the identity, and $u \in U^-$. Then there is a unique $n' \in U_s^-$ such that $p(n) := s^{-1}nn' \in \widetilde{P}$. Furthermore, $s^{-1}nuw_0 \in \widetilde{P}w_0K_1$ if and only if $u = n'u'$ with $u' \in U^- \cap K_1$.*

Proof. The first claim is an immediate consequence of the Bruhat decomposition in the rank one group M_s . For the latter claim, write $u = n'u'$. Then $s^{-1}nuw_0 \in \widetilde{P}w_0K_1$ if and only if $p(n)u' \in \widetilde{P}K_1$ and this set is equal to $\widetilde{P}(U^- \cap K_1)$ since K_1 admits an Iwasawa decomposition. This finishes the proof. □

As an immediate consequence of this lemma, we can simplify (9.2) to leave ourselves with an expression for the coefficient $\tau_{a,b}^{(s,\chi)}$ with only one integral, namely

$$(9.3) \quad \tau_{a,b}^{(s,\chi)} = c_s(\chi)^{-1} \int_{U_s} \lambda_a^{(s\chi)} f_b(p(n)w_0)\psi(n')dn.$$

Since the subset $\{1\} \subset U_s$ where the integrand is not defined is of measure zero, it may safely be ignored.

Proposition 9.4. *The coefficient $\tau_{a,b}^{(w,\chi)}$ is a rational function on Y .*

Proof. Since the simple reflections generate W , it suffices to consider the case where $w = s$ is a simple reflection. Now we may use (9.3) and our strategy is to argue in an analogous manner to the proof of Theorem 4.2.

We use the same decomposition

$$U_s = (U_s \cap K) \cup \left(\bigcup_{r>0} C_r \right)$$

of the domain of integration in (9.3) induced by the valuation of root datum.

If $n \in C_r$ with $r > 0$, then $n' \in K$ and hence $\psi^{-1}(n') = 1$. Furthermore $p(\varpi^{-n\alpha}n\varpi^{n\alpha}) = \varpi^{2n\alpha}p(n) \pmod{U}$. Thus in passing from C_r to C_{r+2n} , the integrand in (9.3) is multiplied by $(\delta^{1/2}\chi)(\varpi^{2n\alpha})$ while the volume of C_{r+2n} is $\delta^{-1}(\varpi^{n\alpha})$ times that of C_r .

Therefore the integral over C_{r+2n} is equal to $\chi(\varpi^{2n\alpha})$ times the integral over C_r . As in the proof of Theorem 4.2, computing the integral (9.3) reduces to adding an integral over a compact piece times a geometric series. The compact piece is dealt with in exactly the same manner as before, completing the proof. \square

10. DIGRESSION ON SU_3

From now until the end of the paper we will assume that G is an unramified group. For any simple reflection s , let G_s be the simply connected cover of the derived group of M_s . The group G_s is a simply-connected, semisimple unramified group of rank one, and such groups are completely classified. There are two possibilities, either G_s is isomorphic to $\text{Res}_{E/F} SL_2$ for an unramified extension E of F , or is isomorphic to $\text{Res}_{E/F} SU_3$, where the special unitary group SU_3 over E is defined in terms of an unramified quadratic extension L of E , which again is unramified over F .

Of these two possibilities, the group SL_2 will be familiar to most readers. We pause to collate some facts about the less well-known SU_3 that will prove to be of use later on.

We use \bar{z} to denote the image of z under the nontrivial element of $\text{Gal}(L/E)$. The special unitary group $SU_3(E)$ is defined to be the subgroup of $SL_3(L)$ preserving the Hermitian form $x_1\bar{y}_3 + x_2\bar{y}_2 + x_3\bar{y}_1$. Explicitly, if J is the matrix with ones on the off-diagonal and zeroes elsewhere, then $X \in SL_3(L)$ is in $SU_3(E)$ if and only if ${}^t\bar{X}JX = J$.

These coordinates are chosen such that the intersection of $SU_3(E)$ with the set of upper-triangular matrices constitutes a Borel subgroup. Its unipotent radical consists of all matrices of the form

$$(10.1) \quad u = \begin{pmatrix} 1 & x & y \\ 0 & 1 & -\bar{x} \\ 0 & 0 & 1 \end{pmatrix},$$

where x and y are elements of L with $x\bar{x} + y + \bar{y} = 0$.

We take our maximal compact subgroup K to be the subgroup consisting of all matrices in $SU_3(E)$ with entries in O_L .

Let v denote the valuation on E . For any $r \in \mathbb{R}$, the set of $u \in U$ with $v(y) \geq r$ forms a subgroup of U . (This is the filtration induced by a valuation of root datum

in Bruhat-Tits theory introduced in the proof of Theorem 4.2.) Let us denote this subgroup by U_r .

The following equation is fundamental, and explicitly realises the first part of Lemma 9.3 in $SU_3(E)$:

$$(10.2) \quad \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 & x & y \\ 0 & 1 & -\bar{x} \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1/\bar{y} & x/y & 1 \\ 0 & \bar{y}/y & \bar{x} \\ 0 & 0 & y \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ \bar{x}/\bar{y} & 1 & 0 \\ \bar{y}^{-1} & -x/y & 1 \end{pmatrix}^{-1}.$$

In the next section we will see how to lift this to an equation in the metaplectic cover.

Let α be the positive generator of $X_*(S)$. We can, and do, write $\alpha = \alpha_1 + \alpha_2$, where α_1 and α_2 are simple coroots of SL_3 .

11. THE DESCENT PROCESS

Since we are only working with unramified groups, Galois descent for an unramified extension of local fields shall be the only descent process that shall concern us.

We will first consider how descent behaves under restriction of scalars for semi-simple groups. Let E be an unramified field extension of F , \mathbf{G}' a reductive group over F and $\mathbf{G} = \text{Res}_{E/F} \mathbf{G}'$.

Let Γ be the Galois group $\text{Gal}(E/F)$. Let T' be a maximal torus of \mathbf{G}' and $T = \text{Res}_{E/F}(T')$, which is a maximal torus of \mathbf{G} . As Galois modules we have $X_*(T) = X_*(T') \otimes \mathbb{Z}[\Gamma]$.

Consider $T'(E) = T(F) \hookrightarrow T(E)$. At the level of cocharacter lattices this corresponds to the diagonal embedding $X_*(T') \hookrightarrow X_*(T') \otimes \mathbb{Z}[\Gamma] = X_*(T)$, namely $y \mapsto \sum_{\gamma \in \Gamma} y \otimes \gamma$. Let us write Q' for the restriction of Q to the image of $X_*(T')$ in $X_*(T)$.

The precise claim about the behaviour of the central extension under descent in this situation is the following:

Proposition 11.1 ([BD, Proposition 12.9]). *The central extension of $\mathbf{G}(F)$ by μ_n associated with the quadratic form Q and the central extension of $\mathbf{G}'(E)$ by μ_n associated with the quadratic form Q' are isomorphic.*

The other descent calculation we need to study in detail is descent from SL_3 to SU_3 , which will enable us to perform computations in the metaplectic cover of SU_3 . Our strategy is to realise $\widetilde{SU_3(E)}$ as a subgroup of the $n(q+1)$ -fold cover of $SL_3(L)$, with the same quadratic form characterising the extension in each case. There is an explicit cocycle for the cover of $SL_3(L)$ given to us by Banks, Levi and Sepanski. Their result [BLS, Theorem 7], together with the equations appearing in its proof provide an algorithmic method to multiply in $\widetilde{SL_3(L)}$. It provides us with a section \mathbf{s} and a 2-cocycle σ for which multiplication is given by $\mathbf{s}(g_1g_2) = s(g_1)s(g_2)\sigma(g_1, g_2)$. We caution the reader that upon restriction of \mathbf{s} to $SU_3(E)$, the image does not lie in the n -fold cover $\widetilde{SU_3(E)}$.

Our strategy for circumventing this problem to find explicit elements of $\widetilde{SU_3(E)}$ is to use Theorem 2.3 which states that all unipotent subgroups are uniquely split in central extensions. We will use this fact to lift the identity (10.2) into an identity in the metaplectic cover.

By construction the section \mathfrak{s} canonically splits the group U of upper-triangular unipotent matrices in SL_3 . Hence, the splitting of the lower-triangular unipotent subgroup U^- must be given by $u \mapsto \mathfrak{s}(w_0)\mathfrak{s}(w_0uw_0)\mathfrak{s}(w_0)$, where $w_0 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$.

Let

$$n_1 = \begin{pmatrix} 1 & x & y \\ 0 & 1 & -\bar{x} \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad n_2 = \begin{pmatrix} 1 & -x/y & \bar{y}^{-1} \\ 0 & 1 & \bar{x}/\bar{y} \\ 0 & 0 & 1 \end{pmatrix}.$$

Using the results of Banks, Levi and Sepanski referenced above, it may be computed that $\sigma(n_1w_0n_2, w_0) = (x, \bar{y}/y)^{Q(\alpha)}$ and $\sigma(w_0, p(n)) = (y, \bar{y})^{Q(\alpha)}$. Thus we have

$$\mathfrak{s}(w_0)\mathfrak{s}(n_1)\mathfrak{s}(w_0)\mathfrak{s}(n_2)\mathfrak{s}(w_0) = (x, y/\bar{y})^{Q(\alpha)}(y, \bar{y})^{Q(\alpha)}\mathfrak{s}(p(n)).$$

There is a subtlety that needs to be taken care of. According to our construction in Section 2, the choice of representative for the simple reflection s is not the same as the element w_0 we have been using so far in this computation. Instead, it differs by a factor of θ^α , where $\theta \in L$ is such that $|\theta| = 1$ and $\theta + \bar{\theta} = 0$.

Now let us write $y = a\theta\varpi^m$. Then after one more (simpler) cocycle computation, we find that

$$(11.1) \quad s^{-1}nn' = (x, y/\bar{y})^{Q(\alpha)}(y, \bar{y})^{Q(\alpha)}(a, \varpi)^{-mQ(\alpha)}a^\alpha\varpi^{m\alpha} \pmod{U}.$$

12. GINDIKIN-KARPELEVIC FORMULA

Recall that for a simple reflection s , G_s is defined to be the simply connected cover of the derived group of the corresponding Levi subgroup M_s . Let q denote the cardinality of the residue field of E , where E is as in the classification of G_s . For a simple coroot α , let $x_\alpha^{n_\alpha} = \chi(\varpi^{n_\alpha\alpha})$, $n_\alpha = n/\text{gcd}(n, Q(\alpha))$ and $\epsilon = (-1)^{n_\alpha}$. We have the following refinement of Theorem 4.2. Since $c_{w_1w_2}(\chi) = c_{w_1}(w_2\chi)c_{w_2}(\chi)$ whenever $\ell(w_1w_2) = \ell(w_1) + \ell(w_2)$, this is enough to evaluate $c_w(\chi)$ for all Weyl group elements w .

Theorem 12.1. *Suppose that G is unramified, and the simple reflection s corresponds to the simple coroot α . Then*

$$c_s(\chi) = \begin{cases} \frac{1 - q^{-1}x_\alpha^{n_\alpha}}{1 - x_\alpha^{n_\alpha}} & \text{if } G_s \cong SL_2(E), \\ \frac{(1 + \epsilon q^{-1}x_\alpha^{n_\alpha})(1 - \epsilon q^{-2}x_\alpha^{n_\alpha})}{1 - x_\alpha^{2n_\alpha}} & \text{if } G_s \cong SU_3(E). \end{cases}$$

Proof. The author has already written a proof in the case where $G_s \cong SL_2(E)$ [Mc1, Theorem 6.4], so we will not repeat the argument here. When $G_s \cong SU_3(E)$, we present the proof as a warm up for the more challenging computation of $\tau_{a,b}^{(s,\chi)}$ that will appear in Section 14. We follow the strategy from the proof of Theorem 4.2. Hence we have to evaluate the integral in the formula

$$(12.1) \quad c_s(\chi)v_0 = \int_{U_s} \phi_K(s^{-1}u)du.$$

The integral over $U_s \cap K$ is trivially equal to v_0 . We focus our attention on the remainder of the integral.

As in (10.1), we will write

$$(12.2) \quad u = \begin{pmatrix} 1 & x & y \\ 0 & 1 & -\bar{x} \\ 0 & 0 & 1 \end{pmatrix},$$

where x and y are elements of L with $x\bar{x} + y + \bar{y} = 0$. Note that this last equation implies that $y \in O_L$ if and only if $u \in K$.

Let $-v(y) = m$. First let us suppose that $2v(x) = v(y)$. Then m is even; let $k = m/2$. Recall $y = a\theta\varpi^{-m}$. Then using the formula (2.2) for an unramified Hilbert symbol, the product $(x, y/\bar{y})(y, \bar{y})(a, \varpi)^{-m}$ is equal to

$$\left(\frac{\bar{a}}{a}\right)^k a^m \bar{a}^{-m} a^{-m} = (a\bar{a})^{-k},$$

where by abuse of notation take the reductions modulo ϖ of all a 's and \bar{a} 's in the above equation, and include $\mathbb{F}_{q^2}^\times$ into \mathbb{C}^\times .

If instead $2v(x) \neq v(y)$, then $y/\bar{y} \equiv -1 \pmod{\varpi}$, so under these circumstances, $(x, y/\bar{y}) = 1$, using $q \equiv 1 \pmod{2n}$. Thus the root of unity $(x, y/\bar{y})(y, \bar{y})(a, \varpi)^{-m}$ is equal to $(\varpi, a)^m$.

Therefore, for all $u \in C_m$, we have $\phi_K(s^{-1}u) = (a\bar{a}, \varpi)^{mQ(\alpha)/2} \phi_K(\varpi^{m\alpha})$. So (12.1) becomes

$$c_s(\chi) = v_0 + \sum_{m=1}^\infty \int_{C_m} (a\bar{a}, \varpi)^{mQ(\alpha)/2} \phi_K(\varpi^{m\alpha}).$$

From this equation it is evident that the expression does not depend on the ambient group G , only on G_s . If we choose to take $G = SU_5$, then there is a coroot β such that $B(\alpha, \beta) = Q(\alpha)$. By (2.1), the only powers of ϖ^α which lie in H are powers of ϖ^{n_α} . So all terms in the above sum where m is not divisible by n_α must be zero.

First consider the case when n_α is odd. Then n_α divides $2k$ if and only if it divides k . For such $m = 2k$, we have $(a\bar{a}, \varpi)^{mQ(\alpha)/2} = (a\bar{a}, \varpi)^{kQ(\alpha)}$ and this exponent is divisible by n , hence the root of unity is 1. Therefore the contribution to (12.1) from the terms where m is even is

$$\begin{aligned} \sum_{l=1}^\infty \text{vol}(C_{2ln_\alpha})(q^{-2}x_\alpha)^{2ln_\alpha} v_0 &= \sum_{l=1}^\infty q^{4ln_\alpha} (1 - q^{-3})(q^{-2}x_\alpha)^{2ln_\alpha} v_0 \\ &= (1 - q^{-3}) \frac{x_\alpha^{2n_\alpha}}{1 - x_\alpha^{2n_\alpha}} v_0. \end{aligned}$$

When m is odd and divisible by n_α , write $m = (2l+1)n_\alpha$. Necessarily $v(y) \neq 2v(x)$ so the root of unity is simply $(\varpi, a)^{mQ(\alpha)}$, again the exponent is divisible by n and so the root of unity is 1. So the contribution to (12.1) from the terms where m is odd is

$$\begin{aligned} \sum_{l=0}^\infty \text{vol}(C_{(2l+1)n_\alpha})(q^{-2}x_\alpha)^{(2l+1)n_\alpha} v_0 &= \sum_{l=0}^\infty (q-1)q^{(2l+1)n_\alpha-2} (q^{-2}x_\alpha)^{(2l+1)n_\alpha} v_0 \\ &= q^{-2}(q-1) \frac{x_\alpha^{n_\alpha}}{1 - x_\alpha^{2n_\alpha}} v_0. \end{aligned}$$

This completes the proof when n_α is odd. Now let us assume that n_α is even.

As before, we only need to consider the integral over C_m where m is divisible by n_α . This implies that m is even. Let $m = 2k$.

If n_α divides k , then $(a\bar{a}, \varpi)^{kQ(\alpha)} = 1$. The contribution to (12.1) from such terms is again

$$(1 - q^{-3}) \frac{x_\alpha^{2n_\alpha}}{1 - x_\alpha^{2n_\alpha}} v_0.$$

Now suppose that $m = 2k$ is such that n_α divides m but does not divide $2k$.

The fibres of the map from C_m to \mathbb{F}_q^\times given by $u \mapsto a \pmod{\varpi}$ do not all have the same size. In particular the volume of the fibre over a point not in \mathbb{F}_q is $q + 1$ times the volume of the fibre over a point in \mathbb{F}_q .

Note that for $a \in \mathbb{F}_q^\times$, we have $(a\bar{a}, \varpi)^{kQ(\alpha)} = 1$ while on the larger group $\mathbb{F}_{q^2}^\times$, the homomorphism $a \mapsto (a\bar{a}, \varpi)^{kQ(\alpha)}$ is nontrivial.

We compute

$$\begin{aligned} \int_{C_m} (a\bar{a}, \varpi)^{kQ(\alpha)} &= q^{2m-3} \left((q+1) \sum_{a \in \mathbb{F}_{q^2}^\times} (a\bar{a}, \varpi)^{kQ(\alpha)} - q \sum_{a \in \mathbb{F}_q^\times} (a\bar{a}, \varpi)^{kQ(\alpha)} \right) \\ &= -q^{2m-2}(q-1). \end{aligned}$$

Therefore the contribution to (12.1) from such terms is (setting $m = (2l + 1)n_\alpha$)

$$\sum_{l=0}^\infty -q^{(4l+2)n_\alpha-2}(q-1)(q^{-2}x_\alpha)^{(2l+1)n_\alpha} = \frac{-q^{-2}(q-1)x_\alpha^{n_\alpha}}{1-x_\alpha^{2n_\alpha}}.$$

This covers all possible cases, and a simple addition of the rational functions we have obtained completes the proof. □

13. THE ACTION WHEN $G_s \cong \text{Res } SL_2$

Let s be a simple reflection in W . Suppose $G_s \cong SL_2(E)$, q is the cardinality of the residue field of E and α is the unique positive coroot. The following theorem is equivalent to [KP, Lemma I.3.3]. We give a different and more direct proof, which will serve as a template for the more involved computation in the following section. Given any integer t , define the Gauss sum $\mathfrak{g}_{SL_2(E)}(t)$ by

$$\mathfrak{g}_{SL_2(E)}(t) = \int_{O_E^\times} (v, \varpi)^t \psi\left(\frac{v}{\varpi}\right) dv$$

with a choice of Haar measure such that the total volume of O_E^\times is $q - 1$.

Recall the coefficients $\tau_{a,b}^{(s,\chi)}$ defined in (9.1). For two elements $\lambda, \mu \in X_*(S)$, we write $\lambda \sim \mu$ if $\lambda - \mu \in \Lambda$.

Theorem 13.1. *Suppose that $a = \varpi^\nu$ and $b = \varpi^\mu$. Then we can write $\tau_{a,b}^{(s,\chi)} = \tau_{a,b}^1 + \tau_{a,b}^2$ where*

$$\begin{aligned} \tau_{a,b}^1 &= 0 \quad \text{unless } \nu \sim \mu, \\ \tau_{a,b}^2 &= 0 \quad \text{unless } s\nu \sim \mu - \alpha. \end{aligned}$$

If $\nu = \mu$, then

$$\tau_{a,b}^1 = (1 - q^{-1}) \frac{x_\alpha^{n_\alpha \lceil \frac{B(\alpha,\mu)}{n_\alpha Q(\alpha)} \rceil}}{1 - q^{-1} x_\alpha^{n_\alpha}}.$$

If $\nu = s\mu + \alpha$, then

$$\tau_{a,b}^2 = q^{-1} \mathfrak{g}_{SL_2(E)}(B(\alpha, \mu) - Q(\alpha)) \frac{1 - x_\alpha^{n_\alpha}}{1 - q^{-1} x_\alpha^{n_\alpha}}.$$

Proof. Write $n = \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \in G_s$ and $x = \varpi^{-m}v^{-1}$, where $v \in O_E^\times$ and $m \in \mathbb{Z}$. The analogous statement to (11.1) is $p(n) = (v, \varpi)^{mQ(\alpha)}v^\alpha\varpi^{m\alpha}u'$ for some $u' \in U$. We now calculate

$$\begin{aligned}
 f_b(p(n)w_0) &= (v, \varpi)^{mQ(\alpha)}\delta^{1/2}(\varpi^{m\alpha})\pi_\chi(v^\alpha\varpi^{m\alpha})f_b(w_0) \\
 &= q^{-m}(v, \varpi)^{mQ(\alpha)}\pi_\chi(bv^\alpha\varpi^{m\alpha})v_0 \\
 &= q^{-m}(v, \varpi)^{mQ(\alpha)}[b, v^\alpha]\pi_\chi(v^\alpha b\varpi^{m\alpha})v_0 \\
 (13.1) \qquad &= q^{-m}(v, \varpi)^{mQ(\alpha)+B(\alpha,\mu)}\pi_\chi(\varpi^{\mu+m\alpha})v_0.
 \end{aligned}$$

In the second line we use $f_b(w_0) = \pi_\chi(b)v_0$ and the b ends up on the left of $v^\alpha\varpi^{m\alpha}$ since the π_χ -action on functions in $I(\chi)$ is by $\pi_\chi(x)f(y) = f(yx)$. In the last line we have used the commutator relation and the fact that $\phi_K(v^\alpha) = \phi_K(1) = v_0$ as ϕ_K is spherical.

Recall that we are trying to evaluate

$$\tau_{a,b}^{(s,\chi)} = c_s(\chi)^{-1} \int_{U_s} \lambda_a^{(s\chi)} f_b(p(n)w_0) \psi^{-1}(n') dn.$$

We decompose U_s into shells where $|x|$ is constant on each shell, and the above calculations show that

$$\tau_{a,b}^{(s,\chi)} = c_s(\chi)^{-1} q^{-1} \lambda_a^{(s\chi)} \sum_{m \in \mathbb{Z}} \pi_\chi(\varpi^{\mu+m\alpha}) \phi_K(1) \int_{O_E^\times} (v, \varpi)^{mQ(\alpha)+B(\alpha,\mu)} \psi(-\varpi^m v) dv,$$

where a normalisation of Haar measure on O_E^\times is chosen such that the group has volume $q - 1$.

If $m \leq -2$, then the corresponding integral over O_E^\times vanishes. Let us now consider the contribution from the summand with $m = -1$. Here, the presence of the term $\lambda_a^{(s\chi)}\pi_\chi(\varpi^{\mu+m\alpha})v_0$ implies that this contribution is nonzero only when $s\nu \sim \mu - \alpha$. When $s\mu = \nu + \alpha$, the $m = -1$ term in the sum gives a contribution of $c_s(\chi)^{-1}q^{-1}\mathfrak{g}_{SL_2(E)}(B(\alpha, \mu) - Q(\alpha))$ to $\tau_{a,b}^{(s,\chi)}$. This corresponds to the $\tau_{a,b}^2$ part of the proposition.

Now we turn to the terms where $m \geq 0$. Here, the argument of ψ is in O_E , so the character ψ automatically takes the value 1. Hence, by Lemma 2.10, the integral over O_E^\times vanishes unless the integrand is identically equal to 1. This occurs if and only if $B(\alpha, \mu) + mQ(\alpha) \equiv 0 \pmod{n}$.

It was shown in the proof of [Mc2, Theorem 11.1] that $B(\alpha, \mu)$ is divisible by $Q(\alpha)$. Thus the condition for nonvanishing of the integral now becomes $m \equiv -B(\alpha, \mu)/Q(\alpha) \pmod{n_\alpha}$. Let us write $m = kn_\alpha - B(\alpha, \mu)/Q(\alpha)$. Then k runs over all integers greater than or equal to $\lceil \frac{B(\alpha,\mu)}{n_\alpha Q(\alpha)} \rceil$. By Lemma 2.6, for integers m in this family, all elements of the form $\varpi^{\mu+m\alpha}$ lie in the same H -coset.

Therefore the sum over $m \geq 0$ is zero unless $s\nu \sim \mu - \frac{B(\alpha,\mu)}{Q(\alpha)}\alpha$. By the definition of the action of W on the cocharacter lattice, this is equivalent to $\nu \sim \mu$.

If indeed we do have $\nu = \mu$, then we obtain a contribution to $\tau_{a,b}^{(s,\chi)}$ from the terms $m \geq 0$ equal to

$$c_s(\chi)^{-1}(1 - q^{-1}) \sum_{k \geq \lceil \frac{B(\alpha,\mu)}{n_\alpha Q(\alpha)} \rceil} x_\alpha^{kn_\alpha}.$$

This corresponds to the $\tau_{a,b}^1$ part of the proposition and the only thing remaining to do is to sum this geometric series and substitute the evaluation of $c_s(\chi)$ from Theorem 12.1. \square

14. THE ACTION WHEN $G_s \cong \text{Res } SU_3$

Now that we have warmed up by proving a Gindikin-Karpelevic formula and covered the computation of the coefficients $\tau_{a,b}^{(s,\chi)}$ when $G_s \cong \text{Res } SL_2$, we present the analogue of Theorem 13.1 when $G_s \cong \text{Res } SU_3$.

So assume that $G_s \cong \text{Res}_{E/F} SU_3$, let q be the cardinality of the residue field of E and let α be the unique positive rational coroot. Given any integer t , we define the SU_3 Gauss sum $\mathfrak{g}_{SU_3(E)}(t)$ by

$$\mathfrak{g}_{SU_3(E)}(t) = \sum_{u \in U(\mathbb{F}_q) \setminus \{1\}} (y\bar{y}, \varpi)^t \psi\left(\frac{x}{\varpi y}\right),$$

where $u = \begin{pmatrix} 1 & x & y \\ 0 & 1 & -\bar{x} \\ 0 & 0 & 1 \end{pmatrix} \in U(\mathbb{F}_q)$, the unipotent radical of $SU_3(\mathbb{F}_q)$.

Theorem 14.1. *Suppose that $c = \varpi^\nu$ and $b = \varpi^\mu$. Then we can write $\tau_{c,b}^{(s,\chi)} = \tau_{c,b}^1 + \tau_{c,b}^2$, where*

$$\begin{aligned} \tau_{c,b}^1 &= 0 \quad \text{unless } \nu \sim \mu, \\ \tau_{c,b}^2 &= 0 \quad \text{unless } s\nu \sim \mu - 2\alpha. \end{aligned}$$

If $\nu = \mu$, then

$$\tau_{c,b}^1 = \frac{(1 - q^{-3})x_\alpha^{2n_\alpha \lceil \frac{B(\alpha,\mu)}{2n_\alpha Q(\alpha)} \rceil} + (q^{-1} - q^{-2})q^{(1-\epsilon)/2}x_\alpha^{(2\lceil \frac{B(\alpha,\mu) + n_\alpha Q(\alpha) - Q(\alpha)}{2n_\alpha Q(\alpha)} \rceil - 1)n_\alpha}}{(1 - \epsilon q^{-1}x_\alpha^{n_\alpha})(1 + \epsilon q^{-2}x_\alpha^{n_\alpha})}.$$

If $\nu = s\mu + 2\alpha$, then

$$\tau_{c,b}^2 = q^{-3} \mathfrak{g}_{SU_3(E)}(B(\alpha,\mu)/2 - Q(\alpha)) \frac{1 - x_\alpha^{2n_\alpha}}{(1 - \epsilon q^{-1}x_\alpha^{n_\alpha})(1 + \epsilon q^{-2}x_\alpha^{n_\alpha})}.$$

Proof. We begin by collecting notation, mostly from previous sections, which we will use in this proof.

The unramified quadratic extension of E used to define SU_3 is denoted L , we use a bar to denote the action of the nontrivial element of $\text{Gal}(L/E)$ and $\theta \in O_L^\times$ is such that $\bar{\theta} = -\theta$.

The coroot α can be written as $\alpha_1 + \alpha_2$, where α_1 and α_2 are simple roots for the geometric cocharacter lattice of SU_3 that form an orbit under the action of $\text{Gal}(L/E)$.

An element in U_s is written $u = \begin{pmatrix} 1 & x & y \\ 0 & 1 & -\bar{x} \\ 0 & 0 & 1 \end{pmatrix}$, where $x, y \in L$ are such that

$$x\bar{x} + y + \bar{y} = 0.$$

For nonzero $u \in U_s$, we let $m = -v(y)$, $y = a\theta\varpi^{-m}$ and $z = x/y$. Then $1/y = -z\bar{z}/2 + h\theta$ for some $h \in E$.

The element $p(u)$ is equal to $(a, \bar{a})^{mQ(\alpha)/2} a^{\alpha_1} \bar{a}^{\alpha_2} \varpi^{m\alpha}$ times an element of U . The commutator of $a^{\alpha_1} \bar{a}^{\alpha_2}$ and ϖ^μ is $(a\bar{a}, \varpi)^{B(\alpha,\mu)/2}$. Thus the evaluation of the

function f_b analogous to (13.1) is

$$f_b(p(u)w_0) = q^{-2m}(a\bar{a}, \varpi)^{(mQ(\alpha)+B(\alpha,\mu))/2}\pi_\chi(\varpi^{\mu+m\alpha})v_0.$$

After substitutions, equation (9.3) becomes (14.1)

$$\tau_{c,b}^{(s,\chi)} = \sum_{m \in \mathbb{Z}} \frac{q^{-2m}}{c_s(\chi)} \lambda_c^{(s\chi)}(\pi_\chi(\varpi^{\mu+m\alpha})v_0) \int_{C_m} (a\bar{a}, \varpi)^{(mQ(\alpha)+B(\alpha,\mu))/2} \psi(z) du.$$

First consider the terms in the sum where $m \leq -3$ and is odd. Parametrising C_m by the coordinates (z, h) , the integrand factors into a piece $(a\bar{a}, \varpi)^{(mQ(\alpha)+B(\alpha,\mu))/2}$ which only depends on h and a piece $\psi(z)$ which only depends on z . Therefore the integral over C_m contains a factor equal to $\int_{\varpi^{(m+1)/2}\mathcal{O}_L} \psi(z) dz$. This is zero by Lemma 2.10.

Now suppose that $m \leq 3$ and is even. Pick $u \in O_L^\times$. Consider the measure-preserving homeomorphism $(z, h) \mapsto (z + \varpi^{-1}u, h)$ of C_m . The effect of this homeomorphism on the corresponding term of (14.1) is to multiply the integrand by $\psi(u/\varpi)$. As u can be chosen such that $\psi(u/\varpi) \neq 1$, these terms with $m \leq 3$ are all zero.

Consider the term where $m = -2$. It is

$$c_s(\chi)^{-1} q^4 \lambda_c^{(s\chi)}(\pi_\chi(\varpi^{\mu-2\alpha})v_0) \int_{C_{-2}} (a\bar{a}, \varpi)^{B(\alpha,\mu)/2-Q(\alpha)} \psi(x/y) du.$$

The presence of the factor $\lambda_c^{(s\chi)}\pi_\chi(\varpi^{\mu-2\alpha})v_0$ implies that this term vanishes unless $s\nu \sim \mu - 2\alpha$. When $\nu = s\mu + 2\alpha$, by definition of the SU_3 Gauss sum, we get

$$c_s(\chi)^{-1} q^{-3} \mathfrak{g}_{SU_3(E)}(B(\alpha, \mu)/2 - Q(\alpha)).$$

This corresponds to the $\tau_{a,b}^2$ term in the theorem.

Now suppose that $m \geq -1$. Then $z \in \mathcal{O}_L$ so $\psi(z) = 1$. Therefore we have to evaluate the integral

$$(14.2) \quad \int_{C_m} (a\bar{a}, \varpi)^{(mQ(\alpha)+B(\alpha,\mu))/2} du.$$

First consider the case where m is odd. Then $a \equiv \bar{a} \pmod{\varpi}$. The integral (14.2) thus reduces to a constant factor times

$$\sum_{a \in \mathbb{F}_q^\times} (a, \varpi)^{mQ(\alpha)+B(\alpha,\mu)}.$$

We evaluate it by Lemma 2.10 to see that when m is odd, (14.2) evaluates to

$$\int_{C_m} (a\bar{a}, \varpi)^{(mQ(\alpha)+B(\alpha,\mu))/2} du = \begin{cases} \text{vol}(C_m) & \text{if } n \mid mQ(\alpha) + B(\alpha, \mu), \\ 0 & \text{otherwise.} \end{cases}$$

Now suppose that m is even. We note that $B(\alpha, \mu) \in 2\mathbb{Z}$ and $B(\alpha, \mu)/2$ is divisible by $Q(\alpha)$. This is because $\alpha = \alpha_1 + \alpha_2$, where α_1 and α_2 are in the same Galois orbit. Thus by linearity of B and Galois-invariantness of μ , we have $B(\alpha, \mu) = 2B(\alpha_1, \mu)$, and we already know that $B(\alpha_1, \mu)$ is divisible by $Q(\alpha)$. Let us write $m = 2k$ and $B(\alpha, \mu) = 2lQ(\alpha)$.

By the same volume computation that appeared in Section 12, we have

$$\int_{C_m} (a\bar{a}, \varpi)^{(mQ(\alpha)+B(\alpha,\mu))/2} du = q^{2m-3} \left((q+1) \sum_{a \in \mathbb{F}_{q^2}^\times} (a\bar{a}, \varpi)^{(k+l)Q(\alpha)} - \sum_{a \in \mathbb{F}_q^\times} (a, \varpi)^{2(k+l)Q(\alpha)} \right).$$

Again, this can be evaluated using Lemma 2.10 to obtain, for m even,

$$\int_{C_m} (a\bar{a}, \varpi)^{(mQ(\alpha)+B(\alpha,\mu))/2} du = \begin{cases} \text{vol}(C_m) & \text{if } n_\alpha \mid k+l, \\ -q^{2m-2}(q-1) & \text{if } n_\alpha \nmid k+l \text{ and } n_\alpha \mid 2(k+l), \\ 0 & \text{if } n_\alpha \nmid 2(k+l). \end{cases}$$

In all cases, we see that we get zero unless n divides $mQ(\alpha) + B(\alpha, \mu)$. This is equivalent to $m \equiv -B(\alpha, \mu)/Q(\alpha) \pmod{n_\alpha}$. As $n_\alpha \alpha \in \Lambda$, under such circumstances we have

$$\mu + m\alpha \sim \mu - \frac{B(\alpha, \mu)}{Q(\alpha)}\alpha = s\mu.$$

The factor $\lambda_c^{(s\chi)}(\pi_\chi(\varpi^{\mu+m\alpha})v_0)$ in (14.1) is zero unless $s\nu \sim \mu + m\alpha$. Combining these observations shows that we only have a contribution to (14.1) from terms with $m \geq -1$ when $\nu \sim \mu$.

To explicitly evaluate the term $\tau_{c,b}^1$ when $\nu = \mu$ is now a routine matter of evaluating the geometric series obtained as a consequence of the integral evaluations performed above. □

15. COMPARISON

We conclude this paper by comparing our above results with those of Chinta and Gunnells [CG] on the construction of the local part of a Weyl group multiple Dirichlet series. To do so, we suppose that G is split, simple and simply connected, and that Q is chosen to take the value 1 on short coroots.

The group homomorphism from $X_*(S)$ to \tilde{G} , $\lambda \mapsto \mathfrak{s}(\varpi^\lambda)$ provides us with a splitting of the short exact sequence (3.1), and hence determines for us an isomorphism $\mathcal{O}(Y) \cong \mathbb{C}[\Lambda]$. Let Σ be the multiplicative subset of $\mathbb{C}[\Lambda]$ such that $\mathcal{O}(V) \cong \Sigma^{-1}\mathbb{C}[\Lambda]$.

Make a choice of left coset representatives Γ for H in \widetilde{M} consisting of elements of the form $\mathfrak{s}(\varpi^\lambda)$. The sections $\{\lambda_a^{(\chi)}\}_{a \in \Gamma}$ of \mathcal{E}^* provide us with a trivialisation of \mathcal{E}^* and hence an isomorphism

$$\Gamma(V, \mathcal{E}^*) \cong \bigoplus_{a \in \Gamma} \Sigma^{-1}\mathbb{C}[\Lambda].$$

Define an isomorphism $\varphi : \Gamma(V, \mathcal{E}^*) \rightarrow \Sigma^{-1}\mathbb{C}[X_*(S)]$ by

$$\varphi \left(\sum_{\mathfrak{s}(\varpi^\mu) \in \Gamma} f_\mu \cdot \lambda_{\mathfrak{s}(\varpi^\mu)}^{(\chi)} \right) = \sum_{\mathfrak{s}(\varpi^\mu) \in \Gamma} f_\mu \cdot \mu,$$

where $f_\mu \in \mathcal{O}(V)$.

As the definition of the W -action on $\mathcal{E}^*|_V$ is by transporting the action of W on $\mathcal{F}|_V$, we have, for $w \in W$,

$$w \cdot \lambda_a^{(\chi)} = \sum_{b \in \Gamma} \tau_{a,b}^{(w, w^{-1}\chi)} \lambda_b^{(w^{-1}\chi)}.$$

At this point, let us introduce a change in notation. We shall write $\tau_{\mu, \nu}^i$ for $\tau_{\mathfrak{s}(\varpi^\mu), \mathfrak{s}(\varpi^\nu)}$ and $i = 1, 2$. For $\mu \in X_*(S)$, we now write x^μ for the corresponding element of the group algebra $\mathbb{C}[X_*(S)]$.

When $w = s$ is the simple reflection corresponding to the simple coroot α and $a = \mathfrak{s}(\varpi^\mu)$, Theorem 13.1 yields

$$(15.1) \quad s \cdot \lambda_a^{(\chi)} = \tau_{\mu, \mu}^1 \lambda_a^{(s\chi)} + \tau_{\mu, s\mu + \alpha}^2 \lambda_{s\mu + \alpha}^{(s\chi)}.$$

Translating (15.1) under the isomorphism φ gives the following expression for the action of a simple reflection on $\Sigma^{-1}\mathbb{C}[X_*(S)]$:

$$s \cdot x^\mu = \tau_{\mu, \mu}^1 x^\mu + \tau_{\mu, s\mu + \alpha}^2 x^{s\mu + \alpha}.$$

Substituting in the values of τ^1 and τ^2 from Theorem 13.1 yields the following explicit formula:

$$s_\alpha \cdot x^\lambda = \frac{x^{s\lambda}}{1 - q^{-1}x_\alpha^{n_\alpha}} \left((1 - q^{-1})x_\alpha^{n_\alpha \lceil \frac{B(\alpha, \lambda)}{n_\alpha Q(\alpha)} \rceil - \frac{B(\alpha, \lambda)}{Q(\alpha)}} + q^{-1}x_\alpha^{-1}(1 - x_\alpha^{n_\alpha})\mathfrak{g}_{SL_2(F)}(B(\alpha, \lambda) - Q(\alpha)) \right).$$

We compare this to the action introduced in [CG] for the purpose of constructing Weyl group multiple Dirichlet series. Our lattice $X_*(S)$ corresponds to the lattice denoted Λ in [CG]. As in [CO, §9], we need to make a minor change of variables from the Chinta-Gunnells paper to eliminate extraneous powers of q . Also as per [CO, §9], we need to modify the action by multiplication with a rather innocuous factor in order to directly compare this action derived from intertwining operators with that from [CG].

In [CG, Definition 3.1], an action is defined for any dominant λ . We write $f \mapsto f|_\lambda w$ for this action. Then $f||w := x^\lambda(x^{-\lambda}f|_\lambda w)$ is independent of λ and defines an action of W on $\Sigma^{-1}\mathbb{C}[X_*(S)]$.

Let Φ denote the set of coroots, and $\Phi = \Phi^+ \sqcup \Phi^-$ the decomposition into positive and negative coroots. For $w \in W$, let $\Phi(w) = \{\alpha \in \Phi^+ | w\alpha \in \Phi^-\}$. Now consider the two Weyl group actions \circ_1 and \circ_2 , under which the actions of $w \in W$ on $f \in \Sigma^{-1}\mathbb{C}[X_*(S)]$ are

$$w \circ_1 f = \frac{c_{w_0}(\chi)}{c_{w_0}(w\chi)}(w \cdot f),$$

$$w \circ_2 f = \text{sgn}(w) \left(\prod_{\alpha \in \Phi(w)} x_\alpha^{n_\alpha} \right) (f||w).$$

For any $w \in W$, $f \in \Sigma^{-1}\mathbb{C}[X_*(S)]$ and $g \in \Sigma^{-1}\mathbb{C}[\Lambda]$, the actions \cdot and $||$ satisfy the transformation property

$$w \cdot (gf) = (wg)w \cdot f, \quad (gf)||w = (wg)(f||w),$$

where wg denotes the usual action of W on $\Sigma^{-1}\mathbb{C}[\Lambda]$. Therefore \circ_1 and \circ_2 are indeed Weyl group actions. There is an explicit formula for $c_w(\chi)$, namely

$$c_w(\chi) = \prod_{\alpha \in \Phi(w)} \frac{1 - q^{-1}x_\alpha^{n_\alpha}}{1 - x_\alpha^{n_\alpha}}.$$

This formula follows from Theorem 12.1 and the cocycle formula $c_{w_1 w_2}(\chi) = c_{w_1}(w_2 \chi) c_{w_2}(\chi)$ whenever $\ell(w_1 w_2) = \ell(w_1) + \ell(w_2)$.

Proposition 15.1. *The two Weyl group actions \circ_1 and \circ_2 are the same action.*

Proof. To prove this proposition, it suffices to consider the action of a simple reflection. We have explicit formulae for the action of a simple reflection on a monomial x^λ in each case, and, upon comparison, we see that the actions are the same. The case of general f then follows by linearity. \square

Chinta and Gunnells use their action to construct the p -part of a Weyl group multiple Dirichlet series. This requires the construction of an auxiliary polynomial

$$N(\chi, \lambda) = \prod_{\alpha > 0} \frac{1 - q^{-1}x_\alpha^{n_\alpha}}{1 - x_\alpha^{n_\alpha}} \sum_{w \in W} \operatorname{sgn}(w) \left(\prod_{\alpha \in \Phi(w)} x_\alpha^{n_\alpha} \right) (1|_\lambda w)(\chi).$$

Now the culmination of the above leads to our final result. Informally, this states that the value of the metaplectic Whittaker function on a torus element is equal to the p -part of a Weyl group multiple Dirichlet series.

Theorem 15.2. *Let λ be dominant. The following identity holds:*

$$(\delta^{-1/2} \mathcal{W}_\chi)(\varpi^\lambda) = \chi(\varpi^\lambda) N(\chi, \lambda).$$

The results of this section suggest how to extend the results in [CG] beyond the split case. We note that some of the content of this paper would accomplish some of the necessary work to achieve this aim. For example, we have obtained an independent proof of [CG, Theorem 3.2]. Although this proof is more indirect than that in [CG], it has the advantage of being more conceptual.

REFERENCES

- [BD] Jean-Luc Brylinski and Pierre Deligne, *Central extensions of reductive groups by \mathbf{K}_2* , Publ. Math. Inst. Hautes Études Sci. **94** (2001), 5–85, DOI 10.1007/s10240-001-8192-2. MR1896177 (2004a:20049)
- [BLS] William D. Banks, Jason Levy, and Mark R. Sepanski, *Block-compatible metaplectic cocycles*, J. Reine Angew. Math. **507** (1999), 131–163, DOI 10.1515/crll.1999.011. MR1670203 (99m:20100)
- [BT] Armand Borel and Jacques Tits, *Groupes réductifs* (French), Inst. Hautes Études Sci. Publ. Math. **27** (1965), 55–150. MR0207712 (34 #7527)
- [BZ] I. N. Bernstein and A. V. Zelevinsky, *Induced representations of reductive p -adic groups. I*, Ann. Sci. École Norm. Sup. (4) **10** (1977), no. 4, 441–472. MR0579172 (58 #28310)
- [C] W. Casselman, *The unramified principal series of p -adic groups. I. The spherical function*, Compositio Math. **40** (1980), no. 3, 387–406. MR571057 (83a:22018)
- [CG] Gautam Chinta and Paul E. Gunnells, *Constructing Weyl group multiple Dirichlet series*, J. Amer. Math. Soc. **23** (2010), no. 1, 189–215, DOI 10.1090/S0894-0347-09-00641-9. MR2552251 (2011g:11100)
- [CO] Gautam Chinta and Omer Offen, *A metaplectic Casselman-Shalika formula for GL_r* , Amer. J. Math. **135** (2013), no. 2, 403–441, DOI 10.1353/ajm.2013.0013. MR3038716
- [CS] W. Casselman and J. Shalika, *The unramified principal series of p -adic groups. II. The Whittaker function*, Compositio Math. **41** (1980), no. 2, 207–231. MR581582 (83i:22027)

- [GG] W. T. Gan and F. Gao, *The Langlands-Weissman Program for Brylinski-Deligne extensions*. <http://arxiv.org/abs/1409.4039>.
- [KP] D. A. Kazhdan and S. J. Patterson, *Metaplectic forms*, Inst. Hautes Études Sci. Publ. Math. **59** (1984), 35–142. MR743816 (85g:22033)
- [Mc1] Peter J. McNamara, *Metaplectic Whittaker functions and crystal bases*, Duke Math. J. **156** (2011), no. 1, 1–31, DOI 10.1215/00127094-2010-064. MR2746386 (2012a:11060)
- [Mc2] Peter J. McNamara, *Principal series representations of metaplectic groups over local fields*, Multiple Dirichlet series, L-functions and automorphic forms, Progr. Math., vol. 300, Birkhäuser/Springer, New York, 2012, pp. 299–327, DOI 10.1007/978-0-8176-8334-4_13. MR2963537
- [T] J. Tits, *Normalisateurs de tores. I. Groupes de Coxeter étendus* (French), J. Algebra **4** (1966), 96–116. MR0206117 (34 #5942)
- [W] Martin H. Weissman, *Split metaplectic groups and their L-groups*, J. Reine Angew. Math. **696** (2014), 89–141, DOI 10.1515/crelle-2012-0111. MR3276164

SCHOOL OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF SYDNEY, NEW SOUTH WALES
2006, AUSTRALIA

E-mail address: `mcnamara@maths.usyd.edu.au`

Current address: School of Mathematics and Physics, The University of Queensland, Brisbane
St. Lucia, Queensland 4072, Australia