

PULLBACK INVARIANTS OF THURSTON MAPS

SARAH KOCH, KEVIN M. PILGRIM, AND NIKITA SELINGER

ABSTRACT. Associated to a Thurston map $f : S^2 \rightarrow S^2$ with postcritical set P are several different invariants obtained via pullback: a relation $S_P \xleftarrow{f} S_P$ on the set S_P of free homotopy classes of curves in $S^2 \setminus P$, a linear operator $\lambda_f : \mathbb{R}[S_P] \rightarrow \mathbb{R}[S_P]$ on the free \mathbb{R} -module generated by S_P , a virtual endomorphism $\phi_f : \text{PMod}(S^2, P) \dashrightarrow \text{PMod}(S^2, P)$ on the pure mapping class group, an analytic self-map $\sigma_f : \mathcal{T}(S^2, P) \rightarrow \mathcal{T}(S^2, P)$ of an associated Teichmüller space, and an analytic self-correspondence $X \circ Y^{-1} : \mathcal{M}(S^2, P) \rightrightarrows \mathcal{M}(S^2, P)$ of an associated moduli space. Viewing these associated maps as invariants of f , we investigate relationships between their properties.

CONTENTS

| | |
|--|------|
| 1. Introduction | 4621 |
| 2. Fundamental identifications | 4624 |
| 3. Pullback invariants | 4627 |
| 4. When $\sigma_f : \partial\mathcal{T}_B \rightarrow \partial\mathcal{T}_A$ | 4636 |
| 5. When $\sigma_f : \mathcal{T}_B \rightarrow \mathcal{T}_A$ is constant | 4637 |
| 6. Noninjectivity of the virtual homomorphism | 4640 |
| 7. Finite global attractor for the pullback relation | 4644 |
| 8. Shadowing | 4648 |
| 9. Obstructed twists and repelling fixed points in $\partial\mathcal{M}_P$ | 4648 |
| Acknowledgements | 4653 |
| References | 4653 |

1. INTRODUCTION

Thurston maps are orientation-preserving branched covers $f : S^2 \rightarrow S^2$ from the oriented topological two-sphere to itself that satisfy certain properties. They were introduced by Thurston as combinatorial invariants associated to postcritically finite rational functions $R : \mathbb{P}^1 \rightarrow \mathbb{P}^1$, regarded as dynamical systems on the Riemann sphere. A fundamental theorem of complex dynamics is Thurston’s characterization and rigidity theorem [DH], which (i) characterizes which Thurston maps f arise from rational functions, and (ii) says that apart from a well-understood family of ubiquitous exceptions, the Möbius conjugacy class of R is determined by the combinatorial class of its associated Thurston map f . The proof of this theorem transforms the question of determining whether f arises from a rational function R to the question of whether an associated *pullback map* $\sigma_f : \mathcal{T}_P \rightarrow \mathcal{T}_P$, an analytic self-map of a certain Teichmüller space, has a fixed point.

Received by the editors April 2, 2013 and, in revised form, May 10, 2014.
 2010 *Mathematics Subject Classification*. Primary 30F60, 32G15, 37F20.

As combinatorial (as opposed to analytic or algebraic) objects, Thurston maps, in principle, should be easier to classify than postcritically finite rational maps. For many years, the lack of suitable invariants for general Thurston maps frustrated attempts toward this goal. Kameyama [Kam] developed algebraic methods geared specifically to the classification of combinatorial classes of maps arising as *twists* of a given Thurston map f : these are maps of the form $h_0 \circ f \circ h_1$ where h_0, h_1 are homeomorphisms. In a ground-breaking paper [BN1], Bartholdi and Nekrashevych introduced several new general tools. One innovation was to apply the general theory of self-similar groups to study both the dynamics of a given Thurston map f and the classification of twists of f . Another was to exploit the existence of an associated analytic correspondence on the moduli space \mathcal{M}_P covered by the graph of σ_f . These new tools have led to much better invariants for Thurston maps and to a better understanding of the pullback map σ_f ; see [Nek1], [CFPP], [Koc1], [Koc2], [Pil3], [BEKP], [HP1], [HP2], [Kel], [Nek2], [BN2], [Lod].

In this work, we deepen the investigations of the relationship between dynamical, algebraic, and analytic invariants associated to Thurston maps.

Fundamental invariants. Let S^2 denote a topological two-sphere equipped with an orientation. Fix an identification $S^2 = \mathbb{P}^1 = \widehat{\mathbb{C}}$; we use S^2 for topological objects, \mathbb{P}^1 for holomorphic objects, and $\widehat{\mathbb{C}}$ when formulas are required. Let $P \subseteq S^2$ be a finite set with $\#P \geq 3$ (in the case $\#P = 3$, the groups and spaces are trivial, so it is helpful to imagine at first that $\#P \geq 4$).

The following objects are basic to our study:

- (1) \mathcal{S}_P , the set of free homotopy classes of simple, unoriented, essential, non-peripheral, closed curves in $S^2 \setminus P$; the symbol o denotes the union of the sets of free homotopy classes of unoriented inessential and peripheral curves. A curve representing an element of \mathcal{S}_P we call *nontrivial*. A multicurve Γ on $S^2 \setminus P$ is a possibly empty subset of \mathcal{S}_P represented by nontrivial, pairwise disjoint, pairwise nonhomotopic curves. Let \mathcal{MS}_P denote the set of possibly empty multicurves on $S^2 \setminus P$.
- (2) $\mathbb{R}[\mathcal{S}_P]$, the free \mathbb{R} -module generated by \mathcal{S}_P ; this arises naturally in the statement of Thurston’s characterization theorem.
- (3) $G_P := \text{PMod}(S^2, P)$, the pure mapping class group (that is, orientation-preserving homeomorphisms $g : (S^2, P) \rightarrow (S^2, P)$ fixing P pointwise, up to isotopy through homeomorphisms fixing P pointwise). Each nontrivial element of G_P has infinite order. The group G_P contains a distinguished subset Tw whose elements are multitwists; that is

$$\text{Tw} := \bigcup_{\Gamma \text{ a multicurve on } S^2 \setminus P} \text{Tw}(\Gamma),$$

where $\text{Tw}(\Gamma)$ is the subgroup of G_P generated by Dehn twists around components of Γ . A multitwist is *positive* if it is a composition of positive powers of right Dehn twists. Let

$$\text{Tw}^+ := \bigcup_{\Gamma \text{ a multicurve on } S^2 \setminus P} \text{Tw}^+(\Gamma),$$

where $\text{Tw}^+(\Gamma)$ is the subgroup of G_P generated by positive Dehn twists around components of Γ . The *support* of a multitwist is the smallest multicurve about which the twists comprising it occur; this is well defined.

- (4) $\mathcal{T}_P := \mathcal{T}(S^2, P)$, the Teichmüller space of (S^2, P) , as in [DH]. It comes equipped with two natural metrics, the Teichmüller Finsler metric $d_{\mathcal{T}_P}$ and the Weil-Petersson (WP) inner product metric $d_{\mathcal{T}_P}^{WP}$. The metric space $(\mathcal{T}_P, d_{\mathcal{T}_P})$ is complete, and any pair of points are joined by a unique geodesic. In contrast, $(\mathcal{T}_P, d_{\mathcal{T}_P}^{WP})$ is incomplete. The WP-completion $\overline{\mathcal{T}}_P$ has a rich geometric structure [Mas] (see also [HK]); it is a stratified space where each stratum \mathcal{T}_P^Γ corresponds to a multicurve Γ and consists of noded Riemann surfaces whose nodes correspond to pinching precisely those curves comprising Γ to points. We denote by $\partial\mathcal{T}_P$ the WP boundary of \mathcal{T}_P .
- (5) $\mathcal{M}_P := \mathcal{M}(S^2, P)$ is the corresponding moduli space. It is a complex manifold, isomorphic to a complex affine hyperplane complement. The natural projection $\pi : \mathcal{T}_P \rightarrow \mathcal{M}_P$ is a universal cover with deck group G_P .

Given a basepoint $\tau_\otimes \in \mathcal{T}_P$, there is a natural identification of G_P with $\pi_1(\mathcal{M}_P, m_\otimes)$ where $m_\otimes := \pi(\tau_\otimes)$; the isomorphism $\pi_1(\mathcal{M}_P, m_\otimes) \rightarrow G_P$ proceeds via isotopy extension, while the isomorphism $G_P \rightarrow \pi_1(\mathcal{M}_P, m_\otimes)$ is induced by composing the evaluation map at those points marked by P with an isotopy to the identity through maps fixing three points of \mathcal{M}_P ; see [Lod] for details.

A subgroup $L < G_P$ is *purely parabolic* if $L \subseteq \text{Tw}$. In terms of the identification $G_P \leftrightarrow \pi_1(\mathcal{M}_P, m_\otimes)$, L is purely parabolic if and only if the displacements of each of its elements satisfies $\inf_{\tau \in \mathcal{T}_P} d_{\mathcal{T}_P}(\tau, g.\tau) = 0$ for all $g \in L$. A purely parabolic subgroup is *complete* if for each multitwist $M \in L$ and each $\gamma \in \mathcal{S}_P$, we have $\gamma \in \text{supp}(M) \Rightarrow T_\gamma \in L$, where T_γ is the right Dehn twist about γ . A complete parabolic subgroup is necessarily of the form $\text{Tw}(\Gamma)$, the set of all multitwists about elements of Γ , where Γ is some nonempty multicurve.

Let $f : S^2 \rightarrow S^2$ be an orientation-preserving branched covering map with critical set Ω_f . Its *postcritical set* is

$$P_f := \bigcup_{n>0} f^{\circ n}(\Omega_f);$$

we assume throughout this work that $\#P_f < \infty$. Suppose $P \subseteq S^2$ is finite, $P_f \subseteq P$, and $f(P) \subseteq P$; we then call the map of pairs $f : (S^2, P) \rightarrow (S^2, P)$ a *Thurston map*. Throughout this work, we refer to a Thurston map by the symbol f , and often suppress mention of the noncanonical subset P .

The following objects are associated to a Thurston map $f : (S^2, P) \rightarrow (S^2, P)$ via pullback:

- (1) a relation $\beta_f : \mathcal{S}_P \cup \{o\} \rightarrow \mathcal{S}_P \cup \{o\}$,
- (2) a nonnegative linear operator $\lambda_f : \mathbb{R}[\mathcal{S}_P] \rightarrow \mathbb{R}[\mathcal{S}_P]$,
- (3) a virtual endomorphism $\phi_f : G_P \dashrightarrow G_P$,
- (4) an analytic map $\sigma_f : \mathcal{T}_P \rightarrow \mathcal{T}_P$; this extends continuously to $\sigma_f : \overline{\mathcal{T}}_P \rightarrow \overline{\mathcal{T}}_P$ by [Sel1, §4],
- (5) an analytic correspondence $X \circ Y^{-1} : \mathcal{M}_P \rightrightarrows \mathcal{M}_P$; the double-arrow notation reflects our view that this is a one-to-many “function”.

Precise definitions and a summary of basic properties of these associated maps will be given in the next two sections.

Main results. The goal of this work is to examine how properties of the objects in (1) through (5) are related. That such relationships should exist is expected

since there are fundamental identifications between various elements associated to the domains of these maps; see §2. Our main results include the following:

- We characterize when the map on Teichmüller space $\sigma_f : \mathcal{T}_P \rightarrow \mathcal{T}_P$ sends the Weil-Petersson boundary to itself (Theorem 4.1).
- We characterize when the map on Teichmüller space $\sigma_f : \mathcal{T}_P \rightarrow \mathcal{T}_P$ is constant (Theorem 5.1); the proof we give corrects an error in [BEKP].
- We show that the virtual endomorphism on the pure mapping class group $\phi_f : G_P \dashrightarrow G_P$ is essentially never injective (Theorem 6.3).
- It is natural to investigate whether a Thurston map f induces an action on projectivized measured foliations, i.e. on the Thurston boundary of \mathcal{T}_P . Using Theorem 6.3, however, we show (Theorem 6.4) that each nonempty fiber of the pullback map σ_f accumulates on the entire Thurston boundary. This substantially strengthens the conclusion of [Sel1, Theorem 9.4].
- We give sufficient analytic criteria (Theorem 7.2) for the existence of a finite global attractor for the pullback relation $\mathcal{S}_P \xleftarrow{f} \mathcal{S}_P$ on curves—equivalently, for the action of the pullback map $\sigma_f : \overline{\mathcal{T}}_P \rightarrow \overline{\mathcal{T}}_P$ on strata.
- We prove an orbit lifting result (Proposition 8.1) which asserts that under the hypothesis that the virtual endomorphism ϕ_f is surjective, finite orbit segments of the pullback correspondence $X \circ Y^{-1}$ on moduli space can be lifted to finite orbit segments of σ_f .
- In the case when $\#P = 4$, we relate fixed points of the various associated maps (Theorem 9.1). If in addition the inverse of the pullback correspondence is actually a function (i.e. the map X is injective), sharper statements are possible (Theorem 9.2). Our results generalize, clarify, and put into context the algebraic, analytic, and dynamical findings in the analysis of twists of $z \mapsto z^2 + i$ given in [BN1, §6]. Using the shadowing result from Proposition 8.1, these results also demonstrate that for certain unobstructed Thurston maps, one can build finite orbits of the pullback map whose underlying surfaces behave in prescribed ways. For example, one can arrange it so that the length of the systole can become shorter and shorter for a while before stabilizing.

While motivated by the attempt at a combinatorial classification of dynamical systems, many of the results we obtain are actually more naturally phrased for a nondynamical branched covering map $f : (S^2, A) \rightarrow (S^2, B)$. Where possible, we first phrase and prove more general results (Proposition 3.1, Theorem 4.1, Theorem 5.1, Proposition 6.2, Theorem 6.3, and Theorem 6.4); the aforementioned theorems then become corollaries of these more general results.

Conventions. All branched covers and homeomorphisms are orientation-preserving. Throughout, f denotes a branched covering of $S^2 \rightarrow S^2$ of degree $d \geq 2$. The symbols A, B, P denote finite subsets of S^2 , which contain at least three points.

2. FUNDAMENTAL IDENTIFICATIONS

2.1. Preliminaries.

Teichmüller and moduli spaces. The Teichmüller space $\mathcal{T}_P = \mathcal{T}(S^2, P)$ is the space of equivalence classes of orientation-preserving homeomorphisms $\varphi : S^2 \rightarrow \mathbb{P}^1$,

whereby $\varphi_1 \sim \varphi_2$ if there is a Möbius transformation $\mu : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ so that

- $\varphi_1 = \mu \circ \varphi_2$ on the set P and
- φ_1 is isotopic to $\mu \circ \varphi_2$ relative to the set P .

The moduli space $\mathcal{M}_P = \mathcal{M}(S^2, P)$ is the set of all injective maps $\varphi : P \hookrightarrow \mathbb{P}^1$ modulo postcomposition by Möbius transformations. The Teichmüller space and the moduli space are complex manifolds of dimension $\#P - 3$, and the map $\pi_P : \mathcal{T}_P \rightarrow \mathcal{M}_P$ given by $\pi_P : [\varphi] \mapsto [\varphi|_P]$ is a holomorphic universal covering map.

Note that since we have identified $S^2 = \mathbb{P}^1$, both \mathcal{M}_P and \mathcal{T}_P have natural basepoints represented by the classes of the inclusion and identity maps, respectively.

Teichmüller metric. Equipped with the Teichmüller Finsler metric $d_{\mathcal{T}_P}$, the space \mathcal{T}_P becomes a complete uniquely geodesic metric space, homeomorphic to the open ball $B^{\#P-3}$; it is not, however, nonpositively curved.

WP metric. In contrast, when equipped with the WP metric $d_{\mathcal{T}_P}^{WP}$, the space \mathcal{T}_P is negatively curved but incomplete. The completion $\overline{\mathcal{T}}_P$ is a stratified space whose strata are indexed by (possibly empty) multicurves. Each stratum is homeomorphic to the product of the Teichmüller spaces of the components of the noded surface obtained by collapsing each curve of Γ to a point. Indeed, this completion coincides with the augmented Teichmüller space parametrizing noded Riemann surfaces marked by P (see [HK]). It is noncompact, coarsely negatively curved, and quasi-isometric to the pants complex; for an extensive overview, see [Wol].

Thurston compactification. For $\gamma \in \mathcal{S}_P$ and $\tau \in \mathcal{T}_P$ let $\ell_\gamma(\tau)$ denote the length of the unique geodesic in the hyperbolic surface associated to τ . The map $\tau \mapsto (\ell_\gamma(\tau))_{\gamma \in \mathcal{S}_P}$ defines an embedding $\mathcal{T}_P \rightarrow \mathbb{R}_{\geq 0}^{\mathcal{S}_P}$ which projects to an embedding $\mathcal{T}_P \rightarrow \mathbb{P}\mathbb{R}_{\geq 0}^{\mathcal{S}_P}$. Thurston showed that the closure of the image is homeomorphic to the closed ball $\overline{B}^{\#P-3}$, and that the boundary points may be identified with *projective measured foliations* on (S^2, P) . A comprehensive reference is [FLP]; cf. also the book by Ivanov [Iva1].

Fix a basepoint $\tau_\otimes \in \mathcal{T}_P$; this gives rise to a basepoint $m_\otimes := \pi(\tau_\otimes) \in \mathcal{M}_P$. Recall that we then have an identification $G_P \leftrightarrow \pi_1(\mathcal{M}_P, m_\otimes)$. The following folklore theorem is well known.

Theorem 2.1 (Fivefold way). *There are natural bijections between the following sets of objects:*

- (1) multicurves Γ on $S^2 \setminus P$;
- (2) “purely atomic” measured foliations $\mathcal{F}(\Gamma) := \sum_{\gamma \in \Gamma} \nu_\gamma$, where ν_γ is the delta-mass at γ , which counts the number of intersections of a curve with γ ;
- (3) complete purely parabolic subgroups L of G_P ;
- (4) strata $\mathcal{T}_P^\Gamma \subseteq \overline{\mathcal{T}}_P$;
- (5) certain subgroups of loops in moduli space (thought of as generated by certain pure braids) and corresponding via “pushing” to complete purely parabolic subgroups L of $\pi_1(\mathcal{M}_P, m_\otimes)$.

The subgroups arising in (5) will be described shortly. The bijections are given as follows:

- (1) \rightarrow (2) Take the zero measured foliation if Γ is empty. Otherwise, take a foliation of a regular neighborhood of Γ with the width (transverse measure) of

each neighborhood equal to 1; this gives the so-called *normal form* of a measured foliation \mathcal{F} obtained by collapsing the unfoliated regions onto their spines ([FLP, §6.5]). The result is unique, up to the natural Whitehead equivalence on foliations. Alternatively, choosing a complex structure on $S^2 \setminus P$. One may apply a well-known result of Jenkins and Strebel (see e.g. [DH, Prop. 4.2]) to obtain a corresponding quadratic differential whose closed horizontal trajectories provide the associated measured foliation.

(2) \rightarrow (1) If the foliation is the zero foliation, take $\Gamma = o$, the empty multicurve; otherwise, take one core curve from each cylinder in the normal form of the foliation.

(1) \rightarrow (3) Take the subgroup $L := \text{Tw}(\Gamma)$ generated by Dehn twists about elements of Γ .

(3) \rightarrow (1) Take the union Γ of the core curves of the representing twists; this is well defined.

(1) \rightarrow (4) We take those marked noded spheres in which precisely all the curves comprising Γ correspond to nodes, and to nothing else.

(3) \rightarrow (4) The stratum \mathcal{T}_P^Γ is the unique stratum in $\text{Fix}(L)$ of maximum dimension.

(4) \rightarrow (3) Given the stratum \mathcal{T}_P^Γ , take the pointwise stabilizer $L := \text{Tw}(\Gamma) = \text{Stab}_{G_P}(\mathcal{T}_P^\Gamma)$.

(3) \rightarrow (5) We now describe this correspondence.

Let $P := \{p_1, \dots, p_n\} \subseteq S^2$, $n := \#P \geq 4$. Recall that the *configuration space* $\text{Config}(S^2, P)$ is the space of injections $P \hookrightarrow S^2$; the inclusion $P \subseteq S^2$ gives a natural basepoint. Let $\pi : \text{Config}(S^2, P) \rightarrow \mathcal{M}_P$ be the natural projection and π_* the induced map on fundamental groups. Let Γ be a nonempty multicurve represented by curves $\gamma \subseteq S^2 \setminus P$. Below, we describe a procedure for producing, for each $\gamma \in \Gamma$, a loop ℓ_γ in $\text{Config}(S^2, P)$ based at P which projects to the right Dehn twist $\text{Tw}(\gamma)$. This procedure will have the additional property that as elements of $\pi_1(\text{Config}(S^2, P))$, given $\gamma_1, \gamma_2 \in \Gamma$, the elements represented by $\ell_{\gamma_1}, \ell_{\gamma_2}$ will commute.

The idea is based on the simultaneous “pushing” (or Birman spin) of points; see Figure 1. Suppose $z_1, \dots, z_m \in \mathbb{D} := \{|z| < 1\}$ are nonzero complex numbers, pick r with $\max_i \{|z_i|\} < r < 1$, set $z_0 = 0$, and consider the motion $t \mapsto z_i(t) := \exp(2\pi it)z_i, t \in [0, 1], i = 0, \dots, m$. This motion extends to an isotopy of $\overline{\mathbb{D}}$ fixing the origin and the boundary. Taking the result of this extension when $t = 1$, the resulting “multi-spin” homeomorphism of the plane is homotopic, through homeomorphisms fixing z_0, z_1, \dots, z_m , to the right-hand Dehn twist about the circle $\gamma := \{|z| = r\}$ surrounding the z_i ’s. To see this, note that the extension of each individual motion $z_i(t)$ yields a “spin” which is the composition of a left Dehn twist about the left-hand component of a regular neighborhood of the circle traced by $z_i(t)$ with the right Dehn twist about the corresponding right-hand component. The left twist resulting from the motion of $z_1(t)$ is trivial, since this left boundary component is peripheral, and for $i = 1, \dots, m - 1$ the right twist from the motion of z_i cancels the left twist from the motion of z_{i+1} .

Now suppose Γ is a nonempty multicurve. Choose an element $\delta \in \Gamma$ among the possibly several components of Γ such that δ does not separate a pair of elements of Γ . Let V be a component of $S^2 \setminus \delta$ whose closure contains Γ (the component V is unique if $\#\Gamma > 1$). Pick $\gamma \in \Gamma$. Then γ bounds a distinguished Jordan domain $D_\gamma \subseteq V$. Let $\{p_{j_0}, p_{j_1}, \dots, p_{j_m}\} = D_\gamma \cap P$ where $1 \leq j_0 < j_1 < \dots < j_m \leq n$, so




FIGURE 1. Simultaneously pushing the points labelled 4 and 5 around the point labelled 3 yields the right Dehn twist about the bold curve on the outer side of the thin curve passing through point 5.

that j_0 is the smallest index of an element of P in D_γ . Up to postcomposition with rotations about the origin, there is a unique Riemann map $\phi : (D_\gamma, p_{j_0}) \rightarrow (\mathbb{D}, 0)$. Set $z_i = \phi(p_{j_i}), i = 0, \dots, m$, and transport the motion of the z_i constructed in the previous paragraph to a motion of $\{p_{i_0}, p_{i_1}, \dots, p_{i_m}\}$ in S^2 ; it is supported in the interior of D_γ . This motion gives a loop ℓ_γ in the space $\text{Config}(S^2, P)$.

By construction, ℓ_γ projects to the right Dehn twist about γ in the pure mapping class group. Suppose γ_1, γ_2 are distinct elements of Γ . If $D_{\gamma_1} \cap D_{\gamma_2} = \emptyset$, then clearly the elements represented by ℓ_{γ_1} and ℓ_{γ_2} commute. Otherwise, we may assume $D_{\gamma_1} \subseteq D_{\gamma_2}$. By construction, the loop ℓ_{γ_2} , thought of as a motion of points in the disk D_{γ_2} , represents the central element in the braid group on $\#P \cap D_{\gamma_2}$ strands, so it commutes with the element represented by ℓ_{γ_1} .

Remark. By choosing arbitrarily an element $p_\infty \in P \cap (S^2 - \overline{D_\gamma})$ and upon identifying $S^2 - \{p_\infty\}$ with the plane \mathbb{R}^2 , we may view the paths constructed above as loops in the configuration space $\text{Config}(\mathbb{R}^2, P - \{p_\infty\})$ whose fundamental group is then isomorphic to the pure braid group PB_{n-1} .

3. PULLBACK INVARIANTS

Many of the objects we are concerned with arise nondynamically. We first define them in the nondynamical setting (§3.1), and then consider the same objects in the dynamical case (§3.2).

3.1. No dynamics.

Admissible covers. Suppose $A, B \subseteq S^2$ are finite sets, each containing at least three points. A branched covering $f : (S^2, A) \rightarrow (S^2, B)$ is *admissible* if (i) $B \supseteq V_f$, the set of branch values of f , and (ii) $f(A) \subseteq B$; we do not require $A = f^{-1}(B)$.

For the remainder of this subsection, we fix an admissible cover $f : (S^2, A) \rightarrow (S^2, B)$. To (f, A, B) we associate the following objects. Though they depend upon all three elements of the triple, we indicate this dependency by referring only to f , for brevity.

Pullback relation on curves. The *pullback relation*

$$\mathcal{S}_B \cup \{o\} \xleftarrow{f} \mathcal{S}_A \cup \{o\}$$

is defined by setting $o \xleftarrow{f} o$ and $\gamma_1 \xleftarrow{f} \gamma_2$ if and only if γ_2 is homotopic in $S^2 \setminus A$ to a connected component of the preimage of $\gamma_1 \subseteq S^2 \setminus B$ under f . Thus $\gamma \xleftarrow{f} o$ if and only if some preimage of γ is inessential or peripheral in $S^2 \setminus A$. The pullback relation induces a *pullback function* $f^{-1} : \mathcal{MS}_B \rightarrow \mathcal{MS}_A$ by sending the empty multicurve on (S^2, A) to the empty multicurve on (S^2, B) and $\Gamma \mapsto f^{-1}(\Gamma) := \{\tilde{\gamma} \in \mathcal{S}_A : \exists \gamma \in \Gamma, \gamma \xleftarrow{f} \tilde{\gamma}\}$; note that $f^{-1}(\Gamma)$ might be empty.

The associated linear transformation. There is a linear transformation

$$\lambda_f : \mathbb{R}[\mathcal{S}_B] \rightarrow \mathbb{R}[\mathcal{S}_A]$$

defined on basis vectors by

$$\lambda_f(\gamma) = \sum_{\gamma \xleftarrow{f} \gamma_i} d_i \gamma_i, \quad \text{where } d_i := \sum_{f^{-1}(\gamma) \supset \delta \simeq \gamma_i} \frac{1}{\deg(f : \delta \rightarrow \gamma)}.$$

The pullback map. There is a map $\sigma_f : \mathcal{T}_B \rightarrow \mathcal{T}_A$ associated to an admissible cover $f : (S^2, A) \rightarrow (S^2, B)$. Let $\varphi : (S^2, B) \rightarrow (\mathbb{P}^1, \varphi(B))$ be an orientation-preserving homeomorphism. Then there is an orientation-preserving homeomorphism $\psi : (S^2, A) \rightarrow (\mathbb{P}^1, \psi(A))$ and a rational map $F : (\mathbb{P}^1, \psi(A)) \rightarrow (\mathbb{P}^1, \varphi(B))$ so that the following diagram commutes:

$$\begin{CD} (S^2, A) @>\psi>> (\mathbb{P}^1, \psi(A)) \\ @VfVV @V F VV \\ (A^2, B) @>\varphi>> (\mathbb{P}^1, \varphi(B)) \end{CD}$$

We set $\sigma_f : [\varphi] \mapsto [\psi]$. It is well known that this map is well defined, holomorphic, and distance-nonincreasing for the corresponding Teichmüller metrics on the domain and range [DH].

The virtual homomorphism. In dynamical settings, the virtual endomorphism was first introduced in [Kam, §4]; cf. also [BN1].

Proposition 3.1. *Let $f : (S^2, A) \rightarrow (S^2, B)$ be an admissible branched covering. Then the subset*

$$H_f := \{[h] \in G_B : \exists \tilde{h}, h \circ f = f \circ \tilde{h}, \text{ and } \tilde{h}|_A = \text{id}_A\}$$

is a finite-index subgroup of G_B , and the function

$$\phi_f : H_f \rightarrow G_A, [h] \mapsto [\tilde{h}]$$

induced by lifting representatives is well defined and a homomorphism.

Proof. The proof is complicated by the possibility that the associated unramified covering map $f : (S^2, f^{-1}(B)) \rightarrow (S^2, B)$ might admit deck transformations.

We first set up some notation that will be needed later. Let $\mathcal{H}_B := \text{Homeo}^+(S^2, B)$ be the group of orientation-preserving homeomorphisms $(S^2, B) \rightarrow (S^2, B)$ which fix B pointwise, and define $\mathcal{H}_A := \text{Homeo}^+(S^2, A)$ analogously. Let $\mathcal{H}_{\{f^{-1}(B)\}}$ denote the group of orientation-preserving homeomorphisms $(S^2, f^{-1}(B)) \rightarrow (S^2, f^{-1}(B))$ that send the set $f^{-1}(B)$ to itself, possibly via a nontrivial permutation.

Next, we will make use of the following fact from algebra. If A, B, C are groups with $A < B$, $n := [A : B] < \infty$, and $p : B \rightarrow C$ is a surjective homomorphism, then the image $p(A)$ has finite index in C . To see this, write $B = b_1A \sqcup \dots \sqcup b_nA$ and apply the homomorphism p . We will also use the fact that the intersection of a finite collection of finite-index subgroups is again finite-index.

Let $\mathcal{Q} < \mathcal{H}_B \times \mathcal{H}_{\{f^{-1}(B)\}}$ be the subgroup of those ordered pairs of homeomorphisms (h, \tilde{h}) such that $h \circ f = f \circ \tilde{h}$. Let Lift_f denote the image of \mathcal{Q} under the projection to \mathcal{H}_B .


The group \mathcal{Q} acts on the set $f^{-1}(B)$. The intersection

$$\mathcal{Q}_A := \bigcap_{a \in A} \mathcal{Q}_a$$

of the stabilizers $\mathcal{Q}_a, a \in A$, is the subgroup of \mathcal{Q} consisting of pairs (h, \tilde{h}) for which $h \circ f = f \circ \tilde{h}$ and for which the upstairs map \tilde{h} fixes A pointwise. Thus $[\mathcal{Q} : \mathcal{Q}_A] < \infty$. Let $\text{Lift}_{f,A}$ denote the image of \mathcal{Q}_A under the projection to \mathcal{H}_B .

Note that both Lift_f and $\text{Lift}_{f,A}$ are “saturated” with respect to homotopy; that is, for $h \in \text{Lift}_f, g \in \text{Lift}_{f,A}$, and for all $\iota_0 \in \mathcal{H}_B$ isotopic to the identity relative to B , we have $\iota_0 \circ h \in \text{Lift}_f$ and $\iota_0 \circ g \in \text{Lift}_{f,A}$. This follows from the homotopy lifting property for the covering map $f : (S^2, f^{-1}(B)) \rightarrow (S^2, B)$. Indeed, there is a homeomorphism $\iota : (S^2, f^{-1}(B)) \rightarrow (S^2, f^{-1}(B))$ which is isotopic to the identity $(S^2, f^{-1}(B)) \rightarrow (S^2, f^{-1}(B))$ relative to $f^{-1}(B)$, so that $\iota_0 \circ f = f \circ \iota$. Clearly, $(\iota_0 \circ h, \iota_0 \circ \tilde{h}) \in \mathcal{Q}$ and $(\iota_0 \circ g, \iota_0 \circ \tilde{g}) \in \mathcal{Q}_A$ (where \tilde{h} and \tilde{g} are the respective lifts of h and g); therefore $\iota_0 \circ h \in \text{Lift}_f$ and $\iota_0 \circ g \in \text{Lift}_{f,A}$.

Now consider the following commutative diagram:



The right-pointing horizontal arrows are inclusions. The left-pointing horizontal arrow at top left is the composition of (i) projection onto the second factor $\mathcal{H}_{\{f^{-1}(B)\}}$ (by construction, this yields an element of \mathcal{H}_A), with (ii) recording the isotopy class (yielding an element of G_A). The vertical arrows in the top row are projections onto the first factor \mathcal{H}_B . The vertical arrows in the bottom row record the isotopy class as an element of G_B . Thus the vertical arrows are surjective by the definition of their images.

Since $[\mathcal{Q} : \mathcal{Q}_A] < \infty$ we conclude using the above algebraic fact that $[\text{Lift}_f/\text{Homeo}_0^+(S^2, B) : H_f] < \infty$. In the remainder of this paragraph, we prove that $[G_B : \text{Lift}_f/\text{Homeo}_0^+(S^2, B)] < \infty$, from which it follows that $[G_B : H_f]$ is finite. Indeed, let $d := \deg f$, and consider all subgroups of $\pi_1(S^2, B)$ (up to conjugacy) of index d . Since $\pi_1(S^2, B)$ is finitely generated, there are finitely many such subgroups and hence finitely many such conjugacy classes. The restriction $f : S^2 \setminus f^{-1}(B) \rightarrow S^2 \setminus B$ is a covering map, and so determines a conjugacy class ξ of subgroups of index d in $\pi_1(S^2, B)$ corresponding to loops in $S^2 \setminus B$ that lift to loops in $S^2 \setminus f^{-1}(B)$. A homeomorphism $h \in \mathcal{H}_B$ determines an outer automorphism of $\pi_1(S^2, B)$, so we obtain a homomorphism $G_B \rightarrow \text{Out}(\pi_1(S^2, B))$ and a corresponding action of G_B on the finite set of conjugacy classes of subgroups of index d . The stabilizer $(G_B)_\xi$ of ξ is therefore a finite index subgroup of G_B . From elementary covering space theory, a sufficient (and, actually, necessary) condition for a homeomorphism $h \in \mathcal{H}_B$ to lie in the subgroup Lift_f is that the class $[h]$ of h in G_B lie in the stabilizer $(G_B)_\xi$. We conclude that $\text{Lift}_f/\text{Homeo}_0^+(S^2, B) = (G_B)_\xi$ has finite index in G_B .

Next, we show that the vertical arrow labelled $*$ is an isomorphism. It is surjective by definition. By definition, an element of the kernel is a pair of homeomorphisms of the form $(\text{id}_{S^2}, \tilde{h})$ with $f = f \circ \tilde{h}$ (so that \tilde{h} is a deck transformation of the covering $f : S^2 \setminus f^{-1}(B) \rightarrow S^2 \setminus B$) and with $\tilde{h}|_A = \text{id}_A$. We claim that this implies $\tilde{h} = \text{id}_{S^2}$. Since nontrivial deck transformations have no fixed points, we will show that \tilde{h} has a fixed point. We do this by applying the Lefschetz fixed point formula. To deal with compact spaces, let $U = S^2 \setminus N_\varepsilon(B)$ (where $N_\varepsilon(B)$ is an ε -neighborhood of B) and $\tilde{U} := f^{-1}(U)$; then $\tilde{h}|_{\tilde{U}}$ is a deck transformation of the covering of compact spaces $f : \tilde{U} \rightarrow U$. Since $\#A \geq 3$ and by assumption \tilde{h} fixes each element of A , we have that the trace of \tilde{h}_* on $H_1(\tilde{U}, \mathbb{Z})$ is at least 2. By the Lefschetz fixed point formula, this implies \tilde{h} has a fixed point.

Next, we define the virtual homomorphism ϕ_f . Its domain is the subgroup H_f . Given a class $\theta \in H_f$, to define $\phi_f(\theta) \in G_A$ choose a representing homeomorphism $h \in \text{Lift}_{f,A}$. Since the vertical arrow ($*$) is an isomorphism, there is a unique lift \tilde{h} of h under f with the property that it fixes A pointwise. We define $\phi_f(\theta) := [\tilde{h}] \in G_A$.

We now show in two ways that ϕ_f is well defined. First: given two pairs $(h_1, \tilde{h}_1), (h_2, \tilde{h}_2) \in \mathcal{Q}_A$ for which $[h_1] = [h_2]$ in G_B , we must show $[\tilde{h}_1] = [\tilde{h}_2] \in G_A$. Consider the product $(g, \tilde{g}) := (h_2^{-1} \circ h_1, \tilde{h}_2^{-1} \circ \tilde{h}_1)$. Since up to isotopy $f \circ \tilde{g} = f$, we have

$$\sigma_{f \circ \tilde{g}} = \sigma_f \quad \text{as maps } \mathcal{T}_B \rightarrow \mathcal{T}_A.$$

So for all $\tau \in \mathcal{T}_B$,

$$\sigma_{\tilde{g}}(\sigma_f(\tau)) = \sigma_f(\tau),$$

which implies that $\sigma_{\tilde{g}}$ has a fixed point. The action of $\sigma_{\tilde{g}}$ on \mathcal{T}_A depends only on the class $[\tilde{g}] \in G_A$, and this action is via a deck transformation of the covering $\mathcal{T}_A \rightarrow \mathcal{M}_A$. Again using the fact that a deck transformation with a fixed point is the identity, this implies $[\tilde{g}] = \text{id} \in G_A$. Second: an isotopy connecting h_1 and h_2 , which is constant and equal to the identity on B , lifts by f to an isotopy which is constant on $f^{-1}(B)$, and which connects \tilde{h}_1 to a lift \tilde{h}'_2 of h_2 . Since \tilde{h}_1 fixes $A \subseteq f^{-1}(B)$ pointwise, so does \tilde{h}'_2 . We have shown that such a lift is unique, and hence $\tilde{h}'_2 = \tilde{h}_2$ and $[\tilde{h}_1] = [\tilde{h}_2] \in G_A$.

Finally, we claim that ϕ_f is a homomorphism. This follows from the definition of ϕ_f and the fact that $\text{Lift}_{f,A}$ is a group under coordinatewise composition. \square

Restricted virtual homomorphism and other definitions. Here, we briefly comment on the difference between the definition of the virtual homomorphism given here and that of the virtual endomorphism given in [Pil3].

In [Pil3], the discussion treats only the dynamical setting of Thurston maps, and the corresponding virtual endomorphism is defined differently. Here is the connection.

Given an admissible cover $f : (S^2, A) \rightarrow (S^2, B)$, the *restricted* virtual homomorphism, $\phi'_f : G_B \dashrightarrow G_A$ is defined as follows. Set $A' = f^{-1}(B)$, and let $\phi''_f : G_B \rightarrow G_{A'}$ be the virtual homomorphism given by Proposition 3.1; denote its domain H'_f . The restricted virtual homomorphism ϕ'_f is defined as the composition

$$H'_f \xrightarrow{\phi''_f} G_{A'} \rightarrow G_A,$$

where $G_{A'} \rightarrow G_A$ is the map induced by forgetting points in $A' \setminus A$.

Now suppose $A = B = P \supset P_f$, where f is a Thurston map. The virtual endomorphism $G_P \dashrightarrow G_P$ of [Pil3] coincides with the restricted virtual endomorphism $\phi'_f : G_P \dashrightarrow G_P$.

The virtual homomorphism changes in a predictable way under pre- and post-composition by homeomorphisms.

Lemma 3.2. *Let $f : (S^2, A) \rightarrow (S^2, B)$ be an admissible covering map, let $i : (S^2, B) \rightarrow (S^2, B)$ be an orientation-preserving homeomorphism which maps the set B to itself (not necessarily pointwise), and let $j : (S^2, A) \rightarrow (S^2, A)$ be an orientation-preserving homeomorphism which maps the set A to itself (not necessarily pointwise). Then $i \circ f \circ j : (S^2, A) \rightarrow (S^2, B)$ is an admissible cover with associated virtual homomorphism*

$$\phi_{i \circ f \circ j} : G_B \dashrightarrow G_A \text{ with domain } H_{i \circ f \circ j},$$

and

$$H_{i \circ f \circ j} = i \circ H_f \circ i^{-1} \quad \text{and} \quad \phi_{i \circ f \circ j}(h) = j^{-1} \circ \phi_f(i^{-1} \circ h \circ i) \circ j, \quad h \in H_{i \circ f \circ j}.$$

Proof. This follows immediately from the definitions. \square

Functional identities. The following are two fundamental functional identities relating the virtual homomorphism, the linear map, and the pullback map.

We begin with some notation. If $\Gamma = \{\gamma_1, \dots, \gamma_m\}$ is a multicurve and $a_1, \dots, a_m \in \mathbb{Z}$, we set

$$w := \sum_{i=1}^m a_i \gamma_i \quad \text{and} \quad M_w := T_1^{a_1} \cdots T_m^{a_m},$$

where T_i is the right Dehn twist about γ_i . This is well defined since the T_j 's commute pairwise.

The result [Pil3, Thm. 1.2], phrased and proved for Thurston maps, generalizes in a completely straightforward way to the following result; note that the more restrictive domain is referred to below.

Theorem 3.3. *For any multitwist $M_w \in \text{dom}(\phi'_f)$, we have $\lambda_f(w) \in \mathbb{Z}[\mathcal{S}_A]$, and*

$$(1) \quad \phi'_f(M_w) = \phi_f(M_w) = M_{\lambda_f(w)}.$$

From the definitions (see also [BEKP] and [Koc2]) we have

$$(2) \quad \sigma_f(h \cdot \tau) = \phi_f(h) \cdot \sigma_f(\tau), \quad \forall h \in H_f.$$

The Hurwitz space \mathcal{W}_f . This discussion is extracted from [Koc2]. Consider the space

$$\text{Rat}_d \times (\mathbb{P}^1)^B \times (\mathbb{P}^1)^A,$$

where Rat_d denotes the space of rational maps of degree d , $(\mathbb{P}^1)^B$ denotes the space of all *injective* maps $\varphi : B \hookrightarrow \mathbb{P}^1$, and $(\mathbb{P}^1)^A$ denotes the space of all *injective* maps $\varphi : A \hookrightarrow \mathbb{P}^1$ (we are abusing notation). This space is a smooth affine variety.

The group $\text{Aut}(\mathbb{P}^1) \times \text{Aut}(\mathbb{P}^1)$ acts on the space in the following way:

$$(\mu, \nu) \cdot (F, i_B, j_A) \mapsto (\mu \circ F \circ \nu^{-1}, \mu \circ i_B, \nu \circ j_A).$$

This action is free, and we consider the geometric quotient

$$\left(\text{Rat}_d \times (\mathbb{P}^1)^B \times (\mathbb{P}^1)^A \right) / (\text{Aut}(\mathbb{P}^1) \times \text{Aut}(\mathbb{P}^1)).$$

It follows from geometric invariant theory that this is a complex manifold of dimension $2d + 1 + \#A + \#B - 6$ (for details, see [Koc2]).

Consider the map

$$\omega_f : \mathcal{T}_B \rightarrow \left(\text{Rat}_d \times (\mathbb{P}^1)^B \times (\mathbb{P}^1)^A \right) / (\text{Aut}(\mathbb{P}^1) \times \text{Aut}(\mathbb{P}^1))$$

given by $\omega_f : [\phi] \mapsto [F, \phi|_B, \psi|_A]$, where

- $\phi : (S^2, B) \rightarrow (\mathbb{P}^1, \phi(B))$ and $\psi : (S^2, A) \rightarrow (\mathbb{P}^1, \psi(A))$ are orientation-preserving homeomorphisms and
- $F := \phi \circ f \circ \psi^{-1} : (\mathbb{P}^1, \psi(A)) \rightarrow (\mathbb{P}^1, \phi(B))$ is the rational map at right in the diagram defining the pullback map σ_f .

In other words, ω_f records the algebraic data of σ_f , up to appropriate equivalence.

Following [Koc2], the *Hurwitz space* associated to $f : (S^2, A) \rightarrow (S^2, B)$ is $\mathcal{W}_f := \omega_f(\mathcal{T}_B)$. It is a complex manifold of dimension $\#B - 3$, and the map $\omega_f : \mathcal{T}_B \rightarrow \mathcal{W}_f$ is a holomorphic covering map.

The moduli space correspondence. The Hurwitz space \mathcal{W}_f associated to $f : (S^2, A) \rightarrow (S^2, B)$ is equipped with two holomorphic maps (see Figure 2):

$$X : \mathcal{W}_f \rightarrow \mathcal{M}_A \quad \text{given by} \quad [F, \phi|_B, \psi|_A] \mapsto [\psi|_A] \quad \text{and}$$

$$Y : \mathcal{W}_f \rightarrow \mathcal{M}_B \quad \text{given by} \quad [F, \phi|_B, \psi|_A] \mapsto [\phi|_B].$$




FIGURE 2. The fundamental diagram associated to any admissible covering map $f : (S^2, A) \rightarrow (S^2, B)$; this diagram commutes.

We may think of these as comprising one multivalued map $X \circ Y^{-1} : \mathcal{M}_B \rightrightarrows \mathcal{M}_A$. Note that the direction of this correspondence is the same as that of $\sigma_f : \mathcal{T}_B \rightarrow \mathcal{T}_A$, i.e. it corresponds to pulling-back complex structures. We call the pair of maps

$$(X : \mathcal{W}_f \rightarrow \mathcal{M}_A, \quad Y : \mathcal{W}_f \rightarrow \mathcal{M}_B)$$

the *moduli space correspondence* associated with $f : (S^2, A) \rightarrow (S^2, B)$. The following results are proved in [Koc2].

Proposition 3.4. *The map π_B factors as $\pi_B = Y \circ \omega_f$, and the space \mathcal{W}_f is isomorphic (as a complex manifold) to \mathcal{T}_B/H_f .*

Thus $\deg(Y) = [G_P : H_f]$, the index of $\text{dom}(\phi_f)$ in the pure mapping class group.

Corollary 3.5. *The map $Y : \mathcal{W}_f \rightarrow \mathcal{M}_B$ is a finite covering map.*

Proof. This follows from Proposition 3.1. □

On the one hand, the map $Y : \mathcal{W}_f \rightarrow \mathcal{M}_B$ is always a covering map. On the other hand, the map $X : \mathcal{W}_f \rightarrow \mathcal{M}_A$ may have diverse properties: it can be e.g. constant, injective but not surjective, or a surjective ramified covering; see [BEKP].

We say that admissible covers $f : (S^2, A) \rightarrow (S^2, B)$ and $g : (S^2, A) \rightarrow (S^2, B)$ are (A, B) -Hurwitz equivalent if there are homeomorphisms $h \in \mathcal{H}_B$, and $\tilde{h} \in \mathcal{H}_A$ so that $h \circ f = g \circ \tilde{h}$.

While σ_f depends on the admissible cover $f : (S^2, A) \rightarrow (S^2, B)$, the correspondence on moduli space depends only on the (A, B) -Hurwitz class of f .

Proposition 3.6. *The Hurwitz space \mathcal{W}_f is equal to the Hurwitz space \mathcal{W}_g if and only if f and g are (A, B) -Hurwitz equivalent.*

Remark. The classical notion of Hurwitz equivalence requires only that $h \circ f = g \circ \tilde{h}$ for some pair of orientation-preserving homeomorphisms h, \tilde{h} which are not required to fix any finite set pointwise. Thus, (A, B) -Hurwitz equivalence is a finer equivalence relation than classical Hurwitz equivalence. For example, any pair of quadratic maps f, g are Hurwitz equivalent, but it is possible to arrange a suitable choice of $A = B = P_f = P_g$ so that f, g are not (A, B) -Hurwitz equivalent; see [Koc1, Remark 7.3.3] and [Koc2, Example 2.9].

Two definitions of the virtual homomorphism. Choose a basepoint $w_\otimes \in \mathcal{W}_f$ and set $\mu_\otimes := X(w_\otimes), \tilde{\mu}_\otimes := Y(w_\otimes)$. Using the identity maps to define corresponding basepoints for $\mathcal{T}_A, \mathcal{T}_B$, we obtain identifications (see [Lod, Section 2.2]) $G_A \leftrightarrow \pi_1(\mathcal{M}_A, \tilde{\mu}_\otimes)$ and $G_B \leftrightarrow \pi_1(\mathcal{M}_B, \mu_\otimes)$. It can be shown (cf. [Lod, Theorem 2.6]) that the virtual homomorphism $\phi_f : G_B \dashrightarrow G_A$, under these identifications, coincides with the induced maps $X_* \circ (Y_*)^{-1}$ on fundamental groups.

Augmented Teichmüller space and the extension of σ_f . The proof of [Sel1, Prop. 4.3] shows that σ_f is uniformly Lipschitz with respect to the WP metric, and therefore it extends to the WP completion. The action of the deck group corresponding to the covering map $\pi_B : \mathcal{T}_B \rightarrow \mathcal{M}_B$ (respectively $\pi_A : \mathcal{T}_A \rightarrow \mathcal{M}_A$)

extends to the space $\overline{\mathcal{T}}_B$ (respectively $\overline{\mathcal{T}}_A$). The fundamental diagram becomes

$$(3) \quad \begin{array}{ccc} \overline{\mathcal{T}}_B & \xrightarrow{\sigma_f} & \overline{\mathcal{T}}_A \\ \pi_B \downarrow & \searrow \omega_f & \downarrow \pi_A \\ & \overline{\mathcal{W}}_f & \\ Y \swarrow & & \searrow X \\ \overline{\mathcal{M}}_B & & \overline{\mathcal{M}}_A \end{array}$$

where

$$\overline{\mathcal{W}}_f := \overline{\mathcal{T}}_B/H_f, \quad \overline{\mathcal{M}}_B := \overline{\mathcal{T}}_B/G_B, \quad \text{and} \quad \overline{\mathcal{M}}_A := \overline{\mathcal{T}}_A/G_A.$$

The proof of [Sel1, Prop. 6.1] shows

$$(4) \quad \sigma_f(\mathcal{T}_B^\Gamma) \subseteq \mathcal{T}_A^{f^{-1}(\Gamma)}.$$

3.2. Dynamics. We briefly mention the relevant objects in the dynamical setting.

Thurston maps. We will often be concerned with the special case where $A = B = P \supset P_f$. We will require the notion of the orbifold associated to a Thurston map $f : (S^2, P) \rightarrow (S^2, P)$. Following [DH], we note that there is a smallest function ν_f over all functions $\nu : S^2 \rightarrow \mathbb{N} \cup \{\infty\}$ such that

- $\nu(x) = 1$ if $x \notin P$, and
- $\nu(x)$ is a multiple of $\nu(y) \cdot \deg(f, y)$ for all $y \in f^{-1}(x)$; here the second factor is the local degree of f at y .

Indeed, one simply sets $\nu(x)$ to be the least common multiple of the set of local degrees of iterates of f at all points that are iterated preimages of x . The orbifold of f is $O_f(S^2, \nu_f)$; it is *hyperbolic* if the Euler characteristic

$$\chi(O_f) = 2 - \sum_{x \in P} \left(1 - \frac{1}{\nu_f(x)} \right)$$

is negative. We say that f has *Euclidean orbifold* if it does not have hyperbolic orbifold. If $\#P_f \geq 5$, then the orbifold of f is necessarily hyperbolic. Maps with Euclidean orbifolds are classified in [DH].

Pullback relation on curves. Just as in the nondynamical setting, the pullback relation induces a *pullback function* $f^{-1} : \mathcal{MS}_P \rightarrow \mathcal{MS}_P$. A multicurve is *invariant* if $f^{-1}(\Gamma) \subseteq \Gamma$; it is *completely invariant* if $f^{-1}(\Gamma) = \Gamma$.

The Thurston linear transformation. The *Thurston linear transformation*

$$\lambda_f : \mathbb{R}[S_P] \rightarrow \mathbb{R}[S_P]$$

is defined just as before, with $A = B = P$.

There is one key difference in the dynamical setting, which is *Thurston’s Characterization Theorem* [DH, Theorem 1].

Theorem 3.7 (Thurston’s characterization). *If \mathcal{O}_f is hyperbolic, then f is equivalent to a rational map R if and only if for every invariant (equivalently, every) multicurve Γ , the spectrum of the linear map*

$$\lambda_{f,\Gamma} := \lambda_f|_{\mathbb{R}[\Gamma]} : \mathbb{R}[\Gamma] \rightarrow \mathbb{R}[\Gamma]$$

lies in the open unit disk. In this case, R is unique, up to conjugation by Möbius transformations.

If there is an invariant multicurve Γ for which the criterion in Theorem 3.7 fails, then Γ is called a *Thurston obstruction*.

The pullback map. The pullback map $\sigma_f : \mathcal{T}_P \rightarrow \mathcal{T}_P$ is defined as before.

When \mathcal{O}_f is hyperbolic, the criterion in Theorem 3.7 is equivalent to the existence of a fixed point of σ_f in \mathcal{T}_P . Some k th iterate σ_f^k is a strict, nonuniform contraction (when $P = P_f$, [DH, Prop. 3.3(b)] asserts that $k = 2$ will do). So if a fixed point exists, it is unique, the map f is equivalent to a rational map R (unique up to conjugacy by Möbius transformations), and the projection of the fixed point to moduli space corresponds to the geometry of the corresponding subset of the dynamical plane of R . When \mathcal{O}_f is Euclidean, the relationship between the existence of Thurston obstructions and the dynamics on Teichmüller space is more subtle; cf. [Sel2]. However, with the exception of the well-known *integral Lattés maps* induced by the endomorphisms $z \mapsto n \cdot z$ on complex tori, a fixed point of σ_f , if it exists, is unique.

The virtual endomorphism. There are a finite index subgroup $H_f < G_P$ and a virtual endomorphism

$$\phi_f : G_P \dashrightarrow G_P \quad \text{given by} \quad \phi_f : [h] \mapsto [h'],$$

where $h, h' \in \text{Homeo}^+(S^2, P)$, and $h \circ f = f \circ h'$.

The Hurwitz space \mathcal{W}_f . We have the same fundamental diagram as in Figure 2, with $A = B = P$, and we have maps

$$X : \mathcal{W}_f \rightarrow \mathcal{M}_P \quad \text{and} \quad Y : \mathcal{W}_f \rightarrow \mathcal{M}_P.$$

In the dynamical setting, we can iterate the moduli space correspondence $X \circ Y^{-1} : \mathcal{M}_P \rightrightarrows \mathcal{M}_P$.

Augmented Teichmüller space and the extension of σ_f . Considering diagram (3), the proof of [Sel1, Prop. 6.1] shows again that $\sigma_f(\mathcal{T}_P^\Gamma) \subseteq \mathcal{T}_P^{f^{-1}(\Gamma)}$. In particular, completely invariant multicurves (those satisfying $f^{-1}(\Gamma) = \Gamma$) correspond to strata invariant under σ_f . Selinger [Sel1, Thm. 10.4] shows that if f is obstructed, then under suitable hypotheses (all pieces in the corresponding canonical decomposition having hyperbolic orbifold), again there is a unique fixed point $\hat{\tau} \in \overline{\mathcal{T}}_P$ to which all iterates converge. However, there exist both obstructed and unobstructed maps f for which σ_f has periodic points in $\partial\mathcal{T}_P$. Indeed, understanding the dynamical behavior of $\sigma_f : \overline{\mathcal{T}}_P \rightarrow \overline{\mathcal{T}}_P$ is a main motivation for this work.

4. WHEN $\sigma_f : \partial\mathcal{T}_B \rightarrow \partial\mathcal{T}_A$

We return to a nondynamical discussion. The main result of this section is the following characterization of when the pullback map preserves the WP boundary.

Theorem 4.1. *Let $f : (S^2, A) \rightarrow (S^2, B)$ be an admissible cover. The following are equivalent:*

- (1) *For each nonempty $\Gamma \in \mathcal{MS}_B$, $f^{-1}(\Gamma) \neq \emptyset$.*
- (2) *For each $\gamma \in \mathcal{S}_B$, $\gamma \notin \ker(\lambda_f)$.*
- (3) *$\ker(\phi_f) \cap \text{Tw}^+ \subseteq G_B$ is trivial.*
- (4) *The pullback map σ_f on Teichmüller space satisfies $\sigma_f : \partial\mathcal{T}_B \rightarrow \partial\mathcal{T}_A$.*
- (5) *The pullback correspondence $X \circ Y^{-1} : \mathcal{M}_B \rightrightarrows \mathcal{M}_A$ on moduli space is proper.*

If $\#A = \#B$ and if any of the conditions above hold, the pullback correspondence $X \circ Y^{-1} : \mathcal{M}_B \rightrightarrows \mathcal{M}_A$ is surjective.

In (5), properness means that for each compact $K \subseteq \mathcal{M}_A$, the set $Y(X^{-1}(K)) \subseteq \mathcal{M}_B$ is compact. Since Y is a finite covering map, this is equivalent to the properness of the map X .

Proof. The equivalence of (1) and (2) is immediate from the definitions.

The equivalence of (2) and (3) follows immediately from the definitions, equation (1), the nonnegativity of λ_f , and the fact that ϕ_f preserves the positivity of twists. Indeed, from [Pil3, Thm. 1.2] we have the following. Suppose $\Gamma = \{\gamma_1, \dots, \gamma_n\}$ is a nonempty multicurve on $S^2 \setminus B$, $a_1, \dots, a_n \in \mathbb{Z}_{>0}$, $w := \sum a_i \gamma_i \in \mathbb{Z}_+[\mathcal{S}_B]$, and $M_w := T_1^{a_1} \cdot \dots \cdot T_n^{a_n}$ is the corresponding positive multitwist; here T_i is the right Dehn twist about γ_i . If $M_w \in H'_f$, then by equation (1),

$$\phi_f(M_w) = M_{\lambda_f(w)}.$$

So

$$\phi_f(M_w) = \text{id} \iff \lambda_f(w) = 0.$$

Since λ_f is a nonnegative linear operator and $a_i > 0$ for each i , we conclude from the definition of λ_f that

$$\phi_f(M_w) = \text{id} \iff f^{-1}(\Gamma) = o.$$

We now show (1) \iff (4). Recall that the WP boundary $\partial\mathcal{T}_B$ is a union of strata corresponding to nonempty multicurves on $S^2 \setminus B$. By equation (4)

$$\sigma_f(\mathcal{T}_B^\Gamma) \subseteq \mathcal{T}_A^{f^{-1}(\Gamma)}.$$

In particular,

$$\sigma_f(\mathcal{T}_B^\Gamma) \subseteq \partial\mathcal{T}_A \iff f^{-1}(\Gamma) \neq o,$$

which yields (1) \iff (4).

Now we show that failure of (4) implies failure of (5). Suppose for some nonempty multicurve Γ there exist $\tau_n \rightarrow \bar{\tau} \in \mathcal{T}_B^\Gamma$ while $\tilde{\tau}_n := \sigma_f(\tau_n) \rightarrow \tilde{\tau} := \sigma_f(\bar{\tau}) \in \mathcal{T}_A$. But then in moduli space $\mu_n := \pi_B(\tau_n) \rightarrow \infty$ while $\tilde{\mu}_n := \pi_A(\sigma_f(\tau_n)) \subseteq X \circ Y^{-1}(\mu_n)$ does not, showing that properness of the pullback correspondence fails.

Now suppose that (5) fails. Then there exist $\tilde{\mu}_n \rightarrow \tilde{\mu}$ in \mathcal{M}_A and $\mu_n \rightarrow \bar{\mu} \in \partial\mathcal{M}_B$ such that $\tilde{\mu}_n \in X \circ Y^{-1}(\mu_n)$. Consider diagram (3). There exists $\bar{w} \in \overline{\mathcal{W}}_f$ with $Y(\bar{w}) = \bar{\mu}$, $X(w) = \tilde{\mu}$. We see from diagram (3) that for any $\bar{\tau} \in \partial\mathcal{T}_B$ with $w_f(\bar{\tau}) = \bar{w}$, $\sigma_f(\bar{\tau}) \in \pi_A^{-1}(\tilde{\mu}) \in \mathcal{T}_A$ and (4) fails.

It remains to prove surjectivity of $X \circ Y^{-1}$ under the assumptions $\#A = \#B$ and that e.g. condition (5) holds. Since $X \circ Y^{-1}$ is proper, it is not constant.

Recall that \mathcal{M}_B is isomorphic to a hyperplane complement; in particular, it is a Stein manifold and, hence, the finite cover \mathcal{W}_f of \mathcal{M}_B also is a Stein manifold of the same dimension. Therefore, the preimage under X of any point is totally disconnected and, by properness, compact. Since X is analytic, we conclude that the preimage is discrete as well, and hence finite. Because $\#A = \#B$, the complex manifolds \mathcal{W}_f and \mathcal{M}_A have equal dimension, so the map $X : \mathcal{W}_f \rightarrow \mathcal{M}_A$ is open because it has discrete fibers. A continuous proper map between locally compact Hausdorff spaces is closed. Hence the image $X(\mathcal{W}_f)$ is both open and closed in \mathcal{M}_A , and so $X : \mathcal{W}_f \rightarrow \mathcal{M}_A$ is surjective. \square

5. WHEN $\sigma_f : \mathcal{T}_B \rightarrow \mathcal{T}_A$ IS CONSTANT

Theorem 5.1. *Let $f : (S^2, A) \rightarrow (S^2, B)$ be an admissible cover. The following are equivalent:*

- (1) \leftarrow^f is constant.
- (2) $\lambda_f : \mathbb{R}[\mathcal{S}_B] \rightarrow \mathbb{R}[\mathcal{S}_A]$ is constant.
- (3) $\phi_f : G_B \dashrightarrow G_A$ is constant.
- (4) $\sigma_f : \mathcal{T}_B \rightarrow \mathcal{T}_A$ is constant.
- (5) $X \circ Y^{-1} : \mathcal{M}_B \rightrightarrows \mathcal{M}_A$ is constant.

In [BEKP], the previous theorem is proved in the dynamical setting for f a Thurston map. The same ideas for the proof apply in the nondynamical case. However, in [BEKP], there is a mistake in the proof that (2) \Rightarrow (3). The assumption (2) is equivalent to the assumption that every curve, when lifted under f , becomes inessential or peripheral. Even if this holds, it need not be the case that every Dehn twist lifts under f to a pure mapping class element. We give an explicit example after the proof of Theorem 5.1.

Proof. In [BEKP] the logic was: (1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4), and failure of (1) implies failure of (4). Also, condition (1) is stated as “ \leftarrow^f is empty” and condition (5) is omitted.

Here is the revised logic:

- (4) \iff (5),
- (1) \iff (2),
- (3) \Rightarrow (2),
- (3) \iff (4), and
- failure of (4) implies failure of (1).

We immediately have that (4) \iff (5) from the fundamental diagram, Figure 2, and (1) \iff (2) follows immediately from the definitions.

To show that (3) \Rightarrow (2), we show failure of (2) implies failure of (3). If λ_f is not constant, then there exists a simple closed curve $\gamma \in \mathcal{S}_B$ which has an essential, nonperipheral simple closed curve $\delta \in \mathcal{S}_A$ as a preimage under f . Some power of the Dehn twist about γ lifts under f to a product of nontrivial Dehn twists. The hypothesis implies that the lifted map is homotopically nontrivial, so ϕ_f is not constant.

For the remaining implications, we will make use of the following facts. First recall the functional identity from (2),

$$\sigma_f(h \cdot \tau) = \phi_f(h) \cdot \sigma_f(\tau), \quad \forall h \in H_f,$$

recall that W_f is isomorphic (as a complex manifold) to \mathcal{T}_B/H_f (Proposition 3.4), and recall Corollary 3.5 which states that $Y : \mathcal{W}_f \rightarrow \mathcal{M}_B$ is a finite cover.

Second, note that a bounded holomorphic function on a finite cover of \mathcal{M}_B is constant. To see this, recall that \mathcal{M}_B is isomorphic to the complement of a finite set of hyperplanes in \mathbb{C}^n where $n = \#B - 3$. Let $L \in \mathbb{C}^n$ be any complex line which intersects \mathcal{M}_B . This intersection is isomorphic to a compact Riemann surface punctured at finitely many points. If \tilde{L} is any component of the preimage of L under a finite covering, then \tilde{L} is also isomorphic to a compact Riemann surface punctured at finitely many points. By Liouville’s theorem, the function is constant on \tilde{L} . Since L is arbitrary, the function is locally constant, and hence constant.

To establish (3) \Rightarrow (4), suppose (3) holds. Then $\sigma_f : \mathcal{T}_B \rightarrow \mathcal{T}_A$ descends to a holomorphic map

$$\bar{\sigma}_f : \mathcal{W}_f \rightarrow \mathcal{T}_A.$$

But it is well known (see e.g. [Hub]) that \mathcal{T}_A is isomorphic as a complex manifold to a bounded domain of \mathbb{C}^n , so the discussion above implies that σ_f is constant.

To establish that (4) \Rightarrow (3), suppose $h \in H_f$. If $\sigma_f \equiv \tau$ is constant, the deck transformation defined by $\phi_f(h)$ fixes the point τ , and hence must be the identity, so ϕ_f is constant.

To establish that not(4) \Rightarrow not(1), we first prove a lemma of perhaps independent interest. Below, by a bounded set we mean a set with compact closure.

Lemma 5.2. *Let $f : (S^2, A) \rightarrow (S^2, B)$ be an admissible covering map. Then the image of $\sigma_f : \mathcal{T}_B \rightarrow \mathcal{T}_A$ is either a point or unbounded in \mathcal{T}_A . In the latter case, actually $\pi_A(\sigma_f(\mathcal{T}_B))$ is unbounded in \mathcal{M}_A .*

Proof. The fundamental diagram in Figure 2 implies that

$$\begin{aligned} \sigma_f(\mathcal{T}_B) \text{ is bounded in } \mathcal{T}_A &\iff \pi_A(\sigma_f(\mathcal{T}_B)) \text{ is bounded in } \mathcal{M}_A \\ &\iff X(\mathcal{W}_f) \text{ is bounded in } \mathcal{M}_A. \end{aligned}$$

Because a bounded holomorphic function on \mathcal{W}_f is constant, $X(\mathcal{W}_f)$ is bounded in \mathcal{M}_A if and only if it is a single point, or (by Figure 2) if and only if the image of σ_f is a single point. □

Suppose now that $\sigma_f : \mathcal{T}_B \rightarrow \mathcal{T}_A$ is not constant (i.e., failure of (4)). Lemma 5.2 implies that $X(\mathcal{W}_f)$ is not contained in any compact subset of \mathcal{M}_A . Consider the diagram in equation (3). This means that the image $X(\overline{\mathcal{W}_f})$ contains at least one point on the boundary $\partial\mathcal{M}_A$, and hence $\sigma_f(\overline{\mathcal{T}_B})$ contains at least one point in $\partial\mathcal{T}_A$. We see that there exists a boundary stratum \mathcal{T}_B^Γ that is mapped to a boundary stratum of \mathcal{T}_A . Hence, $f^{-1}(\Gamma) \neq \emptyset$, which means \leftarrow^f is not constant.

Remark. One may prove the last implication in a more direct and elementary way by observing that if $X(\mathcal{W}_f)$ is unbounded, then there are points $\tilde{\tau} = \sigma_f(\tau)$ on whose underlying Riemann surfaces $X_{\tilde{\tau}}$ exist annuli of arbitrarily large modulus. Pushing such an annulus forward under the natural map $X_{\tilde{\tau}} \rightarrow X_\tau$ yields a fat annulus on X_τ . The core curves of these annuli must be nontrivial and provide a pair $\gamma \leftarrow^f \tilde{\gamma}$.

Example. Consider the postcritically finite rational map $f : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ defined by

$$f(z) = 2i \left(z^2 - \frac{1+i}{2} \right)^2,$$

with postcritical set $P = \{0, 1, -1, \infty\}$. It factors as $f = g \circ s$, where $s(z) = z^2$. Its mapping properties are shown in Figure 3. In this case, $\sigma_f : \mathcal{T}_P \rightarrow \mathcal{T}_P$ is constant (as proved in [BEKP]), $Y : \mathcal{W}_f \rightarrow \mathcal{M}_P$ is a degree 2 covering map, and $X : \mathcal{W}_f \rightarrow \mathcal{M}_P$ is constant. Because Y has degree 2, the subgroup H_f has index 2 in G_P , so there is a Dehn twist $h \in \text{Homeo}^+(S^2, B)$ that does not lift to a homeomorphism $h' \in \text{Homeo}^+(S^2, A)$. Thus even if $\sigma_f : \mathcal{T}_P \rightarrow \mathcal{T}_P$ is constant, the index $[G_P : H_f]$ may be greater than 1.

One may see this directly. Let γ_0 be the boundary of a small regular neighborhood D of the segment $[0, 1] \subseteq \mathbb{C}$ and let $h_0 : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be the right Dehn twist about γ_0 .

Claim. If $h_1 : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ satisfies $h_0 \circ f = f \circ h_1$ (i.e. h_1 is a lift of h_0 under f), then $h_1 \notin G_P$.




FIGURE 3. The mapping properties of f . The points in grey are $-1, 0, +1$. The annulus A_0 is shown in dark gray at right.

Proof. We argue by contradiction. We may assume h_0 is supported on an annulus A_0 (see Figure 3) surrounding a bounded Jordan domain D_0 whose boundary is γ_0 , and an unbounded region U_0 . Easy calculations show that the inverse image of D_0 under f consists of two bounded Jordan domains D_1^\pm each mapping as a quadratic branched cover onto D_0 and ramified at the points $c_\pm := \pm\sqrt{\frac{1+i}{2}}$ (the positive sign corresponding to the root with positive real part), both of which map to the origin under f . The domain D_1^+ contains two preimages of the point 1, namely $+1$ and $+\frac{1+i}{\sqrt{2}}$, while its twin D_1^- also contains two preimages of the point 1, namely -1 and $-\frac{1+i}{\sqrt{2}}$. The points $\pm 1 \in D_1^\pm$ belong to P , so if $h_1 : (\mathbb{P}^1, P) \rightarrow (\mathbb{P}^1, P)$ is a homeomorphism which is a lift of h_0 , fixing P pointwise, then $h_1(1) = 1$ and $h_1(-1) = -1$.

Since $f : D_1^\pm - \{c_\pm\} \rightarrow D_0 - \{0\}$ are both unramified coverings and $h_0 : D_0 - \{0\} \rightarrow D_0 - \{0\}$ is the identity map, we conclude $h_1 : D_1^\pm - \{c_\pm\} \rightarrow D_1^\pm - \{c_\pm\}$ is a deck transformation of this covering fixing a point, and hence is the identity on D_1^\pm .

The preimage of the annulus A_0 is a pair of disjoint, nonnested annuli A_1^\pm with an inner boundary component γ_1^\pm equal to ∂D_1^\pm . Since $f : A_1^\pm \rightarrow A_0$ is quadratic and unramified, and, by the previous paragraph, the restriction $h_1|_{D_1^\pm} = \text{id}$, we must have $h_1 \neq \text{id}$ on the outer boundary components of A_1^\pm ; indeed, h_1 there effects a half-twist.

The preimage of U_0 under f is a single unbounded region U_1 , which is homeomorphic to the plane minus two disks and three points; it maps in a four-to-one fashion, ramified only at the origin. The restriction $f : U_1 - \{f^{-1}(0)\} \rightarrow U_0 - \{-1\}$ is an unramified covering map, so $h_1 : U_1 - \{f^{-1}(-1)\} \rightarrow U_1 - \{f^{-1}(-1)\}$ is a deck transformation of this covering. By the previous paragraph, it is distinct from the identity.

We will obtain a contradiction by proving that $h_1 : U_1 - \{f^{-1}(-1)\} \rightarrow U_1 - \{f^{-1}(-1)\}$ has a fixed point; this is impossible for deck transformations other than the identity. We use the Lefschetz fixed point formula. By removing a neighborhood of ∞ and of -1 , and lifting these neighborhoods, we place ourselves in the setting of compact planar surfaces with boundary, so that this theorem will apply. Under h_1 , the boundary component near infinity is sent to itself, as are the outer boundaries of A_1^\pm and the boundary component surrounding the origin (since the origin is the uniquely ramified point of f over U_0). The remaining pair of boundary components are permuted amongst themselves. The action of $h_1 : U_1 - \{f^{-1}(-1)\} \rightarrow U_1 - \{f^{-1}(-1)\}$ on rational homology has trace equal to either 3 or 5. A fixed point thus exists, and the proof is complete. \square

Remark. There exists a lift h_1 of h_0 under f . First, there is a lift h' of h_0 under g , obtained by setting $h' = \text{id}$ on the preimage of U_0 . This extends to a half-twist on the preimage A'_0 of A_0 under g , which then in turn extends to a homeomorphism fixing the preimage D'_0 of D_0 under g ; inside D'_0 , this homeomorphism interchanges the points $1, i$ which are the preimages of 1 . It is then straightforward to show that h' lifts under s by setting $h_1 = \text{id}$ on U_1 and extending similarly over the annuli A_1^\pm and the domains D_1^\pm .

6. NONINJECTIVITY OF THE VIRTUAL HOMOMORPHISM

In this section, we begin with a nondynamical discussion about the injectivity of the virtual homomorphism $\phi_f : G_B \dashrightarrow G_A$ associated to an admissible covering map $f : (S^2, A) \rightarrow (S^2, B)$.

It follows from Lemma 3.2 that $\phi_f : G_B \dashrightarrow G_A$ is injective if and only if $\phi_{i \circ f \circ j} : G_B \dashrightarrow G_A$ is injective.

Lemma 6.1. *Let $f : (S^2, A) \rightarrow (S^2, B)$ be an admissible cover. If $\sigma_f : \mathcal{T}_B \rightarrow \mathcal{T}_A$ is injective, then $\phi_f : G_B \dashrightarrow G_A$ is injective.*

Proof. Suppose $h \in \ker(\phi_f)$. Then from the functional identity (2) we have $\sigma_f(h \cdot \tau) = \sigma_f(\tau)$ for all $\tau \in \mathcal{T}_B$, and so non-injectivity of ϕ_f implies non-injectivity of σ_f . \square

Remark. We are not aware of any examples of an admissible covering map $f : (S^2, A) \rightarrow (S^2, B)$ where $\phi_f : G_B \dashrightarrow G_A$ is injective but $\sigma_f : \mathcal{T}_B \rightarrow \mathcal{T}_A$ is not.

Our discussion naturally breaks up into three cases depending on $\#A$ and $\#B$. We treat the first two cases in the proposition below. The remaining case where $\#A = \#B$ is treated in Theorem 6.3.

Proposition 6.2. *Let $f : (S^2, A) \rightarrow (S^2, B)$ be an admissible covering map with virtual homomorphism $\phi_f : G_B \dashrightarrow G_A$.*

- (1) *If $\#B > \#A$, then ϕ_f is not injective.*
- (2) *If $\#B < \#A$, then ϕ_f is injective if $A = f^{-1}(B)$.*

Proof. We begin with a proof of (1) due to D. Margalit. Let $G_{\max} \leq G_B$ be a maximal abelian subgroup of G_B . The rank of G_{\max} is $\#B - 3$ by Theorem A in [BLM]. Let H_{\max} be a maximal abelian subgroup of H_f , the domain of ϕ_f . Because H_f has finite index in G_B , the rank of H_{\max} is also equal to $\#B - 3$ (Theorem 6.4C, [Iva2]). If ϕ_f were injective, then $\phi_f(H_{\max})$ would be an abelian subgroup in G_A of rank $\#B - 3$. But the rank of any abelian subgroup of G_A is bounded above by $\#A - 3$, and since $\#B > \#A$ by hypothesis, the virtual homomorphism ϕ_f cannot be injective.

We now prove (2). The Teichmüller spaces $\mathcal{T}_A, \mathcal{T}_B$ are each canonically isomorphic to the Teichmüller spaces where the points in A and B represent punctures. Under the hypothesis that $A = f^{-1}(B)$, the pullback map $\sigma_f : \mathcal{T}_B \rightarrow \mathcal{T}_A$ is induced by lifting complex structures under an unramified covering map. It is well known that in this case σ_f is a global isometry with respect to the Teichmüller metrics (a Teichmüller mapping ψ corresponding to a quadratic differential q lifts to a Teichmüller mapping corresponding to a lifted quadratic differential) and is therefore injective. By Lemma 6.1, ϕ_f is injective. □

Remark. Condition (2) is sufficient but not necessary. For example, in the dynamical setting, if $A = B = P_f$, if $f : (S^2, P_f) \rightarrow (S^2, P_f)$ is a Thurston map, and if f has Euclidean orbifold, then ϕ_f is injective. This is immediately clear in the case where $\#P_f = 3$. In the case where $\#P_f = 4$, the result follows from Lemma 6.1 and from the fact that $\sigma_f : \mathcal{T}_{P_f} \rightarrow \mathcal{T}_{P_f}$ is an automorphism [DH].

For the remaining case, we will require a nondynamical version of the notion of hyperbolic orbifold (see Section 3.2). Let $f : (S^2, A) \rightarrow (S^2, B)$ be an admissible cover with $\#A = \#B$. We say that f has *potentially hyperbolic orbifold* if there are orientation-preserving homeomorphisms i, j with

$$j^{-1} : (S^2, A) \rightarrow (S^2, j^{-1}(A))$$

and

$$i : (S^2, B) \rightarrow (S^2, i(B)),$$

so that $P := j^{-1}(A) = i(B)$, and $i \circ f \circ j : (S^2, P) \rightarrow (S^2, P)$ is a Thurston map with hyperbolic orbifold. Note that if $\#A = \#B$, then it is easy to find homeomorphisms j and i so that $f : (S^2, P) \rightarrow (S^2, P)$ is a Thurston map. The content of this definition is that some such Thurston map has hyperbolic orbifold.

Theorem 6.3. *Let $f : (S^2, A) \rightarrow (S^2, B)$ be an admissible cover with $\#A = \#B \geq 4$, and suppose that f has potentially hyperbolic orbifold. Then the virtual homomorphism $\phi_f : G_B \dashrightarrow G_A$ is not injective: its kernel contains a pseudo-Anosov element.*

Because f has potentially hyperbolic orbifold, there are homeomorphisms i, j with $j^{-1} : (S^2, B) \rightarrow (S^2, P)$ and $i : (S^2, A) \rightarrow (S^2, P)$ so that $F := i \circ f \circ j : (S^2, P) \rightarrow (S^2, P)$ is a Thurston map with distinguished set P , and this Thurston map has hyperbolic orbifold. We will prove that the virtual endomorphism $\phi_F : G_P \dashrightarrow G_P$ associated to F is not injective. It will then follow from Lemma 3.2 that the virtual homomorphism $\phi_f : G_B \dashrightarrow G_A$ is not injective, concluding the proof.

The proof of this theorem will use both analytic and algebraic arguments. From this, we will derive

Theorem 6.4. *Let $f : (S^2, A) \rightarrow (S^2, B)$ be an admissible cover with $\#A = \#B$, and suppose that f has potentially hyperbolic orbifold. Then the closure in the Thurston compactification of each nonempty fiber of $\sigma_f : \mathcal{T}_B \rightarrow \mathcal{T}_A$ map contains the Thurston boundary of \mathcal{T}_B .*

6.1. Proof of Theorem 6.3. We begin with some preliminary results of independent interest.

Suppose $f : (S^2, A) \rightarrow (S^2, B)$ is an admissible covering map with virtual homomorphism $\phi_f : G_B \dashrightarrow G_A$; as before, let $H_f < G_B$ be the domain of ϕ_f . Let d_B be the Teichmüller distance on \mathcal{T}_B , and let d_A be the Teichmüller distance on \mathcal{T}_A .

Recall that every element g of G_B has a *minimal displacement* given by

$$\delta_B(g) := \inf_{\tau \in \mathcal{T}_B} d_B(\tau, g \cdot \tau) \in [0, \infty).$$

Define the analogous object δ_A for G_A . Also recall that we have the functional identity

$$\sigma(h \cdot \tau) = \phi_f(h) \cdot \sigma(\tau), \quad \forall h \in H_f.$$

The following results are related. However, Proposition 6.5 and Corollary 6.7 involve nondynamical statements, while Corollary 6.6 and Corollary 6.8 involve dynamical statements.

Proposition 6.5. *Let $f : (S^2, A) \rightarrow (S^2, B)$ be an admissible cover. For each $h \in H_f$ we have*

$$\delta_A(\phi_f(h)) \leq \delta_B(h).$$

Proof. Fix $\varepsilon > 0$. We have

$$\begin{aligned} \delta_B(h) &= \inf_{\xi} d_B(\xi, h \cdot \xi) && \text{definition} \\ &> d_B(\tau, h \cdot \tau) - \varepsilon && \text{for some } \tau \in \mathcal{T}_B \\ &\geq d_A(\sigma_f(\tau), \sigma_f(h \cdot \tau)) - \varepsilon && \sigma_f \text{ is distance nonincreasing} \\ &= d_A(\sigma_f(\tau), \phi_f(h) \cdot \sigma(\tau)) - \varepsilon && \text{functional identity} \\ &\geq \inf_{\xi} d_A(\xi, \phi_f(h) \cdot \xi) - \varepsilon && \text{definition} \\ &= \delta_A(\phi_f(h)) - \varepsilon && \text{definition.} \end{aligned}$$

□

Corollary 6.6. *Let $f : (S^2, P) \rightarrow (S^2, P)$ be a Thurston map with hyperbolic orbifold.*

(1) *There exists $k \in \mathbb{N}$ such that if $h \in H_{f \circ k}$ is pseudo-Anosov, then*

$$\delta_P(\phi_f^{\circ k}(h)) < \delta_P(h).$$

(2) *If $\sigma_f : \mathcal{T}_P \rightarrow \mathcal{T}_P$ is strictly distance-decreasing and $h \in H_f$, then*

$$\delta_P(\phi_f(h)) < \delta_P(h).$$

Proof. Being pseudo-Anosov implies we can take τ to realize the infimum in the definition of $\delta_P(h)$, so that ε in the previous proof is equal to 0. By [DH, Cor. 3.4], there exists a $k \in \mathbb{N}$ such that the k th iterate $\sigma_f^{\circ k}$ strictly decreases Teichmüller distances. □

Corollary 6.7. *Let $f : (S^2, A) \rightarrow (S^2, B)$ be an admissible cover. Then the virtual homomorphism $\phi_f : G_B \dashrightarrow G_A$ sends multitwists to multitwists.*

Proof. There are no elliptic pure mapping class elements, so twists h are characterized by the condition that $\delta_B(h) = 0$ (or $\delta_A(h) = 0$ if $h \in G_A$). But $\delta_B(h) \geq \delta_A(\phi_f(h))$, so $\delta_B(h) = 0 \Rightarrow \delta_A(\phi_f(h)) = 0$. \square

Corollary 6.8. *Let $f : (S^2, P) \rightarrow (S^2, P)$ be a Thurston map, and let k be as in Corollary 6.6. If the orbifold of f is hyperbolic and $h \in H_{f \circ k}$ is a pseudo-Anosov element, then the mapping class $\phi_f^{\circ k}(h)$ cannot be conjugate to h in G_P . In particular, $\phi_f^{\circ k}(h) \neq h$ and $\phi_f(h) \neq h$. If $\sigma_f : \mathcal{T}_P \rightarrow \mathcal{T}_P$ is distance decreasing, then the mapping class $\phi_f(h)$ cannot be conjugate to h ; in particular, $\phi_f(h) \neq h$.*

Remark. One can show Proposition 6.5 analytically in another way, via the correspondence on moduli space and the definition of the virtual endomorphism in terms of the induced map on the fundamental group determined by the correspondence. As before, choose basepoints $w_{\otimes}, \mu_{\otimes}, \tilde{\mu}_{\otimes}$ with $Y(w_{\otimes}) = \mu_{\otimes}, X(w_{\otimes}) = \tilde{\mu}_{\otimes}$. Represent the mapping class $h \in G_B$ by a loop γ in \mathcal{M}_B based at μ_{\otimes} . By a suitable choice of basepoints we may assume the length of the loop γ is very close to $\delta_B(h)$. If $h \in H_f$, this loop lifts to a loop $\tilde{\gamma}$ based at w_{\otimes} , so the length of $\tilde{\gamma}$ and of γ are the same. The map $X : \mathcal{W}_f \rightarrow \mathcal{M}_A$ cannot increase lengths, so we conclude that $X(\tilde{\gamma})$ in \mathcal{M}_A under this projection has length which is at most that of γ in \mathcal{M}_B .

Proof of Theorem 6.3. Let S_m denote the m -times-punctured sphere. We will need a few results about its extended (nonpure) mapping class group, $\text{Mod}(S_m)$. The first is the following result of Bell and Margalit [BM, Thm. 1].

Theorem 6.9. *Let $m \geq 5$. If H is a finite index subgroup of $\text{Mod}(S_m)$ and $\rho : H \rightarrow \text{Mod}(S_m)$ is an injective homomorphism, then there is a unique $g \in \text{Mod}(S_m)$ so that $\rho(h) = ghg^{-1}$ for all $h \in H$.*

Recall we are proving that if $f : (S^2, P) \rightarrow (S^2, P)$ is a Thurston map with hyperbolic orbifold, then the kernel of the virtual endomorphism $\phi_f : G_P \dashrightarrow G_P$ contains a pseudo-Anosov element. Note that G_P is canonically identified with a finite index subgroup of the full mapping class group $\text{Mod}(S_m)$.

Assume first that $\#P \geq 5$.

Suppose to the contrary that ϕ_f is injective. Let k be the integer of Corollary 6.6. Then $\phi_f^{\circ k}$ is also injective. By Theorem 6.9 applied to $\rho = \phi_f^{\circ k}$ and $H := H_{f \circ k}$, there is a unique $g \in G_P$ so that $\phi_f^{\circ k}(h) = ghg^{-1}$ for all $h \in H$. Since H is a finite-index subgroup of G_P , it contains a pseudo-Anosov element, h . But then $\phi_f^{\circ k}(h) = ghg^{-1}$, contradicting Corollary 6.8. Let $N := \ker(\phi_f)$ and let H now denote the domain of ϕ_f . We have just shown that N is nontrivial.

Now assume that $\#P = 4$ so that \mathcal{M}_P and \mathcal{W}_f are each Riemann surfaces of finite type. Assuming that ϕ_f is injective, Theorem 4.1 implies $X : \mathcal{W}_f \rightarrow \mathcal{M}_P$ is proper and surjective; it is also open. It follows that $X : \mathcal{W}_f \rightarrow \mathcal{M}_P$ is a finite branched covering. Let $w_{\otimes} \in \mathcal{W}_f$, let $m_{\otimes} := X(w_{\otimes})$ be basepoints, and consider $X_* : \pi_1(\mathcal{W}_f, w_{\otimes}) \rightarrow \pi_1(\mathcal{M}_P, m_{\otimes})$. Suppose X_* is injective.

Let $p : \mathcal{Z} \rightarrow \mathcal{M}_P$ be the covering of \mathcal{M}_P induced by the image subgroup $X_*(\pi_1(\mathcal{W}_f, w_{\otimes}))$. By elementary covering space theory, there exists a lift $\tilde{X} : \mathcal{W}_f \rightarrow \mathcal{Z}$ of X satisfying $p \circ \tilde{X} = X$. Since degrees of branched coverings are multiplicative, \tilde{X} is a finite branched covering. By construction, the induced map

on fundamental groups \tilde{X}_* is injective and surjective, hence an isomorphism. Since \mathcal{M}_P is a thrice-punctured sphere, \mathcal{Z} and \mathcal{W}_f are punctured surfaces of strictly negative Euler characteristic, say $-k_{\mathcal{Z}}, -k_{\mathcal{W}_f}$, respectively. The fundamental groups of \mathcal{Z} and \mathcal{W}_f are then free groups of ranks, respectively $1 + k_{\mathcal{Z}}, 1 + k_{\mathcal{W}_f}$. Since they are isomorphic (via X_*), these ranks coincide, hence so do the Euler characteristics of \mathcal{Z} and \mathcal{W}_f . By the Riemann-Hurwitz formula, the degree of \tilde{X} must be 1. It follows that \tilde{X} is an unramified cover and hence that $X = p \circ \tilde{X}$ is an unramified cover as well.¹ But then local inverse branches of $X \circ Y^{-1}$ are isometries with respect to the hyperbolic (equivalently, Teichmüller) metric. Taking a composition of two such branches, we again obtain an isometry. This is impossible, since the orbifold of F is assumed hyperbolic, so that σ_f and, hence, $X \circ Y^{-1}$ have second iterates that are contractions.

We conclude that in both cases N is nontrivial. Since S_m is torsion-free, N is infinite. The next results we need concern the notion of irreducibility: a subgroup $H < \text{Mod}(S_m)$ is *reducible* if there exists a nonempty multicurve which is stabilized by every element of H . A subgroup containing a pseudo-Anosov element cannot be reducible. Hence any finite-index subgroup $H < \text{Mod}(S_m)$ is irreducible, since given any pseudo-Anosov element, some power will be a pseudo-Anosov element which lies in H . By [Iva1, Cor. 7.13], the subgroup N is irreducible, and by [Iva1, Cor. 7.14], N contains a pseudo-Anosov element.² \square

6.2. Proof of density, Theorem 6.4. We will prove that if $f : (S^2, A) \rightarrow (S^2, B)$ is an admissible map with $\#A = \#B$ and potentially hyperbolic orbifold, then the closure in the Thurston compactification of each nonempty fiber of $\sigma_f : \mathcal{T}_B \rightarrow \mathcal{T}_A$ contains the Thurston boundary of \mathcal{T}_B .

Let $N = \ker(\phi_f)$; by Theorem 6.3, N contains pseudo-Anosov elements. Suppose $\tau \in \mathcal{T}_P$ is an element of a nonempty fiber E of σ_f . Given a subgroup of G_B , recall that its *limit set* in the Thurston boundary of \mathcal{T}_B is defined as the closure of the set of fixed points of its pseudo-Anosov elements. For $n \in N$ we have $\sigma_f(n \cdot \tau) = \sigma_f(\tau)$, so that E is N -invariant. Since pseudo-Anosov elements have north-south dynamics on the Thurston compactification of \mathcal{T}_P , the accumulation set of E in the Thurston compactification contains the limit set $\Lambda(N)$. But N is a normal subgroup of H_f , so if $x \in \Lambda(N)$ is a fixed point of some $n \in N$, then $h \cdot x$ is a fixed point of $h \circ n \circ h^{-1} \in N$ for all $h \in H_f$. Thus $\Lambda(N)$ is H_f -invariant and it follows that $\Lambda(N) = \Lambda(H_f)$. Since H_f is of finite index in G_B , we conclude $\Lambda(N) = \Lambda(H_f) = \Lambda(G_B)$, which is equal to the Thurston boundary of \mathcal{T}_B . \square

7. FINITE GLOBAL ATTRACTOR FOR THE PULLBACK RELATION

For the remainder of the article, we restrict ourselves to the dynamical setting; that is, $f : (S^2, P) \rightarrow (S^2, P)$ is a Thurston map with a postcritical set P . In this section, we establish conditions on f under which there exists a finite set $\mathcal{A} \subseteq S_P$ such that $\mathcal{A} \xleftarrow{f} \mathcal{A}$ and the orbit of each curve eventually falls into \mathcal{A} ; we call \mathcal{A} a *finite global attractor*.

In this paragraph, we state a known algebraic condition for the existence of a finite global attractor. Suppose S is a finite generating set for a group G . We denote by $\|g\|$ the word length of g in the generators S .

¹The authors acknowledge Allan Edmonds for providing the above arguments in this paragraph.
²The authors acknowledge Dan Margalit for this reference.




FIGURE 4. (Image by W. Floyd.) The horizontal curve is fixed under pullback, as are all elements of its orbit under iteration of a double Dehn twist about the vertical curve.

Definition 7.1. The virtual endomorphism $\phi : G \dashrightarrow G$ is called *contracting* if the *contraction ratio*

$$\rho := \limsup_{n \rightarrow \infty} \left(\limsup_{\|g\| \rightarrow \infty} \frac{\|\phi^n(g)\|}{\|g\|} \right)^{1/n} < 1.$$

The contraction ratio of the virtual endomorphism ϕ is independent of the choice of the generating set and is an asymptotic property: ϕ is contracting if and only if some iterate of ϕ is contracting. In [Pil3, Thm. 1.4], it is shown that the algebraic criterion of contraction of the virtual endomorphism $\phi_f : G_P \dashrightarrow G_P$ implies the existence of a finite global attractor for $\mathcal{S}_P \xleftarrow{f} \mathcal{S}_P$. It is easy to construct examples of obstructed maps f for which $\mathcal{S}_P \xleftarrow{f} \mathcal{S}_P$ does not have a finite global attractor— for example, there could be a large essential subsurface $\Sigma \subseteq S^2 \setminus P$ for which $f|_\Sigma = \text{id}_\Sigma$, leading to e.g. infinitely many fixed curves. It is perhaps somewhat more surprising that one can achieve this with an expanding map.

Example. Let f be the Thurston map obtained by starting with the degree four integral Lattès example and applying the “blowing up” surgery along a single vertical arc joining two critical points (see Figure 4). The domain and range spheres are each the union of the two squares A, B at the left side of Figure 4 along their common boundary. The arrows go in the direction of pulling back. The map f

sends small topological quadrilaterals labelled A, B to the large squares labelled A, B , respectively; their boundary edges e_i map to the boundary edge with the same label e_i . As usual, set $P := P_f$.

It is therefore natural to restrict our attention to the case when f is a rational map, in which case $\sigma_f : \mathcal{T}_P \rightarrow \mathcal{T}_P$ has a unique fixed point τ_{\otimes} ; let $m_{\otimes} = \pi_P(\tau_{\otimes})$. Below, we give two different analytic conditions on the correspondence $X \circ Y^{-1} : \mathcal{M}_P \rightrightarrows \mathcal{M}_P$ which imply that $\mathcal{S}_P \xleftarrow{f} \mathcal{S}_P$ has a finite global attractor. These properties depend only on the (A, B) -Hurwitz class of f and not on the choice of base fixed point.

In preparation for stating our results, the following two paragraphs give definitions.

A nonempty subset $K \subseteq \mathcal{M}_P$ is *invariant* under the correspondence $X \circ Y^{-1}$ if $X \circ Y^{-1}(K) \subseteq K$. Consider the case when there exists a nonempty compact invariant subset, K . This condition is quite strong. If K is such a set, then so is any R -neighborhood of K , defined using the Teichmüller metric, since lengths of paths do not increase under application $X \circ Y^{-1}$. In this case it follows that \mathcal{M}_P is exhausted by compact invariant connected sets. Informally, then, the condition of having a nonempty invariant compact subset may be thought of as asserting that the ends of moduli space are repellers of $X \circ Y^{-1}$. The relationship between the property of having a nonempty invariant compact subset and other topological, dynamical, and algebraic properties is investigated in [KPS2].

We will call a length metric ℓ on \mathcal{M}_P *WP-like* if its lift $\tilde{\ell}$ to \mathcal{T}_P has the property that the identity map defines a homeomorphism between the completions of \mathcal{T} with respect to $\tilde{\ell}$ and with respect to the WP metric. This implies, in particular, that (\mathcal{M}, ℓ) has finite diameter. We say that the pullback correspondence $X \circ Y^{-1}$ on \mathcal{M}_P is *uniformly contracting* with respect to a WP-like length metric ℓ on \mathcal{M}_P if there is a constant $0 \leq \lambda < 1$ such that for each curve $\gamma : [0, 1] \rightarrow \mathcal{M}_P$ of finite length, and each lift $\tilde{\gamma}$ of γ under Y , we have $\ell(X \circ \tilde{\gamma}) \leq \lambda \cdot \ell(\gamma)$, where $\ell(\gamma)$ is the length of γ .

Theorem 7.2. *Suppose $f : (\mathbb{P}^1, P) \rightarrow (\mathbb{P}^1, P)$ is a rational Thurston map with hyperbolic orbifold.*

- (1) *If the correspondence $X \circ Y^{-1} : \mathcal{M}_P \rightrightarrows \mathcal{M}_P$ has a nonempty compact invariant subset, then ϕ_f is contracting, and the pullback relation \xleftarrow{f} has a finite global attractor.*
- (2) *If the correspondence $X \circ Y^{-1} : \mathcal{M}_P \rightrightarrows \mathcal{M}_P$ is uniformly contracting with respect to a WP-like length metric ℓ , then under iteration of the pullback relation \xleftarrow{f} , every curve becomes trivial, i.e. the set $\{o\}$ consisting of the trivial curves (inessential and peripheral) is a finite global attractor.*

Examples. (1) The correspondence induced by the Rabbit polynomial $f(z) = z^2 + c_R$ satisfies the hypothesis in (1). As shown in [BN1, §5.1], the map X is an inclusion, and in coordinates identifying \mathcal{M}_P with $\mathbb{P}^1 \setminus \{0, 1, \infty\}$, the map $Y \circ X^{-1}$ is given by the restriction of the critically finite hyperbolic rational function $g(w) = 1 - \frac{1}{w^2}$ to the complement of $\{0, \infty, \pm 1\}$. The Julia set K of g is then a nonempty compact invariant subset for $X \circ Y^{-1} = g^{-1}$. Theorem 7.2(1) implies that the pullback relation on curves has a finite global attractor. This is computed in [Pil3, §8] using algebraic techniques.

The same arguments apply in the case of preperiod 2, period 1 quadratic polynomials; see [BN1, §7.2] and [Pil3, §9].

- (2) The correspondence induced by the dendrite Julia set polynomial $f(z) = z^2 + i$ satisfies the hypothesis in (2). As shown in [BN1, §6.2], the map X is an inclusion, and in coordinates identifying \mathcal{M}_P with $\mathbb{P}^1 \setminus \{0, 1, \infty\}$, the map $Y \circ X^{-1}$ is given by the restriction of the critically finite Lattès-type subhyperbolic (but not hyperbolic) rational function $g(w) = (1 - 2/w)^2$ to the complement of $\{0, \infty, 1, 2\}$. The map g defines a self-orbifold-cover of the Euclidean $(2, 4, 4)$ -orbifold (see [DH, §9]) which uniformly expands the corresponding Euclidean metric ℓ by the factor $|1 + i| = \sqrt{2}$. The identity map of \mathcal{M}_P defines a homeomorphism between the completions with respect to the Weil-Petersson and Euclidean metrics, and it follows easily that ℓ is WP-like. Theorem 7.2(2) implies that under iterated pullback, each curve becomes trivial. This is established in [Pil3, §9] using algebraic techniques. Guizhen Cui (personal communication) has pointed out that there is an elementary proof of this fact: it is easy to see that any curve which does not surround the unique 2-cycle in P_f must become trivial, and any curve which does must, by the Schwarz-Pick lemma, have a geodesic representative which strictly shrinks under pullback.
- (3) The correspondence induced by $f(z) = \frac{3z^2}{2z^3+1}$, studied in [BEKP, §4], satisfies the hypotheses of neither (1) nor (2) above. The analysis in [Lod], however, shows that nonetheless the pullback relation on curves has a finite global attractor.

Proof. Since f is assumed rational, $\sigma_f : \mathcal{T}_P \rightarrow \mathcal{T}_P$ has a fixed point which we denote by τ_{\otimes} ; it projects to $w_{\otimes} \in \mathcal{W}_f$ and $m_{\otimes} \in \mathcal{M}_P$.

We first prove (1) by showing that the virtual endomorphism $\phi_f : G_P \dashrightarrow G_P$ is contracting and then appealing to [Pil3, Theorem 1.4]. To see that ϕ_f is contracting, recall that ϕ_f is also the virtual endomorphism on $\pi_1(\mathcal{M}_P, m_{\otimes})$ given by $(X \circ Y^{-1})_* = X_* \circ Y^*$, where $Y^*(\gamma)$ is the lift under Y of a loop $\gamma \in \pi_1(\mathcal{M}_P, m_{\otimes})$ based at $w_{\otimes} \in \mathcal{W}_f$. Equip \mathcal{M}_P with the Teichmüller metric (equivalently, by Royden’s theorem, the Kobayashi metric) and \mathcal{W}_f with the lift of this metric. Given a smooth loop in \mathcal{M}_P based at m_{\otimes} in the domain of the virtual endomorphism, it lifts under Y to a loop of the same length in \mathcal{W}_f and projects to a loop in \mathcal{M}_P , whose length is strictly shorter. The space \mathcal{M}_P is a hyperplane complement, so its fundamental group is generated by a finite collection S of loops. The union of these loops lies in some invariant compact subset K ; we may assume K is a compact submanifold with smooth boundary. In K , the word length $\|g\|_S$ with respect to S is comparable to the infimum of lengths of loops in K representing g . Since K is compact and invariant, and since the orbifold of f is hyperbolic, the second iterate of the correspondence is uniformly contracting on K . It follows that lengths of such liftable loops in K , and hence the word lengths of the corresponding group elements, are uniformly contracted under iteration and so ϕ_f is contracting.

We now prove (2). Let $\tilde{\ell}$ denote the lifted WP-like metric. Results of [Sel1, §4] imply that the pullback map σ_f extends to the Weil-Petersson completion $\overline{\mathcal{T}}_P$, which, by the WP-like property of ℓ , coincides with the $\tilde{\ell}$ completion of \mathcal{T}_P . The hypothesis implies that σ_f is uniformly contracting with respect to $\tilde{\ell}$, and hence that under iteration, each point in $\overline{\mathcal{T}}_P$ converges exponentially fast to τ_{\otimes} under iteration. Now let Γ be a multicurve on $S^2 \setminus P$. Proposition [Sel1, Prop. 6.1]

implies that for each multicurve Γ , the pullback map σ_f sends the stratum \mathcal{T}_P^Γ to the stratum $\mathcal{T}_P^{f^{-1}(\Gamma)}$.

Define the *complexity* of Γ , denoted $\kappa(\Gamma)$, to be the distance between τ_\otimes and the stratum \mathcal{T}_P^Γ with respect to the lifted metric $\tilde{\ell}$. The admissibility of ℓ implies

- (1) for all Γ , $\kappa(\Gamma) < \infty$, and
- (2) there exists $r_0 > 0$ such that for all $\Gamma \neq \emptyset$, $\kappa(\Gamma) > r_0$.

The hypothesis, the admissibility of ℓ , and the fact that $\sigma_f(\mathcal{T}_P^\Gamma) \subseteq \mathcal{T}_P^{f^{-1}(\Gamma)}$ imply

- (3) there exists $\lambda < 1$ such that for all Γ , $\kappa(f^{-1}(\Gamma)) \leq \lambda \cdot \kappa(\Gamma)$.

From (1), (2), and (3), it follows that given any multicurve Γ , there exists $n \in \mathbb{N}$ such that $\kappa(f^{-n}(\Gamma)) = 0$ and hence that $f^{-n}(\Gamma)$ is trivial.

Hence under iterated pullback, each multicurve becomes trivial. It follows that every curve becomes trivial under iterated pullback. □

8. SHADOWING

In this section, we briefly consider one consequence of surjectivity of the virtual endomorphism; this will be used in §9.2 below.

Proposition 8.1. *Let $f : (S^2, P) \rightarrow (S^2, P)$ be a Thurston map. Suppose that $\phi_f : G_P \dashrightarrow G_P$ is surjective. Then any finite orbit segment of the correspondence $X \circ Y^{-1} : \mathcal{M}_P \rightrightarrows \mathcal{M}_P$ is covered by an orbit segment of the pullback map $\sigma_f : \mathcal{T}_P \rightarrow \mathcal{T}_P$.*

Proof. Let $\mu_1 \in \mathcal{M}_P$, and let $\mu_2 \in X(Y^{-1}(\mu_1)) \subseteq \mathcal{M}_P$. By the fundamental diagram, Figure 2, there exist $\tau'_1, \tau'_2 \in \mathcal{T}_P$ so that $\sigma_f(\tau'_1) = \tau'_2$, and $\pi_P(\tau'_1) = \mu_1$ and $\pi_P(\tau'_2) = \mu_2$. Let $\tau_2 \in \pi_P^{-1}(\mu_2)$. Then there exists $g \in G_P$ so that $\tau_2 = g \cdot \tau'_2$. Since $\phi_f : G_P \dashrightarrow G_P$ is surjective, there exists $h \in G_P$ so that $\phi_f(h) = g$. Define $\tau_1 := h \cdot \tau'_1$. Then

$$\sigma_f(\tau_1) = \sigma_f(h \cdot \tau'_1) = \phi_f(h) \cdot \sigma_f(\tau'_1) = g \cdot \tau'_2 = \tau_2.$$

The claim now readily follows by induction.

9. OBSTRUCTED TWISTS AND REPELLING FIXED POINTS IN $\partial\mathcal{M}_P$

If f is a Thurston map with $\#P = 4$, then the projection maps $Y, X : \mathcal{W}_f \rightarrow \mathcal{M}_P$ extend to holomorphic maps $Y, X : \overline{\mathcal{W}}_f \rightarrow \overline{\mathcal{M}}_P \simeq \mathbb{P}^1$ yielding a holomorphic correspondence on the compactified moduli space $X \circ Y^{-1} : \mathbb{P}^1 \rightarrow \mathbb{P}^1$.

9.1. Obstructed twists.

Theorem 9.1. *Let $f : (\mathbb{P}^1, P) \rightarrow (\mathbb{P}^1, P)$ be a rational Thurston map with hyperbolic orbifold. Suppose $\#P = 4$.*

Then the following conditions are equivalent:

- (1) *There exist $g \in G_P$, a twist T , and a nonzero $k \in \mathbb{Z}$ such that $f_* := g \circ f$ commutes up to homotopy with T^k .*
- (2) *There exist $g \in G_P$ such that $f_* := g \circ f$ has an obstruction $\Gamma := \{\gamma\}$ for which $\lambda_f(\gamma) = \gamma$.*
- (3) *There exist an element $g \in G_P$, a twist T , and a nonzero integer k such that $(T^k)^g \in H_f$ and $\phi_f((T^k)^g) = T^k$.*
- (4) *There exist curves γ_1, γ_2 in $\mathbb{P}^1 \setminus P$ such that (i) $g(\gamma_1) = \gamma_2$ for some $g \in G_P$, and (ii) $\lambda_f(\gamma_1) = \gamma_2$.*

- (5) *There exist a point $0 \in \partial\mathcal{M}_P$ and a single-valued branch $\mu \mapsto \beta(\mu)$ of the pullback correspondence $X \circ Y^{-1}$ near 0 such that $\beta(0) = 0$ and, in suitable local holomorphic coordinates for which 0 is the origin, $\beta(\mu) = a\mu + O(\mu^2)$, where $0 < |a| < 1$.*

Proof. (1) \iff (2) We take the elements g to be the same.

Condition (1) is equivalent to the condition that $T^k \circ f_* = f_* \circ T^k$ up to homotopy, i.e. that $T^k \in \text{dom}(\phi_{f_*})$ and $\phi_{f_*}(T^k) = T^k$. In turn this is equivalent to the condition that for some k' with $|k'| \geq |k|$ we have $T^{k'} \in \text{dom}(\phi'_f)$ and $\phi_{f_*}(T^{k'}) = T^{k'}$. Denote by γ the core curve of T . Put $w = k'\gamma$ so that the multitwist $M_w := T^{k'}$.

If condition (1) holds, then equation (1) implies

$$\phi_{f_*}(M_w) = M_w \Rightarrow \lambda_{f_*}(w) = w \Rightarrow \lambda_{f_*}(k'\gamma) = k'\gamma \Rightarrow \lambda_{f_*}(\gamma) = \gamma$$

by linearity, and so γ is an obstruction for f_* with eigenvalue 1.

Now suppose condition (2) holds, with the obstruction being given by γ . Choose $k' \in \mathbb{Z}$ so $T^{k'} \in \text{dom}(\phi'_{f_*})$. Again by equation (1) we have

$$\lambda_{f_*}(\gamma) = \gamma \Rightarrow \lambda_f(k'\gamma) = k'\gamma \Rightarrow \phi_{f_*}(T^{k'}) = T^{k'},$$

as required.

(1) \iff (3) By applying Lemma 3.2 with $i = g$ and $j = \text{id}$, for any $g \in G$ we have

$$\phi_f((T^k)^g) = T^k \iff \phi_{g \circ f}(T^k) = T^k.$$

(4) \iff (2) This follows immediately from the observation that $\lambda_{g \circ f}(\gamma) = \lambda_f(g^{-1}(\gamma))$.


(1) \Rightarrow (5) If $T^k \circ f_* = f_* \circ T^k$ up to homotopy, then $\sigma_{T^k} \circ \sigma_{f_*} = \sigma_{f_*} \circ \sigma_{T^k}$. Let γ be the core curve of the twist T , let $\Gamma = \{\gamma\}$, and consider the quotient

$$\mathbb{H}/\langle z \mapsto z + 2k \rangle \simeq \mathcal{T}_P/k\text{Tw}(\Gamma) \simeq \mathbb{D}^*$$

which we identify with the punctured disk. Then σ_f descends to an analytic map

$$\bar{\sigma}_{f_*} : \mathbb{D}^* \rightarrow \mathbb{D}^*.$$


The fundamental diagram becomes



where the vertical arrows are covering maps. Since Γ is f_* -invariant, the induced map $\bar{\sigma}_{f_*}$ extends over the origin, yielding an analytic map

$$\bar{\sigma}_{f_*} : (\mathbb{D}, 0) \rightarrow (\mathbb{D}, 0).$$

For convenience of notation, we denote by 0 the corresponding end of \mathcal{M}_P and of \mathcal{W}_f . We obtain a commutative diagram of analytic maps:



Since f has hyperbolic orbifold, $\bar{\sigma}_{f_*} \notin \text{Aut}(\mathbb{D})$. By the Schwarz lemma, its derivative at the origin therefore satisfies $|(\bar{\sigma}_{f_*})'(0)| < 1$. Taking inverse branches near the origin, we have

$$(X \circ Y^{-1})'(0) = (\pi_P \circ \bar{\sigma}_{f_*} \circ \pi_P^{-1})'(0) = \sigma'_{f_*}(0),$$

and so the pullback correspondence $X \circ Y^{-1}$ has an attracting fixed point in the boundary of moduli space.

In the remaining paragraphs, we prove that $\sigma'_{f_*}(0) \neq 0$ and that (5) \Rightarrow (3).

We first discuss some specializations of the results in the “fivefold way” Theorem 2.1 to the case $\#P = 4$; cf. [Lod]. We may identify $P = P_f = \{0, 1, \infty, m_{\otimes}\}$, where $m_{\otimes} \in X \circ Y^{-1}(m_{\otimes})$ (since f is rational). Recall that there is a natural identification $G_P \leftrightarrow \pi_1(\mathcal{M}_P, m_{\otimes})$ such that $\phi_f : G_P \dashrightarrow G_P$ coincides with the virtual endomorphism $X_* \circ Y^*$ of $\pi_1(\mathcal{M}_P, m_{\otimes})$ induced by the correspondence $X \circ Y^{-1} : (\mathcal{M}_P, m_{\otimes}) \rightrightarrows (\mathcal{M}_P, m_{\otimes})$. Under this identification, a simple oriented loop γ_{loop} based at m_{\otimes} in \mathcal{M}_P which is peripheral and which encloses 0 on its left-hand side corresponds to the right Dehn twist T about the right-hand boundary component γ_{curve} of a regular neighborhood of γ_{loop} in $\mathbb{P}^1 \setminus \{0, 1, \infty, m_{\otimes}\}$. Denote by λ the group element of $\pi_1(\mathcal{M}, m_{\otimes})$ corresponding to g . We see that

$$\phi_{g \circ f}(T^k) = T^k \iff \phi_f((T^k)^g) = (T^k)^g \iff (X_* \circ Y^*)(\lambda \cdot \gamma_{loop}^k \cdot \lambda^{-1}) = \gamma_{loop}^k,$$

where \cdot denotes concatenation of paths, λ^{-1} is the path λ traversed in the opposite direction, and in the right-hand expression the path λ is traversed first.

Now suppose (1) holds, so that $(X_* \circ Y^*)(\lambda \cdot \gamma_{loop}^k \cdot \lambda^{-1}) = \gamma_{loop}^k$. The assumption that $\lambda \cdot \gamma_{loop}^k \cdot \lambda^{-1}$ is in the domain of (Y^*) implies that the k th power of a tiny peripheral simple closed curve surrounding $0 \in \mathcal{M}_P \cup \{0\}$ lifts under Y to a tiny simple closed curve surrounding $0 \in \mathcal{W}_f \cup \{0\}$. Since $(X_* \circ Y^*)(\lambda \cdot \gamma_{loop}^k \cdot \lambda^{-1}) = \gamma_{loop}^k$, the tiny curve surrounding $0 \in \mathcal{W}_f \cup \{0\}$, obtained by lifting, maps by X to the k th power of a simple closed curve surrounding $0 \in \mathcal{M}_P \cup \{0\}$. This shows that the local degree of the branch of $X \circ Y^{-1}$ at 0 is 1 , and, hence, the derivative at this point does not vanish. This finishes the proof of (1) \Rightarrow (5).

The proof of (5) \Rightarrow (3) proceeds similarly. If (5) holds, then the local degrees of X and Y at 0 are the same, say k . Thus under the correspondence a k th power of a tiny simple peripheral curve about 0 lifts under Y and projects under X to a k th power of a tiny simple peripheral curve about 0 . Since free homotopy classes of curves correspond to conjugacy classes of loops based at m_{\otimes} , we conclude that $(X_* \circ Y^*)(\lambda \cdot \gamma_{loop}^k \cdot \lambda^{-1}) = \gamma_{loop}^k$ for some loop γ_{loop} surrounding 0 and some loop λ based at m_{\otimes} . Thus (3) holds, and the proof is complete. \square

9.2. Dynamical consequences.

Theorem 9.2. *Under the hypotheses and notation of Theorem 9.1, assume further that the map X of the correspondence on moduli space is injective, so that an inverse of the pullback map descends and extends to a holomorphic self-map³ $Y \circ X^{-1} : \mathbb{P}^1 \rightarrow \mathbb{P}^1$. Then the following further properties hold:*

- I. *For any Thurston map $F : (S^2, P) \rightarrow (S^2, P)$ which is (P, P) -Hurwitz equivalent to f , the associated virtual endomorphism ϕ_F on G_P is surjective.*
- II. *The end 0 is a repelling fixed point of $Y \circ X^{-1}$.*
- III. *Conditions (1) and (3) hold with $k = 1$.*
- IV. *The univalent pullback relation on curves induced by the rational map f has arbitrarily long nontrivial orbits (necessarily comprised of distinct curves) whose elements are all homologous (and hence differ by elements of G_P).*
- V. *The collection of obstructed Thurston maps $T^n \circ f_*$, $n \in \mathbb{Z}$, are pairwise combinatorially inequivalent.*
- VI. *For any $\delta > 0$ and any sequence $\varepsilon_1 > \varepsilon_2 > \dots > \varepsilon_k$ there exists $\tau_0 \in \mathcal{T}_P$ and iterates $n_1 > m_1 > n_2 > m_2 > \dots > n_k > m_k$ such that (i) $\pi_P(\tau_0) = m_\otimes$; (ii) for each $i = 1, \dots, k$,*

$$\ell(\tau_{n_i}) < \varepsilon_i, \quad d(\pi_P(\tau_{m_i}), m_\otimes) < \delta,$$

where $\ell(\tau)$ is the length of the systole on the underlying Riemann surface.

Proof. I. Different choices of representatives in the (P, P) -Hurwitz class of f lead to virtual endomorphisms which differ by inner automorphisms. So (just to fix ideas) it suffices to verify this for the case of the rational map f , corresponding to the fixed point m_\otimes . But this is clear, since the nonconstant inclusion $X : \mathcal{W}_f \hookrightarrow \mathcal{M}_P$ of finitely punctured spheres must induce a surjection on fundamental groups.

- II. This follows immediately from condition (5) in Theorem 9.1.
- III. The previous conclusion implies that Y is locally injective near 0, and hence that a simple peripheral loop γ_{loop} in \mathcal{M}_P based at m_\otimes surrounding the end 0 in \mathcal{M}_P on its left-hand side lifts under Y to a simple loop in \mathcal{W}_f based at w_\otimes surrounding the end 0 in \mathcal{W}_f . This implies that the corresponding twist T lies in H_f . Since X is injective and the end 0 in \mathcal{W}_f maps to the end 0 in \mathcal{M}_P , we have that $\phi_f(T)$ is conjugate to T via some $g \in G_P$, i.e. $\phi_f(T^g) = T$, and so condition (3) of Theorem 9.1 holds with $k = 1$. The proof of the equivalence of (1) and (3) in Theorem 9.1 shows that the integers k in (1) and in (3) are the same.
- IV. The argument given below is due to L. Bartholdi. By the previous assertion, we may suppose condition (3) in Theorem 9.1 holds with $k = 1$, and let f, g, T, ϕ_f be as in the statement there. Recall that G_P is free, generated by any pair of Dehn twists whose core curves intersect minimally. Choose such a twist $S \in G_P$ so that $G_P = \langle S, T \rangle$. Since ϕ_f is surjective, there exists $\tilde{T} := T^g$ and $\tilde{S} \in G$ such that $\phi_f(\tilde{T}) = T, \phi_f(\tilde{S}) = S$. Since G_P is free on the generators S, T , the homomorphism defined by sending $T \mapsto \tilde{T}, S \mapsto \tilde{S}$ is an injection $\sigma : G_P \rightarrow G_P$ giving a section of ϕ_f , i.e. $\phi_f \circ \sigma = \text{id}_{G_P}$. For $n \in \mathbb{N}$

³A so-called “ g -map”, in the language of the first author.

let

$$w_n = g \cdot \sigma(g) \cdot \dots \cdot \sigma^{\circ(n-1)}(g).$$

Then an easy induction argument shows

$$\phi_f(T^{w_n}) = T^{w_{n-1}}.$$

The elements T^{w_n} are primitive Dehn twists whose core curve C_n lifts under f to the core curve C_{n-1} of $T^{w_{n-1}}$. These curves must be pairwise distinct (otherwise f is obstructed) and pairwise homologous (since the twists are pairwise conjugate). This proves (4).

V. We again suppose that condition (3) in Theorem 9.1 holds with $k = 1$, and we let f, g, T, ϕ_f be as in the statement there.

Lemma 9.3. *Suppose F is a Thurston map with hyperbolic orbifold for which $\#P_F = 4$. If F is obstructed, it has a unique Thurston obstruction $\Gamma = \{\gamma\}$. If $h \circ F = F \circ h$ up to homotopy for some $h \in G_P$, then $h(\gamma) = \gamma$, preserving orientation.*

Cf. [BN1, Prop. 6.10].

Proof. By the main result of [Pil], there is an obstruction (called the *canonical obstruction* of F) that is disjoint from all other obstructions. But on a sphere with four marked points, any two distinct essential nonperipheral curves intersect. A homeomorphism representing a pure mapping class that yields an automorphism of F up to homotopy must send an obstruction to an obstruction, and hence must fix the unique obstruction, preserving orientation since (by pureness) each disk it bounds must be mapped to itself. \square

Suppose $n_1, n_2 \in \mathbb{Z}$ and $T^{n_1} \circ f_*$ is combinatorially equivalent to $T^{n_2} \circ f_*$ via $g \in G_P$, so that up to homotopy relative to P ,

$$g \circ T^{n_1} \circ f_* \circ g^{-1} = T^{n_2} \circ f_*.$$

By construction, both $T^{n_1} \circ f_*$ and $T^{n_2} \circ f_*$ have a common obstruction, $\Gamma = \{\gamma\}$, which is the core curve of the twist, T . By Lemma 9.3, $g(\gamma) = \gamma$, preserving orientation. Hence $g = T^l$ for some $l \in \mathbb{Z}$ and

$$g \circ T^{n_1} \circ f_* \circ g^{-1} = T^l \circ T^{n_1} \circ f_* \circ T^{-l} = T^{n_2} \circ f_*.$$

Since f_* commutes with T up to homotopy, we have

$$f_* \circ T^{n_1} = f_* \circ T^{n_2},$$

and so $n_1 = n_2$ since the right action of G_P on the set \mathcal{F} of homotopy classes of Thurston maps with postcritical set P is free ([Pil3, §3], [Kam, Prop. 3.1]).

VI. Since Y is a covering map, the critical values of the map $Y \circ X^{-1}$ lie in the set of ends of \mathcal{M}_P . Since these ends must map to themselves, we conclude that $Y \circ X^{-1}$ is analytically conjugate to a subhyperbolic rational map R with a repelling fixed point at the origin, 0; the point m_{\otimes} is also fixed and repelling. In particular, $Y \circ X^{-1}$ is uniformly expanding near its Julia set with respect to a complete orbifold length metric.

Let $V \subseteq \mathcal{M}_P \cup \{0\}$ be a simply-connected domain containing 0 and m_{\otimes} . The expanding property implies that for $r \in \mathbb{N}$ large enough, there exist domains $U_0, U_{m_{\otimes}}$ compactly contained in V , and univalent inverse branches of R^r giving analytic isomorphisms $R_0^{-r} : V \rightarrow U_0$ and $R_{m_{\otimes}}^{-r} : V \rightarrow U_{m_{\otimes}}$

that fix 0 and m_{\otimes} , respectively. The maps $R_0^{-r}, R_{m_{\otimes}}^{-r}$ are branches of the r th iterate of $X \circ Y^{-1}$; together with their domains, we have a so-called iterated function system. By contraction, the fixed points of these branches are unique.

In \mathcal{M}_P , the loci defined by the conditions $\ell(\tau) < \varepsilon$ and $d(\pi_P(\tau), m_{\otimes}) < \delta$ contain neighborhoods of the end 0 and the basepoint m_{\otimes} of \mathcal{M}_P . Define a sequence of points $\mu_n \in \mathcal{M}_P$ comprising an orbit under the pullback correspondence as follows. Set $\mu_0 := m_{\otimes}$. Apply the branch $R_0^{-r} : V \rightarrow U_0$ enough times so that the length of the systole becomes less than ε_1 . Now apply the branch $R_{m_{\otimes}}^{-r}$ enough times so that the resulting point lies in the neighborhood $d(\pi_P(\tau), m_{\otimes}) < \delta$. Now again apply the branch $R_0^{-r} : V \rightarrow U_0$ enough times so that the length of the systole becomes less than ε_2 , etc. In this fashion, we obtain a finite orbit segment of the pullback correspondence in \mathcal{M}_P . By conclusion I, the virtual endomorphism ϕ_f on G_P is surjective. Hence Proposition 8.1 applies. We conclude that the orbit in \mathcal{M}_P can be lifted to an orbit in \mathcal{T}_P having the desired properties. \square

Example. Consider Example (2) from Section 7. As shown in [BN1, § 6.2], the dendrite Julia set polynomial $f(z) = z^2 + i$ satisfies the hypothesis of the previous theorem. Therefore all conclusions of the theorem follow; moreover, the following is true. Recall that $Y \circ X^{-1}$ is conjugate to the rational map $g(w) = (1 - 2/w)^2$, the Julia set of which is the entire Riemann sphere. Therefore, the forward orbit of any point in \mathcal{M}_P under the pullback correspondence, which is the same as the backward orbit of $Y \circ X^{-1}$, is dense in \mathcal{M}_P . In other words, for any point $\tau \in \mathcal{T}_P$, the projection $\pi_P(\bigcup_{g \in G_P, n \in \mathbb{N}} \{\sigma_f^n(g \cdot \tau)\})$ of the union of σ_f -orbits of all points in \mathcal{T}_P , that are in the same fiber as τ over \mathcal{M}_P , is dense in \mathcal{M}_P . This situation is somewhat surprising as we know that every particular orbit $\sigma_f^n(g \cdot \tau)$ converges geometrically fast to the unique attracting fixed point τ_{\otimes} , which corresponds to f . For this example, the statement of the conclusion VI in the last theorem can be further strengthened. We can find a point τ_0 in the fiber of m_{\otimes} that follows an arbitrary itinerary in the moduli space with an arbitrary precision before converging to the fixed point.

ACKNOWLEDGEMENTS

The authors thank Indiana University for supporting the visits of the first and third authors. The first author was supported by an NSF postdoctoral fellowship and the second author by a Simons collaboration grant. We thank A. Edmonds, D. Margalit, and V. Turaev for useful conversations.

REFERENCES

- [BN1] Laurent Bartholdi and Volodymyr Nekrashevych, *Thurston equivalence of topological polynomials*, Acta Math. **197** (2006), no. 1, 1–51, DOI 10.1007/s11511-006-0007-3. MR2285317 (2008c:37072)
- [BN2] Laurent Bartholdi and Volodymyr V. Nekrashevych, *Iterated monodromy groups of quadratic polynomials. I*, Groups Geom. Dyn. **2** (2008), no. 3, 309–336, DOI 10.4171/GGD/42. MR2415302 (2009d:37074)
- [BM] Robert W. Bell and Dan Margalit, *Injections of Artin groups*, Comment. Math. Helv. **82** (2007), no. 4, 725–751, DOI 10.4171/CMH/108. MR2341838 (2009b:20062)

- [BLM] Joan S. Birman, Alex Lubotzky, and John McCarthy, *Abelian and solvable subgroups of the mapping class groups*, Duke Math. J. **50** (1983), no. 4, 1107–1120, DOI 10.1215/S0012-7094-83-05046-9. MR726319 (85k:20126)
- [BEKP] Xavier Buff, Adam Epstein, Sarah Koch, and Kevin Pilgrim, *On Thurston's pullback map*, Complex dynamics, A K Peters, Wellesley, MA, 2009, pp. 561–583, DOI 10.1201/b10617-20. MR2508269 (2010g:37071)
- [CFP] J. W. Cannon, W. J. Floyd, and W. R. Parry, *Finite subdivision rules*, Conform. Geom. Dyn. **5** (2001), 153–196 (electronic), DOI 10.1090/S1088-4173-01-00055-8. MR1875951 (2002j:52021)
- [CFPP] J. W. Cannon, W. J. Floyd, W. R. Parry, and K. M. Pilgrim, *Subdivision rules and virtual endomorphisms*, Geom. Dedicata **141** (2009), 181–195, DOI 10.1007/s10711-009-9352-7. MR2520071 (2010d:37086)
- [DH] Adrien Douady and John H. Hubbard, *A proof of Thurston's topological characterization of rational functions*, Acta Math. **171** (1993), no. 2, 263–297, DOI 10.1007/BF02392534. MR1251582 (94j:58143)
- [FLP] Albert Fathi, François Laudenbach, and Valentin Poénaru, *Thurston's work on surfaces*, Mathematical Notes, vol. 48, Princeton University Press, Princeton, NJ, 2012. Translated from the 1979 French original by Djun M. Kim and Dan Margalit. MR3053012
- [HP1] Peter Haïssinsky and Kevin M. Pilgrim, *Finite type coarse expanding conformal dynamics*, Groups Geom. Dyn. **5** (2011), no. 3, 603–661, DOI 10.4171/GGD/141. MR2813529
- [HP2] Peter Haïssinsky and Kevin M. Pilgrim, *An algebraic characterization of expanding Thurston maps*, J. Mod. Dyn. **6** (2012), no. 4, 451–476. MR3008406
- [HK] John H. Hubbard and Sarah Koch, *An analytic construction of the Deligne-Mumford compactification of the moduli space of curves*, J. Differential Geom. **98** (2014), no. 2, 261–313. MR3263519
- [Hub] John Hamal Hubbard, *Teichmüller theory and applications to geometry, topology, and dynamics. Vol. 1*, Teichmüller theory. With contributions by Adrien Douady, William Dunbar, Roland Roeder, Sylvain Bonnot, David Brown, Allen Hatcher, Chris Hruska and Sudeb Mitra. With forewords by William Thurston and Clifford Earle. Matrix Editions, Ithaca, NY, 2006. MR2245223 (2008k:30055)
- [Iva1] Nikolai V. Ivanov, *Subgroups of Teichmüller modular groups*, Translations of Mathematical Monographs, vol. 115, American Mathematical Society, Providence, RI, 1992. Translated from the Russian by E. J. F. Primrose and revised by the author. MR1195787 (93k:57031)
- [Iva2] Nikolai V. Ivanov, *Mapping class groups*, <http://www.math.msu.edu/~ivanov/m99.ps>, 1998.
- [Kam] Atsushi Kameyama, *The Thurston equivalence for postcritically finite branched coverings*, Osaka J. Math. **38** (2001), no. 3, 565–610. MR1860841 (2002h:57004)
- [Kel] Gregory A. Kelsey, *Mapping schemes realizable by obstructed topological polynomials*, Conform. Geom. Dyn. **16** (2012), 44–80, DOI 10.1090/S1088-4173-2012-00239-1. MR2893472
- [Koc1] Sarah Koch, *Teichmüller theory and endomorphisms of \mathbb{P}^n* , PhD thesis, University of Provence, 2007.
- [Koc2] Sarah Koch, *Teichmüller theory and critically finite endomorphisms*, Adv. Math. **248** (2013), 573–617, DOI 10.1016/j.aim.2013.08.019. MR3107522
- [KPS2] Sarah Koch, Kevin M. Pilgrim, and Nikita Selinger, *Limit sets of Thurston pullback semigroups*, Manuscript, 2014.
- [Lod] Russell Lodge, *Boundary values of the Thurston pullback map*, ProQuest LLC, Ann Arbor, MI, 2012. Thesis (Ph.D.)—Indiana University. MR3054977
- [Mas] Howard Masur, *Extension of the Weil-Petersson metric to the boundary of Teichmüller space*, Duke Math. J. **43** (1976), no. 3, 623–635. MR0417456 (54 #5506)
- [Nek1] Volodymyr Nekrashevych, *Combinatorics of polynomial iterations*, Complex dynamics, A K Peters, Wellesley, MA, 2009, pp. 169–214, DOI 10.1201/b10617-5. MR2508257 (2011k:37082)
- [Nek2] Volodymyr Nekrashevych, *Combinatorial models of expanding dynamical systems*, Ergodic Theory Dynam. Systems **34** (2014), no. 3, 938–985, DOI 10.1017/etds.2012.163. MR3199801

- [Pil1] Kevin M. Pilgrim, *Canonical Thurston obstructions*, Adv. Math. **158** (2001), no. 2, 154–168, DOI 10.1006/aima.2000.1971. MR1822682 (2001m:57004)
- [Pil2] Kevin M. Pilgrim, *Combination, decomposition, and structure theory for postcritically finite branched coverings of the two-sphere to itself*, to appear, Springer Lecture Notes in Math., 2003.
- [Pil3] Kevin M. Pilgrim, *An algebraic formulation of Thurston’s characterization of rational functions* (English, with English and French summaries), Ann. Fac. Sci. Toulouse Math. (6) **21** (2012), no. 5, 1033–1068, DOI 10.5802/afst.1361. MR3088266
- [Sel1] Nikita Selinger, *Thurston’s pullback map on the augmented Teichmüller space and applications*, Invent. Math. **189** (2012), no. 1, 111–142, DOI 10.1007/s00222-011-0362-3. MR2929084
- [Sel2] Nikita Selinger, *Topological characterization of canonical Thurston obstructions*, J. Mod. Dyn. **7** (2013), no. 1, 99–117, DOI 10.3934/jmd.2013.7.99. MR3071467
- [Wol] Scott A. Wolpert, *The Weil-Petersson metric geometry*, Handbook of Teichmüller theory. Vol. II, IRMA Lect. Math. Theor. Phys., vol. 13, Eur. Math. Soc., Zürich, 2009, pp. 47–64, DOI 10.4171/055-1/2. MR2497791 (2010i:32012)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 530 CHURCH STREET, ANN ARBOR, MICHIGAN 48109

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, 831 E. THIRD STREET, BLOOMINGTON, INDIANA 47405

INSTITUTE FOR MATHEMATICAL SCIENCES, STONY BROOK UNIVERSITY, STONY BROOK, NEW YORK 11794-3660