

LOCAL MAPS AND THE REPRESENTATION THEORY OF OPERATOR ALGEBRAS

ELIAS G. KATSOULIS

ABSTRACT. Using representation theory techniques we prove that various spaces of derivations or one-sided multipliers over certain operator algebras are reflexive. A sample result: any bounded local derivation (local left multiplier) on an automorphic semicrossed product $C(\Omega) \times_{\sigma} \mathbb{Z}^+$ is a derivation (resp. left multiplier). In the process we obtain various results of independent interest. In particular, we show that the finite dimensional nest representations of the tensor algebra of a topological graph separate points.

1. INTRODUCTION

One of the early results of Barry Johnson [17] asserts that if \mathcal{A} is a semisimple Banach algebra¹ and $S : \mathcal{A} \rightarrow \mathcal{A}$ is an operator that leaves invariant all closed left ideals of \mathcal{A} , then S is a left multiplier of \mathcal{A} . This result has been the source of inspiration for subsequent work, including its recent use by the author [23] to cast new light on a familiar behavior for the adjointable operators on a Hilbert C^* -module.

It is easy to see that the maps preserving all closed left ideals of a Banach algebra \mathcal{A} coincide with the approximately local left multipliers on \mathcal{A} . If \mathcal{X} is a right \mathcal{A} -module, then a map $S : \mathcal{A} \rightarrow \mathcal{X}$ is said to be an *approximately local left multiplier* iff for any $A \in \mathcal{A}$ there exists a sequence $\{X_{A,n}\}_n$ in \mathcal{X} so that $S(A) = \lim_n X_{A,n}A$. One of the motivating questions for the present work is to what extent Johnson's Theorem is valid beyond semisimple operator algebras, i.e., for which operator algebras all approximately local left multipliers are left multipliers. Equivalently, we ask for which operator algebras \mathcal{A} , the algebra $LM(\mathcal{A})$ consisting of all left multipliers on \mathcal{A} is reflexive, i.e., $\text{alg lat } LM(\mathcal{A}) = LM(\mathcal{A})$. (Here we view $LM(\mathcal{A})$ as a subalgebra of all bounded operators on \mathcal{A} .)

This line of investigation is not new. Don Hadwin, Jiankui Li and their collaborators [18–21, 31] have been investigating questions of this type for the past 20 years for various reflexive operator algebras through the use of idempotents and their separating spaces. In particular, Hadwin and Li [21] have shown that Johnson's Theorem holds for all CSL algebras. Also Brešar, Šemrl and others have been investigating local multipliers (and derivations) in various other settings, including the purely algebraic, again through an intricate use of idempotents [2, 3]. The

Received by the editors June 11, 2014 and, in revised form, June 19, 2014.

2010 *Mathematics Subject Classification*. Primary 46L08, 47B49, 47L40, 47L65.

Key words and phrases. Local derivation, local multiplier, reflexivity, topological graph, tensor algebra, C^* -correspondence, Cuntz-Pimsner C^* -algebra.

¹In this paper all Banach algebras have an approximate identity and all linear mappings are necessarily bounded.

present paper advocates the more systematic use of representation theory on this line of investigation and adds new examples which were previously inaccessible to the list of operator algebras for which Johnson's Theorem is valid. In particular, Theorem 3.4 shows that all approximately local left multipliers for the tensor algebra of a topological graph are actually left multipliers. This applies to various algebras that are currently under investigation, including Peters' semicrossed products, the tensor algebras of multivariable dynamics, subalgebras of Stacey's and Exel's crossed products and the tensor algebras of graphs. Theorem 3.4 is proven with the use of representation theory and a crucial step in its proof is a result of independent interest: the finite dimensional nest representations of the tensor algebra of a topological graph separate points (Theorem 3.2). This generalizes one of the main results of an earlier work of Davidson and the author [11, Theorem 4.7].

It turns out that the problem of deciding whether approximately local multipliers are actually multipliers is intimately related to the study of local derivations. The concept of a local derivation was introduced by Dick Kadison in his seminal paper [22] and was further studied by Johnson [16], Larson [29], Larson and Sourour [30] and many others. If \mathcal{X} is a bimodule of a Banach algebra \mathcal{A} , then a map $\delta : \mathcal{A} \rightarrow \mathcal{X}$ is said to be a *local derivation* iff for every $A \in \mathcal{A}$, there exists derivation δ_A so that $\delta(A) = \delta_A(A)$. (The obvious approximate version of this definition establishes the concept of approximately local derivation.) The question of whether all approximately local derivations of \mathcal{A} are actually derivations is equivalent to the reflexivity (in the sense of [29]) of the space $\mathcal{Z}_1(\mathcal{A})$ of all derivations and includes as a special case the corresponding one for local derivations. Johnson established the much stronger result that the space $\mathcal{Z}_1(\mathcal{A}, \mathcal{X})$ of all derivations from a C^* -algebra \mathcal{A} into an \mathcal{A} -module is reflexive thus generalizing an earlier result of Kadison [22]. Hadwin and Li [20] and Crist [8] have established the reflexivity for $\mathcal{Z}_1(\mathcal{A})$ in the case where \mathcal{A} is a CD CSL algebra or a direct limit of finite dimensional CSL algebras respectively. Beyond these two classes, very little is known regarding the reflexivity of $\mathcal{Z}_1(\mathcal{A})$ for a non-selfadjoint operator algebra \mathcal{A} . (Note however [31].)

The other central result of this paper, Theorem 4.5, establishes that approximately local derivations are derivations for a large class of tensor algebras of topological graphs. The proof is more involved than that of Theorem 3.4 and introduces new tools, including that of an acyclic discrete graph and its associated representations. It turns out that if there are sufficiently many acyclic or transitive discrete graphs associated with a topological graph, then one can build a nice representation theory for its tensor algebra (see Lemmas 4.2 and 4.4). Our theory applies in particular to Peters' semicrossed products $C_0(\Omega) \times_{\sigma} \mathbb{Z}^+$ when σ is a homeomorphism on Ω [1, 10, 34], to the tensor subalgebras of Exel and Vershik's crossed products by topologically free coverings [4, 5, 14], to tensor algebras of Exel's crossed products by partial automorphisms [13] and to various tensor algebras of multivariable dynamical systems [9]. In all these cases, approximately local derivations are derivations. We also obtain a similar result for various non-selfadjoint algebras which do not come from C^* -correspondences. In particular, Proposition 4.13 implies that any local derivation on the universal non-selfadjoint algebra generated by n contractions is a derivation.

Unlike the case of C^* -algebras [16] or CSL algebras [21], a local derivation on a tensor algebra \mathcal{A} with range in a Banach \mathcal{A} -bimodule need not be a derivation. Counterexamples to demonstrate that behavior are presented in the last section of

the paper. The author welcomes this phenomenon and believes that it will lead to computable isomorphism invariants for these algebras. This will be explored in a subsequent work.

2. REVISITING THE GENERAL THEORY

In this section we gather various results needed for the rest of the paper. Theorems 2.1 and 2.3 are due to Don Hadwin, Jiankui Li and their collaborators [18–21, 31]. Both results have appeared repeatedly in the literature in various forms and degrees of generality. It turns out that in the form needed here, both have quite elementary proofs (even in the non-unital case) which we include. Examples 2.4 and 2.5 at the end of the section appear to be new; these are quite crucial for building the counterexamples of the subsequent sections.

Theorem 2.1. *Let \mathcal{A} be a Banach algebra generated (as a Banach space) by its idempotents and \mathcal{X} be a right Banach \mathcal{A} -module. Then any approximately local left multiplier from \mathcal{A} into \mathcal{X} is a multiplier. Hence $LM(\mathcal{A}, \mathcal{X})$ is reflexive.*

Proof. Let $S : \mathcal{A} \rightarrow \mathcal{X}$ be an approximate left multiplier. Note that for any $A, B, P \in \mathcal{A}$ with $P = P^2$, we have $S(ABP) \in \overline{\mathcal{X}BP}$ and $S(AB(I - P)) \in \overline{\mathcal{X}(B - BP)}$. Therefore,

$$\begin{aligned} S(AB)P &= S(ABP)P + S(AB(I - P))P \\ &= S(ABP)P = S(ABP). \end{aligned}$$

Letting B range over an approximate unit for \mathcal{A} , we obtain $S(AP) = S(A)P$. Since \mathcal{A} is generated by its idempotents, S is a left multiplier, as desired. □

One can easily see that a slight modification of the above proof also implies that the space $\text{End}_{\mathcal{A}}(\mathcal{X})$ of right \mathcal{A} -module operators on \mathcal{X} is also reflexive. We leave the details to the reader.

Lemma 2.2. *Let \mathcal{X} be a Banach \mathcal{A} -bimodule for a unital Banach algebra \mathcal{A} and let $\delta : \mathcal{A} \rightarrow \mathcal{X}$ be an approximately local derivation. Then*

$$\delta(PQ) = \delta(P)Q + P\delta(Q)$$

for any idempotents $P, Q \in \mathcal{A}$.

Proof. Since δ is an approximately local derivation,

$$P^\perp \delta(PQ)Q^\perp = P\delta(P^\perp Q)Q^\perp = 0,$$

where $P^\perp \equiv I - P$ and similarly for Q^\perp . Hence

$$(1) \quad \delta(PQ)Q^\perp = P\delta(PQ)Q^\perp = P\delta(Q)Q^\perp$$

and similarly

$$\delta(PQ^\perp)Q = P\delta(Q^\perp)Q.$$

By replacing in the above equation Q^\perp with $I - Q$ and then distributing, we obtain

$$\delta(P)Q - \delta(PQ) = P\delta(I)Q - P\delta(Q)Q$$

and since $\delta(I) = 0$,

$$(2) \quad \delta(PQ)Q = \delta(P)Q + P\delta(Q)Q.$$

The conclusion follows by adding (1) and (2). □

Theorem 2.3. *Let \mathcal{A} be a Banach algebra generated by its idempotents and \mathcal{X} a Banach \mathcal{A} -bimodule. Then $\mathcal{Z}_1(\mathcal{A}, \mathcal{X})$ is reflexive.*

Proof. Extend δ by linearity to an approximate derivation $\hat{\delta}$ of the unitization \mathcal{A}_1 of \mathcal{A} , by setting $\hat{\delta}(I) = 0$. Then the extension $\hat{\delta}$, and therefore δ itself, satisfies the conclusion of the above result for $P, Q \in \mathcal{A}$. Since \mathcal{A} is generated by its idempotents, the result follows. □

In the papers [18–21, 31, 36] the reader will find more general results regarding local maps on Banach algebras. We now present some counterexamples which demonstrate that even in the simplest of situations, local maps may fail the corresponding global property.

Example 2.4. If

$$\mathfrak{A} = \left\{ \begin{pmatrix} \lambda & \mu \\ 0 & \lambda \end{pmatrix} \mid \lambda, \mu \in \mathbb{C} \right\} = \{ \lambda I + \mu E_{12} \mid \lambda, \mu \in \mathbb{C} \},$$

then

$$S_{\mathfrak{A}} : \mathfrak{A} \longrightarrow \mathfrak{A} : \lambda I + \mu E_{12} \longmapsto \lambda I + 2\mu E_{12}$$

is a local multiplier which is not a multiplier.

Indeed if $\lambda \neq 0$, then $S_{\mathfrak{A}}(\lambda I + \mu E_{12}) = (I + \mu/\lambda E_{12})(\lambda I + \mu E_{12})$ or otherwise $S_{\mathfrak{A}}(\mu E_{12}) = 2I(\mu E_{12})$. This shows that $S_{\mathfrak{A}}$ is a local multiplier. It is easy to see that in the case $\lambda \neq 0$, the factor $(I + \mu/\lambda E_{12})$ is uniquely determined by $\lambda I + \mu E_{12}$ and so $S_{\mathfrak{A}}$ cannot be a multiplier.

Example 2.5. Let $\{E_{ij}\}_{i,j=1}^3$ be the standard matrix unit system of $M_3(\mathbb{C})$ and let $N = E_{12} + E_{23}$.

If

$$\mathfrak{B} = \left\{ \begin{pmatrix} \lambda & \mu & \nu \\ 0 & \lambda & \mu \\ 0 & 0 & \lambda \end{pmatrix} \mid \lambda, \mu, \nu \in \mathbb{C} \right\} = \{ \lambda I + \mu N + \nu N^2 \mid \lambda, \mu, \nu \in \mathbb{C} \},$$

then

$$\delta_{\mathfrak{B}} : \mathfrak{B} \longrightarrow \mathfrak{B} : \lambda I + \mu N + \nu N^2 \longmapsto \nu N^2$$

is a local derivation which is not a derivation.

Indeed, it is easy to see that the mappings

$$\delta_1 : \mathfrak{B} \longrightarrow \mathfrak{B} : \lambda I + \mu N + \nu N^2 \longmapsto \mu N^2$$

and

$$\delta_2 : \mathfrak{B} \longrightarrow \mathfrak{B} : X \longmapsto (E_{11} + E_{22})X - X(E_{11} + E_{22})$$

are derivations. Furthermore, if $X = \lambda I + \mu N + \nu N^2$, then

$$\delta_{\mathfrak{B}}(X) = \begin{cases} (\nu/\mu)\delta_1(X) & \text{if } \mu \neq 0, \\ \delta_2(X) & \text{otherwise.} \end{cases}$$

Therefore, $\delta_{\mathfrak{B}}$ is a local derivation.

However, $\delta_{\mathfrak{B}}$ is not a derivation. Indeed, if $X = \lambda I + \mu N + \nu N^2$ and $X' = \lambda' I + \mu' N + \nu' N^2$, then the (1, 3)-entry of

$$\delta_{\mathfrak{B}}(XX') - \delta_{\mathfrak{B}}(X)X' - X\delta_{\mathfrak{B}}(X')$$

is equal to $\mu\mu'$, which is not equal to 0 in general.

Example 2.6 (Crist [8]). Let $\{E_{ij}\}$ be the standard matrix unit system for $M_3(\mathbb{C})$. If

$$\mathfrak{C} = \mathbb{C}I + [E_{12}, E_{13}, E_{23}]$$

and

$$\delta_{\mathfrak{C}} : \mathfrak{C} \longrightarrow \mathfrak{C} : \sum x_{ij}E_{ij} \longmapsto (2x_{13} - x_{12} + x_{23})E_{13},$$

then $\delta_{\mathfrak{C}}$ is a local derivation which is not a derivation.

3. TENSOR ALGEBRAS, NEST REPRESENTATIONS AND LOCAL MULTIPLIERS

Davidson and the author have shown in [11, Theorem 4.7] that the finite dimensional nest representations of the tensor algebra of a countable graph separate points. In this section we extend this result to all tensor algebras of topological graphs. This applies in particular to algebras associated to one or more dynamical systems such as Peters’ semicrossed products. Combined with the results of the previous section, this result allows us to obtain definitive information regarding the local multipliers on these algebras.

A C^* -correspondence $(X, \mathcal{A}, \varphi_X)$ consists of a C^* -algebra \mathcal{A} , a Hilbert \mathcal{A} -module $(X, \langle \cdot, \cdot \rangle)$ and a non-degenerate $*$ -homomorphism $\varphi_X : \mathcal{A} \rightarrow \mathcal{L}(X)$.

A (Toeplitz) representation (π, t) of a C^* -correspondence into a C^* -algebra \mathcal{B} , is a pair of a $*$ -homomorphism $\pi : \mathcal{A} \rightarrow \mathcal{B}$ and a linear map $t : X \rightarrow \mathcal{B}$, such that

- (i) $\pi(a)t(\xi) = t(\varphi_X(a)(\xi))$,
- (ii) $t(\xi)^*t(\eta) = \pi(\langle \xi, \eta \rangle)$,

for $a \in \mathcal{A}$ and $\xi, \eta \in X$. A representation (π, t) is said to be *injective* iff π is injective; in that case t is an isometry.

The C^* -algebra generated by a representation (π, t) equals the closed linear span of $t^n(\bar{\xi})t^m(\bar{\eta})^*$, where for simplicity $\bar{\xi} \equiv (\xi^1, \dots, \xi^{(n)}) \in X^n$ and $t^n(\bar{\xi}) \equiv t(\xi_1) \dots t(\xi_n)$. For any representation (π, t) there exists a $*$ -homomorphism $\psi_t : \mathcal{K}(X) \rightarrow \mathcal{B}$, such that $\psi_t(\Theta_{\xi, \eta}^X) = t(\xi)t(\eta)^*$.

It is easy to see that for a C^* -correspondence $(X, \mathcal{A}, \varphi_X)$ there exists universal Toeplitz representation, denoted as (π_∞, t_∞) , so that any other representation of $(X, \mathcal{A}, \varphi_X)$ is equivalent to a direct sum of subrepresentations of (π_∞, t_∞) . We define the Cuntz-Pimsner-Toeplitz C^* -algebra \mathcal{T}_X as the C^* -algebra generated by all elements of the form $\pi_\infty(a), t_\infty(\xi)$, $a \in \mathcal{A}$, $\xi \in X$. The algebra \mathcal{T}_X satisfies the following universal property: for any Toeplitz representation (π, t) of X , there exists a representation $\pi \times t$ of \mathcal{T}_X so that $\pi(a) = ((\pi \times t) \circ \pi_\infty)(a)$, $\forall a \in \mathcal{A}$, and $t(\xi) = ((\pi \times t) \circ t_\infty)(\xi)$, $\forall \xi \in X$.

Definition 3.1. The *tensor algebra* \mathcal{T}_X^+ of a C^* -correspondence $(X, \mathcal{A}, \varphi_X)$ is the norm-closed subalgebra of \mathcal{T}_X generated by all elements of the form $\pi_\infty(a), t_\infty(\xi)$, $a \in \mathcal{A}$, $\xi \in X$.

It is worth mentioning here that \mathcal{T}_X^+ also sits naturally inside the Cuntz-Pimsner algebra \mathcal{O}_X associated with the C^* -correspondence X ; this follows from [24, 32]. As we will not be making essential use of that theory here, we skip the pertinent definitions and results and instead direct the reader to [24, 32] for more details.

The tensor algebras for C^* -correspondences were pioneered by Muhly and Solel in [32]. They form a broad class of non-selfadjoint operator algebras which includes as special cases Peters’ semicrossed products [34], Popescu’s non-commutative disc algebras [35], the tensor algebras of graphs (introduced in [32] and further studied

in [25]) and the tensor algebras for multivariable dynamics [9], to mention but a few.

Due to its universality, the Cuntz-Pimsner-Toeplitz C^* -algebra \mathcal{T}_X admits a gauge action that leaves $\pi_\infty(A)$ elementwise invariant and “twists” each $t_\infty(\xi)$, $\xi \in X$, by a unimodular scalar. Using this action, and reiterating a familiar trick with the Fejer kernel, one can verify that each element $T \in \mathcal{T}_X^+$ admits a Fourier series expansion

$$(3) \quad T = \pi_\infty(a) + \sum_{n=1}^\infty t_\infty^n(\bar{\xi}_n), \quad a \in A, \bar{\xi}_n \in \mathcal{X}^n, n = 1, 2, \dots,$$

where the summability is in the Cesaro sense. (See [26, 32] for more details.)

A broad class of C^* -correspondences arises naturally from the concept of a topological graph. A topological graph $\mathcal{G} = (\mathcal{G}^0, \mathcal{G}^1, r, s)$ consists of two σ -locally compact² spaces $\mathcal{G}^0, \mathcal{G}^1$, a continuous proper map $r : \mathcal{G}^1 \rightarrow \mathcal{G}^0$ and a local homeomorphism $s : \mathcal{G}^1 \rightarrow \mathcal{G}^0$. The set \mathcal{G}^0 is called the base (vertex) space and \mathcal{G}^1 the edge space. When \mathcal{G}^0 and \mathcal{G}^1 are both equipped with the discrete topology, we have a discrete countable graph (see below).

With a topological graph $\mathcal{G} = (\mathcal{G}^0, \mathcal{G}^1, r, s)$ there is a C^* -correspondence $X_{\mathcal{G}}$ over $C_0(\mathcal{G}^0)$. The right and left actions of $C_0(\mathcal{G}^0)$ on $C_c(\mathcal{G}^1)$ are given by

$$(fFg)(e) = f(r(e))F(e)g(s(e))$$

for $F \in C_c(\mathcal{G}^1)$, $f, g \in C_0(\mathcal{G}^0)$ and $e \in \mathcal{G}^1$. The inner product is defined for $F, G \in C_c(\mathcal{G}^1)$ by

$$\langle F | G \rangle (v) = \sum_{e \in s^{-1}(v)} \overline{F(e)}G(e)$$

for $v \in \mathcal{G}^0$. Finally, $X_{\mathcal{G}}$ denotes the completion of $C_c(\mathcal{G}^1)$ with respect to the norm

$$(4) \quad \|F\| = \sup_{v \in \mathcal{G}^0} \langle F | F \rangle (v)^{1/2}.$$

When \mathcal{G}^0 and \mathcal{G}^1 are both equipped with the discrete topology, then the tensor algebra $\mathcal{T}_{\mathcal{G}}^+ \equiv \mathcal{T}_{X_{\mathcal{G}}}^+$ associated with \mathcal{G} coincides with the quiver algebra of Muhly and Solel [32]. In that case, $\mathcal{T}_{\mathcal{G}}^+$ has a natural presentation which we now describe.

Let $\mathcal{G} = (\mathcal{G}^0, \mathcal{G}^1, r, s)$ be a countable discrete graph and let \mathcal{G}^∞ be the path space of \mathcal{G} . This consists of all vertices $v \in \mathcal{G}^0$ and all paths $p = e_k e_{k-1} \dots e_1$, where the e_i are edges satisfying $s(e_i) = r(e_{i-1})$, $i = 1, 2, \dots, k$, $k \in \mathbb{N}$. (Paths of the form $p = e_k e_{k-1} \dots e_1$ are said to have length $|p| = k$ and vertices are called paths of length 0.) The maps r and s extend to \mathcal{G}^∞ in the obvious way, two paths p_1 and p_2 are composable whenever $s(p_2) = r(p_1)$ and in that case, the composition $p_2 p_1$ is just the concatenation of p_1 and p_2 . Let $\{\xi_p\}_{p \in \mathcal{G}^\infty}$ denote the usual orthonormal basis of the Fock space $\mathcal{H}_{\mathcal{G}} \equiv l^2(\mathcal{G}^\infty)$, where ξ_p is the characteristic function of $\{p\}$. The left creation operator L_q , $q \in \mathcal{G}^\infty$, is defined by

$$L_q \xi_p = \begin{cases} \xi_{qp} & \text{if } s(q) = r(p), \\ 0 & \text{otherwise.} \end{cases}$$

By [15, Corollary 2.2], the algebra generated by $\{L_p \mid p \in \mathcal{G}^\infty\}$ is (completely isometrically) isomorphic to $\mathcal{T}_{\mathcal{G}}^+$. Its weak closure, denoted as $\mathcal{L}_{\mathcal{G}}$, is the familiar free semigroupoid algebra of \mathcal{G} , first studied by Kribs and Power [28].

²Due to this assumption, all discrete graphs appearing in this paper are countable.

Another class of examples for topological graphs arises from multivariable dynamical systems. If Ω is a σ -locally compact Hausdorff space equipped with n proper continuous self maps $\sigma = (\sigma_1, \sigma_2, \dots, \sigma_n)$, then let $\mathcal{G}^0 = \Omega$, $\mathcal{G}^1 = \{1, 2, \dots, n\} \times \Omega$, let $s : \mathcal{G}^1 \rightarrow \mathcal{G}^0$ be the natural projection and let $r(i, x) = \sigma_i(x)$. This defines a topological graph denoted as $\mathcal{G}(\Omega, \sigma)$ and such graphs are naturally associated to various familiar operator algebras. Indeed in the case of a single variable dynamical system (Ω, σ) , the tensor algebra $\mathcal{T}_{\mathcal{G}(\Omega, \sigma)}^+$ is completely isometrically isomorphic to Peters' semicrossed product $C_0(\Omega) \times_{\sigma} \mathbb{Z}^+$ [34], which was first studied by Arveson [1]. This is the universal non-selfadjoint operator algebra generated by a copy of $C_0(\Omega)$ and an isometry V , subject to the covariance relations $fV = V(f \circ \sigma)$, $f \in C_0(\Omega)$. For more general dynamical systems (Ω, σ) , the tensor algebra $\mathcal{T}_{\mathcal{G}(\Omega, \sigma)}^+$, usually denoted as $\mathcal{T}^+(\Omega, \sigma)$, was first studied by Davidson and the author in [9].

The following generalizes one of the main results of an earlier work of Davidson and the author [11, Theorem 4.7] and it is essential for the main result of this section, Theorem 3.4.

Theorem 3.2. *If $\mathcal{G} = (\mathcal{G}^0, \mathcal{G}^1, r, s)$ is a topological graph, then the finite dimensional nest representations of its tensor algebra $\mathcal{T}_{\mathcal{G}}^+$ separate points.*

Proof. We will construct a family of representations (π_v, t_v) , $v \in \mathcal{G}^0$, of the correspondence $X_{\mathcal{G}}$ so that the integrated representations $\varphi_v \equiv \pi_v \times t_v$ satisfy:

- (i) for each $v \in \mathcal{G}^0$ there exists a discrete graph \mathcal{G}_v so that

$$(5) \quad \overline{\varphi_v(\mathcal{T}_{\mathcal{G}}^+)}^{\text{wot}} = \mathcal{L}_{\mathcal{G}_v},$$

where $\mathcal{L}_{\mathcal{G}_v}$ is the free semigroupoid algebra associated with \mathcal{G}_v ,

- (ii) the representations φ_v , $v \in \mathcal{G}^0$, separate the points in $\mathcal{T}_{\mathcal{G}}^+$.

Once this is done, the proof follows from the result of Davidson and Katsoulis [11, Theorem 4.7] that the weakly continuous finite dimensional nest representations of the free semigroupoid algebras separate points.

Fix a $v \in \mathcal{G}^0$ and define inductively

$$\begin{aligned} \mathcal{G}_{v, n+1}^0 &= r(\mathcal{G}_{v, n}^1), \\ \mathcal{G}_{v, n+1}^1 &= s^{-1}(\mathcal{G}_{v, n+1}^0), \quad n = 0, 1, \dots, \end{aligned}$$

where $\mathcal{G}_{v, 0}^0 = \{v\}$ and $\mathcal{G}_{v, 0}^1 = s^{-1}(\{v\})$. If

$$\mathcal{G}_v^i = \bigcup_{n \geq 0} \mathcal{G}_{v, n}^i, \quad i = 0, 1,$$

then we define $\mathcal{G}_v \equiv (\mathcal{G}_v^0, \mathcal{G}_v^1, r, s)$, where r and s are the restrictions on \mathcal{G}_v^1 of the corresponding maps coming from $\mathcal{G} = (\mathcal{G}^0, \mathcal{G}^1, r, s)$. (The family of discrete graphs \mathcal{G}_v , $v \in \mathcal{G}^0$, appearing in this proof is said to be the *family of discrete graphs associated with \mathcal{G}* .)

Let \mathcal{G}_v^∞ be the (finite) path space of \mathcal{G}_v and let $\mathcal{H}_{\mathcal{G}_v}$ be the Fock Hilbert space associated with \mathcal{G}_v . For $f \in C_0(\mathcal{G}^0)$ and $F \in C_c(\mathcal{G}^1)$ we define

$$\pi_v(f)\xi_p = f(r(p))\xi_p$$

and

$$t_v(F)\xi_p = \sum_{e \in s^{-1}(r(p))} \overline{F(e)}\xi_{ep}, \quad p \in \mathcal{G}_v^\infty,$$

with the understanding that $r(p) = p$, when $p \in \mathcal{G}_v^0$.

Clearly,

$$\|t_v(F)\xi_p\|^2 = \|\langle F | F \rangle (r(p))\| \leq \|F\|$$

for any $p \in \mathcal{G}_v^\infty$. Furthermore, if $p, p' \in \mathcal{G}_v^\infty$ are distinct paths, then the vectors $t_v(F)\xi_p$ and $t_v(F)\xi_{p'}$ are orthogonal. Hence $t_v(F)$ extends to a bounded operator on $\mathcal{H}_{\mathcal{G}_v}$. Its adjoint satisfies $t_v^*(F)\xi_w = 0$, when $w \in \mathcal{G}_v^0$, and

$$t_v^*(F)\xi_{ep} = \overline{F(e)}\xi_p$$

otherwise. Therefore,

$$\begin{aligned} t_v(F)^*t_v(G)\xi_p &= t_v(F)^* \left(\sum_{e \in s^{-1}(r(p))} G(e)\xi_{ep} \right) \\ &= \sum_{e \in s^{-1}(r(p))} \overline{F(e)}G(e)\xi_p \\ &= \langle F | G \rangle (r(p))\xi_p = \pi_v(\langle F | G \rangle)\xi_p. \end{aligned}$$

A simple calculation also shows that $\pi_v(f)t_v(F)\pi_v(g) = t_v(fFg)$, for all $f, g \in C_0(\mathcal{G}^0)$ and $F \in C_c(\mathcal{G}^1)$. Hence (π_v, t_v) extends to an isometric covariant representation of the correspondence $X_{\mathcal{G}}$.

Let $\pi \equiv \bigoplus_{v \in \mathcal{G}^0} \pi_v$ and $t \equiv \bigoplus_{v \in \mathcal{G}^0} t_v$. We will show now that the C^* -algebra $C^*(\pi, t)$ generated by $\pi(C_0(\mathcal{G}^0))$ and $t(X_{\mathcal{G}})$ is isomorphic to the Toeplitz-Cuntz-Pimsner C^* -algebra $\mathcal{T}_{\mathcal{G}}$. From this it will follow that the family of representations $\varphi_v, v \in \mathcal{G}^0$, separates the points in $\mathcal{T}_{\mathcal{G}}^+$, since $\mathcal{T}_{\mathcal{G}}^+ \subseteq \mathcal{T}_{\mathcal{G}}$.

By the Gauge Invariant Uniqueness Theorem of Katsura [27, Theorem 6.2], it suffices to show that π is faithful on $C_0(\mathcal{G}^0)$, the C^* -algebra $C^*(\pi, t)$ admits a gauge action and finally that

$$I(\pi, t) \equiv \{f \in C_0(\mathcal{G}^0) \mid \pi(f) \in \psi_t(K(X_{\mathcal{G}}))\} = 0.$$

Since $\pi_v(f)\xi_v = f(v)\xi_v, v \in \mathcal{G}^0$, it is clear that π is faithful. To define a gauge action on $C^*(\pi, t)$ let

$$\alpha_\lambda : C^*(\pi, t) \longrightarrow C^*(\pi, t) : A \longmapsto U_\lambda^*AU_\lambda, \quad \lambda \in \mathbb{T},$$

where $U_\lambda\xi_p = \lambda^{|p|}\xi_p, p \in \mathcal{G}_v^\infty$ and $|p|$ equals the length of the path p . Finally, for any $G \in X_{\mathcal{G}}$, we have $t_v(G)^*\xi_v = 0$, for all $v \in \mathcal{G}^0$. However if $\pi(f) \in I(\pi, t)$, then $\pi_v(f)$ can be approximated by linear combinations of elements of the form

$$t_v(F_1) \dots t_v(F_k)t_v(G_1)^* \dots t_v(G_k)^*, \quad k = 1, 2, \dots$$

Since $t_v(G)^*\xi_v = 0$, for all $G \in X_{\mathcal{G}}$, we have

$$\|f(v)\| = \|f(v)\xi_v\| = \|\pi_v(f)\xi_v\| = 0,$$

for all $v \in \mathcal{G}^0$. Hence $f = 0$, as desired.

It remains to verify the validity of (5). Let v_1, v_2, \dots be an enumeration of \mathcal{G}_v^0 . Fix an $i \in \mathbb{N}$; for each $n \in \mathbb{N}$ use Urysohn's Lemma to obtain $f_n = f_{n,i} \in C_0(\mathcal{G}_v^0)$ with

$$f_n(v_i) = 1, \quad f_n(v_j) = 0, \quad \text{for } 1 \leq j \leq n, j \neq i,$$

and $|f_n(v_j)| \leq 1$, for all other j . If \hat{L}_{v_i} is any weak limit of $\{\pi_v(f_n)\}_n$, then $\hat{L}_{v_i}\xi_p = \xi_p$, if $r(p) = v_i$, or otherwise $\hat{L}_{v_i}\xi_p = 0$. Hence, $\hat{L}_{v_i} = L_{v_i}$, for all i . Furthermore,

$$(6) \quad \overline{\varphi_v(\mathcal{T}_{\mathcal{G}}^+)}^{\text{wot}} \ni t_v(F)L_{v_i} = \sum_{e \in s^{-1}(v_i)} F(e)L_e \in \mathcal{L}_{\mathcal{G}}.$$

However $\sum_{i=1}^\infty L_{v_i} = I$ and so (6) implies that

$$t_v(F) = \sum_{i=1}^\infty t_v(F)L_{v_i} \in \mathcal{L}_{\mathcal{G}}, \quad F \in C_c(\mathcal{G}_v^1),$$

i.e., $\overline{\varphi_v(\mathcal{T}_{\mathcal{G}}^+)}^{\text{wot}} \subseteq \mathcal{L}_{\mathcal{G}}$. To obtain the reverse inclusion, let e_1, e_2, \dots be an enumeration of \mathcal{G}_v^1 . Fix an $i \in \mathbb{N}$; for each $n \in \mathbb{N}$ use Urysohn’s Lemma to obtain $F_n \in C_c(\mathcal{G}_v^1)$ with

$$F_n(e_i) = 1, \quad F_n(e_j) = 0, \quad \text{for } 1 \leq j \leq n, j \neq i,$$

and $|F_n(e_j)| \leq 1$, for all other j . If $v_{j_i} = s(e_i)$, then again by (6) any weak limit of $\{t_v(F_n)L_{v_{j_i}}\}_n$ will equal L_{e_i} and the conclusion follows. \square

Remark 3.3. The representations φ_v in the proof of Theorem 3.2 were inspired by the familiar orbit representations (see [9, 34]). However, it is frequently the case that these two classes of representations differ.

Indeed, let (Ω, σ) be a dynamical system on a locally compact Hausdorff space Ω and recall that Peters’ semicrossed product $C_0(\Omega) \times_\sigma \mathbb{Z}^+$ is the tensor algebra corresponding to the topological graph $\mathcal{G} = (\mathcal{G}^0, \mathcal{G}^1, r, s)$ with $\mathcal{G}^0 = \mathcal{G}^1 = \Omega$, $s = \text{id}$ and $r = \sigma$. Let $v \in \Omega$ and let \mathcal{H} be a separable Hilbert space with orthonormal base $\{\xi_n\}_{n=0}^\infty$. The orbit representation λ_v of $C_0(\Omega) \times_\sigma \mathbb{Z}^+$ is defined by

$$\begin{aligned} \lambda_v(f)\xi_n &= f(\sigma^{(n)}(v))\xi_n, \quad f \in C_0(\Omega), \quad n = 0, 1, 2, \dots, \\ \lambda_v(V) &= S, \end{aligned}$$

where $V \in C_0(\Omega) \times_\sigma \mathbb{Z}^+$ is the universal isometry and S the forward shift on $\{\xi_n\}_{n=0}^\infty$.

It is easy to see that when $v \in \Omega$ is aperiodic (Definition 4.6), then $\lambda_v(C_0(\Omega) \times_\sigma \mathbb{Z}^+)$ is SOT-dense in the nest algebra of all lower triangular infinite matrices with respect to $\{\xi_n\}_{n=1}^\infty$. Therefore, $\lambda_v(C_0(\Omega) \times_\sigma \mathbb{Z}^+)'' = B(\mathcal{H})$. On the other hand, the range of any of the $\varphi_{v'}$, $v' \in \Omega$, constructed above is SOT-dense in a free semigroupoid algebra and so [28, Corollary 4.5] implies that $\varphi_{v'}(C_0(\Omega) \times_\sigma \mathbb{Z}^+)''$ is a free semigroupoid algebra. Since $B(\mathcal{H})$ is not a free semigroupoid algebra, no such $\varphi_{v'}$, $v' \in \Omega$, can be unitarily equivalent to λ_v , when v is aperiodic.

In [31] it was observed that the representation theory of Davidson and the author [11] has definitive applications on the theory of local maps for graph algebras. The following extends the results of [31] to a much broader class of operator algebras.

Theorem 3.4. *If $\mathcal{G} = (\mathcal{G}^0, \mathcal{G}^1, r, s)$ is a topological graph, then any approximately local left multiplier on $\mathcal{T}_{\mathcal{G}}^+$ is actually a left multiplier, i.e., the algebra $LM(\mathcal{T}_{\mathcal{G}}^+)$ of left multipliers on $\mathcal{T}_{\mathcal{G}}^+$ is reflexive.*

Proof. Let S be an approximately local multiplier on $\mathcal{T}_{\mathcal{G}}^+$. By the previous theorem, there exists a separating family of representations

$$\rho_i : \mathcal{T}_{\mathcal{G}}^+ \rightarrow B(\mathcal{H}_i), \quad i \in \mathbb{I},$$

on finite dimensional Hilbert space so that each $\rho_i(\mathcal{T}_{\mathcal{G}}^+)$ is a finite dimensional nest algebra. Furthermore, $\rho_i(\mathcal{T}_{\mathcal{G}}^+)$ is a right $\mathcal{T}_{\mathcal{G}}^+ / \ker \rho_i$ -module, with the right action coming from ρ_i . Since S preserves closed left ideals we obtain

$$S_i : \mathcal{T}_{\mathcal{G}}^+ / \ker \rho_i \longrightarrow \rho_i(\mathcal{T}_{\mathcal{G}}^+); A + \ker \rho_i \longmapsto \rho_i(S(A)), \quad i \in \mathbb{I}.$$

It is easy to verify that S_i is an approximate left multiplier and so Theorem 2.1 implies that S_i is actually a left multiplier. Hence

$$S_i(AB + \ker \rho_i) = S_i(A + \ker \rho_i)\rho_i(B)$$

and so

$$\rho_i(S(AB) - S(A)B) = 0, \quad \text{for all } i \in \mathbb{I}.$$

Since $\bigcap_i \ker \rho_i = \{0\}$, the conclusion follows. □

Corollary 3.5. *Let (Ω, σ) be a multivariable dynamical system and let $\mathcal{T}^+(\Omega, \sigma)$ be its tensor algebra. Then any approximately local left multiplier on $\mathcal{T}^+(\Omega, \sigma)$ is a left multiplier.*

There is another class of operator algebras associated to a multivariable dynamical system (Ω, σ) , the so-called semicrossed product $C_0(\Omega) \times_{\sigma} \mathbb{F}_n^+$, which was introduced and first studied by Davidson and the author in [9]. We do not know whether approximately local left multipliers are left multipliers in general but the following provides an answer in a particular situation.

Proposition 3.6. *Let (\mathcal{A}, σ) be an automorphic multivariable C^* -dynamical system and assume that the crossed product C^* -algebra $\mathcal{A} \times_{\sigma} \mathbb{F}_n$ is residually finite dimensional (RFD). Then any approximately local left multiplier on $\mathcal{A} \times_{\sigma} \mathbb{F}_n^+$ is a left multiplier.*

Proof. Let U_1, U_2, \dots, U_n be the free unitaries in $\mathcal{A} \times_{\sigma} \mathbb{F}_n$ implementing the covariance relations. Let $\pi_i, i \in \mathbb{I}$, be a family of irreducible representations that separates the points in $\mathcal{A} \times_{\sigma} \mathbb{F}_n$ and let $\hat{\pi}_i, i \in \mathbb{I}$, be the restriction of that family on $\mathcal{A} \times_{\sigma} \mathbb{F}_n^+$. Since $\pi_i(\mathcal{A} \times_{\sigma} \mathbb{F}_n^+)$ is an operator algebra acting on a finite dimensional space, it is inverse closed and therefore it contains the adjoints of the unitary operators $\pi_i(U_j), i = 1, 2, \dots, n$. Hence $\pi_i(\mathcal{A} \times_{\sigma} \mathbb{F}_n^+)$ equals $\pi_i(\mathcal{A} \times_{\sigma} \mathbb{F}_n)$ and so $\hat{\pi}_i, i \in \mathbb{I}$, is a separating family of finite dimensional irreducible representations for $\mathcal{A} \times_{\sigma} \mathbb{F}_n^+$. As with Theorem 3.4, this suffices to prove the result. □

Proposition 3.6 applies in particular to the case where $\mathcal{A} = \mathbb{C}$ and σ consists of n -copies of the identity map. Indeed

Corollary 3.7. *Let $C^*(\mathbb{F}_n)$ be the full C^* -algebra of the free group on n -generators and let $A(\mathbb{F}_n^+)$ be the non-selfadjoint subalgebra of $C^*(\mathbb{F}_n)$ generated by the generators. Then any approximately local multiplier on $A(\mathbb{F}_n^+)$ is a multiplier.*

Proof. The result follows from a well-known result of Choi [7, Theorem 1] and Proposition 3.6. □

An interesting result of Samei [36] asserts that if all approximately local one-sided multipliers from a Banach algebra \mathcal{A} into its one-sided modules are one-sided multipliers, then all approximately local derivations on \mathcal{A} -bimodules are derivations. This approach was used by Hadwin and Li [21] to prove that all local derivations on CSL algebras are actually derivations by proving the corresponding result for approximately one-sided multipliers.

It turns out that this approach is not applicable here. Neither Theorem 3.4 nor Corollary 3.5 extend to approximate local left multipliers on a right \mathcal{T}_G^+ -module, instead of \mathcal{T}_G^+ itself. The following provides a broad class of counterexamples.

Example 3.8. Let (Ω, σ) be a dynamical system on a locally compact Hausdorff space and assume that σ has a fixed point $x \in \Omega$. Then there exists a finite dimensional $C_0(\Omega) \times_{\sigma} \mathbb{Z}^+$ -module \mathfrak{A} and a local left multiplier $S : C_0(\Omega) \times_{\sigma} \mathbb{Z}^+ \rightarrow \mathfrak{A}$ which is not a left multiplier.

Indeed, let \mathfrak{A} and $S_{\mathfrak{A}}$ be as in Example 2.4. Since $x \in \Omega$ is a fixed point for σ the mapping

$$\pi : C_0(\Omega) \times_{\sigma} \mathbb{Z}^+ \longrightarrow \mathfrak{A}; \sum_{n=0}^{\infty} V^n f_n \longmapsto \begin{pmatrix} f_0(x) & f_1(x) \\ 0 & f_0(x) \end{pmatrix}$$

is a representation that makes \mathfrak{A} a right $C_0(\Omega) \times_{\sigma} \mathbb{Z}^+$ -module with the natural action coming from π . If

$$S : C_0(\Omega) \times_{\sigma} \mathbb{Z}^+ \longrightarrow \mathfrak{A}; A \longmapsto S_{\mathfrak{A}}(\pi(A)),$$

then S is a local left multiplier which is not a left multiplier (otherwise $S_{\mathfrak{A}}$ would be a multiplier as well).

4. LOCAL DERIVATIONS

In this section we investigate whether an analogue of Theorem 3.4 is valid for approximately local derivation instead of multipliers. This is a harder problem and the following illustrates where the difficulty lies.

Proposition 4.1. *Let \mathcal{A} be an operator algebra and assume that*

- (i) *the finite dimensional nest representations separate points in \mathcal{A} ,*
- (ii) *any derivation $\delta : \mathcal{A} \rightarrow \mathcal{A}$ leaves invariant any closed ideal.*

Then any approximately local derivation $\delta : \mathcal{A} \rightarrow \mathcal{A}$ is a derivation.

Proof. Let δ be an approximately local derivation on \mathcal{A} and let $\{\varphi_i\}_{i \in \mathbb{I}}$ be a family of finite dimensional nest representations that separate points in \mathcal{A} . Since locally δ can be approximated by derivations, it leaves all closed ideals invariant. Hence we obtain well-defined approximately local derivations

$$\delta_i : \mathcal{A}/\ker \varphi_i \longrightarrow \varphi_i(\mathcal{A}); A + \ker \varphi_i \mapsto \varphi_i(\delta(A)), \quad i \in \mathbb{I},$$

with the right action on $\varphi_i(\mathcal{A})$ coming from φ_i . Since $\mathcal{A}/\ker \varphi_i$ is generated by its idempotents (it is a finite dimensional nest algebra), Theorem 2.3 implies that the δ_i are actually derivations. Hence

$$\varphi_i(\delta(AB) - \delta(A)B - A\delta(B)) = 0,$$

for all $i \in \mathbb{I}$. Since $\bigcap_{i \in \mathbb{I}} \ker \varphi_i = \{0\}$, the conclusion follows. □

If we knew that any derivation on the tensor algebra of a topological graph leaves closed ideals invariant, e.g., it is inner, then the result above combined with Theorem 3.2 would imply that local derivations are derivations. With the exception of [12], it seems that nothing is known in that direction. We therefore adopt a different approach: using the concept of an acyclic graph, we build a more flexible representation theory than that of Theorem 3.2. The drawback is that it is not applicable in all situations.

We begin by recalling some further results regarding the structure of the tensor algebra of a topological graph $\mathcal{G} = (\mathcal{G}^0, \mathcal{G}^1, r, s)$. Let

$$\mathcal{G}^n \equiv \{e_n e_{n-1} \dots e_1 \mid e_i \in \mathcal{G}^1, s(e_{i+1}) = r(e_i), 1 \leq i \leq n\}$$

be the space of paths of length n equipped with the topology it inherits as a subset of $\mathcal{G}^1 \times \mathcal{G}^1 \times \dots \times \mathcal{G}^1$. Equip \mathcal{G}^n with the obvious domain and range maps $s : \mathcal{G}^n \rightarrow \mathcal{G}^0$ and $r : \mathcal{G}^n \rightarrow \mathcal{G}^0$, i.e., $s(e_n e_{n-1} \dots e_1) = s(e_1)$ and $r(e_n e_{n-1} \dots e_1) = r(e_n)$. Then [26, Lemma 1.25] shows that the quadruple $(\mathcal{G}^n, \mathcal{G}^0, r, s)$ becomes a topological graph which we simply denote as \mathcal{G}^n . Furthermore, [26, Proposition 1.27] shows that $X_{\mathcal{G}^n} \simeq X_{\mathcal{G}} \otimes X_{\mathcal{G}} \otimes \dots \otimes X_{\mathcal{G}}$, via the identification

$$(7) \quad (F_n \otimes F_{n-1} \otimes \dots \otimes F_1)(e_n e_{n-1} \dots e_1) = F_n(e_n) F_{n-1}(e_{n-1}) \dots F_1(e_1),$$

where $F_i \in X_{\mathcal{G}}$, $i = 1, 2, \dots, n$, $e_n e_{n-1} \dots e_1 \in \mathcal{G}^n$ and \otimes denotes the internal tensor product of C^* -correspondences.

Now fix a $v \in \mathcal{G}^0$ so that the graph \mathcal{G}_v is acyclic, i.e., no paths of length higher than zero start and end at the same vertex, and recall the representation $\varphi_v = \pi_v \times t_v$ of $\mathcal{T}_{\mathcal{G}}^+$, as it appears in the proof of Theorem 3.2. Using φ_v , we construct now a family of finite dimensional representations for $\mathcal{T}_{\mathcal{G}}^+$ whose kernels satisfy a useful property.

For each $n = 0, 1, 2, \dots$, let

$$\mathcal{F}_{v,n} = \left\{ p \in \mathcal{G}_v^\infty \mid s(p) \in \bigcup_{i=0}^n \mathcal{G}_{v,i}^0 \text{ and } r(p) \in \bigcup_{i=0}^n \mathcal{G}_{v,i}^0 \right\}$$

and

$$\mathcal{H}_{v,n} = \bigvee \{ \xi_p \mid p \in \mathcal{F}_{v,n} \} \subseteq \mathcal{H}_{\mathcal{G}_v}.$$

It is easy to verify that $\mathcal{H}_{v,n}$ is co-invariant for $\varphi_v(\mathcal{T}_{\mathcal{G}}^+)$ and so the compression of φ_v on $\mathcal{H}_{v,n}$ defines a representation of $\mathcal{T}_{\mathcal{G}}^+$ denoted as $\varphi_{v,n}$. Since $\bigvee_n \mathcal{H}_{v,n} = \mathcal{H}_{\mathcal{G}_v}$, the family $\{\varphi_{v,n}\}_{n \in \mathbb{N}}$ separates points in $\mathcal{T}_{\mathcal{G}}^+ / \ker \varphi_v$.

In order to ensure that the representations appearing in the next lemma are finite dimensional we need to assume that the topological graph $\mathcal{G} = (\mathcal{G}^0, \mathcal{G}^1, r, s)$ is *source finite*, i.e., $s^{-1}(\{v\})$ is a finite set for any $v \in \mathcal{G}^0$. This assumption is automatically satisfied when \mathcal{G}^1 is compact and in all applications considered in this paper.

Lemma 4.2. *If $\varphi_{v,n}$ is as above, with \mathcal{G}_v acyclic, then $\varphi_{v,n}(\mathcal{T}_{\mathcal{G}}^+)$ is a finite dimensional operator algebra which is generated by its idempotents.*

Proof. Since \mathcal{G}_v is acyclic, $\mathcal{G}_{v,i}^0 \cap \mathcal{G}_{v,j}^0 = \emptyset$, for $i \neq j$. This implies that the cardinality of $\mathcal{F}_{v,n}$ is finite. Hence $\mathcal{H}_{v,n}$ is finite dimensional and the algebra $\varphi_{v,n}(\mathcal{T}_{\mathcal{G}}^+)$ is generated as a linear space by the (finitely many) compressions $\hat{L}_p \equiv P_{\mathcal{H}_{v,n}} L_p |_{\mathcal{H}_{v,n}}$, $p \in \mathcal{F}_{v,n}$.

If $p \in \mathcal{F}_{v,n} \cap \mathcal{G}_v^0$, then \hat{L}_p is an idempotent (actually an orthogonal projection). Otherwise, $\hat{L}_p = \hat{L}_{r(p)}\hat{L}_p\hat{L}_{s(p)}$ and since \mathcal{G}_v is acyclic we have

$$\hat{L}_{r(p)}\hat{L}_{s(p)} = 0.$$

Hence the identity

$$2\hat{L}_p = (\hat{L}_{r(p)} + \hat{L}_p) - (\hat{L}_{r(p)} - \hat{L}_p)$$

shows that \hat{L}_p belongs to the span of two idempotents and the conclusion follows. □

If $\mathfrak{G} = (\mathfrak{G}^0, \mathfrak{G}^1, r, s)$ is a topological graph and $S \subseteq \mathfrak{G}^1$, then $N(S)$ denotes the collection of continuous functions $F \in X_{\mathfrak{G}}$ with $F|_S = 0$, i.e., vanishing at S .

Lemma 4.3. *Let $\mathfrak{G} = (\mathfrak{G}^0, \mathfrak{G}^1, r, s)$ be a topological graph.*

(i) *If $S_1, S_2 \subseteq \mathfrak{G}^1$ are closed, then*

$$N(S_1 \cap S_2) = \overline{N(S_1) + N(S_2)}.$$

(ii) *If $S_1 \subseteq \mathfrak{G}^0, S_2 \subseteq \mathfrak{G}^1$ are closed, then*

$$N(r^{-1}(S_1) \cup S_2) = \overline{[(f \circ r)F \mid f|_{S_1} = 0, F|_{S_2} = 0]}.$$

(iii) *If $S_1 \subseteq \mathfrak{G}^0, S_2 \subseteq \mathfrak{G}^1$ are closed, then*

$$N(s^{-1}(S_1) \cup S_2) = \overline{[(f \circ s)F \mid f|_{S_1} = 0, F|_{S_2} = 0]}.$$

Proof. Let $K \subseteq \mathfrak{G}^1$ be compact. We claim that for any sequence $\{F_n\}_n$ with $\text{supp } F_n \subseteq K, n \in \mathbb{N}$, convergence in $X_{\mathfrak{G}}$ is equivalent to convergence in the usual supremum norm $\| \cdot \|_{\infty}$ of $C_0(\mathfrak{G}^1)$.

Indeed, this will follow if we show that there exists $n \in \mathbb{N}$ so that $\|F\| \leq n\|F\|_{\infty}$, for any F with $\text{supp } F \subseteq K$. Let U_1, U_2, \dots, U_n be an open cover of K so that $s|_{U_i}$ is injective for all $i = 1, 2, \dots, n$. Notice that for any function G , with $\text{supp } G \subseteq U_i$ for some i , we have $\|G\| = \|G\|_{\infty}$. Hence, if $\{H_i\}_{i=1}^n$ is a partition of unity for K subordinated by the cover U_1, U_2, \dots, U_n , then,

$$\|F\| \leq \sum_{i=1}^n \|H_i F\| = \sum_{i=1}^n \|H_i F\|_{\infty} \leq n\|F\|_{\infty}$$

and the conclusion follows.

(i) Let $F \in N(S_1 \cap S_2)$. By [26, Lemma 1.6] and its proof, F can be approximated by functions of the form HF , where $H \in C_c(\mathfrak{G}^1)$. Clearly such functions belong to $N(S_1 \cap S_2)$ and so without loss of generality we may assume that $\text{supp } F = K$ is compact. The proof now follows familiar lines as we only need to approximate F by elements of $N(S_1) + N(S_2)$, in the usual supremum norm, while staying inside K .

Let $\epsilon > 0$ and let $K_{\epsilon} = \{e \in K \mid |F(e)| \geq \epsilon\}$. Since K_{ϵ} is disjoint from $S_1 \cap S_2$, we can cover K_{ϵ} with finitely many U_1, U_2, \dots, U_m so that each of the U_i is disjoint from one of the S_1, S_2 . Let $H_i, i = 1, 2, \dots, m$, be a partition of unity for K_{ϵ} which is subordinate to U_1, U_2, \dots, U_m . Then $\sum_{i=1}^m H_i F \in N(S_1) + N(S_2)$ and is ϵ -close to F .

(ii) Let $F \in N(r^{-1}(S_1) \cup S_2)$. Once again, by [26, Lemma 1.6] there is no loss of generality assuming that F has compact support. If $\{f_i\}_{i \in \mathbb{I}}$ is an increasing approximate unit for $C_0(\mathfrak{G}^0 \setminus S_1) \subseteq C_0(\mathfrak{G}^0)$, then Dini's Theorem implies that the family $\{(f_i \circ r)F\}_{i \in \mathbb{I}}$ approximates F (in both norms) and proves the lemma.

(iii) The proof is similar to that of (ii). □

Lemma 4.3 is now being used in the following with $\mathcal{G}^i, i \in \mathbb{N}$, in the place of \mathfrak{G} .

Lemma 4.4. *If $\varphi_{v,n}$ is as above, with \mathcal{G}_v acyclic, then*

$$\ker \varphi_{v,n} = \overline{[(\ker \varphi_{v,n})^2]}.$$

Proof. From the Fourier series expansion of (3) and the results of Katsura discussed in the beginning of this section, it follows that each element of $\mathcal{T}_{\mathcal{G}}^+$ admits a Fourier series expansion of the form

$$(8) \quad \pi_{\infty}(f) + \sum_{i=1}^{\infty} t_{\infty}^i(F_i),$$

with $f \in C_0(\mathcal{G}^0)$ and $F_i \in X_{\mathcal{G}^i}, i = 1, 2, \dots$. It is easy to see now that $\ker \varphi_{v,n}$ consists precisely of all elements of $\mathcal{T}_{\mathcal{G}}^+$, whose Fourier series (8) satisfies

$$f \in N(\{p \in \mathcal{F}_{v,n} \mid |p| = 0\})$$

and

$$F_i \in N(\{p \in \mathcal{F}_{v,n} \mid |p| = i\}), \quad i = 1, 2, \dots$$

Note that in the case where $i > n$, the set $\{p \in \mathcal{F}_{v,n} \mid |p| = i\}$ is empty and so the above simply says that F_i ranges over $X_{\mathcal{G}^i}$.

To prove the lemma, fix an i and notice that any finite sum of the following form belongs to $[(\ker \varphi_{v,n})^2]$:

$$(9) \quad \sum_m \left(\pi_{\infty}(f_m)t_{\infty}^i(F_m) + t_{\infty}^i(F'_m)\pi_{\infty}(f'_m) \right) \\ = t_{\infty}^i \left(\sum_m (f_m \circ r)F_m + \sum_m (f'_m \circ s)F'_m \right),$$

where $f_m, f'_m \in N(\bigcup_{i=0}^n \mathcal{G}_{v,i}^0)$ and $F_m, F'_m \in N(\{p \in \mathcal{F}_{v,n} \mid |p| = i\})$, for all m . However, Lemma 4.3 shows that sums of the form $\sum_m (f_m \circ r)F_m$ are dense in

$$(10) \quad N \left(r^{-1} \left(\bigcup_{j=0}^n \mathcal{G}_{v,j}^0 \right) \cup \{p \in \mathcal{F}_{v,n} \mid |p| = i\} \right),$$

and similarly, sums of the form $\sum_m (f'_m \circ s)F'_m$ are dense in

$$(11) \quad N \left(\{p \in \mathcal{F}_{v,n} \mid |p| = i\} \cup s^{-1} \left(\bigcup_{j=0}^n \mathcal{G}_{v,j}^0 \right) \right),$$

where r, s denotes the range and source functions on paths of length i . Since \mathcal{G}_v is acyclic

$$r^{-1} \left(\bigcup_{j=0}^n \mathcal{G}_{v,j}^0 \right) \cap s^{-1} \left(\bigcup_{j=0}^n \mathcal{G}_{v,j}^0 \right) = \{p \in \mathcal{F}_{v,n} \mid |p| = i\}$$

and so Lemma 4.3(i) shows that the closed linear space generated by (10) and (11) equals $\{p \in \mathcal{F}_{v,n} \mid |p| = i\}$. This shows that the sums as in (9) can approximate the i -th Fourier coefficient of any element in $\ker \varphi_{v,n}$. □

Another possibility that we will be considering here is that the graph \mathcal{G}_v may be *transitive*, i.e., given any two vertices $v', v'' \in \mathcal{G}_v^0$ there exists a path starting at v' and ending at v'' . In that case, Davidson and the author have shown [11, Theorem 4.4] that $\mathcal{L}_{\mathcal{G}_v}$ admits a separating family of finite dimensional irreducible representations. Composing these representations with the representation φ_v of Theorem 3.2, we obtain a separating family $\varphi_{v,n}$, $n \in \mathbb{N}$, of finite dimensional irreducible representations of $\mathcal{T}_{\mathcal{G}^+} / \ker \varphi_v$.

Theorem 4.5. *Let $\mathcal{G} = (\mathcal{G}^0, \mathcal{G}^1, r, s)$ be a source finite topological graph and let $\{\mathcal{G}_v\}_{v \in \mathcal{G}^0}$ be the family of discrete graphs associated with \mathcal{G} . Assume that the set of all points $v \in \mathcal{G}^0$, for which \mathcal{G}_v is either acyclic or transitive, is dense in \mathcal{G}^0 . Then any approximately local derivation on $\mathcal{T}_{\mathcal{G}^+}$ is a derivation.*

Proof. Let δ be an approximately local derivation. Consider the representations $\varphi_{v,n}$, discussed above, where v ranges over all points for which \mathcal{G}_v is either acyclic or transitive and $n \in \mathbb{N}$. Because of the density assumption this family separates points.

Claim. For any approximately local derivation $\hat{\delta} : \mathcal{T}_{\mathcal{G}^+} \rightarrow \mathcal{T}_{\mathcal{G}^+}$, we have

$$\hat{\delta}(\ker \varphi_{v,n}) \subseteq \ker \varphi_{v,n}.$$

Indeed let $A \in \ker \rho_{x,n}$ and let δ_A be a derivation of $T_{\mathcal{G}^+}$ so that $\hat{\delta}(A) = \delta_A(A)$. We distinguish two cases.

If \mathcal{G}_v is transitive, then we have noted in the discussion preceding this theorem that $\varphi_{v,n}$ is a finite dimensional irreducible representation and so $\ker \varphi_{v,n}$ is a primitive ideal. Since derivations of Banach algebras leave primitive ideals invariant [33, Proposition 6.4.16], $\hat{\delta}(A) = \delta_A(A) \in \ker \varphi_{v,n}$.

On the other hand, if \mathcal{G}_v is acyclic, then Lemma 4.4 shows that $(\ker \varphi_{v,n})^2 = \ker \varphi_{v,n}$. Hence any derivation of $\mathcal{T}_{\mathcal{G}^+}$ leaves $\ker \rho_{x,n}$ invariant. In particular, this applies to δ_A and so $\hat{\delta}(A) = \delta_A(A) \in \ker \varphi_{v,n}$ in this case as well.

The rest of the proof follows now familiar lines. By the Claim above, δ preserves $\ker \varphi_{v,n}$ and so we obtain a map

$$\delta_{x,n} : \mathcal{T}_{\mathcal{G}^+} / \ker \varphi_{v,n} \longrightarrow \varphi_{v,n}(\mathcal{T}_{\mathcal{G}^+}); A + \ker \varphi_{v,n} \mapsto \varphi_{v,n}(\delta(A)).$$

It is easy to see, again from the Claim above, that $\delta_{x,n}$ is an approximate local derivation. Furthermore, by Lemma 4.2, $\mathcal{T}_{\mathcal{G}^+} / \ker \varphi_{v,n}$ is generated by its idempotents and so Theorem 2.3 implies that $\delta_{x,n}$ is actually a derivation. Hence

$$\varphi_{v,n}(\delta(AB) - \delta(A)B - A\delta(B)) = 0.$$

Since $\bigcap_{v,n} \ker \varphi_{v,n} = \{0\}$, the conclusion follows. □

The first significant application of the above result is to Peters' semicrossed products, where nothing was previously known regarding local derivations.

Definition 4.6. Let (Ω, σ) be a dynamical system on a locally compact Hausdorff space. A point $x \in \Omega$ is said to be *periodic* if there exists $i \in \mathbb{N}$ so that $\sigma^{(i)}(x) = x$. The point $x \in \Omega$ is said to be *eventually periodic* if x is not periodic but one of its iterations $\sigma^{(i)}(x)$, $i \in \mathbb{N}$, is periodic. Otherwise x is said to be *aperiodic*.

Let (Ω, σ) be a dynamical system on a locally compact Hausdorff space. The semicrossed product $C(\Omega) \times_{\sigma} \mathbb{Z}^+$ is the tensor algebra corresponding to the topological graph $\mathcal{G} = (\mathcal{G}^0, \mathcal{G}^1, r, s)$ with $\mathcal{G}^0 = \mathcal{G}^1 = \Omega$, $s = \text{id}$ and $r = \sigma$. If $v \in \Omega$, then

$$\mathcal{G}_v^0 = \mathcal{G}_v^1 = \mathcal{O}_v \equiv \{v, \sigma(v), \sigma^{(2)}(v), \dots\},$$

i.e., the orbit of v . If v is periodic with period n , then the corresponding graph \mathcal{G}_v is the n -cycle graph \mathcal{C}_n that has been studied quite extensively in the graph algebra theory; this is a transitive graph. On the other hand, if v is aperiodic, then \mathcal{G}_v is an infinite countable graph with exactly one edge starting from each $\sigma^{(n)}(v)$ (and ending at $\sigma^{(n+1)}(v)$), $n = 0, 1, 2, \dots$. This graph is acyclic.

There is also a third possibility that the point $v \in \Omega$ is eventually periodic for σ . In that case the graph \mathcal{G}_v is a combination of the above graphs. If $k \in \mathbb{N}$ is the least positive integer so that $\sigma^{(k)}(v)$ is periodic, then at $\sigma^{(k)}(v)$ the graph \mathcal{G}_v supports an n -cycle graph (n is the period of $\sigma^{(k)}(v)$) and receives a tail starting at v . We do not know how to deal with that case as it is not covered by Theorem 4.5 and so in the next result we assume that such points are topologically insignificant.

Corollary 4.7. *Let (Ω, σ) be a dynamical system on a locally compact Hausdorff space Ω . Assume that the eventually periodic points of σ have empty interior, e.g., σ is a homeomorphism. Then any approximately local derivation on $C(\Omega) \times_{\sigma} \mathbb{Z}^+$ is a derivation. Hence $\mathcal{Z}^1(C(\Omega) \times_{\sigma} \mathbb{Z}^+)$ is reflexive.*

Let Ω be a compact Hausdorff space and let $\sigma : \Omega \rightarrow \Omega$ be a covering map. Consider the topological graph $\mathcal{G}_{\text{EV}}(\Omega, \sigma) = (\mathcal{G}^0, \mathcal{G}^1, r, s)$, where $\mathcal{G}^0 = \mathcal{G}^1 = \Omega$, $r = \text{id}$ and $s = \sigma$. Assume further that the system (Ω, σ) is *topologically free*, i.e., the sets $\Omega_{m,n} \equiv \{v \in \Omega \mid \sigma^{(m)}(v) = \sigma^{(n)}(v)\}$ have empty interior for all $m \neq n$. It is known [5, Theorem 6.1] that the Cuntz-Pimsner C^* -algebra associated with $\mathcal{G}_{\text{EV}}(\Omega, \sigma)$ is isomorphic to the Exel crossed product C^* -algebra $C(\Omega) \rtimes_{\sigma, L} \mathbb{N}$, where L is the transfer operator that averages over inverse images of points. The condition of topological freeness was isolated by Exel and Vershik in their original work [14] and was further studied in [4–6]. It is a consequence of these works that the tensor algebra $\mathcal{T}_{\text{EV}}^+(\Omega, \sigma)$ for $\mathcal{G}_{\text{EV}}(\Omega, \sigma)$ admits the following elegant description.

Example 4.8. A faithful representation for $\mathcal{T}_{\text{EV}}^+(\Omega, \sigma)$, provided that (Ω, σ) is topologically free.

Let \mathcal{H} be a Hilbert space with orthonormal base $\{e_v\}_{v \in \Omega}$. For $f \in C(\Omega)$ let M_f be the multiplication operator $M_f(e_v) = f(v)e_v$, $v \in \Omega$, and

$$S(e_v) = (L(1_{\Omega})(v))^{-1/2} \sum_{w \in \sigma^{-1}(\{v\})} e_w.$$

In [6, Theorem 6] it is shown that the C^* -algebra generated by M_f , $f \in C(\Omega)$, and S is canonically isomorphic to $C(\Omega) \rtimes_{\sigma, L} \mathbb{N}$, i.e., with generators going to generators. Furthermore, it is a consequence of [5, Proposition 3.1] and [5, Theorem 6.1] that $C(\Omega) \rtimes_{\sigma, L} \mathbb{N}$ is once again canonically isomorphic to $\mathcal{O}_{\text{EV}}(\Omega, \sigma)$. This implies that $\mathcal{T}_{\text{EV}}^+(\Omega, \sigma)$ is isomorphic to the non-selfadjoint algebra generated by M_f , $f \in C(\Omega)$, and S .

Corollary 4.9. *Let Ω be a compact Hausdorff space and let $\sigma : \Omega \rightarrow \Omega$ be a topologically free covering map. If $\mathcal{G}_{\text{EV}}(\Omega, \sigma) = (\mathcal{G}^0, \mathcal{G}^1, r, s)$, where $\mathcal{G}^0 = \mathcal{G}^1 = \Omega$, $r = \text{id}$ and $s = \sigma$, then any approximately local derivation on $\mathcal{T}_{\text{EV}}^+(\Omega, \sigma)$ is a derivation.*

Proof. It is not difficult to see that for a point $v \in \Omega$, the corresponding discrete graph \mathcal{G}_v is acyclic iff $v \notin \bigcup_{m \neq n} \Omega_{m,n}$. By the Baire Category Theorem, $\bigcup_{m \neq n} \Omega_{m,n}$ has empty interior and the conclusion follows from Theorem 4.5 \square

Another application of Theorem 4.5 takes place for certain tensor algebras that materialize as subalgebras of Exel’s crossed products by partial automorphisms. Let Ω be a locally compact space and σ a partial homeomorphism, i.e., $\sigma : U \rightarrow V$ is a homeomorphism between two open subsets U, V of Ω . Let $\theta : C_0(U) \rightarrow C_0(V)$ be the $*$ -automorphism induced by σ . The triple $(\theta, C_0(U), C_0(V))$ is called a partial automorphism of $C_0(\Omega)$. In [13, Definition 3.7] Exel associates a C^* -algebra $C_0(\Omega) \times_{\theta} \mathbb{Z}$ with the partial automorphism $(\theta, C_0(U), C_0(V))$. This algebra is defined through a universal property but Katsura has shown that it corresponds to the Cuntz-Pimsner algebra corresponding to the graph $\mathcal{G} = (\mathcal{G}^0, \mathcal{G}^1, r, s)$, where $\mathcal{G}_0 = \Omega, \mathcal{G}_1 = U, r = \sigma$ and s is the natural embedding.

Corollary 4.10. *Let $C_0(\Omega) \times_{\theta} \mathbb{Z}$ be Exel’s crossed product by the partial automorphism $(\theta, C_0(U), C_0(V))$ of Ω and let $C_0(\Omega) \times_{\theta} \mathbb{Z}^+$ be the associated tensor algebra. Then any approximately local derivation on $C_0(\Omega) \times_{\theta} \mathbb{Z}^+$ is a derivation.*

We now focus on tensor algebras for multivariable dynamical systems. If (Ω, σ) is such a system, with $\sigma = (\sigma_1, \sigma_2, \dots, \sigma_n)$, and $u = i_k i_{k-1} \dots i_1 \in \mathbb{F}_n^+$, then we write $\sigma_u \equiv \sigma_{i_k} \circ \sigma_{i_{k-1}} \circ \dots \circ \sigma_{i_1}$.

Corollary 4.11. *Let (Ω, σ) be a multivariable system on a locally compact space Ω and assume that the set*

$$\Omega_0 = \left\{ v \in \Omega \mid \sigma_u(v) = \sigma_w(v), \text{ for some } u, w \in \mathbb{F}_n^+ \text{ with } |u| \neq |w| \right\}$$

has empty interior. Then any approximately local derivation on $\mathcal{T}^+(\Omega, \sigma)$ is a derivation.

Proof. It is easy to verify that \mathcal{G}_v is acyclic provided that $v \in \Omega \setminus \Omega_0$. \square

The assumptions of the above corollary hold in particular when $\Omega = \mathbb{T}$, and $\sigma_k(z) = e^{2\pi i \theta_k} z, k = 1, 2$, with θ_1, θ_2 irrational satisfying $\theta_1/\theta_2 \notin \mathbb{Q}$.

Corollary 4.12. *Let \mathfrak{G} be a finite group acting on a locally compact Hausdorff space Ω via a family of automorphisms $\sigma_{\mathfrak{G}} \equiv \{\sigma_g\}_{g \in \mathfrak{G}}$. Then any approximately local derivation on $\mathcal{T}^+(\Omega, \sigma_{\mathfrak{G}})$ is a derivation.*

Proof. We claim that for any $v \in \Omega$, the graph \mathcal{G}_v of Theorem 3.2 is transitive, i.e., given any two vertices $v', v'' \in \mathcal{G}_v^0$ there exists a path starting at v' and ending at v'' . Indeed by construction for any $v' \in \mathcal{G}_v^0$ there is a path of the form

$$(g_i, \sigma_{g_i^{-1}}(v')), \dots, (g_2, \sigma_{g_1}(v))(g_1, v), \quad g_k \in \mathfrak{G}, k = 1, 2, \dots, i,$$

starting at v and ending at v' . Since \mathfrak{G} is a group, the opposite path

$$(g_1^{-1}, g_1(v)) \dots (g_{i-1}^{-1}, g_i^{-1}(v'))(g_i^{-1}, v')$$

exists thus establishing the existence of paths between v and any v' in both directions. Thus \mathcal{G}_v is transitive. The result follows now from Theorem 4.5. \square

We observed in Proposition 3.6 that the semicrossed product $\mathcal{A} \times_{\sigma} \mathbb{F}_n^+$ for a multivariable C^* -dynamical system (\mathcal{A}, σ) admits a separating family $\hat{\pi}_i, i \in \mathbb{I}$ of finite dimensional irreducible representations. Since the kernels of these representations

are primitive ideals of $\mathcal{A} \times_{\sigma} \mathbb{F}_n^+$, any approximately local derivation leaves them invariant. Therefore by following the same steps as in the proof of Corollary 4.7 we obtain

Proposition 4.13. *Let (\mathcal{A}, σ) be an automorphic multivariable C^* -dynamical system and assume that the crossed product C^* -algebra $\mathcal{A} \times_{\sigma} \mathbb{F}_n$ is residually finite dimensional (RFD). Then any approximately local derivation on $\mathcal{A} \times_{\sigma} \mathbb{F}_n^+$ is a derivation.*

In particular, Proposition 4.13 applies to the universal algebra $A(\mathbb{F}_n^+) \subseteq C^*(\mathbb{F}_n)$ generated by n contractions (see the proof of Corollary 3.7). Here is another application coming from [9].

Corollary 4.14. *Let $\sigma = (\sigma_1, \sigma_2)$ be a multivariable system on the two-point space $\Omega = \{0, 1\}$, with $\sigma_1 = \text{id}$ and $\sigma_2(i) = i + 1 \pmod 2, i = 0, 1$. Then any approximately local derivation on $C(\Omega) \times_{\sigma} \mathbb{F}_2^+$ is a derivation.*

Proof. In [9, Example 3.24], we observed that $C(\Omega) \times_{\sigma} \mathbb{F}_2 \simeq M_2(C^*(\mathbb{F}_3))$, which is RFD, and so the conclusion follows from Proposition 4.13. □

5. CONCLUDING REMARKS AND OPEN PROBLEMS

Theorem 4.5 does not extend to local derivations with range on a bimodule rather than the tensor algebra itself. The counterexample is actually on a semicrossed product bimodule.

Example 5.1. Let (Ω, σ) be a dynamical system on a locally compact Hausdorff space and assume that σ has a fixed point $x \in \Omega$. Then there exists a finite dimensional $C_0(\Omega) \times_{\sigma} \mathbb{Z}^+$ -module \mathfrak{B} and a local derivation $\delta : C_0(\Omega) \times_{\sigma} \mathbb{Z}^+ \rightarrow \mathfrak{B}$ which is not a derivation.

Indeed, let \mathfrak{B} and $\delta_{\mathfrak{B}}$ be as in Example 2.5. Since $x \in \Omega$ is a fixed point for σ the mapping

$$\pi : C_0(\Omega) \times_{\sigma} \mathbb{Z}^+ \longrightarrow \mathfrak{B}; \sum_{n=0}^{\infty} V^n f_n \longmapsto \begin{pmatrix} f_0(x) & f_1(x) & f_2(x) \\ 0 & f_0(x) & f_1(x) \\ 0 & 0 & f_0(x) \end{pmatrix}$$

is a representation that makes \mathfrak{B} a right $C_0(\Omega) \times_{\sigma} \mathbb{Z}^+$ -module with the natural action coming from π . If

$$\delta : C_0(\Omega) \times_{\sigma} \mathbb{Z}^+ \longrightarrow \mathfrak{B}; A \longmapsto \delta_{\mathfrak{B}}(\pi(A)),$$

then δ is a local derivation which is not a derivation (otherwise $\delta_{\mathfrak{B}}$ would be a derivation as well).

A modification of the above example also works for the non-commutative disc algebra \mathcal{A}_n , i.e., the tensor algebra of the graph with one vertex and n -loops, thus showing the failure of Theorem 4.5 for local derivations on \mathcal{A}_n -modules as well. It turns out that using Example 2.6, we can construct yet another \mathcal{A}_n -module to demonstrate that failure.

Example 5.2. Let $\mathcal{A}_n, n \geq 2$, be the non-commutative disc algebra with generators A_1, A_2, \dots, A_n and let $\mathfrak{C}, \delta_{\mathfrak{C}}$ be as in Example 2.6. From the dilation theory for row contractions, it follows that there exists a representation $\pi : \mathcal{A}_n \rightarrow M_3(\mathbb{C})$ so that

$$\pi(A_1) = E_{12}, \pi(A_2) = E_{23} \text{ and } \pi(A_i) = 0, \text{ for all other } i.$$

If

$$\delta : \mathcal{A}_n \longrightarrow \mathfrak{C}; A \longmapsto \delta_{\mathfrak{C}}(\pi(A)),$$

then δ is a local derivation from \mathcal{A}_n onto the \mathcal{A}_n -module $\mathfrak{C} = \pi(\mathcal{A}_n)$ which is not a derivation.

The above example shows that \mathcal{A}_2 admits local derivations on modules of dimension 3 and 4, which are not derivations. Does this persist on higher dimensions? One is tempted to guess that the answer is affirmative but we do not have a systematic way of producing such examples. What about semicrossed product modules? Can one construct such counterexamples as Example 5.1 with modules of dimension higher than 3? Also we wonder whether any counterexamples do exist in the case where σ is free.

It goes without saying that the following two open problems are tantalizing.

Problem 1. Prove or disprove that a local left multiplier on the tensor algebra of a C^* -correspondence is actually a left multiplier.

Problem 2. Prove or disprove that an approximately local derivation on an arbitrary semicrossed product is actually a derivation.

ACKNOWLEDGEMENTS

A good part of this paper materialized while the author was visiting the Chinese Academy of Sciences in May 2014. The author would like to thank his hosts, Professors Liming Ge and Wei Yuan, for the stimulating conversations and their hospitality during his stay.

REFERENCES

- [1] William B. Arveson, *Operator algebras and measure preserving automorphisms*, Acta Math. **118** (1967), 95–109. MR0210866 (35 #1751)
- [2] Matej Brešar, *Characterizing homomorphisms, derivations and multipliers in rings with idempotents*, Proc. Roy. Soc. Edinburgh Sect. A **137** (2007), no. 1, 9–21, DOI 10.1017/S0308210504001088. MR2359769 (2008j:16097)
- [3] Matej Brešar and Peter Šemrl, *Mappings which preserve idempotents, local automorphisms, and local derivations*, Canad. J. Math. **45** (1993), no. 3, 483–496, DOI 10.4153/CJM-1993-025-4. MR1222512 (94k:47054)
- [4] Nathan Brownlowe and Iain Raeburn, *Exel's crossed product and relative Cuntz-Pimsner algebras*, Math. Proc. Cambridge Philos. Soc. **141** (2006), no. 3, 497–508, DOI 10.1017/S030500410600956X. MR2281412 (2007i:46067)
- [5] Nathan Brownlowe, Iain Raeburn, and Sean T. Vittadello, *Exel's crossed product for non-unital C^* -algebras*, Math. Proc. Cambridge Philos. Soc. **149** (2010), no. 3, 423–444, DOI 10.1017/S030500411000037X. MR2726727 (2011k:46092)
- [6] Toke Meier Carlsen and Sergei Silvestrov, *On the Exel crossed product of topological covering maps*, Acta Appl. Math. **108** (2009), no. 3, 573–583, DOI 10.1007/s10440-008-9372-6. MR2563500 (2010k:46071)
- [7] Man Duen Choi, *The full C^* -algebra of the free group on two generators*, Pacific J. Math. **87** (1980), no. 1, 41–48. MR590864 (82b:46069)
- [8] Randall L. Crist, *Local derivations on operator algebras*, J. Funct. Anal. **135** (1996), no. 1, 76–92, DOI 10.1006/jfan.1996.0004. MR1367625 (96m:46128)
- [9] Kenneth R. Davidson and Elias G. Katsoulis, *Operator algebras for multivariable dynamics*, Mem. Amer. Math. Soc. **209** (2011), no. 982, viii+53, DOI 10.1090/S0065-9266-10-00615-0. MR2752983 (2012c:47194)
- [10] Kenneth R. Davidson and Elias G. Katsoulis, *Isomorphisms between topological conjugacy algebras*, J. Reine Angew. Math. **621** (2008), 29–51, DOI 10.1515/CRELLE.2008.057. MR2431249 (2010b:47226)

- [11] Kenneth R. Davidson and Elias Katsoulis, *Nest representations of directed graph algebras*, Proc. London Math. Soc. (3) **92** (2006), no. 3, 762–790, DOI 10.1017/S0024611505015662. MR2223543 (2009m:47202)
- [12] Benton L. Duncan, *Derivations for a class of matrix function algebras*, Linear Algebra Appl. **422** (2007), no. 1, 67–76, DOI 10.1016/j.laa.2006.09.006. MR2298996 (2008c:46069)
- [13] Ruy Exel, *Circle actions on C^* -algebras, partial automorphisms, and a generalized Pimsner-Voiculescu exact sequence*, J. Funct. Anal. **122** (1994), no. 2, 361–401, DOI 10.1006/jfan.1994.1073. MR1276163 (95g:46122)
- [14] R. Exel and A. Vershik, *C^* -algebras of irreversible dynamical systems*, Canad. J. Math. **58** (2006), no. 1, 39–63, DOI 10.4153/CJM-2006-003-x. MR2195591 (2007d:46058)
- [15] Neal J. Fowler and Iain Raeburn, *The Toeplitz algebra of a Hilbert bimodule*, Indiana Univ. Math. J. **48** (1999), no. 1, 155–181, DOI 10.1512/iumj.1999.48.1639. MR1722197 (2001b:46093)
- [16] B. E. Johnson, *Local derivations on C^* -algebras are derivations*, Trans. Amer. Math. Soc. **353** (2001), no. 1, 313–325 (electronic), DOI 10.1090/S0002-9947-00-02688-X. MR1783788 (2002c:46132)
- [17] B. E. Johnson, *Centralisers and operators reduced by maximal ideals*, J. London Math. Soc. **43** (1968), 231–233. MR0223890 (36 #6937)
- [18] The Hadwin Lunch Bunch, *Local multiplications on algebras spanned by idempotents*, Linear and Multilinear Algebra **37** (1994), no. 4, 259–263, DOI 10.1080/03081089408818328. MR1310968 (96f:47070)
- [19] Don Hadwin and Jeanne Wald Kerr, *Local multiplications on algebras*, J. Pure Appl. Algebra **115** (1997), no. 3, 231–239, DOI 10.1016/S0022-4049(96)00013-8. MR1431834 (98b:16006)
- [20] Don Hadwin and Jiankui Li, *Local derivations and local automorphisms on some algebras*, J. Operator Theory **60** (2008), no. 1, 29–44. MR2415555 (2009e:47056)
- [21] Don Hadwin and Jiankui Li, *Local derivations and local automorphisms*, J. Math. Anal. Appl. **290** (2004), no. 2, 702–714, DOI 10.1016/j.jmaa.2003.10.015. MR2033052 (2004j:47071)
- [22] Richard V. Kadison, *Local derivations*, J. Algebra **130** (1990), no. 2, 494–509, DOI 10.1016/0021-8693(90)90095-6. MR1051316 (91f:46092)
- [23] E. G. Katsoulis, *The reflexive closure of the adjointable operators*, Illinois J. Math. **58** (2014), no. 2, 359–367. MR3367652
- [24] Elias G. Katsoulis and David W. Kribs, *Tensor algebras of C^* -correspondences and their C^* -envelopes*, J. Funct. Anal. **234** (2006), no. 1, 226–233, DOI 10.1016/j.jfa.2005.12.013. MR2214146 (2007a:46061)
- [25] Elias Katsoulis and David W. Kribs, *Isomorphisms of algebras associated with directed graphs*, Math. Ann. **330** (2004), no. 4, 709–728, DOI 10.1007/s00208-004-0566-6. MR2102309 (2005i:47114)
- [26] Takeshi Katsura, *A class of C^* -algebras generalizing both graph algebras and homeomorphism C^* -algebras. I. Fundamental results*, Trans. Amer. Math. Soc. **356** (2004), no. 11, 4287–4322 (electronic), DOI 10.1090/S0002-9947-04-03636-0. MR2067120 (2005b:46119)
- [27] Takeshi Katsura, *On C^* -algebras associated with C^* -correspondences*, J. Funct. Anal. **217** (2004), no. 2, 366–401, DOI 10.1016/j.jfa.2004.03.010. MR2102572 (2005e:46099)
- [28] David W. Kribs and Stephen C. Power, *Free semigroupoid algebras*, J. Ramanujan Math. Soc. **19** (2004), no. 2, 117–159. MR2076898 (2005c:47106)
- [29] David R. Larson, *Reflexivity, algebraic reflexivity and linear interpolation*, Amer. J. Math. **110** (1988), no. 2, 283–299, DOI 10.2307/2374503. MR935008 (89d:47096)
- [30] David R. Larson and Ahmed R. Sourour, *Local derivations and local automorphisms of $\mathcal{B}(X)$* , Operator theory: operator algebras and applications, Part 2 (Durham, NH, 1988), Proc. Sympos. Pure Math., vol. 51, Amer. Math. Soc., Providence, RI, 1990, pp. 187–194, DOI 10.1090/pspum/051.2/1077437. MR1077437 (91k:47106)
- [31] Jiankui Li and Zhidong Pan, *Annihilator-preserving maps, multipliers, and derivations*, Linear Algebra Appl. **432** (2010), no. 1, 5–13, DOI 10.1016/j.laa.2009.06.002. MR2566454 (2010j:47055)
- [32] Paul S. Muhly and Baruch Solel, *Tensor algebras over C^* -correspondences: representations, dilations, and C^* -envelopes*, J. Funct. Anal. **158** (1998), no. 2, 389–457, DOI 10.1006/jfan.1998.3294. MR1648483 (99j:46066)

- [33] Theodore W. Palmer, *Banach algebras and the general theory of *-algebras. Vol. I*, Encyclopedia of Mathematics and its Applications, vol. 49, Cambridge University Press, Cambridge, 1994. Algebras and Banach algebras. MR1270014 (95c:46002)
- [34] Justin Peters, *Semicrossed products of C^* -algebras*, J. Funct. Anal. **59** (1984), no. 3, 498–534, DOI 10.1016/0022-1236(84)90063-6. MR769379 (86e:46063)
- [35] Gelu Popescu, *Non-commutative disc algebras and their representations*, Proc. Amer. Math. Soc. **124** (1996), no. 7, 2137–2148, DOI 10.1090/S0002-9939-96-03514-9. MR1343719 (96k:47077)
- [36] Ebrahim Samei, *Approximately local derivations*, J. London Math. Soc. (2) **71** (2005), no. 3, 759–778, DOI 10.1112/S0024610705006496. MR2132382 (2006a:46061)

DEPARTMENT OF MATHEMATICS, EAST CAROLINA UNIVERSITY, GREENVILLE, NORTH CAROLINA 27858

E-mail address: `katsoulise@ecu.edu`