

## CURVATURE DECAY ESTIMATES OF GRAPHICAL MEAN CURVATURE FLOW IN HIGHER CODIMENSIONS

KNUT SMOCZYK, MAO-PEI TSUI, AND MU-TAO WANG

ABSTRACT. We derive pointwise curvature estimates for graphical mean curvature flows in higher codimensions for a flat ambient space. To the best of our knowledge, these are the first such estimates without assuming smallness of first derivatives of the defining map. An immediate application is a convergence theorem of the mean curvature flow of the graph of an area decreasing map between flat Riemann surfaces.

### 1. INTRODUCTION

Let  $\Sigma_1$  and  $\Sigma_2$  be two compact Riemannian manifolds and  $M = \Sigma_1 \times \Sigma_2$  be the product manifold. We consider a smooth map  $f : \Sigma_1 \rightarrow \Sigma_2$  and denote the graph of  $f$  by  $\Sigma$ ;  $\Sigma$  is a submanifold of  $M$  by the embedding  $id \times f$ . We study the deformation of  $f$  by the mean curvature flow. The idea is to deform  $\Sigma$  along its mean curvature vector field in  $M$  with the hope that  $\Sigma$  will remain a graph. This is the negative gradient flow of the volume functional, and a stationary point is a “minimal map” introduced by Schoen in [S1].

To describe previous results, we recall the differential of  $f$ ,  $df$ , at each point of  $\Sigma_1$  is a linear map between the tangent spaces. The Riemannian structures enable us to define the adjoint of  $df$ . Let  $\{\lambda_i\}$  denote the eigenvalues of  $\sqrt{(df)^T df}$ , or the singular values of  $df$ , where  $(df)^T$  is the adjoint of  $df$ . Note that  $\lambda_i$  is always nonnegative. We say  $f$  is an *area decreasing map* if  $\lambda_i \lambda_j < 1$  for any  $i \neq j$  at each point. In particular,  $f$  is area decreasing if  $df$  has rank one everywhere.

In [TW] (see also [W4, W5, W6]), it was proved that the area decreasing condition is preserved along the mean curvature flow and that the following global existence and convergence theorem holds.

**Theorem** ([TW], 2004). *Let  $\Sigma_1$  and  $\Sigma_2$  be compact Riemannian manifolds of constant sectional curvatures  $k_1$  and  $k_2$  respectively. Suppose  $k_1 \geq |k_2|$ ,  $k_1 + k_2 \geq 0$  and  $\dim(\Sigma_1) \geq 2$ . If  $f$  is a smooth area decreasing map from  $\Sigma_1$  to  $\Sigma_2$ , the mean curvature flow of the graph of  $f$  remains the graph of an area decreasing map and exists for all time. Moreover, if  $k_1 + k_2 > 0$ , then it converges smoothly to the graph of a constant map.*

---

Received by the editors January 23, 2014 and, in revised form, November 28, 2014.

2010 *Mathematics Subject Classification*. Primary 53C44.

*Key words and phrases*. Mean curvature flow.

The first author was supported by the DFG (German Research Foundation).

The second author was partially supported by a Collaboration Grant for Mathematicians from the Simons Foundation, #239677.

The third author was partially supported by National Science Foundation grants DMS 1105483 and DMS 1405152.

This result has been generalized to allow more general curvature conditions [LL, SHS1]. For example, the convergence part can be established when  $k_1 + k_2 = 0$  and  $k_1 \geq |k_2| > 0$  in [LL]. An important ingredient of these proofs is to use the positivity of  $k_1$  to show that the gradient of  $f$  approaches zero as  $t \rightarrow \infty$ . In [SHS1] the convergence follows if the sectional curvatures  $\text{sec}_{\Sigma_1}, \text{sec}_{\Sigma_2}$  of  $\Sigma_1$  and  $\Sigma_2$  are not necessarily constant and satisfy

$$\text{sec}_{\Sigma_1} > -\sigma, \quad \text{Ric}_{\Sigma_1} \geq (n-1)\sigma \geq (n-1)\text{sec}_{\Sigma_2}$$

for some positive constant  $\sigma$ , where  $n = \dim(\Sigma_1)$ . In this case the positivity of  $\text{Ric}_{\Sigma_1}$  is important to get the convergence. However, in all cases mentioned above the convergence part in the case  $k_1 = k_2 = 0$  remains an open standing problem.

In general, the global existence and convergence of a mean curvature flow relies on the boundedness of the second fundamental form. In the above theorem, the boundedness of the second fundamental form is obtained by an indirect blow-up argument; see [W1, W2, W3, TW]. The idea of the proof of convergence is to use the positivity of  $k_1 + k_2$  (or  $k_1$ , resp.  $\text{Ric}_{\Sigma_1}$ ) to show that the gradient of  $f$  is approaching zero, which in turn gives the boundedness of the second fundamental form when the flow exists for sufficiently long time. In [SHS2] mean curvature estimates are shown in case of length decreasing maps ( $\lambda_i < 1$ ). Other curvature estimates for higher codimensional graphical mean curvature flows have been obtained under various conditions [CCH, CCY]. However, to the best of our knowledge, there is no direct pointwise curvature estimate for higher codimensional mean curvature flow without assuming smallness conditions on the Lipschitz norm of the first derivatives. In this paper, we prove pointwise estimates assuming only a weaker condition (area decreasing) on the gradient of the map. As a result, the convergence of the flow can be established in dimension two when  $k_1 = k_2 = 0$ .

**Theorem 1.** *Let  $(\Sigma_1, g_1)$  and  $(\Sigma_2, g_2)$  be complete flat Riemann surfaces,  $\Sigma_1$  being compact. Suppose  $\Sigma \subset (\Sigma_1 \times \Sigma_2, g_1 \times g_2)$  is the graph of an area decreasing map  $f : \Sigma_1 \rightarrow \Sigma_2$  and let  $\Sigma_t$  denote its mean curvature flow with initial surface  $\Sigma_0 = \Sigma$ . Then  $\Sigma_t$  remains the graph of an area decreasing map  $f_t$  along the mean curvature flow. The flow exists smoothly for all time, and  $\Sigma_t$  converges smoothly to a totally geodesic submanifold as  $t \rightarrow \infty$ . Moreover, we have the mean curvature decay estimate*

$$t|H|^2 \leq \frac{2}{\alpha}$$

where  $\alpha = \inf_{\Sigma_0} \frac{2(1-\lambda_1^2\lambda_2^2)}{(1+\lambda_1^2)(1+\lambda_2^2)} > 0$  and  $\lambda_1$  and  $\lambda_2$  are the singular values of  $df$ .

*Remark 1.1.* Let  $f : \Sigma_1 \rightarrow \Sigma_2$  be an arbitrary smooth map between flat Riemann surfaces  $(\Sigma_1, g_1)$ ,  $(\Sigma_2, g_2)$  and suppose  $\Sigma_1$  is compact. Then there exists a constant  $c > 0$  such that all singular values of  $f$  satisfy  $\lambda_i\lambda_j < c^2$ . The map  $f : (\Sigma_1, g_1) \rightarrow (\Sigma_2, c^{-2}g_2)$  becomes area decreasing, and we can apply Theorem 1 to this case since the new metric  $\tilde{g}_2 = c^{-2}g_2$  is still flat. Such a procedure only applies in the flat case and changes the Lipschitz norm of the map. Nevertheless, the geometry of the graph of  $f$  is different.

As in [SW, Sm3], consider the symplectic structure  $dx^1 \wedge dy^1 + dx^2 \wedge dy^2$  on  $(T^2, \{x^i\}_{i=1,2}) \times (T^2, \{y^j\}_{j=1,2})$  and suppose  $\Sigma$  is Lagrangian with respect to this symplectic structure. The following stronger decay estimate on the second fundamental can be obtained in this case.

**Theorem 2.** *Let  $f : T^2 \rightarrow T^2$  be an area decreasing map such that its graph  $\Sigma$  is a Lagrangian submanifold in  $T^2 \times T^2$  with respect to the above symplectic structure. Then the same conclusion as in Theorem 1 holds and*

$$t|A|^2 \leq C_\alpha,$$

where  $C_\alpha$  is a positive constant that only depends on  $\alpha$ .

We first revisit the curvature estimates in codimension one by Ecker and Huisken [EH1, EH2]. A direct generalization of their estimate only works in the higher codimensional case when the gradient of the defining function is small enough. However, we were able to reformulate their estimates in a different way that can be adapted to the higher codimensional case. It turns out in higher codimensions a more sophisticated approach has to be developed to accommodate the complexity of the normal bundle.

### 2. ECKER AND HUISKEN’S ESTIMATES IN CODIMENSION ONE

In this section, we slightly rewrite the estimate in [EH1] so it can be adapted to the higher codimensional situation in later sections. Consider the mean curvature flow of the graph of a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  and let  $v = \sqrt{1 + |Df|^2}$ . Recall that the evolution equations of  $v$  and  $|A|^2$  are

$$\begin{aligned} \left(\frac{d}{dt} - \Delta\right)v &= -|A|^2v - 2\frac{|\nabla v|^2}{v} \text{ and} \\ \left(\frac{d}{dt} - \Delta\right)|A|^2 &= -2|\nabla A|^2 + 2|A|^4. \end{aligned}$$

We obtain the evolution equation of  $\ln v^2$  as

$$(2.1) \quad \left(\frac{d}{dt} - \Delta\right)\ln(v^2) = -2|A|^2 - \frac{1}{2}|\nabla \ln(v^2)|^2.$$

Using  $|\nabla A|^2 \geq |\nabla |A||^2 = \frac{|A|^2}{4}|\nabla \ln |A|^2|^2$ , we have

$$\left(\frac{d}{dt} - \Delta\right)|A|^2 \leq -\frac{|A|^2}{2}|\nabla \ln |A|^2|^2 + 2|A|^4.$$

Taking  $\ln$  of  $|A|^2$ , we obtain

$$(2.2) \quad \left(\frac{d}{dt} - \Delta\right)\ln(|A|^2) \leq 2|A|^2 + \frac{1}{2}|\nabla \ln |A|^2|^2.$$

As in [EH1], equations (2.1) and (2.2) together imply a sup norm bound for  $|A|^2v^2$ .

The following differential inequality for  $\ln(\delta t|A|^2 + \epsilon)$ , which is similar to equation (2.2), gives a decay estimate of  $|A|^2$ .

**Lemma 2.1.** *Given any  $\epsilon > 0$  and  $\delta < 2\epsilon$ ,*

$$\left(\frac{d}{dt} - \Delta\right)\ln(\delta t|A|^2 + \epsilon) \leq 2|A|^2 + \frac{1}{2}|\nabla \ln(\delta t|A|^2 + \epsilon)|^2.$$

*Proof.* Using  $\Delta \ln(\delta t|A|^2 + \epsilon) = \frac{\delta t \Delta |A|^2}{\delta t|A|^2 + \epsilon} - \frac{\delta^2 t^2 |\nabla |A|^2|^2}{(\delta t|A|^2 + \epsilon)^2}$ , we compute the evolution equation of  $\ln(\delta t|A|^2 + \epsilon)$ :

$$\begin{aligned} & \left(\frac{d}{dt} - \Delta\right) \ln(\delta t|A|^2 + \epsilon) \\ &= \frac{1}{\delta t|A|^2 + \epsilon} \left(\frac{d}{dt} - \Delta\right) (\delta t|A|^2 + \epsilon) + |\nabla \ln(\delta t|A|^2 + \epsilon)|^2 \\ &\leq \frac{1}{\delta t|A|^2 + \epsilon} \left(\delta |A|^2 + \delta t(2|A|^4 - \frac{|\nabla |A|^2|^2}{2|A|^2})\right) + |\nabla \ln(\delta t|A|^2 + \epsilon)|^2 \\ &\leq \frac{\delta |A|^2 + 2[\delta t|A|^2 + \epsilon]|A|^2 - 2\epsilon|A|^2}{\delta t|A|^2 + \epsilon} - \frac{1}{2} \frac{\delta t |\nabla |A|^2|^2}{(\delta t|A|^2 + \epsilon)|A|^2} \\ &\quad + |\nabla \ln(\delta t|A|^2 + \epsilon)|^2 \\ &\leq 2|A|^2 + \frac{\delta |A|^2 - 2\epsilon|A|^2}{\delta t|A|^2 + \epsilon} - \frac{1}{2} \frac{\delta t |\nabla |A|^2|^2}{(\delta t|A|^2 + \epsilon)|A|^2} + |\nabla \ln(\delta t|A|^2 + \epsilon)|^2 \\ &\leq 2|A|^2 + \frac{1}{2} |\nabla \ln(\delta t|A|^2 + \epsilon)|^2. \end{aligned}$$

Here we use  $\delta - 2\epsilon < 0$  and

$$-\frac{1}{2} \frac{\delta t |\nabla |A|^2|^2}{(\delta t|A|^2 + \epsilon)|A|^2} + \frac{1}{2} |\nabla \ln(\delta t|A|^2 + \epsilon)|^2 \leq 0. \quad \square$$

**Theorem 3.**  $\sup_{\Sigma_t}(t|A|^2) \leq v_0^2$  where  $v_0 = \sup_{\Sigma_0} v > 0$ .

*Proof.* From the evolution equation of  $v$ , we have  $\sup_{\Sigma_t} v^2 \leq v_0^2$ . Choosing  $\epsilon = 1$  and  $\delta = 1$  in the previous lemma and combining with equation (2.1), we derive

$$\begin{aligned} \left(\frac{d}{dt} - \Delta\right) \ln((t|A|^2 + 1)v^2) &\leq \frac{1}{2} |\nabla \ln(t|A|^2 + 1)|^2 - \frac{1}{2} |\nabla \ln(v^2)|^2 \\ &\leq \frac{1}{2} \nabla \ln\left(\frac{t|A|^2 + 1}{v^2}\right) \cdot \nabla \ln((t|A|^2 + 1)v^2). \end{aligned}$$

The maximum principle implies

$$\sup_{\Sigma_t} ((t|A|^2 + 1)v^2) \leq \sup_{\Sigma_0} v^2.$$

Therefore  $t|A|^2 \leq (t|A|^2 + 1)v^2 \leq v_0^2$ . □

### 3. ESTIMATES IN HIGHER CODIMENSIONS

Our basic set-up here is a mean curvature flow  $F : \Sigma \times [0, T] \rightarrow M$  of an  $n$ -dimensional submanifold  $\Sigma$  inside an  $(n+m)$ -dimensional flat Riemannian manifold  $M$ . Given any tensor on  $M$ , we may consider the pull-back tensor by  $F_t$  and consider the evolution equation with respect to the time-dependent induced metric on  $F_t(\Sigma) = \Sigma_t$ . For the purpose of applying the maximum principle, it suffices to derive the equation at a space-time point. We write all geometric quantities in terms of orthonormal frames, keeping in mind all quantities are defined independently of choices of frames. At any point  $p \in \Sigma_t$ , we choose any orthonormal frames  $\{e_i\}_{i=1, \dots, n}$  for  $T_p \Sigma_t$  and  $\{e_\alpha\}_{\alpha=n+1, \dots, n+m}$  for the normal space  $N_p \Sigma_t$ . The second fundamental form  $h_{\alpha ij}$  is denoted by  $h_{\alpha ij} = \langle \nabla_{e_i}^M e_j, e_\alpha \rangle$ , and the mean curvature vector is denoted by  $H_\alpha = \sum_i h_{\alpha ii}$ . For any  $j, k$ , we pretend

$$h_{n+i, jk} = 0$$

if  $i > m$ . Also we denote  $|A|^2 = \sum_{\alpha, i, j} h_{\alpha ij}^2$  and  $|H|^2 = \sum_\alpha H_\alpha^2$ .

First, we recall the evolution equations for  $|H|^2$  and  $|A|^2$ . The following proposition is taken from Corollary 3.8 and Corollary 3.9 from the survey paper [Sm2].

**Proposition 3.1.** *Suppose  $M$  is a flat Riemannian manifold. For a mean curvature flow  $F : \Sigma \times [0, T) \rightarrow M$  of any dimension, the quantities  $|A|^2$  and  $|H|^2$  satisfy the following equations along the mean curvature flow:*

$$(3.1) \quad \begin{aligned} & \frac{d}{dt}|A|^2 = \Delta|A|^2 - 2|\nabla^\perp A|^2 \\ & + 2 \sum_{\alpha, \gamma, i, m} \left( \sum_k h_{\alpha ik} h_{\gamma mk} - h_{\alpha mk} h_{\gamma ik} \right)^2 + 2 \sum_{i, j, m, k} \left( \sum_\alpha h_{\alpha ij} h_{\alpha mk} \right)^2 \end{aligned}$$

and

$$(3.2) \quad \frac{d}{dt}|H|^2 = \Delta|H|^2 - 2|\nabla^\perp H|^2 + 2 \sum_{i, k} \left( \sum_\alpha H_\alpha h_{\alpha ik} \right)^2.$$

Using Theorem 1 from [AMJ], we have

$$2 \sum_{\alpha, \gamma, i, m} \left( \sum_k h_{\alpha ik} h_{\gamma mk} - h_{\alpha mk} h_{\gamma ik} \right)^2 + 2 \sum_{i, j, m, k} \left( \sum_\alpha h_{\alpha ij} h_{\alpha mk} \right)^2 \leq 3|A|^4.$$

(This improves the prior bound of  $4|A|^4$  used in [W1].) Using

$$2 \sum_{i, k} \left( \sum_\alpha H_\alpha h_{\alpha ik} \right)^2 \leq 2|A|^2|H|^2,$$

we obtain the next lemma.

**Lemma 3.1.** *Suppose  $M$  is a flat Riemannian manifold. For a mean curvature flow  $F : \Sigma \times [0, T) \rightarrow M$  of any dimension, we have the following differential inequalities for  $|A|^2$  and  $|H|^2$ :*

$$(3.3) \quad \begin{aligned} & \left( \frac{d}{dt} - \Delta \right) |A|^2 \leq -2|\nabla^\perp A|^2 + 3|A|^4, \\ & \left( \frac{d}{dt} - \Delta \right) |H|^2 \leq -2|\nabla^\perp H|^2 + 2|A|^2|H|^2. \end{aligned}$$

We note that the term  $3|A|^4$  in the first equation is different from the term  $2|A|^4$  in the codimension one case. It cannot be improved unless the normal bundle is flat. This creates a major difficulty in attempting to generalize the codimension one estimate to the higher codimension case.

In the following, we derive differential inequalities for various geometric quantities which will be used for curvature decay estimates in Section 4.

**Lemma 3.2.** *Suppose  $M$  is a flat Riemannian manifold. For a mean curvature flow  $F : \Sigma \times [0, T) \rightarrow M$  of an  $n$ -dimensional submanifold  $\Sigma$ . Given any  $\epsilon > 0$  and  $0 < \delta \leq \frac{2\epsilon}{n}$ , we have the following differential inequalities:*

$$(3.4) \quad \begin{aligned} & \left( \frac{d}{dt} - \Delta \right) \ln(\delta t |H|^2 + \epsilon) \leq 2|A|^2 + \frac{|\nabla \ln(\delta t |H|^2 + \epsilon)|^2}{2}, \\ & \left( \frac{d}{dt} - \Delta \right) \ln(\delta t |A|^2 + \epsilon) \leq 3|A|^2 + \frac{|\nabla \ln(\delta t |A|^2 + \epsilon)|^2}{2}. \end{aligned}$$

*Proof.* From Lemma 3.1, we have

$$\begin{aligned} \left(\frac{d}{dt} - \Delta\right) \ln(|H|^2) &= \frac{1}{|H|^2} \left(\frac{d}{dt} - \Delta\right) |H|^2 + |\nabla \ln |H|^2|^2 \\ &\leq \frac{1}{|H|^2} \left(2|A|^2 |H|^2 - 2|\nabla |H|^2|^2\right) + |\nabla \ln |H|^2|^2 \\ &\leq 2|A|^2 + \frac{1}{2} |\nabla \ln |H|^2|^2. \end{aligned}$$

In the last step, we have used  $-\frac{2|\nabla |H|^2|^2}{|H|^2} = -\frac{|\nabla \ln |H|^2|^2}{2}$ .

Using  $\left(\frac{d}{dt} - \Delta\right)(\delta t |H|^2 + \epsilon) = \delta |H|^2 + \delta t \left(\frac{d}{dt} - \Delta\right) |H|^2$  and Lemma 3.1, we compute the evolution equation of  $\ln(\delta t |H|^2 + \epsilon)$  to obtain

$$\begin{aligned} &\left(\frac{d}{dt} - \Delta\right) \ln(\delta t |H|^2 + \epsilon) \\ &= \frac{1}{\delta t |H|^2 + \epsilon} \left(\frac{d}{dt} - \Delta\right) (\delta t |H|^2 + \epsilon) + |\nabla \ln(\delta t |H|^2 + \epsilon)|^2 \\ &\leq \frac{1}{\delta t |H|^2 + \epsilon} \left(\delta |H|^2 + \delta t (2|A|^2 |H|^2 - 2|\nabla |H|^2|^2)\right) + |\nabla \ln(\delta t |H|^2 + \epsilon)|^2 \\ &= \frac{\delta |H|^2 + 2(\delta t |H|^2 + \epsilon) |A|^2 - 2\epsilon |A|^2}{\delta t |H|^2 + \epsilon} - \frac{2\delta t |\nabla |H|^2|^2}{\delta t |H|^2 + \epsilon} + |\nabla \ln(\delta t |H|^2 + \epsilon)|^2 \\ &= 2|A|^2 + \frac{\delta |H|^2 - 2\epsilon |A|^2}{\delta t |H|^2 + \epsilon} - \frac{\delta t |\nabla |H|^2|^2}{2(\delta t |H|^2 + \epsilon) |H|^2} + |\nabla \ln(\delta t |H|^2 + \epsilon)|^2. \end{aligned}$$

In the last step, we have used  $|\nabla |H|^2|^2 = \frac{|\nabla |H|^2|^2}{4|H|^2}$ . Since  $|H|^2 \leq n|A|^2$  and  $-\frac{\delta t |\nabla |H|^2|^2}{2(\delta t |H|^2 + \epsilon) |H|^2} + \frac{1}{2} |\nabla \ln(\delta t |H|^2 + \epsilon)|^2 \leq 0$ , we can choose  $\delta \leq \frac{2\epsilon}{n}$  and get

$$\left(\frac{d}{dt} - \Delta\right) \ln(\delta t |H|^2 + \epsilon) \leq 2|A|^2 + \frac{1}{2} |\nabla \ln(\delta t |H|^2 + \epsilon)|^2.$$

The rest of the lemma can be proved in a similar fashion to Lemma 2.1. □

To derive an a priori curvature estimate, we need to find the right geometric quantity to counteract the quadratic growth of the second fundamental forms in Lemma 3.2. When  $n = 2$ , we are able to find the right quantity to establish the a priori mean curvature estimate.

In [TW], a parallel symmetric two tensor  $S$  is introduced to study the area decreasing map. We first recall some basic notation and definitions from Section 3 and Section 4 in [TW].

When  $M = \Sigma_1 \times \Sigma_2$  is the product of  $\Sigma_1$  and  $\Sigma_2$ , we denote the projections by  $\pi_1 : M \rightarrow \Sigma_1$  and  $\pi_2 : M \rightarrow \Sigma_2$ . By abusing notation, we also denote the differentials by  $\pi_1 : T_p M \rightarrow T_{\pi_1(p)} \Sigma_1$  and  $\pi_2 : T_p M \rightarrow T_{\pi_2(p)} \Sigma_2$  at any point  $p \in M$ .

When  $\Sigma$  is the graph of  $f : \Sigma_1 \rightarrow \Sigma_2$ , the equation at each point can be written in terms of the singular values of  $df$  and special bases adapted to  $df$ . Denote the singular values of  $df$ , or eigenvalues of  $\sqrt{(df)^T df}$ , by  $\{\lambda_i\}_{i=1, \dots, n}$ . Let  $r$  denote the rank of  $df$ . We can rearrange them so that  $\lambda_i = 0$  when  $i$  is greater than  $r$ . By singular value decomposition, there exist orthonormal bases  $\{a_i\}_{i=1, \dots, n}$  for  $T_{\pi_1(p)} \Sigma_1$  and  $\{a_\alpha\}_{\alpha=n+1, \dots, n+m}$  for  $T_{\pi_2(p)} \Sigma_2$  such that

$$df(a_i) = \lambda_i a_{n+i}$$

for  $i$  less than or equal to  $r$  and  $df(a_i) = 0$  for  $i$  greater than  $r$ . Moreover,

$$(3.5) \quad e_i = \begin{cases} \frac{1}{\sqrt{1+\lambda_i^2}}(a_i + \lambda_i a_{n+i}) & \text{if } 1 \leq i \leq r, \\ a_i & \text{if } r+1 \leq i \leq n, \end{cases}$$

becomes an orthonormal basis for  $T_p\Sigma$ , and

$$(3.6) \quad e_{n+p} = \begin{cases} \frac{1}{\sqrt{1+\lambda_p^2}}(a_{n+p} - \lambda_p a_p) & \text{if } 1 \leq p \leq r, \\ a_{n+p} & \text{if } r+1 \leq p \leq m, \end{cases}$$

becomes an orthonormal basis for  $N_p\Sigma$ .

The tangent space of  $M = \Sigma_1 \times \Sigma_2$  is identified with  $T\Sigma_1 \oplus T\Sigma_2$ . Let  $\pi_1$  and  $\pi_2$  denote the projection onto the first and second summand in the splitting. We define the parallel symmetric two-tensor  $S$  by

$$(3.7) \quad S(X, Y) = \langle \pi_1(X), \pi_1(Y) \rangle - \langle \pi_2(X), \pi_2(Y) \rangle$$

for any  $X, Y \in TM$ .

Let  $\Sigma$  be the graph of  $f : \Sigma_1 \rightarrow \Sigma_1 \times \Sigma_2$ .  $S$  restricts to a symmetric two-tensor on  $\Sigma$ , and we can represent  $S$  in terms of the orthonormal basis (3.5).

Let  $r$  denote the rank of  $df$ . By (3.5), it is not hard to check that

$$(3.8) \quad \begin{aligned} \pi_1(e_i) &= \frac{a_i}{\sqrt{1+\lambda_i^2}}, \pi_2(e_i) = \frac{\lambda_i a_{n+i}}{\sqrt{1+\lambda_i^2}} \text{ for } 1 \leq i \leq r, \text{ and} \\ \pi_1(e_i) &= a_i, \pi_2(e_i) = 0 \text{ for } r+1 \leq i \leq n. \end{aligned}$$

Similarly, by (3.6) we have

$$(3.9) \quad \begin{aligned} \pi_1(e_{n+p}) &= \frac{-\lambda_p a_p}{\sqrt{1+\lambda_p^2}}, \pi_2(e_{n+p}) = \frac{a_{n+p}}{\sqrt{1+\lambda_p^2}} \text{ for } 1 \leq p \leq r, \text{ and} \\ \pi_1(e_{n+p}) &= 0, \pi_2(e_{n+p}) = a_{n+p} \text{ for } r+1 \leq p \leq m. \end{aligned}$$

From the definition of  $S$ , we have

$$(3.10) \quad S(e_i, e_j) = \frac{1 - \lambda_i^2}{1 + \lambda_i^2} \delta_{ij}.$$

In particular, the eigenvalues of  $S$  are

$$(3.11) \quad \frac{1 - \lambda_i^2}{1 + \lambda_i^2}, \quad i = 1, \dots, n.$$

Now, at each point we express  $S$  in terms of the orthonormal basis  $\{e_i\}_{i=1, \dots, n}$  and  $\{e_\alpha\}_{\alpha=n+1, \dots, n+m}$ . Let  $I_{k \times k}$  denote a  $k$  by  $k$  identity matrix. Then  $S$  can be written in the block form

$$(3.12) \quad S = \left( S(e_k, e_l) \right)_{1 \leq k, l \leq n+m} = \begin{pmatrix} B & 0 & D & 0 \\ 0 & I_{n-r \times n-r} & 0 & 0 \\ D & 0 & -B & 0 \\ 0 & 0 & 0 & -I_{m-r \times m-r} \end{pmatrix}$$

where  $B$  and  $D$  are  $r$  by  $r$  matrices with  $B_{ij} = S(e_i, e_j) = \frac{1 - \lambda_i^2}{1 + \lambda_i^2} \delta_{ij}$  and  $D_{ij} = S(e_i, e_{n+j}) = \frac{-2\lambda_i}{1 + \lambda_i^2} \delta_{ij}$  for  $1 \leq i, j \leq r$ .

Next we recall the evolution equation of parallel two-tensors from [SW, Sm3]. Given a parallel two-tensor  $S$  on  $M$ , we consider the evolution of  $S$  restricted to  $\Sigma_t$ . This is a family of time-dependent symmetric two-tensors on  $\Sigma_t$ .

**Proposition 3.2.** *Suppose  $M$  is a flat Riemannian manifold. For a mean curvature flow  $F : \Sigma \times [0, T) \rightarrow M$  of an  $n$ -dimensional submanifold. Let  $S$  be a parallel two-tensor on  $M$ . Then the pull-back of  $S$  to  $\Sigma_t$  satisfies the following equation:*

$$(3.13) \quad \begin{aligned} \left(\frac{d}{dt} - \Delta\right)S_{ij} &= -h_{\alpha il}H_{\alpha}S_{lj} - h_{\alpha jl}H_{\alpha}S_{li} \\ &+ R_{kik\alpha}S_{\alpha j} + R_{kjk\alpha}S_{\alpha i} \\ &+ h_{\alpha kl}h_{\alpha ki}S_{lj} + h_{\alpha kl}h_{\alpha kj}S_{li} - 2h_{\alpha ki}h_{\beta kj}S_{\alpha\beta} \end{aligned}$$

where  $\Delta$  is the rough Laplacian on two-tensors over  $\Sigma_t$ , and  $S_{\alpha i} = S(e_{\alpha}, e_i)$ ,  $S_{\alpha\beta} = S(e_{\alpha}, e_{\beta})$ , and  $R_{kik\alpha} = R(e_k, e_i, e_k, e_{\alpha})$  is the curvature of  $M$ .

The evolution equations (3.13) of  $S$  can be written in terms of evolving orthonormal frames as in Hamilton [H]. If the orthonormal frames

$$(3.14) \quad F = \{F_1, \dots, F_a, \dots, F_n\}$$

are given in local coordinates by

$$F_a = F_a^i \frac{\partial}{\partial x_i},$$

then to keep them orthonormal, i.e.  $g_{ij}F_a^iF_b^j = \delta_{ab}$ , we evolve  $F$  by the formula

$$\frac{\partial}{\partial t}F_a^i = g^{ij}g^{\alpha\beta}h_{\alpha jl}H_{\beta}F_a^l.$$

Let  $S_{ab} = S_{ij}F_a^iF_b^j$  be the components of  $S$  in  $F$ . Then  $S_{ab}$  satisfies the equation

$$(3.15) \quad \begin{aligned} \left(\frac{d}{dt} - \Delta\right)S_{ab} &= R_{cac\alpha}S_{\alpha b} + R_{cbc\alpha}S_{\alpha a} \\ &+ h_{\alpha cd}h_{\alpha ca}S_{db} + h_{\alpha cd}h_{\alpha cb}S_{da} \\ &- 2h_{\alpha ca}h_{\beta cb}S_{\alpha\beta}. \end{aligned}$$

In the following, we will compute the evolution of  $Tr(S) = \sum_{i,j} g^{ij}S_{ij}$  in the case  $\Sigma_1, \Sigma_2$  are flat Riemann surfaces, i.e. where the curvature tensor  $R = 0$ . By equation (3.10),  $Tr(S) = \frac{1-\lambda_1^2}{1+\lambda_1^2} + \frac{1-\lambda_2^2}{1+\lambda_2^2} = \frac{2(1-\lambda_1^2\lambda_2^2)}{(1+\lambda_1^2)(1+\lambda_2^2)}$ . The next proposition gives a new proof that the area decreasing condition is preserved along the mean curvature. In addition, the equation satisfied by  $\ln Tr(S)$  plays a critical role in the next section’s curvature estimates.

**Proposition 3.3.** *Suppose  $M = \Sigma_1 \times \Sigma_2$  where  $\Sigma_1$  and  $\Sigma_2$  are complete flat Riemann surfaces. Suppose  $\Sigma$  is the graph of a map  $f : \Sigma_1 \rightarrow \Sigma_2$  and let  $\Sigma_t$  denote its mean curvature flow with initial surface  $\Sigma_0 = \Sigma$ . Let  $S$  be given by (3.7); then  $Tr(S)$  on  $\Sigma_t$  satisfies the equation*

$$(3.16) \quad \begin{aligned} &\left(\frac{d}{dt} - \Delta\right) \ln Tr(S) \\ &= 2|A|^2 + \frac{|\nabla \ln(TrS)|^2}{2} + \frac{2\sum_{c=1}^2 (T_{11}h_{4c2} + T_{22}h_{3c1})^2}{Tr(S)^2}, \end{aligned}$$

where  $T_{11} = \frac{2\lambda_1}{(1+\lambda_1^2)}$  and  $T_{22} = \frac{2\lambda_2}{(1+\lambda_2^2)}$ .



*Proof.* Using the evolution equation of  $S$  (with respect to an orthonormal frame) in equation (3.15) and  $S(e_{2+i}, e_{2+j}) = -S_{ij}$ , we derive

$$\begin{aligned} & \left(\frac{d}{dt} - \Delta\right)Tr(S) \\ &= \sum_{a,b} \delta^{ab} \left(\sum_{\alpha,c,d} h_{\alpha cd} h_{\alpha ca} S_{db} + h_{\alpha cd} h_{\alpha cb} S_{da} - 2h_{\alpha ca} h_{\beta cb} S_{\alpha\beta}\right) \\ &= \sum_a \left(\sum_{\alpha,c} 2h_{\alpha ca}^2 S_{aa} - 2h_{\alpha ca}^2 S_{\alpha\alpha}\right) \\ &= \sum_a \left(\sum_{p,c} 2h_{2+p\ ca}^2 S_{aa} + 2h_{2+p\ ca}^2 S_{pp}\right) \\ &= \sum_{p,c} \left[2h_{2+p\ c1}^2 S_{11} + 2h_{2+p\ c1}^2 S_{pp} + 2h_{2+p\ c2}^2 S_{22} + 2h_{2+p\ c2}^2 S_{pp}\right] \\ &= \sum_c \left[(2h_{4c1}^2 + 2h_{3c2}^2)(S_{11} + S_{22}) + 4h_{3c1}^2 S_{11} + 4h_{4c2}^2 S_{22}\right]. \end{aligned}$$

Using  $|A|^2 = \sum_{c=1}^2 (h_{4c1}^2 + h_{3c2}^2 + h_{3c1}^2 + h_{4c2}^2)$ , we obtain

$$(3.17) \quad \left(\frac{d}{dt} - \Delta\right)Tr(S) = 2|A|^2 Tr(S) + 2(S_{11} - S_{22}) \sum_{c=1}^2 (h_{3c1}^2 - h_{4c2}^2).$$

We claim the following relation holds:

$$\begin{aligned} & 4Tr(S)(S_{11} - S_{22}) \sum_{c=1}^2 (h_{3c1}^2 - h_{4c2}^2) + |\nabla Tr(S)|^2 \\ (3.18) \quad &= 4 \sum_{c=1}^2 (T_{11} h_{4c2} + T_{22} h_{3c1})^2. \end{aligned}$$

Equation (3.16) follows from equations (3.17) and (3.18).

In the rest of the proof, we verify equation (3.18). The covariant derivative of the restriction of  $S$  on  $\Sigma$  can be computed by

$$\begin{aligned} & (\nabla_{e_k} S)(e_i, e_j) \\ &= e_k(S(e_i, e_j)) - S(\nabla_{e_k} e_i, e_j) - S(e_i, \nabla_{e_k} e_j) \\ &= S(\nabla_{e_k}^M e_i - \nabla_{e_k} e_i, e_j) - S(e_i, \nabla_{e_k}^M e_j - \nabla_{e_k} e_j) \\ &= h_{\alpha ki} S_{\alpha j} + h_{\beta kj} S_{\beta i}. \end{aligned}$$

Since  $S_{2+p\ l} = -\frac{2\lambda_p \delta_{pl}}{(1+\lambda_p^2)}$ , we derive

$$\begin{aligned} (3.19) \quad S_{ij,k} &= -\frac{2h_{2+p\ ki} \lambda_p \delta_{pj}}{(1+\lambda_p^2)} - \frac{2h_{2+p\ kj} \lambda_p \delta_{pi}}{(1+\lambda_p^2)} \\ &= -h_{2+p\ ki} T_{pp} \delta_{pj} - h_{2+p\ kj} T_{pp} \delta_{pi} \\ &= -h_{2+j\ ki} T_{jj} - h_{2+i\ kj} T_{ii}. \end{aligned}$$

In particular,  $\nabla_k(S_{ii}) = -2T_{ii} h_{2+i\ ki}$  and

$$|\nabla Tr(S)|^2 = 4 \sum_{k=1}^2 (T_{11}^2 h_{3k1}^2 + 2T_{11} T_{22} h_{3k1} h_{4k2} + T_{22}^2 h_{4k2}^2).$$

We compute

$$\begin{aligned}
 & 4(S_{11}^2 - S_{22}^2) \sum_{k=1}^2 (h_{3k1}^2 - h_{4k2}^2) + |\nabla Tr(S)|^2 \\
 &= 4(S_{11}^2 - S_{22}^2) \sum_{k=1}^2 (h_{3k1}^2 - h_{4k2}^2) \\
 &\quad + 4 \sum_{k=1}^2 (T_{11}^2 h_{3k1}^2 + 2T_{11}T_{22}h_{3k1}h_{4k2} + T_{22}^2 h_{4k2}^2) \\
 &= 4 \sum_{k=1}^2 (T_{11}h_{4k2} + T_{22}h_{3k1})^2,
 \end{aligned}$$

where we use the fact that  $S_{11}^2 + T_{11}^2 = S_{22}^2 + T_{22}^2 = 1$  to complete the square in the last equality. This verifies (3.18). □

#### 4. PROOF OF THE THEOREMS

**4.1. Proof of Theorem 1.** Since  $Tr(S) = \frac{2(1-\lambda_1^2\lambda_2^2)}{(1+\lambda_1^2)(1+\lambda_2^2)}$ , by Proposition 3.3 and the maximum principle, we deduce that the area decreasing condition is preserved by the mean curvature flow and  $inf_{\Sigma_t} Tr(S) \geq \alpha$ , where  $\alpha = inf_{\Sigma_0} Tr(S)$ . On any  $\Sigma_t$ ,  $t > 0$ ,  $Tr(S) \geq \alpha > 0$  and  $(1 + \lambda_1^2)(1 + \lambda_2^2) < \frac{2}{\alpha}$ . Thus  $\Sigma_t$  remains as the graph of an area decreasing map.

Next we derive the mean curvature decay estimate. Combining the first equation in Lemma 3.2 (with  $\delta = \epsilon = 1$ ) and Proposition 3.3, we obtain

$$\left(\frac{d}{dt} - \Delta\right) \ln \frac{(t|H|^2 + 1)}{Tr(S)} \leq \frac{1}{2} \nabla \ln \frac{(t|H|^2 + 1)}{Tr(S)} \cdot \nabla \ln [(t|H|^2 + 1)Tr(S)].$$

By the maximum principle, we have

$$sup_{\Sigma_t} \ln \frac{(t|H|^2 + 1)}{Tr(S)} \leq sup_{\Sigma_0} \ln \frac{1}{Tr(S)}.$$

Note that  $f_t$  remains area decreasing and

$$Tr(S) = \frac{2(1 - \lambda_1^2\lambda_2^2)}{(1 + \lambda_1^2)(1 + \lambda_2^2)} < 2.$$

This implies that  $t|H|^2 \leq Tr(S) sup_{\Sigma_0} \frac{1}{Tr(S)} \leq sup_{\Sigma_0} \frac{2}{Tr(S)} = \frac{2}{\alpha}$ . If  $\Sigma_2$  is compact we already have long-time existence of the flow by the methods in [W1, W3, TW]. In case  $\Sigma_2$  is complete and noncompact we can use the mean curvature estimate to obtain a  $C^0$ -estimate on finite time intervals, and then we may proceed as in [SHS2] to get long-time existence.

Now we can prove the  $C^\infty$  convergence using the mean curvature decay estimate. By the Gauss formula,  $\int_{\Sigma_t} |A|^2 d\mu_t = \int_{\Sigma_t} |H|^2 d\mu_t \rightarrow 0$ . Therefore  $\int_{\Sigma_t} |A|^2 d\mu_t$  is sufficiently small when  $t$  is large enough; the  $\epsilon$  regularity theorem in [I] (see also [E]) implies  $sup_{\Sigma_t} |A|^2$  is uniformly bounded. The general convergence theorem of Simon [S2] implies  $C^\infty$  convergence of  $\Sigma_t$  to a minimal submanifold  $\Sigma_\infty$ , which is totally geodesic by the Gauss formula again.

**4.2. Proof of Theorem 2.** Now we prove a decay estimate for the second fundamental form in the case when  $\Sigma$  is a Lagrangian submanifold. It is well known that  $\Sigma_t$  remains as a Lagrangian submanifold in  $T^2 \times T^2$  from [Sm1]. Using the first equation in Lemma 3.1 and Proposition 3.3, we derive

$$\begin{aligned} & \left(\frac{d}{dt} - \Delta\right) \ln\left(\frac{|A|^2}{Tr(S)^2}\right) \\ & \leq \frac{|\nabla \ln(|A|^2)|^2}{2} - |\nabla \ln Tr(S)|^2 - |A|^2 - \frac{4}{Tr(S)^2} \sum_{c=1}^2 (T_{11}h_{4c2} + T_{22}h_{3c1})^2 \\ & = \frac{1}{2} \nabla \ln\left(\frac{|A|^2}{Tr(S)^2}\right) \cdot \nabla \ln(|A|^2 Tr(S)^2) - |A|^2 \\ & \quad + |\nabla \ln Tr(S)|^2 - \frac{4}{Tr(S)^2} \sum_{c=1}^2 (T_{11}h_{4c2} + T_{22}h_{3c1})^2. \end{aligned}$$

We estimate the last two terms on the right hand side. From equation (3.18), we obtain

$$\begin{aligned} & |\nabla \ln Tr(S)|^2 - \frac{4}{Tr(S)^2} \sum_{c=1}^2 (T_{11}h_{4c2} + T_{22}h_{3c1})^2 \\ & = -\frac{4(S_{11} - S_{22})}{Tr(S)} \sum_{c=1}^2 (h_{3c1}^2 - h_{4c2}^2). \end{aligned}$$

Let  $J$  be the standard almost complex structure on  $(T^2, \{x^i\}_{i=1,2}) \times (T^2, \{y^j\}_{j=1,2})$  that maps from the tangent space of the first component to the tangent space of the second one, and vice versa. Suppose  $f$  is the defining map of a Lagrangian surface  $\Sigma$  in  $(T^2, \{x^i\}_{i=1,2}) \times (T^2, \{y^j\}_{j=1,2})$  with respect to  $dx^1 \wedge dy^1 + dx^2 \wedge dy^2$ . By the Lagrangian condition,  $Jdf$  is a self-adjoint map on the tangent space of  $(T^2, \{x^i\}_{i=1,2})$ . Therefore, there exists an orthonormal basis  $\{a_i\}_{i=1,2}$  such that

$$df(a_i) = \lambda_i J(a_i), \quad i = 1, 2.$$

We can then choose  $a_{2+i} = J(a_i)$  in equations (3.5) and (3.6) such that

$$e_i = \frac{1}{\sqrt{1 + \lambda_i^2}}(a_i + \lambda_i J(a_i))$$

and

$$e_{2+i} = \frac{1}{\sqrt{1 + \lambda_i^2}}(J(a_i) - \lambda_i a_i), \quad i = 1, 2.$$

From these expressions, it is easy to check that  $e_3 = J(e_1)$  and  $e_4 = J(e_2)$ . Since  $\Sigma_t$  remains Lagrangian, such orthonormal frames can be picked at any point on  $\Sigma_t$ . Because  $J$  is parallel,

$$\langle \nabla_{e_c}^M e_2, e_3 \rangle = \langle \nabla_{e_c}^M e_1, e_4 \rangle,$$

and we have  $h_{3c2}^2 = h_{4c1}^2$  for  $c = 1, 2$ . Therefore,

$$\left| \sum_{c=1}^2 (h_{3c1}^2 - h_{4c2}^2) \right| \leq |h_{311} - h_{322}| |H_3| + |h_{411} - h_{422}| |H_4| \leq 2\sqrt{2} |A| |H|$$

and

$$\|\nabla \ln \operatorname{Tr}(S)\|^2 - \frac{4}{\operatorname{Tr}(S)^2} \sum_{c=1}^2 (T_{11}h_{4c2} + T_{22}h_{3c1})^2 \leq C|A||H|,$$

where  $C$  depends only on  $\alpha = \inf_{\Sigma_0} \frac{2(1-\lambda_1^2\lambda_2^2)}{(1+\lambda_1^2)(1+\lambda_2^2)} > 0$  on the initial surface. For example,  $C = \frac{16\sqrt{2}}{\alpha}$  suffices. To this end,

$$\left(\frac{d}{dt} - \Delta\right) \ln\left(\frac{|A|^2}{\operatorname{Tr}(S)^2}\right) \leq \frac{1}{2} \nabla \ln\left(\frac{|A|^2}{\operatorname{Tr}(S)^2}\right) \cdot \nabla \ln(|A|^2 \operatorname{Tr}(S)^2) - |A|^2 + C|A||H|.$$

Note that  $-|A|^2 \leq -\alpha^2 \frac{|A|^2}{\operatorname{Tr}(S)^2}$  and  $|H|^2 \leq \frac{2}{t\alpha}$  from Theorem 1, and we derive

$$-|A|^2 + C|A||H| \leq -\frac{1}{2}|A|^2 + \frac{C^2}{2}|H|^2 \leq -\frac{\alpha^2|A|^2}{2\operatorname{Tr}(S)^2} + \frac{C^2}{\alpha t}.$$

Therefore, the quantity  $\frac{|A|^2}{\operatorname{Tr}(S)^2}$  satisfies the following differential inequality:

$$\begin{aligned} & \left(\frac{d}{dt} - \Delta\right) \left(\frac{|A|^2}{\operatorname{Tr}(S)^2}\right) \\ & \leq \frac{1}{2} \nabla \left(\frac{|A|^2}{\operatorname{Tr}(S)^2}\right) \cdot \nabla \ln(|A|^2 \operatorname{Tr}(S)^2) + \left(-\frac{\alpha^2|A|^2}{2\operatorname{Tr}(S)^2} + \frac{C^2}{\alpha t}\right) \frac{|A|^2}{\operatorname{Tr}(S)^2}. \end{aligned}$$

Consider the ODE  $u' = \left(-\frac{\alpha^2}{2}u + \frac{C^2}{\alpha t}\right)u$ . Note that  $w = \frac{2(1+\frac{C^2}{\alpha})}{\alpha^2 t}$  is a solution to  $u' = \left(-\frac{\alpha^2}{2}u + \frac{C^2}{\alpha t}\right)u$  and  $\lim_{t \rightarrow 0^+} w(t) = \infty$ . Thus we have  $\sup_{\Sigma_t} \left(\frac{|A|^2}{\operatorname{Tr}(S)^2}\right) \leq \frac{2(1+\frac{C^2}{\alpha})}{\alpha^2 t}$  and  $|A|^2 \leq \frac{8(1+\frac{C^2}{\alpha})}{\alpha^2 t}$ .

#### ACKNOWLEDGEMENTS

Part of this paper was completed while the authors were visiting Taida Institute of Mathematical Sciences, National Center for Theoretical Sciences, Taipei Office in National Taiwan University, Taipei, Taiwan, and the Riemann Center for Geometry and Physics in Leibniz Universität Hannover. The authors wish to express their gratitude for the excellent support they received during their stay.

#### REFERENCES

- [AMJ] Li An-Min and Li Jimin, *An intrinsic rigidity theorem for minimal submanifolds in a sphere*, Arch. Math. (Basel) **58** (1992), no. 6, 582–594, DOI 10.1007/BF01193528. MR1161925 (93b:53050)
- [CCH] Albert Chau, Jingyi Chen, and Weiyong He, *Lagrangian mean curvature flow for entire Lipschitz graphs*, Calc. Var. Partial Differential Equations **44** (2012), no. 1-2, 199–220, DOI 10.1007/s00526-011-0431-x. MR2898776
- [CCY] Albert Chau, Jingyi Chen, and Yu Yuan, *Lagrangian mean curvature flow for entire Lipschitz graphs II*, Math. Ann. **357** (2013), no. 1, 165–183, DOI 10.1007/s00208-013-0897-2. MR3084345
- [E] Klaus Ecker, *Regularity theory for mean curvature flow*, Progress in Nonlinear Differential Equations and their Applications, 57, Birkhäuser Boston, Inc., Boston, MA, 2004. MR2024995 (2005b:53108)
- [EH1] Klaus Ecker and Gerhard Huisken, *Mean curvature evolution of entire graphs*, Ann. of Math. (2) **130** (1989), no. 3, 453–471, DOI 10.2307/1971452. MR1025164 (91c:53006)
- [EH2] Klaus Ecker and Gerhard Huisken, *Interior estimates for hypersurfaces moving by mean curvature*, Invent. Math. **105** (1991), no. 3, 547–569, DOI 10.1007/BF01232278. MR1117150 (92i:53010)

- [H] Richard S. Hamilton, *Harnack estimate for the mean curvature flow*, J. Differential Geom. **41** (1995), no. 1, 215–226. MR1316556 (95m:53055)
- [I] Tom Ilmanen, *Singularities of mean curvature flow of surfaces*, preprint (1997).
- [LL] Kuo-Wei Lee and Yng-Ing Lee, *Mean curvature flow of the graphs of maps between compact manifolds*, Trans. Amer. Math. Soc. **363** (2011), no. 11, 5745–5759, DOI 10.1090/S0002-9947-2011-05204-9. MR2817407 (2012h:53154)
- [S1] Richard M. Schoen, *The role of harmonic mappings in rigidity and deformation problems*, Complex geometry (Osaka, 1990), Lecture Notes in Pure and Appl. Math., vol. 143, Dekker, New York, 1993, pp. 179–200. MR1201611 (94g:58055)
- [S2] Leon Simon, *Asymptotics for a class of nonlinear evolution equations, with applications to geometric problems*, Ann. of Math. (2) **118** (1983), no. 3, 525–571, DOI 10.2307/2006981. MR727703 (85b:58121)
- [SHS1] Andreas Savas-Halilaj and Knut Smoczyk, *Homotopy of area decreasing maps by mean curvature flow*, Adv. Math. **255** (2014), 455–473, DOI 10.1016/j.aim.2014.01.014. MR3167489
- [SHS2] Andreas Savas-Halilaj and Knut Smoczyk, *Evolution of contractions by mean curvature flow*, Math. Ann. **361** (2015), no. 3-4, 725–740, DOI 10.1007/s00208-014-1090-y. MR3319546
- [Sm1] Knut Smoczyk, *A canonical way to deform a Lagrangian submanifold*, arXiv: dg-ga/9605005 (1996).
- [Sm2] Knut Smoczyk, *Mean curvature flow in higher codimension: introduction and survey*, Global differential geometry, Springer Proc. Math., vol. 17, Springer, Heidelberg, 2012, pp. 231–274, DOI 10.1007/978-3-642-22842-1\_9. MR3289845
- [Sm3] Knut Smoczyk, *Longtime existence of the Lagrangian mean curvature flow*, Calc. Var. Partial Differential Equations **20** (2004), no. 1, 25–46, DOI 10.1007/s00526-003-0226-9. MR2047144 (2004m:53119)
- [SW] Knut Smoczyk and Mu-Tao Wang, *Mean curvature flows of Lagrangians submanifolds with convex potentials*, J. Differential Geom. **62** (2002), no. 2, 243–257. MR1988504 (2004d:53086)
- [TW] Mao-Pei Tsui and Mu-Tao Wang, *Mean curvature flows and isotopy of maps between spheres*, Comm. Pure Appl. Math. **57** (2004), no. 8, 1110–1126, DOI 10.1002/cpa.20022. MR2053760 (2005b:53110)
- [W1] Mu-Tao Wang, *Mean curvature flow of surfaces in Einstein four-manifolds*, J. Differential Geom. **57** (2001), no. 2, 301–338. MR1879229 (2003j:53108)
- [W2] Mu-Tao Wang, *Deforming area preserving diffeomorphism of surfaces by mean curvature flow*, Math. Res. Lett. **8** (2001), no. 5-6, 651–661, DOI 10.4310/MRL.2001.v8.n5.a7. MR1879809 (2003f:53122)
- [W3] Mu-Tao Wang, *Long-time existence and convergence of graphic mean curvature flow in arbitrary codimension*, Invent. Math. **148** (2002), no. 3, 525–543, DOI 10.1007/s002220100201. MR1908059 (2003b:53073)
- [W4] Mu-Tao Wang, *Gauss maps of the mean curvature flow*, Math. Res. Lett. **10** (2003), no. 2-3, 287–299, DOI 10.4310/MRL.2003.v10.n3.a2. MR1981905 (2004m:53121)
- [W5] Mu-Tao Wang, *Subsets of Grassmannians preserved by mean curvature flows*, Comm. Anal. Geom. **13** (2005), no. 5, 981–998. MR2216149 (2006m:53102)
- [W6] Mu-Tao Wang, *Remarks on a class of solutions to the minimal surface system*, Geometric evolution equations, Contemp. Math., vol. 367, Amer. Math. Soc., Providence, RI, 2005, pp. 229–235, DOI 10.1090/conm/367/06758. MR2115762 (2005i:53010)

INSTITUT FÜR DIFFERENTIALGEOMETRIE AND RIEMANN CENTER FOR GEOMETRY AND PHYSICS,  
LEIBNIZ UNIVERSITÄT HANNOVER, WELFENGARTEN 1, 30167 HANNOVER, GERMANY

*E-mail address:* [smoczyk@math.uni-hannover.de](mailto:smoczyk@math.uni-hannover.de)

DEPARTMENT OF MATHEMATICS, NATIONAL TAIWAN UNIVERSITY, TAIPEI 10617, TAIWAN – AND –  
DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF TOLEDO, 2801 W. BANCROFT  
STREET, TOLEDO, OHIO 43606-3390

*E-mail address:* [mao-pei.tsui@utoledo.edu](mailto:mao-pei.tsui@utoledo.edu)

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, 2990 BROADWAY, NEW YORK, NEW  
YORK 10027

*E-mail address:* [mtwang@math.columbia.edu](mailto:mtwang@math.columbia.edu)