

## FIXED-POINTS IN THE CONE OF TRACES ON A $C^*$ -ALGEBRA

MIKAEL RØRDAM

*Dedicated to the memory of John Roe*

ABSTRACT. Nicolas Monod introduced the class of groups with the fixed-point property for cones, characterized by always admitting a nonzero fixed-point when acting (suitably) on proper weakly complete cones. He proved that his class of groups contains the class of groups with subexponential growth and is contained in the class of supramenable groups. In this paper we investigate what Monod’s results say about the existence of invariant traces on (typically nonunital)  $C^*$ -algebras equipped with an action of a group with the fixed-point property for cones. As an application of these results, we provide results on the existence (and nonexistence) of traces on the (nonuniform) Roe algebra.

### 1. INTRODUCTION

Whenever a discrete amenable group acts on a unital  $C^*$ -algebra with at least one tracial state, the  $C^*$ -algebra admits an invariant tracial state, and the crossed product  $C^*$ -algebra admits a tracial state. This, moreover, characterizes amenable groups. The purpose of this paper is to find statements, similar to this well-known fact, about the existence of invariant traces on nonunital  $C^*$ -algebras using the results of the recent paper by Monod [16] in which the class of groups with the *fixed-point property for cones* is introduced and developed.

In [16] a group is said to have the fixed-point property for cones if whenever it acts continuously on a proper weakly complete cone (embedded in a locally convex topological vector space), such that the action is of *cobounded type* and *locally bounded*, then there is a nonzero fixed-point in the cone. Being of cobounded type is an analogue of an action on a locally compact Hausdorff space being *co-compact*; see [16, Definition 2]. The action is locally bounded if there is a nonzero bounded orbit; see [16, Definition 1].

Even the group of integers,  $\mathbb{Z}$ , can fail to leave invariant any nonzero trace when acting on a nonunital  $C^*$ -algebra. For example, the stabilization of the Cuntz algebra  $\mathcal{O}_2$  is the crossed product of the stabilized CAR-algebra  $\mathcal{A}$  with an action of  $\mathbb{Z}$  that scales the traces on  $\mathcal{A}$  by a factor of 2. In particular, there are no nonzero invariant traces on  $\mathcal{A}$ . The cone of (densely defined lower semicontinuous) traces on  $\mathcal{A}$  is isomorphic to the cone  $[0, \infty)$ , and the induced action of  $\mathbb{Z}$  on this cone is multiplication by 2, which of course fails to be locally bounded.

The action of any group  $G$  on a locally compact Hausdorff space  $X$  induces a locally bounded action on the cone of Radon measures on  $X$  (which again is

---

Received by the editors August 4, 2018, and, in revised form, January 1, 2019.

2010 *Mathematics Subject Classification*. Primary 46L35; Secondary 46L05, 37A55.

Supported by the Danish Council for Independent Research, Natural Sciences, and the Danish National Research Foundation (DNRF) through the Centre for Symmetry and Deformation at the University of Copenhagen.

the same as the cone of densely defined lower semicontinuous traces on  $C_0(X)$ . However, any infinite group can act on the locally compact noncompact Cantor set  $\mathbf{K}^*$  in a non-co-compact way so that there are no nonzero invariant Radon measures; see [15, Section 4]. Such an action of  $G$  on the Radon measures on  $\mathbf{K}^*$  fails to be of cobounded type.

The two examples above explain why one must impose conditions on the action, such as being of cobounded type and being locally bounded, to get meaningful results on when invariant traces exist.

Monod proves in [16] that the class of groups with the fixed-point property for cones contains the class of groups of subexponential growth and is contained in the class of supramenable groups introduced by Rosenblatt in [23]. It is not known if there are supramenable groups of exponential growth, so the three classes of groups could coincide, although the common belief seems to be that they all are different. Monod proved a number of permanence properties for his class of groups, prominently including the fact that it is closed under central extensions; see [16, Theorem 8]. He also shows that the property of having the fixed-point property for cones can be recast in several ways, including the property that for each nonzero positive function  $f \in \ell^\infty(G)$  there is a nonzero invariant positive linear functional (called an *invariant integral* in [16]) on the subspace  $\ell^\infty(G, f)$  of all bounded functions  $G$ -dominated by  $f$ .

We begin our paper in Section 2 by recalling properties of possibly unbounded positive traces on (typically nonunital)  $C^*$ -algebras, including when they are lower semicontinuous and when they are singular. By default, all traces in this paper are assumed to be positive. Interestingly, many of the traces predicted by Monod turn out to be singular. Traces on  $C^*$ -algebras were treated systematically already by Dixmier in [7]. Elliott, Robert, and Santiago consider in [8] the cone of lower semicontinuous traces as an invariant of the  $C^*$ -algebra, they derive useful properties of this cone, and they make the point that such traces most conveniently are viewed as maps defined on the positive cone of the  $C^*$ -algebra taking values in  $[0, \infty]$ . While this indeed is a convenient way to describe unbounded traces, and one we in part shall use here, the cone structure from the point of view of this paper is sometimes better portrayed when traces are viewed as linear functionals on a suitable domain: a hereditary symmetric algebraic ideal in the  $C^*$ -algebra.

By a theorem of G. K. Pedersen one can identify the cone of lower semicontinuous densely defined traces on a  $C^*$ -algebra with the cone of traces defined on its Pedersen ideal. In the case in which the primitive ideal space of the  $C^*$ -algebra is compact, we show that there are nonzero lower semicontinuous densely defined traces if and only if the stabilization of the  $C^*$ -algebra contains no full properly infinite projections, thus extending well-known results from both the unital and the simple case. A similar compactness condition appears in our reformulation of coboundedness of the action of the group on the cone of traces.

In Section 3 we explain when a  $C^*$ -algebra equipped with an action of a group  $G$  with the fixed-point property for cones admits an invariant densely defined lower semicontinuous trace. Given that the cone of densely defined lower semicontinuous traces is always proper and weakly complete, all that we have to do is to explain when the action of the group on this cone is of cobounded type and when it is locally bounded, respectively. The former can be translated into a compactness statement,

as mentioned above, while the latter just means that there exists a nonzero trace which is bounded on all  $G$ -orbits.

In Section 4 we examine the situation in which the group  $G$  acts on  $\ell^\infty(G)$  with the aim of describing for which positive  $f \in \ell^\infty(G)$  the invariant integrals on  $\ell^\infty(G)$  normalizing  $f$  are lower semicontinuous and singular, respectively. As it turns out, frequently they must be singular. We use this to give an example of a  $G$ -invariant densely defined trace on a  $C^*$ -algebra that does not extend to a trace on the crossed product.

Finally, in Section 5 we consider the particular example of invariant traces on  $\ell^\infty(G, \mathcal{K})$  and traces on the Roe algebra  $\ell^\infty(G, \mathcal{K}) \rtimes G$ . We show that  $\ell^\infty(G, \mathcal{K})$  has only the “obvious” densely defined lower semicontinuous traces, and hence that it never has nonzero  $G$ -invariant ones, when  $G$  is infinite. In many cases, however, there are invariant lower semicontinuous traces with smaller domains, such as domains defined by projections in  $\ell^\infty(G, \mathcal{K})$ . Specifically we show that any projection in  $\ell^\infty(G, \mathcal{K})$  whose dimension (as a function on  $G$ ) grows subexponentially is normalized by an invariant lower semicontinuous trace if  $G$  has the fixed-point property for cones. In general, for any nonlocally finite group  $G$  there are (necessarily exponentially growing) projections in  $\ell^\infty(G, \mathcal{K})$  not normalized by any invariant trace, and which are properly infinite in the Roe algebra, while the Roe algebra of a locally finite group is always stably finite.

## 2. HEREDITARY IDEALS AND CONES OF TRACES

By a *hereditary ideal*  $\mathcal{I}$  in a  $C^*$ -algebra  $\mathcal{A}$  we shall mean an algebraic two-sided self-adjoint ideal satisfying the hereditary property: If  $0 \leq a \leq b$ ,  $b \in \mathcal{I}$ , and  $a \in \mathcal{A}$ , then  $a \in \mathcal{I}$ . If  $x^*x \in \mathcal{I}$  whenever  $xx^* \in \mathcal{I}$ , for all  $x \in \mathcal{A}$ , then we say that  $\mathcal{I}$  is *symmetric*. All closed two-sided ideals are hereditary and symmetric.

If a group  $G$  acts on the  $C^*$ -algebra  $\mathcal{A}$ , then we refer to an ideal  $\mathcal{I}$  as being  *$G$ -invariant* (or just invariant) if it is invariant under the group action. If  $M$  is a subset of  $\mathcal{A}$ , then let  $\mathcal{I}_{\mathcal{A}}(M)$  (resp.,  $\overline{\mathcal{I}_{\mathcal{A}}}(M)$ ) denote the smallest hereditary ideal in  $\mathcal{A}$  (resp., the smallest closed two-ideal in  $\mathcal{A}$ ) which contains  $M$ , and let similarly  $\mathcal{I}_{\mathcal{A}}^G(M)$  and  $\overline{\mathcal{I}_{\mathcal{A}}^G}(M)$  denote the smallest  $G$ -invariant hereditary ideal and the smallest  $G$ -invariant closed two-ideal in  $\mathcal{A}$  which contain  $M$ , respectively.

**Example 2.1.** The *Pedersen ideal*,  $\text{Ped}(\mathcal{A})$ , of a  $C^*$ -algebra  $\mathcal{A}$  is the (unique) smallest dense ideal in  $\mathcal{A}$ ; see [18, Section 5.6]. It is a hereditary symmetric ideal, and, even better, for each  $x \in \text{Ped}(\mathcal{A})$ , the hereditary sub- $C^*$ -algebra,  $\overline{x\mathcal{A}x^*}$ , of  $\mathcal{A}$  is contained in  $\text{Ped}(\mathcal{A})$ . In particular, the Pedersen ideal of  $\mathcal{A}$  is closed under continuous function calculus on its normal elements, as long as the continuous function vanishes at 0. If  $x \in \mathcal{A}$  is such that  $x^*x \in \text{Ped}(\mathcal{A})$ , then  $x \in \text{Ped}(\mathcal{A})$ , which shows that the Pedersen ideal is also symmetric.

**Example 2.2.** Not all self-adjoint two-sided ideals in a  $C^*$ -algebra are hereditary. Consider, for example, the commutative  $C^*$ -algebra  $\mathcal{A} = C([-1, 1])$ , the element  $f \in C([-1, 1])$  given by  $f(t) = |t|$ , for  $t \in [-1, 1]$ , and the (two-sided) self-adjoint ideal  $\mathcal{I} = \mathcal{A}f$  in  $\mathcal{A}$ . The function  $g(t) = \max\{-\frac{1}{2}t, t\}$ ,  $t \in [-1, 1]$ , then satisfies  $0 \leq g \leq f$ , but  $g \notin \mathcal{I}$ .

Hereditary ideals need not be spanned (or even generated) by their positive elements. Indeed, take again  $\mathcal{A} = C([-1, 1])$ , and let  $\mathcal{I} = \mathcal{A}\iota$ , where  $\iota(t) = t$ , for  $t \in [-1, 1]$ . If  $f \in \mathcal{I} \cap \mathcal{A}^+$ , then  $f = g \cdot \iota$ , for some  $g \in \mathcal{P}$ , where  $\mathcal{P}$  is the set of

functions  $g \in \mathcal{A}$  such that  $g(t) \leq 0$ , for  $t \in [-1, 0]$ , and  $g(t) \geq 0$ , for  $t \in [0, 1]$ . Since  $g(0) = 0$ , for all  $g \in \mathcal{P}$ , it is not possible to write  $\iota \in \mathcal{I}$  as a linear combination of functions in  $\mathcal{I} \cap \mathcal{A}^+$ . To see that  $\mathcal{I}$  is hereditary, suppose that  $0 \leq h \leq f$ , where  $f \in \mathcal{I} \cap \mathcal{A}^+$  and  $h \in \mathcal{A}$ . Then  $f = g \cdot \iota$ , for some  $g \in \mathcal{P}$ . Hence  $|h(t)/t| \leq |g(t)|$ , for all  $t \neq 0$ , from which we see that  $h \in \mathcal{I}$ .

Let us also note that (algebraic) two-sided ideals need not be self-adjoint. Consider the commutative  $C^*$ -algebra  $\mathcal{A} = C(\mathbb{D})$ , where  $\mathbb{D}$  is the closed unit disk in the complex plane, the function  $f(z) = z$ ,  $z \in \mathbb{D}$ , and the (two-sided) ideal  $\mathcal{I} = \mathcal{A}f$  in  $\mathcal{A}$ . Then  $f$  belongs to  $\mathcal{I}$ , but  $f^* = \bar{f}$  does not.

For a  $C^*$ -algebra  $\mathcal{A}$  let  $\tilde{\mathcal{A}}$  denote the unitization of  $\mathcal{A}$  when it is nonunital, or  $\mathcal{A}$  itself when it already is unital.

**Lemma 2.3.** *Let  $\mathcal{A}$  be a  $C^*$ -algebra, and let  $C$  be a subcone of  $\mathcal{A}^+$  satisfying the following:*

- (i) *If  $a \in C$  and  $x \in \tilde{\mathcal{A}}$ , then  $x^*ax \in C$ .*
- (ii)  *$C$  is hereditary: If  $0 \leq a \leq b$ ,  $a \in \mathcal{A}$ , and  $b \in C$ , then  $a \in C$ .*

*Let  $\mathcal{I}$  be the linear span of  $C$ . Then  $\mathcal{I}$  is a hereditary ideal in  $\mathcal{A}$  and  $\mathcal{I} \cap \mathcal{A}^+ = C$ . If  $C$  is symmetric (in the sense that  $x^*x \in C$  implies  $xx^* \in C$ , for all  $x \in \mathcal{A}$ ), then so is  $\mathcal{I}$ .*

*Conversely, if  $\mathcal{I}$  is a hereditary ideal in  $\mathcal{A}$  and if  $C = \mathcal{I} \cap \mathcal{A}^+$ , then  $C$  is a subcone of  $\mathcal{A}^+$  satisfying (i) and (ii) above. The span,  $\mathcal{I}_0$ , of  $C$  is a hereditary subideal of  $\mathcal{I}$ ; and  $\mathcal{I}_0 = \mathcal{I}$  if and only if  $\mathcal{I}$  is generated as a hereditary ideal by its positive elements.*

*Proof.* It follows from (i) that if  $x \in \mathcal{A}$  and  $a \in C$ , then

$$\begin{aligned} xa + ax^* &= (x + 1)a(x + 1)^* - xax^* - a, \\ i(xa - ax^*) &= (x - i)a(x - i)^* - xax^* - a \end{aligned}$$

belong to  $\mathcal{I}$ , whence  $xa$  and  $ax^*$  belong to  $\mathcal{I}$ . This shows that  $\mathcal{I}$  is an ideal in  $\mathcal{A}$ . Clearly,  $\mathcal{I}$  is self-adjoint.

A subcone  $C$  of  $\mathcal{A}^+$  satisfies  $\text{span}(C) \cap \mathcal{A}^+ = C$  if and only if whenever  $a, b \in C$  are such that  $a \leq b$ , then  $b - a \in C$ . Hereditary cones clearly have this property, so  $\mathcal{I} \cap \mathcal{A}^+ = C$ . This also shows that  $\mathcal{I}$  is hereditary (because  $C$  is hereditary).

It is clear that  $C = \mathcal{I} \cap \mathcal{A}^+$  has the stated properties if  $\mathcal{I}$  is a hereditary ideal of  $\mathcal{A}$ ; and  $\mathcal{I}_0$  is a hereditary ideal of  $\mathcal{A}$  by the first part of the lemma. It is contained in  $\mathcal{I}$  and contains by definition all positive elements of  $\mathcal{I}$ . That proves the last claim of the lemma. □

**Corollary 2.4.** *Let  $\mathcal{A}$  be a  $C^*$ -algebra, and let  $M$  be a nonempty subset of  $\mathcal{A}^+$ . Let  $C$  be the set of all elements  $a \in \mathcal{A}^+$  for which there exist  $n \geq 1$ ,  $e_1, e_2, \dots, e_n \in M$ , and  $x_1, x_2, \dots, x_n \in \tilde{\mathcal{A}}$  such that  $a \leq \sum_{j=1}^n x_j^* e_j x_j$ . Then*

- (i)  *$C$  is a subcone of  $\mathcal{A}^+$  satisfying (i) and (ii) of Lemma 2.3,*
- (ii)  *$\mathcal{I}_{\mathcal{A}}(M) = \text{span}(C)$ ,*
- (iii)  *$\mathcal{I}_{\mathcal{A}}(M) \cap \mathcal{A}^+ = C$ .*

*Proof.* It is clear that (i) holds, so Lemma 2.3 implies that  $\mathcal{I} := \text{span}(C)$  is a hereditary ideal in  $\mathcal{A}$  satisfying  $\mathcal{I} \cap \mathcal{A}^+ = C$ . As  $M \subseteq C \subseteq \mathcal{I}$ , we conclude that  $\mathcal{I}_{\mathcal{A}}(M) \subseteq \mathcal{I}$ . Conversely,  $C \subseteq \mathcal{I}_{\mathcal{A}}(M)$ , so  $\mathcal{I} \subseteq \mathcal{I}_{\mathcal{A}}(M)$ . □

**Corollary 2.5.** *Let  $\mathcal{A}$  be a  $C^*$ -algebra, let  $\alpha$  be an action of a group  $G$  on  $\mathcal{A}$ , and let  $M$  be a nonempty subset of  $\mathcal{A}^+$ . Then the following hold:*

- (i)  $\mathcal{I}_{\mathcal{A}}^G(M) = \mathcal{I}_{\mathcal{A}}(G.M)$ , where  $G.M = \{\alpha_t(e) : t \in G, e \in M\}$ .
- (ii) An element  $a \in \mathcal{A}^+$  belongs to  $\mathcal{I}_{\mathcal{A}}^G(M)$  if and only if there exist  $n \geq 1$ ,  $t_1, t_2, \dots, t_n \in G$ ,  $y_1, y_2, \dots, y_n \in \tilde{\mathcal{A}}$ , and  $e_1, e_2, \dots, e_n \in M$  such that  $a \leq \sum_{j=1}^n y_j^* \alpha_{t_j}(e_j) y_j$ .

*Proof.*

(i) As  $G.M \subseteq \mathcal{I}_{\mathcal{A}}^G(M)$ , we see that  $\mathcal{I}_{\mathcal{A}}(G.M) \subseteq \mathcal{I}_{\mathcal{A}}^G(M)$ . Conversely, as  $\mathcal{I}_{\mathcal{A}}(G.M)$  is  $G$ -invariant, it contains  $\mathcal{I}_{\mathcal{A}}^G(M)$ . Part (ii) follows from (i) and from Corollary 2.4. □

For each nonempty subset  $M$  of  $\mathcal{A}^+$  denote by  $\mathcal{J}_{\mathcal{A}}(M)$  the smallest symmetric hereditary ideal in  $\mathcal{A}$  containing  $M$ . If a group  $G$  acts on  $\mathcal{A}$ , then denote by  $\mathcal{J}_{\mathcal{A}}^G(M)$  the smallest symmetric hereditary  $G$ -invariant ideal in  $\mathcal{A}$  containing  $M$ . Since closed two-sided ideals in a  $C^*$ -algebra always are symmetric, we have

$$(2.1) \quad \mathcal{I}_{\mathcal{A}}(M) \subseteq \mathcal{J}_{\mathcal{A}}(M) \subseteq \overline{\mathcal{I}_{\mathcal{A}}}(M), \quad \mathcal{I}_{\mathcal{A}}^G(M) \subseteq \mathcal{J}_{\mathcal{A}}^G(M) \subseteq \overline{\mathcal{I}_{\mathcal{A}}^G}(M).$$

**Lemma 2.6.** *Let  $\mathcal{A}$  be a  $C^*$ -algebra, and let  $M$  be a nonempty subset of projections in  $\mathcal{A}$ . Then  $\mathcal{I}_{\mathcal{A}}(M) = \mathcal{J}_{\mathcal{A}}(M) = \text{Ped}(\mathcal{A}_0)$ , where  $\mathcal{A}_0 = \overline{\mathcal{I}_{\mathcal{A}}}(M)$ . If  $\mathcal{A}$  is equipped with an action of a group  $G$ , then  $\mathcal{I}_{\mathcal{A}}^G(M) = \mathcal{J}_{\mathcal{A}}^G(M) = \text{Ped}(\mathcal{A}_1)$ , where  $\mathcal{A}_1 = \overline{\mathcal{I}_{\mathcal{A}}^G}(M)$ .*

*Proof.* We have  $M \subseteq \text{Ped}(\mathcal{A}_0) \subseteq \mathcal{I}_{\mathcal{A}}(M) \subseteq \mathcal{J}_{\mathcal{A}}(M) \subseteq \mathcal{A}_0$  (the first inclusion holds because each projection in a  $C^*$ -algebra belong to its Pedersen ideal, and second inclusion holds because  $\mathcal{I}_{\mathcal{A}}(M)$  is a dense ideal in  $\mathcal{A}_0$ ). As  $\text{Ped}(\mathcal{A}_0)$  is a hereditary symmetric ideal, which contains  $M$  (cf. Example 2.1),  $\mathcal{J}_{\mathcal{A}}(M) \subseteq \text{Ped}(\mathcal{A}_0)$ , so  $\text{Ped}(\mathcal{A}_0) = \mathcal{I}_{\mathcal{A}}(M) = \mathcal{J}_{\mathcal{A}}(M)$ . The second part of the statement follows from the first part applied to  $G.M$  (instead of  $M$ ). □

**Lemma 2.7.** *Let  $\mathcal{A}$  be a  $C^*$ -algebra, and let  $M$  be a nonempty set of positive elements in  $\mathcal{A}$ . Then  $\mathcal{J}_{\mathcal{A}}(M)$  is the linear span of its positive elements. If  $\mathcal{A}$  is equipped with an action of a group  $G$ , then the same holds for  $\mathcal{J}_{\mathcal{A}}^G(M)$ .*

*Proof.* Set  $C = \mathcal{J}_{\mathcal{A}}^G(M) \cap \mathcal{A}^+$ , and let  $\mathcal{J}_0$  be the linear span of  $C$ . Then  $\mathcal{J}_0$  is a hereditary ideal in  $\mathcal{A}$  by Lemma 2.3. Since  $\mathcal{J}_{\mathcal{A}}^G(M)$  is symmetric and  $G$ -invariant, the same holds for  $C$ , and hence for  $\mathcal{J}_0$ . Moreover,  $M \subseteq C \subseteq \mathcal{J}_0 \subseteq \mathcal{J}_{\mathcal{A}}^G(M)$ . As  $\mathcal{J}_{\mathcal{A}}^G(M)$  is the smallest ideal with these properties,  $\mathcal{J}_0 = \mathcal{J}_{\mathcal{A}}(M)$ . The first claim is proved in a similar manner. □

**Definition 2.8.** Let  $\mathcal{A}$  be a  $C^*$ -algebra. Denote by  $T^+(\mathcal{A})$  the cone of traces on the positive cone of  $\mathcal{A}$ , i.e., the set of additive homogeneous maps  $\tau : \mathcal{A}^+ \rightarrow [0, \infty]$  satisfying  $\tau(x^*x) = \tau(xx^*)$ , for all  $x \in \mathcal{A}$ .

For each hereditary symmetric ideal  $\mathcal{I}$  in  $\mathcal{A}$ , let  $T(\mathcal{I}, \mathcal{A})$  denote the cone of linear traces on  $\mathcal{I}$ , i.e., the set of positive linear maps  $\tau : \mathcal{I} \rightarrow \mathbb{C}$  satisfying  $\tau(x^*x) = \tau(xx^*)$ , whenever  $x \in \mathcal{A}$  is such that  $x^*x$  (and hence  $xx^*$ ) belongs to  $\mathcal{I}$ . We refer to  $\mathcal{I}$  as the *domain* of  $\tau$ . If the domain of  $\tau$  is a dense ideal of  $\mathcal{A}$  (in which case it will contain the Pedersen ideal of  $\mathcal{A}$ ), then  $\tau$  is said to be a *densely defined trace* on  $\mathcal{A}$ .

The cone of traces, here denoted by  $T^+(\mathcal{A})$ , is in [8] denoted by  $T(\mathcal{A})$ . *Note that all traces by default are assumed to be positive.* The following easy fact will be used repeatedly.

**Lemma 2.9.** *Let  $\tau$  be a trace on a  $C^*$ -algebra  $\mathcal{A}$ . Then  $\tau(x^*ax) \leq \|x\|^2\tau(a)$ , for all  $a \in \mathcal{A}^+$  (in the domain of  $\tau$ ) and all  $x$  in the unitization of  $\mathcal{A}$ .*

The “ $\varepsilon$ -cut-down”  $(a - \varepsilon)_+$  of  $a \in \mathcal{A}^+$  appearing in part (i) of the lemma below is defined by applying the continuous positive function  $t \mapsto \max\{t - \varepsilon, 0\}$  to  $a$ .

A trace in  $T^+(\mathcal{A})$ , or a linear trace on  $\mathcal{A}$  defined on a given domain, is said to be *lower semicontinuous* if one of the equivalent conditions in the following lemma holds.

**Lemma 2.10.** *Let  $\mathcal{A}$  be a  $C^*$ -algebra. The following conditions are equivalent for each trace  $\tau$  in  $T^+(\mathcal{A})$  (or for each linear trace on  $\mathcal{A}$ ):*

- (i)  $\tau(a) = \sup_{\varepsilon > 0} \tau((a - \varepsilon)_+)$  for each  $a \in \mathcal{A}^+$  (in the domain of  $\tau$ );
- (ii) whenever  $\{a_n\}_{n=1}^\infty$  is an increasing sequence in  $\mathcal{A}^+$  (in the domain of  $\tau$ ) converging in norm to  $a \in \mathcal{A}^+$  (in the domain of  $\tau$ ), then  $\tau(a) = \lim_{n \rightarrow \infty} \tau(a_n)$ ;
- (iii) whenever  $\{a_n\}_{n=1}^\infty$  is a sequence in  $\mathcal{A}^+$  (in the domain of  $\tau$ ) converging in norm to  $a \in \mathcal{A}^+$  (in the domain of  $\tau$ ), then  $\tau(a) \leq \liminf_{n \rightarrow \infty} \tau(a_n)$ .

*Proof.*

(ii)  $\Rightarrow$  (i) To verify (i), one needs only show that  $\tau(a) = \lim_{n \rightarrow \infty} \tau((a - \varepsilon_n)_+)$  for all sequences  $\{\varepsilon_n\}$  decreasing to 0, but this is just a special case of (ii).

(iii)  $\Rightarrow$  (ii) If  $\{a_n\}_{n=1}^\infty$  is an increasing sequence of positive elements converging to  $a$ , then  $\tau(a_n) \leq \tau(a)$ , for all  $n$ , by positivity of  $\tau$ . If (iii) holds, then this entails that  $\tau(a) = \lim_{n \rightarrow \infty} \tau(a_n)$ .

(i)  $\Rightarrow$  (iii) Let  $\varepsilon > 0$  be given. Choose  $n_0 \geq 1$  such that  $\|a_n - a\| < \varepsilon$ , for all  $n \geq n_0$ . It then follows from [14, Lemma 2.2] that  $(a - \varepsilon)_+ = d_n^* a_n d_n$ , for some contractions  $d_n$  in  $\mathcal{A}$ , for all  $n \geq n_0$ . Hence  $\tau((a - \varepsilon)_+) \leq \tau(a_n)$ , by Lemma 2.9. This shows that  $\liminf_{n \rightarrow \infty} \tau(a_n) \geq \tau((a - \varepsilon)_+)$ . It follows that  $\liminf_{n \rightarrow \infty} \tau(a_n) \geq \sup_{\varepsilon > 0} \tau((a - \varepsilon)_+)$ , which proves that (i) implies (iii).  $\square$

**Theorem 2.11** (Pedersen [19, Corollary 3.2]). *The restriction of any densely defined trace on a  $C^*$ -algebra  $\mathcal{A}$  to the Pedersen ideal of  $\mathcal{A}$  is automatically lower semicontinuous.*

**Definition 2.12.** Denote by  $T_{\text{isc}}(\mathcal{A})$  the cone of linear traces on  $\mathcal{A}$  whose domain is the Pedersen ideal of  $\mathcal{A}$ . In other words,  $T_{\text{isc}}(\mathcal{A}) = T(\text{Ped}(\mathcal{A}), \mathcal{A})$ .

We can identify  $T_{\text{isc}}(\mathcal{A})$  with the set of densely defined lower semicontinuous traces on  $\mathcal{A}$  as follows: Each trace in  $T_{\text{isc}}(\mathcal{A})$  is clearly densely defined, and it is lower semicontinuous by Pedersen’s theorem. Conversely, if  $\tau$  is a lower semicontinuous densely defined trace, then its restriction  $\tau_0$  to the Pedersen ideal of  $\mathcal{A}$  belongs to  $T_{\text{isc}}(\mathcal{A})$ , and  $\tau$  is uniquely determined on its domain by  $\tau_0$  by Lemma 2.10(i), because  $(a - \varepsilon)_+ \in \text{Ped}(\mathcal{A})$  for all positive  $a \in \mathcal{A}$  and all  $\varepsilon > 0$ .

A trace in  $T_{\text{isc}}(\mathcal{A})$  can usually be extended to a lower semicontinuous trace on a larger domain than the Pedersen ideal, and such an extension is unique; see Proposition 2.13 and the subsequent discussion.

It follows from Theorem 2.11 and Lemma 2.6 that each linear trace on a  $C^*$ -algebra  $\mathcal{A}$  with domain  $\mathcal{J}_{\mathcal{A}}(M)$  or  $\mathcal{J}_{\mathcal{A}}^G(M)$  (when  $\mathcal{A}$  has a  $G$ -action) is lower semicontinuous whenever  $M$  is a subset of projections in  $\mathcal{A}$ .

Observe that  $T_{\text{isc}}(B(H)) = \{0\}$ , where  $B(H)$  is the bounded operators on a separable infinite-dimensional Hilbert space  $H$ , while  $T_{\text{isc}}(\mathcal{K}(H))$  and the cone of lower semicontinuous traces in  $T^+(B(H))$ , with both equal to the one-dimensional cone spanned by the canonical trace on  $B(H)$ , in the latter case viewed as a function  $B(H)^+ \rightarrow [0, \infty]$ . The Dixmier trace is an example of a singular trace on  $\mathcal{K}(H)$ . It belongs to  $T^+(B(H))$  and to  $T(\mathcal{I}, \mathcal{K}(H))$ , where  $\mathcal{I} \subset \mathcal{K}(H)$  is its domain, and it is 0 on the finite rank operators.

Consider a general (not necessarily densely defined) linear trace  $\tau$  on  $\mathcal{A}$  with domain  $\mathcal{I}$ . The closure,  $\overline{\mathcal{I}}$ , of  $\mathcal{I}$  is a closed two-sided ideal in  $\mathcal{A}$ , and hence, in particular, a  $C^*$ -algebra; and  $\tau$  is of course densely defined relatively to this  $C^*$ -algebra. We have the following inclusions:

$$\text{Ped}(\overline{\mathcal{I}}) \subseteq \mathcal{I} \subseteq \overline{\mathcal{I}}.$$

The restriction of  $\tau$  to  $\text{Ped}(\overline{\mathcal{I}})$  is lower semicontinuous by Theorem 2.11. If this restriction is 0, then  $\tau$  is said to be *singular*. Each trace  $\tau$  on  $\mathcal{A}$  with domain  $\mathcal{I}$  can in a unique way be written as the sum  $\tau = \tau_1 + \tau_2$  of a lower semicontinuous trace  $\tau_1$  and a singular trace  $\tau_2$ , both with domain  $\mathcal{I}$ . The lower semicontinuous part is obtained by restricting  $\tau$  to the Pedersen ideal (which is lower semicontinuous) and then extending to a lower semicontinuous trace  $\tau_1$  defined on  $\mathcal{I}$  as described in (2.3) and the subsequent comments.

One can smoothly and uniquely pass from a trace in  $T^+(\mathcal{A})$  to a linear trace defined on its natural (maximal) domain.

**Proposition 2.13.** *Let  $\mathcal{A}$  be a  $C^*$ -algebra, and let  $\tau' \in T^+(\mathcal{A})$ . Let  $C$  be the set of positive elements  $a \in \mathcal{A}$  with  $\tau'(a) < \infty$ , and let  $\mathcal{I}$  be the linear span of  $C$ . Then  $\mathcal{I}$  is a hereditary symmetric ideal in  $\mathcal{A}$ ,  $\mathcal{I} \cap \mathcal{A}^+ = C$ , and there is a unique linear trace  $\tau$  with domain  $\mathcal{I}$  that agrees with  $\tau'$  on  $C$ .*

*We can recover  $\tau'$  from  $\tau$  via the formula*

$$(2.2) \quad \tau'(a) = \begin{cases} \tau(a), & a \in C, \\ \infty, & a \in \mathcal{A}^+ \setminus C. \end{cases}$$

*If  $\tau'$  is lower semicontinuous, then so is  $\tau$ .*

*Proof.* Observe that the set  $C$  is a symmetric subcone of  $\mathcal{A}^+$  satisfying conditions (i) and (ii) of Lemma 2.3 (use Lemma 2.9 to see that Lemma 2.3(i) holds). It therefore follows from Lemma 2.3 that  $\mathcal{I}$  is a symmetric hereditary ideal in  $\mathcal{A}$  and that  $\mathcal{I}^+ = \mathcal{I} \cap \mathcal{A}^+ = C$ . By the additivity and homogeneity of  $\tau'$  its restriction to  $C$  extends (uniquely) to a linear map  $\tau: \mathcal{I} \rightarrow \mathbb{C}$ . If  $a \in \mathcal{I}$  is positive, then  $a \in C$ , so  $\tau(a) = \tau'(a) \geq 0$ , which shows that  $\tau$  is positive. Let  $x \in \mathcal{A}$  be such that  $x^*x \in \mathcal{I}$ . Then  $x^*x$  and  $xx^*$  are positive elements in  $\mathcal{I}$ , so both belong to  $C$ , whence  $\tau(x^*x) = \tau'(x^*x) = \tau'(xx^*) = \tau(xx^*)$ , so  $\tau$  is a trace on  $\mathcal{I}$ .

It is clear that (2.2) holds. If  $\tau'$  is lower semicontinuous, then so is its restriction to  $C$ , which shows that  $\tau$  is lower semicontinuous. □

Whenever we talk about a trace on a  $C^*$ -algebra  $\mathcal{A}$ , we shall mean a trace defined on the cone of positive elements of that  $C^*$ -algebra taking values in  $[0, \infty]$ , i.e., a trace in  $T^+(\mathcal{A})$ , and, at the same time, a linear trace on the domain defined in the proposition above, or some other domain to be specified in the context.

As a converse to the proposition above, consider a linear trace  $\tau$  defined on a hereditary symmetric ideal  $\mathcal{I}$  in  $\mathcal{A}$ . Then  $\tau'$  given by (2.2), with  $C = \mathcal{I} \cap \mathcal{A}^+$ ,

belongs to  $T^+(\mathcal{A})$ , and it agrees with  $\tau$  on  $C$ . If we apply Proposition 2.13 to  $\tau'$ , then we get back a new linear trace  $\tau_0$  defined on some symmetric hereditary ideal  $\mathcal{J}_0$  of  $\mathcal{A}$ , which contains the subideal  $\mathcal{I}_0$  of  $\mathcal{I}$  defined in Lemma 2.3 (but perhaps not  $\mathcal{I}$  itself), and  $\tau$  and  $\tau_0$  agree on  $\mathcal{I}_0$ .

However, this extension of  $\tau$  to a trace  $\tau'$  defined on the positive cone of  $\mathcal{A}$  is not unique, and  $\tau'$  need not be lower semicontinuous, even when  $\tau$  is lower semicontinuous. If  $\tau$  is lower semicontinuous, then the map  $\tau': \mathcal{A}^+ \rightarrow [0, \infty]$  defined by

$$(2.3) \quad \tau'(a) = \sup\{\tau(a_0) : 0 \leq a_0 \leq a, a_0 \in \mathcal{I}\}, \quad a \in \mathcal{A}^+,$$

is a lower semicontinuous trace in  $T^+(\mathcal{A})$ , and it is the unique one that extends  $\tau$ . In the sequel, when considering a lower semicontinuous trace, we may wish to view it either as a linear trace defined on its domain or as a trace defined on the positive cone, via (2.3).

There is a canonical way of extending a lower semicontinuous trace  $\tau$  defined on some hereditary symmetric ideal  $\mathcal{I}$  of  $\mathcal{A}$  to its maximal domain: First extend  $\tau$  to a lower semicontinuous trace  $\tau': \mathcal{A}^+ \rightarrow [0, \infty]$  as in (2.3), then take the linearization  $\bar{\tau}$  of  $\tau'$  defined in Proposition 2.13.

We quote the following well-known result for later reference; see, e.g., [20, Lemma 5.3] for a proof.

**Lemma 2.14.** *Let  $\mathcal{A}$  be a  $C^*$ -algebra equipped with an action of a group  $G$ , and let  $\tau$  be a  $G$ -invariant lower semicontinuous trace on  $\mathcal{A}$ . It follows that  $\tau \circ E$  is a lower semicontinuous trace on the (reduced) crossed product  $\mathcal{A} \rtimes G$  that extends  $\tau$ , where  $E: \mathcal{A} \rtimes G \rightarrow \mathcal{A}$  is the standard conditional expectation. If  $\tau$  is densely defined, then so is  $\tau \circ E$ .*

One should here view  $\tau$  and  $\tau \circ E$  as traces defined on the positive cones of  $\mathcal{A}$  and  $\mathcal{A} \rtimes G$ , respectively. For the claim that  $\tau \circ E$  is densely defined when  $\tau$  is, use the fact that  $\tau \circ E$  is finite on the positive cone of  $\text{Ped}(\mathcal{A})$ , and the hereditary ideal in  $\mathcal{A} \rtimes G$  generated by  $\text{Ped}(\mathcal{A})$  is dense in  $\mathcal{A} \rtimes G$ . It is a curious fact that an invariant densely defined trace  $\tau$  on  $\mathcal{A}$  need not in general extend to a trace on the crossed product  $\mathcal{A} \rtimes G$ ; in particular,  $\tau \circ E$  need not be a trace if  $\tau$  is not lower semicontinuous. See Example 4.4.

We end this section by considering when a  $C^*$ -algebra admits a nonzero densely defined trace. Blackadar and Cuntz proved in [2] that a stable *simple*  $C^*$ -algebra either contains a properly infinite projection or admits a nonzero dimension function (defined on its Pedersen ideal). In the latter case, assuming moreover that the  $C^*$ -algebra is exact, it admits a nonzero densely defined trace. (This step follows from the work of Blackadar and Handelman [3], Haagerup [10], and Kirchberg [12], as explained in the last part of the proof of the theorem below.) Also, it is well known that a unital exact  $C^*$ -algebra admits a tracial state if and only if no matrix algebra over it is properly infinite. A common feature of simple and of unital  $C^*$ -algebras is that their primitive ideal spaces are compact. Recall that the primitive ideal space,  $\text{Prim}(\mathcal{A})$ , of a  $C^*$ -algebra  $\mathcal{A}$  is compact if and only if for all upward directed families  $\{\mathcal{I}_\alpha\}$  of closed two-sided ideals in  $\mathcal{A}$  whose union is dense in  $\mathcal{A}$ , there is an  $\alpha$  such that  $\mathcal{A} = \mathcal{I}_\alpha$ .

**Theorem 2.15.** *Let  $\mathcal{A}$  be an exact  $C^*$ -algebra whose primitive ideal space is compact. Then  $\mathcal{A}$  admits a nonzero densely defined lower semicontinuous trace, i.e.,*

$T_{\text{isc}}(\mathcal{A}) \neq \{0\}$ , if and only if the stabilization of  $\mathcal{A}$  does not contain a full properly infinite projection.

*Proof.* The proof is most naturally phrased via dimension functions (as defined by Cuntz in [6]) and the Cuntz semigroup; see, e.g., [5].

Observe first that the cone of densely defined lower semicontinuous traces and the primitive ideal space are not changed by stabilizing the  $C^*$ -algebra, so we may assume that  $\mathcal{A}$  is stable.

The class of (closed two-sided) ideals of the form  $\overline{\mathcal{I}}_{\mathcal{A}}((e - \varepsilon)_+)$ , where  $e \in \mathcal{A}^+$  and  $\varepsilon > 0$ , is upward directed and its union is dense in  $\mathcal{A}$ . Hence  $\mathcal{A} = \overline{\mathcal{I}}_{\mathcal{A}}((e - \varepsilon_0)_+)$ , for some  $e \in \mathcal{A}^+$  and some  $\varepsilon_0 > 0$  by compactness of the primitive ideal space of  $\mathcal{A}$ . Since  $(e - \varepsilon)_+$  belongs to the Pedersen ideal and since  $\mathcal{I}_{\mathcal{A}}((e - \varepsilon)_+)$  is a dense ideal in  $\mathcal{A}$ , for all  $0 < \varepsilon \leq \varepsilon_0$ , it follows that  $\mathcal{I}_{\mathcal{A}}((e - \varepsilon)_+) = \text{Ped}(\mathcal{A})$ , for this  $e \in \mathcal{A}^+$  and for all  $0 < \varepsilon \leq \varepsilon_0$ . Let  $u_\varepsilon = \langle (e - \varepsilon)_+ \rangle$  be the corresponding element, the Cuntz semigroup  $\text{Cu}(\mathcal{A})$  of  $\mathcal{A}$ .

Fix  $0 < \varepsilon \leq \varepsilon_0$ . It follows from Corollary 2.4, and the fact that  $\langle \sum_{j=1}^n x_j^* e_j x_j \rangle \leq \sum_{j=1}^n \langle e_j \rangle$  in  $\text{Cu}(\mathcal{A})$ , for all positive  $e_j$  and all  $x_j$  in  $\tilde{\mathcal{A}}$ , that for each positive  $a$  in  $\text{Ped}(\mathcal{A})$  there exists  $k \geq 1$  such that  $\langle a \rangle \leq k u_\varepsilon$ . In other words,  $u_\varepsilon$  is an order unit for the subsemigroup  $\text{Cu}_0(\mathcal{A})$ , consisting of all classes  $\langle a \rangle$ , where  $a$  is a positive element in  $\text{Ped}(\mathcal{A})$ . In particular,  $u_{\varepsilon_0} \leq u_\varepsilon \leq k u_{\varepsilon_0}$ , for some integer  $k \geq 1$  (that depends on  $\varepsilon$ ).

Consider first the case that  $n u_{\varepsilon_0}$  is properly infinite, for some integer  $n \geq 1$ . Upon replacing  $e$  by an  $n$ -fold direct sum of  $e$  with itself (which is possible since  $\mathcal{A}$  is assumed to be stable), we may assume that  $u_{\varepsilon_0}$  itself is properly infinite, i.e., that  $k u_{\varepsilon_0} \leq u_{\varepsilon_0}$ , for all integers  $k \geq 1$ . By the discussion in the previous paragraph we can then conclude that  $u_\varepsilon$  is properly infinite and that  $x \leq u_\varepsilon$ , for all  $x \in \text{Cu}_0(\mathcal{A})$  and for all  $0 < \varepsilon \leq \varepsilon_0$ .

We can now follow the argument of [17, Proposition 2.7], which uses the notion of *scaling elements* introduced by Blackadar and Cuntz [2], to construct a full properly infinite projection  $p \in \mathcal{A}$ : Fix  $0 \leq \varepsilon < \varepsilon_0$ . Then  $(e - \varepsilon)_+$  is properly infinite, so by [13, Proposition 3.3] there exist positive elements  $b_1, b_2$  in the hereditary sub- $C^*$ -algebra of  $\mathcal{A}$  generated by  $(e - \varepsilon_0)_+$  such that  $b_1 \perp b_1$  and  $(e - \varepsilon_0)_+ \precsim b_j$ , for  $j = 1, 2$ . In particular,  $b_1, b_2 \in \text{Ped}(\mathcal{A})$ . As explained in [17, Remark 2.5], there exists an  $x \in \mathcal{A}$  such that  $x^* x (e - \varepsilon_0)_+ = (e - \varepsilon_0)_+$  and  $x x^*$  belongs to the hereditary sub- $C^*$ -algebra of  $\mathcal{A}$  generated by  $b_1$ . This shows that  $x$  is a scaling element (see [17, Remark 2.4]) satisfying  $x^* x b_2 = b_2$  and  $x x^* b_2 = 0$ . By [2] (see also [17, Remark 2.4]) we get a projection  $p \in \mathcal{A}$  satisfying  $b_2 p = b_2$ . As  $u_{\varepsilon_0} \leq \langle b_2 \rangle \leq \langle p \rangle \leq u_{\varepsilon_0}$ , we conclude that  $\langle p \rangle$  is a properly infinite order unit of  $\text{Cu}_0(\mathcal{A})$ , whence  $p$  is a full properly infinite projection in  $\mathcal{A}$ .

Suppose now that there is no integer  $n \geq 1$  such that  $n u_{\varepsilon_0}$  is properly infinite. We proceed to show that  $T_{\text{isc}}(\mathcal{A}) \neq \{0\}$  in this case. As shown above,  $n u_\varepsilon$  is not properly infinite, for any  $n \geq 1$  and for any  $0 < \varepsilon \leq \varepsilon_0$ . Fix  $0 < \varepsilon_1 < \varepsilon_0$ , and observe that  $n u_{\varepsilon_1} \leq m u_{\varepsilon_1}$  implies that  $n \leq m$ , for all integers  $n, m \geq 0$  (since no multiple of  $u_{\varepsilon_1}$  is properly infinite). The map  $f_0: \mathbb{N}_0 u_{\varepsilon_1} \rightarrow \mathbb{R}^+$ , given by  $f_0(n u_{\varepsilon_1}) = n$ , for  $n \geq 0$ , is therefore a positive additive map on the subsemigroup  $\mathbb{N}_0 u_{\varepsilon_1}$  of  $\text{Cu}_0(\mathcal{A})$  (where  $\mathbb{N}_0 u_{\varepsilon_1}$  is equipped with the relative order arising from  $\text{Cu}_0(\mathcal{A})$ ). By [4, Corollary 2.7] we can extend  $f_0$  to a positive additive map (state)  $f: \text{Cu}_0(\mathcal{A}) \rightarrow \mathbb{R}^+$  (since  $u_{\varepsilon_1}$  is an order unit for  $\text{Cu}_0(\mathcal{A})$ ). Let  $d: \text{Ped}(\mathcal{A})^+ \rightarrow \mathbb{R}^+$  be the associated *dimension function* given by  $d(a) = f(\langle a \rangle)$ , and let  $\bar{d}: \text{Cu}_0(\mathcal{A}) \rightarrow \mathbb{R}^+$  be the corresponding

lower semicontinuous dimension function given by  $\bar{d}(a) = \lim_{\varepsilon > 0} d((a - \varepsilon)_+)$ , for  $a \in \text{Ped}(\mathcal{A})^+$ ; cf. [22, Proposition 4.1]. Then

$$d((e - \varepsilon_0)_+) \leq \bar{d}((e - \varepsilon_1)_+) \leq d((e - \varepsilon_1)_+),$$

and  $d((e - \varepsilon_0)_+) > 0$  because  $d$  is nonzero and  $u_{\varepsilon_0} = \langle (e - \varepsilon_0)_+ \rangle$  is an order unit for  $\text{Cu}_0(\mathcal{A})$ . This shows that  $\bar{d}$  is nonzero.

It follows from Blackadar and Handelman [3, Theorem II.2.2] that the lower semicontinuous dimension function  $\bar{d}$  (called the rank function in [3]) lifts to a lower semicontinuous 2-quasitrace  $\tau$  defined on the “pre- $C^*$ -algebra”  $\text{Ped}(\mathcal{A})$ , i.e.,  $\bar{d} = d_\tau$ , where  $d_\tau(a) = \lim_{n \rightarrow \infty} \tau(a^{1/n})$ , for all positive elements  $a \in \text{Ped}(\mathcal{A})$ . Finally, by Kirchberg’s extension [12] to the nonunital case of Haagerup’s theorem [10], that any 2-quasitrace on an exact  $C^*$ -algebra is a trace,  $\tau$  is a nonzero lower semicontinuous densely defined trace on  $\mathcal{A}$ .  $\square$

It remains unresolved when a  $C^*$ -algebra with noncompact primitive ideal space admits a nonzero densely defined lower semicontinuous trace. Clearly,  $T_{\text{isc}}(\mathcal{A})$  is nonzero for all commutative  $C^*$ -algebras  $\mathcal{A}$ , while the primitive ideal space of a commutative  $C^*$ -algebra is compact only when it is unital. On the other hand, absence of full properly infinite projections is not sufficient to guarantee existence of nonzero lower semicontinuous traces. Take, for example, the suspension (or the cone over) any purely infinite  $C^*$ -algebra; see [13, Proposition 5.1]. In [15, Section 4] it was shown that any infinite group  $G$  admits a (free) action on the locally compact noncompact Cantor set  $\mathbf{K}^*$  with no nonzero invariant Radon measures. Accordingly,  $C_0(\mathbf{K}^*) \rtimes G$  has no nonzero densely defined lower semicontinuous trace, although  $C_0(\mathbf{K}^*) \rtimes G$  admits an approximate unit consisting of projections, and, if  $G$  is supramenable, e.g., if  $G = \mathbb{Z}$ , then no projection in the (stabilization of)  $C_0(\mathbf{K}^*) \rtimes G$  is properly infinite.

The latter example is covered by the proposition below. When  $p$  and  $q$  are projections in a  $C^*$ -algebra  $\mathcal{A}$  and  $n \geq 1$  is an integer, then denote by  $p \otimes 1_n$  the  $n$ -fold direct sum of  $p$  with itself, and write  $p \ll q$  if  $p \otimes 1_n \lesssim q$ , for all  $n \geq 1$ .

**Proposition 2.16.** *Let  $\mathcal{A}$  be a  $C^*$ -algebra admitting an approximate unit consisting of projections. Suppose that for each projection  $p \in \mathcal{A}$  there exists a projection  $q$  in  $\mathcal{A}$  with  $p \ll q$ . Then  $T_{\text{isc}}(\mathcal{A}) = \{0\}$ .*

*Proof.* Suppose that  $\tau \in T_{\text{isc}}(\mathcal{A})$ , let  $p$  be a projection in  $\mathcal{A}$ , and let  $q \in \mathcal{A}$  be another projection such that  $p \ll q$ . Since  $p$  and  $q$  belong to the Pedersen ideal of  $\mathcal{A}$ , and hence to the domain of  $\tau$ , we find that  $\tau(q) < \infty$ , which entails that  $\tau(p) = 0$ . As  $\mathcal{A}$  has an approximate unit consisting of projection, this implies that  $\tau = 0$ .  $\square$

Here is an elementary example of an exact stably finite<sup>1</sup>  $C^*$ -algebra satisfying the conditions of Proposition 2.16, and which accordingly admits no nonzero lower semicontinuous densely defined trace: Let  $\mathcal{A}$  be the inductive limit of the sequence  $\mathcal{A}_1 \rightarrow \mathcal{A}_2 \rightarrow \mathcal{A}_3 \rightarrow \cdots$ , where  $\mathcal{A}_1 = \mathcal{K}$ , the  $C^*$ -algebra of compact operators on a separable Hilbert space, where  $\mathcal{A}_{n+1} = \tilde{\mathcal{A}}_n \otimes \mathcal{K}$ , for  $n \geq 1$ , and where the inclusion  $\mathcal{A}_n \rightarrow \mathcal{A}_{n+1}$  is given by  $a \mapsto a \otimes e \in \mathcal{A}_n \otimes \mathcal{K} \subset \mathcal{A}_{n+1}$ , for some fixed one-dimensional projection  $e \in \mathcal{K}$ .

<sup>1</sup>A  $C^*$ -algebra  $\mathcal{A}$  is said to be *stably finite* if its stabilization  $\mathcal{A} \otimes \mathcal{K}$  contains no infinite projections. This definition is meaningful when the  $C^*$ -algebra has an approximate unit consisting of projections.

3. INVARIANT UNBOUNDED TRACES ON  $C^*$ -ALGEBRAS

We shall here use Monod's characterization of groups with the fixed-point property for cones to say something about the times when a (typically nonunital)  $C^*$ -algebra  $\mathcal{A}$  with an action of a group  $G$  admits an invariant trace. We are mostly interested in the existence of a (nonzero) invariant densely defined lower semicontinuous trace, i.e., an invariant nonzero trace in the cone  $T_{\text{isc}}(\mathcal{A})$  defined in Section 2. But we shall also address the existence of more general traces (including singular traces and not densely defined traces).

Recall from Definition 2.8 that  $T(\mathcal{I}, \mathcal{A})$  is the cone of positive traces on  $\mathcal{A}$  with domain  $\mathcal{I}$  whenever  $\mathcal{I}$  is a hereditary symmetric ideal in  $\mathcal{A}$ . The cone  $T(\mathcal{I}, \mathcal{A})$  is embedded in the complex vector space  $\mathcal{L}(\mathcal{I})$  of all linear functionals on  $\mathcal{I}$  equipped with the locally convex weak topology induced by  $\mathcal{I}$ . The dual space  $\mathcal{L}(\mathcal{I})^*$  of  $\mathcal{L}(\mathcal{I})$  is naturally isomorphic to  $\mathcal{I}$  (cf. [24, 3.14], and as remarked in [16]); i.e.,  $\mathcal{L}(\mathcal{I})^* = \{\varphi_a : a \in \mathcal{I}\}$ , where  $\varphi_a$  denotes the functional  $\varphi_a(\rho) = \rho(a)$ , for  $\rho \in \mathcal{L}(\mathcal{I})$  and  $a \in \mathcal{I}$ . The dual space  $\mathcal{L}(\mathcal{I})^*$  is equipped with the preordering given by  $T^+(\mathcal{I}, \mathcal{A})$ , whereby an element  $\varphi \in \mathcal{L}(\mathcal{I})^*$  is positive if  $\varphi(\tau) \geq 0$ , for all  $\tau \in T(\mathcal{I}, \mathcal{A})$ . Observe that  $\varphi \geq 0$  and  $-\varphi \geq 0$  if and only if  $\varphi(\tau) = 0$ , for all  $\tau \in T(\mathcal{I}, \mathcal{A})$ . The map  $a \mapsto \varphi_a$  is a positive isomorphism, but not necessarily an order embedding, since  $\varphi_a \geq 0$  does not necessarily imply that  $a \geq 0$ .

Monod considers real vector spaces in his paper [16], while our vector spaces are complex by the nature of  $C^*$ -algebras. To translate some properties from Monod's paper to our language, we shall occasionally consider the real vector space of all self-adjoint functionals  $\varphi$  in  $\mathcal{L}(\mathcal{I})^*$ , and we note that  $\varphi_a$  is self-adjoint if and only if  $a \in \mathcal{I}$  is self-adjoint.

The cone  $T(\mathcal{I}, \mathcal{A})$  is said to be *proper* if  $T(\mathcal{I}, \mathcal{A}) \cap -T(\mathcal{I}, \mathcal{A}) = \{0\}$ , or, equivalently, if 0 is the only trace in  $T(\mathcal{I}, \mathcal{A})$  that vanishes on  $\mathcal{I} \cap \mathcal{A}^+$ . This will hold if  $\mathcal{I}$  is the span of its positive elements. Most ideals considered in this paper have this property, including the Pedersen ideal  $\text{Ped}(\mathcal{A})$ , or any of the ideals  $\mathcal{I}_{\mathcal{A}}(M)$ ,  $\mathcal{I}_{\mathcal{A}}^G(M)$ ,  $\mathcal{J}_{\mathcal{A}}(M)$ , or  $\mathcal{J}_{\mathcal{A}}^G(M)$  when  $M$  is any nonempty subset of  $\mathcal{A}^+$ ; cf. Example 2.1, Corollaries 2.4 and 2.5, and Lemma 2.7.

Recall also that  $T_{\text{isc}}(\mathcal{A}) = T(\text{Ped}(\mathcal{A}), \mathcal{A})$ . We allow for the possibility that the cones  $T(\mathcal{I}, \mathcal{A})$  and  $T_{\text{isc}}(\mathcal{A})$  are trivial, that is, equal to  $\{0\}$ , unless otherwise stated.

**Proposition 3.1.** *For each  $C^*$ -algebra  $\mathcal{A}$  and for each hereditary symmetric ideal  $\mathcal{I}$  in  $\mathcal{A}$ , the cone  $T(\mathcal{I}, \mathcal{A})$  is weakly complete. In particular,  $T_{\text{isc}}(\mathcal{A})$  is weakly complete.*

*Proof.* We must show that each weak Cauchy net in  $T(\mathcal{I}, \mathcal{A})$  is weakly convergent; i.e., if  $(\tau_i)_i$  is a net in  $T(\mathcal{I}, \mathcal{A})$  such that  $(\varphi(\tau_i))_i$  is Cauchy in  $\mathbb{C}$ , for all  $\varphi \in \mathcal{L}(\mathcal{I})^*$ , then the net converges weakly in  $T(\mathcal{I}, \mathcal{A})$ . Since  $\varphi_a(\tau_i) = \tau_i(a)$ , for all  $a \in \mathcal{I}$ , being weakly Cauchy implies that  $(\tau_i(a))_i$  is Cauchy, and hence convergent in  $\mathbb{C}$ , for all  $a \in \mathcal{I}$ . Set  $\tau(a) = \lim_i \tau_i(a)$ , for all  $a \in \mathcal{I}$ . It is easy to check that  $\tau: \mathcal{I} \rightarrow \mathbb{C}$  is in fact a trace, so it belongs to  $T(\mathcal{I}, \mathcal{A})$ , and since  $\varphi_a(\tau_i) \rightarrow \varphi_a(\tau)$ , for all  $a \in \mathcal{I}$ ,  $\tau$  is the weak limit of the net  $(\tau_i)_i$ , as desired.  $\square$

Consider an action  $\alpha$  of a (discrete) group  $G$  on  $\mathcal{A}$ . If  $\mathcal{I}$  is a  $G$ -invariant hereditary symmetric ideal in  $\mathcal{A}$ , then  $G$  induces an action of the cone  $T(\mathcal{I}, \mathcal{A})$  by  $t \cdot \tau = \tau \circ \alpha_t^{-1}$ , for  $t \in G$  and  $\tau \in T(\mathcal{I}, \mathcal{A})$ . It is clear that this action of  $G$  on  $T(\mathcal{I}, \mathcal{A})$  is continuous. Each automorphism of  $\mathcal{A}$  leaves the Pedersen ideal invariant, so each group action on  $\mathcal{A}$  induces an action on the cone  $T_{\text{isc}}(\mathcal{A})$ .

The action of  $G$  on  $T(\mathcal{I}, \mathcal{A})$  is in [16] said to be of cobounded type if there exists a positive functional  $\varphi$  in  $\mathcal{L}(\mathcal{I})^*$  which  $G$ -dominates<sup>2</sup> any other self-adjoint functional in  $\mathcal{L}(\mathcal{I})^*$ . This condition is automatically satisfied when  $\mathcal{I} = \mathcal{J}_{\mathcal{A}}^G(e)$ , for some positive element  $e \in \mathcal{A}^+$  (see Corollary 3.4), but not always when  $\mathcal{I}$  is the Pedersen ideal of  $\mathcal{A}$ . However, in the latter case we can reformulate the coboundedness condition into more familiar statements for  $C^*$ -algebras.

A positive element  $e \in \mathcal{I}$  is said to  $G$ -dominate a self-adjoint element  $a \in \mathcal{I}$  if there are group elements  $t_1, \dots, t_n$  such that  $a \leq \sum_{j=1}^n \alpha_{t_j}(e)$ , and  $e$  is said to *tracially  $G$ -dominate*  $a$  if there are group elements  $t_1, \dots, t_n$  such that  $\tau(a) \leq \sum_{j=1}^n \tau(\alpha_{t_j}(e))$ , for all  $\tau \in T(\mathcal{I}, \mathcal{A})$ . The latter holds if and only if  $\varphi_a \leq \sum_{j=1}^n t_j^{-1} \cdot \varphi_e$ —in other words, if  $\varphi_e$   $G$ -dominates  $\varphi_a$ . We can summarize these remarks as follows.

**Lemma 3.2.** *Let  $\mathcal{A}$  be a  $C^*$ -algebra equipped with an action of a group  $G$ , and let  $\mathcal{I}$  be an invariant hereditary symmetric ideal in  $\mathcal{A}$ . The induced action of  $G$  on the cone  $T(\mathcal{I}, \mathcal{A})$  is of cobounded type if and only if there is a positive element  $e \in \mathcal{I}$  which tracially  $G$ -dominates each self-adjoint element  $a \in \mathcal{I}$ .*

**Lemma 3.3.** *Let  $\mathcal{I}$  be a  $G$ -invariant hereditary symmetric ideal in a  $C^*$ -algebra  $\mathcal{A}$ , and let  $e$  be a positive element in  $\mathcal{I}$ . Then the functional  $\varphi_a$  is  $G$ -dominated by  $\varphi_e$ , for each self-adjoint element  $a \in \mathcal{J}_{\mathcal{A}}^G(e)$ .*

*Proof.* Let  $C$  be the set of positive elements  $a \in \mathcal{I}$  such that  $\varphi_a$  is  $G$ -dominated by  $e$ . We claim that  $C$  is a  $G$ -invariant symmetric cone in  $\mathcal{A}^+$ , which satisfies (i) and (ii) of Lemma 2.3. Since  $e$  clearly belongs to  $C$ , it will then follow from Lemma 2.3 that  $\mathcal{J}_{\mathcal{A}}^G(e) \cap \mathcal{A}^+ \subseteq C$ , and this will prove the lemma.

The set of positive  $\varphi \in \mathcal{L}(\mathcal{I})^*$  that are  $G$ -dominated by  $\varphi_e$  is a  $G$ -invariant hereditary cone in the positive cone of  $\mathcal{L}(\mathcal{I})^*$ . As the map  $a \mapsto \varphi_a$  is linear and order preserving and satisfies  $\varphi_{\alpha_t(a)} = t \cdot \varphi_a$ , for  $a \in \mathcal{I}^+$  and  $t \in G$ , we conclude that  $C$  is a hereditary  $G$ -invariant cone in  $\mathcal{A}^+$ . For each  $x \in \mathcal{A}$  for which  $x^*x$  (and hence  $xx^*$ ) belong to  $\mathcal{I}$ , we have  $\varphi_{x^*x} \leq \varphi_{xx^*} \leq \varphi_{x^*x}$ , which implies that  $C$  is symmetric. It remains to show that  $x^*ax$  belongs to  $C$  when  $a$  belongs to  $C$  and  $x$  belongs to  $\tilde{\mathcal{A}}$ . To see this, recall from Lemma 2.9 that  $\tau(x^*ax) \leq \|x\|^2 \tau(a)$ , so  $\varphi_{x^*ax} \leq \|x\|^2 \varphi_a$ , and the latter is  $G$ -dominated by  $\varphi_e$  since  $a \in C$  (and since  $C$  is a cone).  $\square$

The corollary below follows immediately from Lemma 3.3.

**Corollary 3.4.** *The action of a group  $G$  on the cone  $T(\mathcal{J}_{\mathcal{A}}^G(e), \mathcal{A})$  is of cobounded type whenever  $\mathcal{A}$  is a  $C^*$ -algebra with an action of  $G$  and  $e$  is a positive element in  $\mathcal{A}$ .*

The action of a group on the Pedersen ideal of a  $C^*$ -algebra is not always of cobounded type, as illustrated in the proposition below, which covers the case of commutative  $C^*$ -algebras, and which paraphrases and expands a remark in [16, page 71]. We remind the reader that the action of a group  $G$  on a locally compact Hausdorff space is *co-compact* if  $X = G \cdot K$ , for some compact subset  $K$  of  $X$ .

<sup>2</sup>If  $\varphi$  and  $\psi$  are self-adjoint functionals in  $\mathcal{L}(\mathcal{I})^*$ , then  $\psi$  is  $G$ -dominated by  $\varphi$  if  $\psi \leq \sum_{j=1}^n t_j \cdot \varphi$ , for some  $n \geq 1$  and some  $t_1, t_2, \dots, t_n \in G$ .

**Proposition 3.5.** *Let  $X$  be a locally compact Hausdorff space equipped with a continuous action of a group  $G$ . Then the following conditions are equivalent:*

- (i) *The action of  $G$  on  $X$  is co-compact.*
- (ii)  *$X$  is compact in the (non-Hausdorff) topology on  $X$  consisting of the  $G$ -invariant open subsets of  $X$ .*
- (iii) *The action of  $G$  on the cone of Radon measures on  $X$  equipped with the vague topology is of cobounded type.*
- (iv) *The action of  $G$  on the cone  $T_{\text{isc}}(C_0(X))$  is of cobounded type.*

*Proof.*

(i)  $\Rightarrow$  (ii) Let  $K$  be a compact subset of  $X$  witnessing co-compactness of the action. Let  $\{U_i\}_{i \in I}$  be a collection of invariant open sets that covers  $X$ . Select a finite subset  $F \subseteq I$  such that  $\{U_i\}_{i \in F}$  covers  $K$ . Then  $\bigcup_{i \in F} U_i = X$ , being a  $G$ -invariant set that contains  $K$ .

(ii)  $\Rightarrow$  (i) Let  $\{U_i\}_{i \in I}$  be the collection of all open precompact subsets of  $X$ . Then  $X = \bigcup_{i \in I} U_i$  because  $X$  is locally compact. For each  $i \in I$  set  $V_i = \bigcup_{t \in G} t.U_i$ . The families  $\{U_i\}_{i \in I}$  and  $\{V_i\}_{i \in I}$  are both upward directed (both are closed under forming finite unions). It follows by compactness of  $X$  in the topology of invariant open sets that  $X = V_i$ , for some  $i \in I$ . Hence  $X = \bigcup_{t \in G} t.K$  when  $K$  is the (compact) closure of  $U_i$ .

(i)  $\Rightarrow$  (iv) The cone  $T_{\text{isc}}(C_0(X))$  is embedded in the vector space  $\mathcal{L}(C_c(X))$  equipped with the weak topology from  $C_c(X)$ , and the dual space,  $\mathcal{L}(C_c(X))^*$ , is equal to  $C_c(X)$ . By co-compactness of the action we can find sets  $U \subseteq K \subseteq X$  such that  $K$  is compact,  $U$  is open, and  $G.U = X$ . Let  $f \in C_c(X)$  be such that  $1_K \leq f \leq 1$ . Then any real valued function  $g \in C_c(X)$  is  $G$ -dominated by  $f$ . Indeed, if  $F$  is a finite subset of  $G$  such that the support of  $g$  is contained in  $\bigcup_{t \in F} t.U$ , then  $g \leq \|g\|_\infty \sum_{t \in F} t.f$ .

(iv)  $\Rightarrow$  (i) Following the setup of the proof above, we can find a positive function  $f \in C_c(X)$  which  $G$ -dominates any other real valued function in  $C_c(X)$ . Let  $K$  be the support of  $f$ . Let  $x \in X$ , and choose a positive function  $g \in C_c(X)$  such that  $g(x) = 1$ . Then  $g \leq \sum_{t \in F} t.f$  for some finite subset  $F$  of  $G$ . Thus  $t.x \in K$ , for some  $t \in K$ . This shows that  $X = G.K$ .

(iii)  $\Leftrightarrow$  (iv) By Riesz' theorem there is a one-to-one correspondance between Radon measures on  $X$  and positive linear functionals (and hence traces) on  $C_c(X) = \text{Ped}(C_0(X))$ . □

We have previously, in Theorem 2.15, considered  $C^*$ -algebras  $\mathcal{A}$  whose primitive ideal space,  $\text{Prim}(\mathcal{A})$ , is compact, which happens if whenever  $\{\mathcal{I}_\alpha\}$  is an upward directed net of closed two-sided ideals in  $\mathcal{A}$  such that  $\bigcup_\alpha \mathcal{I}_\alpha$  is dense in  $\mathcal{A}$ , then  $\mathcal{A} = \mathcal{I}_\alpha$ , for some  $\alpha$ . Let us say that a  $C^*$ -algebra  $\mathcal{A}$  equipped with an action of a group  $G$  is  $G$ -compact if whenever  $\{\mathcal{I}_\alpha\}$  is an upward directed net of  $G$ -invariant closed two-sided ideals in  $\mathcal{A}$  such that  $\bigcup_\alpha \mathcal{I}_\alpha$  is dense in  $\mathcal{A}$ , then  $\mathcal{A} = \mathcal{I}_\alpha$ , for some  $\alpha$ . In the commutative case this is equivalent to condition (ii) of Proposition 3.5. In the noncommutative case we have the following.

**Proposition 3.6.** *Let  $\mathcal{A}$  be a  $C^*$ -algebra with an action of a group  $G$ , and suppose that  $\mathcal{A}$  is  $G$ -compact. Then the induced action of  $G$  on the cone  $T_{\text{isc}}(\mathcal{A})$  is of cobounded type.*

*Proof.* The family  $\{\overline{\mathcal{J}_\mathcal{A}^G}(e)\}$  of  $G$ -invariant ideals in  $\mathcal{A}$ , where  $e \in \text{Ped}(\mathcal{A})^+$ , is upward directed, and its union is dense in  $\mathcal{A}$ . Hence by the  $G$ -compactness assumption

there is a positive element  $e$  in  $\text{Ped}(\mathcal{A})$  such that the hereditary ideal  $\mathcal{J}_{\mathcal{A}}^G(e)$  is dense in  $\mathcal{A}$ . As  $e \in \text{Ped}(\mathcal{A})$ , this entails that  $\mathcal{J}_{\mathcal{A}}^G(e) = \text{Ped}(\mathcal{A})$ . We can therefore conclude from Lemma 3.3 that each self-adjoint  $\varphi \in \mathcal{L}(\text{Ped}(\mathcal{A}))^*$  is  $G$ -dominated by  $\varphi_e$ .  $\square$

Examples of  $G$ -compact  $C^*$ -algebras include any simple  $C^*$ -algebra and, more generally, any  $C^*$ -algebra with no nontrivial  $G$ -invariant closed two-sided ideals. Also, any  $C^*$ -algebra  $\mathcal{A}$  which contains a  $G$ -full projection, i.e., a projection  $p$  such that  $\overline{\mathcal{I}_{\mathcal{A}}^G(p)} = \mathcal{A}$ , is  $G$ -compact.

It is not hard to show that the conclusion of Proposition 3.6 still holds under the weaker assumption that  $\mathcal{A}/\mathcal{I}_0$  is  $G$ -compact, where  $\mathcal{I}_0$  is the closed two-sided  $G$ -invariant ideal

$$\mathcal{I}_0 = \bigcap_{\tau \in T_{\text{isc}}(\mathcal{A})} \{x \in \mathcal{A} : \tau(x^*x) = 0\}.$$

Following Monod [16], a subset  $M$  of  $T(\mathcal{I}, \mathcal{A})$  is *bounded* if for each open neighborhood  $0 \in \mathcal{U} \subseteq \mathcal{L}(\mathcal{I})$  there is an  $r > 0$  such that  $M \subseteq r\mathcal{U}$ .

**Lemma 3.7.** *Let  $\mathcal{A}$  be a  $C^*$ -algebra, and let  $\mathcal{I}$  be a hereditary symmetric ideal in  $\mathcal{A}$ . A subset  $M$  of  $T(\mathcal{I}, \mathcal{A})$  is bounded if and only if  $\{\tau(a) : \tau \in M\}$  is bounded for all  $a \in \mathcal{I}$ . In particular,  $M \subseteq T_{\text{isc}}(\mathcal{A})$  is bounded if and only if  $\{\tau(a) : \tau \in M\}$  is bounded for all  $a \in \text{Ped}(\mathcal{A})$ .*

*Proof.* For each self-adjoint element  $a \in \mathcal{I}$ , the set  $\mathcal{U}_a := \{\varphi \in \mathcal{L}(\mathcal{I}) : |\varphi(a)| < 1\}$  is an open neighborhood of 0. Hence if  $M$  is bounded, then  $M \subseteq r\mathcal{U}_a$  for some  $r > 0$ , which entails that  $|\tau(a)| < r$ , for all  $\tau \in M$ . Conversely, suppose that  $\{\tau(a) : \tau \in M\}$  is bounded, for all  $a \in \mathcal{I}$ , and let  $\mathcal{U}$  be an open neighborhood of 0. Then there are  $a_1, \dots, a_n$  in  $\mathcal{I}$  such that  $\mathcal{U}_{a_1, \dots, a_n} \subseteq \mathcal{U}$ , where

$$\mathcal{U}_{a_1, \dots, a_n} = \{\varphi \in \mathcal{L}(\mathcal{I}) : |\varphi(a_i)| < 1, \text{ for } i = 1, 2, \dots, n\}.$$

Set  $r_i = \sup\{|\tau(a_i)| : \tau \in M\}$ , and set  $r = 1 + \max_i r_i$ . Then  $|\tau(a_i)| < r$ , for all  $i$  and for all  $\tau \in M$ , whence  $M \subseteq r\mathcal{U}_{a_1, a_2, \dots, a_n} \subseteq r\mathcal{U}$ , as desired.  $\square$

A trace  $\tau \in T(\mathcal{I}, \mathcal{A})$  will be said to be *locally bounded* if it is bounded on the  $G$ -orbit of each  $a \in \mathcal{I}$ . Clearly, any bounded trace is  $G$ -bounded (regardless of the properties of the action). By Monod [16] the action of  $G$  on  $T(\mathcal{I}, \mathcal{A})$  is locally bounded if  $T(\mathcal{I}, \mathcal{A})$  contains a nonzero bounded orbit.

**Lemma 3.8.** *Let  $G$  be a group acting on a  $C^*$ -algebra  $\mathcal{A}$ , let  $\mathcal{I}$  be a hereditary symmetric invariant ideal in  $\mathcal{A}$ , and let  $e$  be a positive element in  $\mathcal{A}$ .*

- (i) *The induced action of  $G$  on  $T(\mathcal{I}, \mathcal{A})$  is locally bounded if and only if there is a nonzero locally bounded trace in  $T(\mathcal{I}, \mathcal{A})$ .*
- (ii) *A trace in  $T(\mathcal{J}_{\mathcal{A}}^G(e), \mathcal{A})$  is locally bounded if it is bounded on the orbit of  $e$ .*

*Proof.*

(i) This is an immediate reformulation of Lemma 3.7. To prove (ii), let  $\tau \in T(\mathcal{J}_{\mathcal{A}}^G(e), \mathcal{A})$  be bounded on the orbit of  $e$ , and let  $C$  be the set of positive elements  $a \in \mathcal{J}_{\mathcal{A}}^G(e)$  such that  $\tau$  is bounded on the orbit of  $a$ . Then  $C$  satisfies conditions (i) and (ii) of Lemma 2.3 (cf. Lemma 2.9),  $e \in C$ , and  $C$  is  $G$ -invariant, so  $C = \mathcal{J}_{\mathcal{A}}^G(e) \cap \mathcal{A}^+$  by Lemma 2.3.  $\square$

Groups with the fixed-point property for cones allow invariant lower semicontinuous densely defined traces on a  $C^*$ -algebra as follows.

**Theorem 3.9.** *Let  $\mathcal{A}$  be a  $C^*$ -algebra with  $T_{\text{isc}}(\mathcal{A}) \neq \{0\}$ , and let  $G$  be a group with the fixed-point property for cones, which acts on  $\mathcal{A}$ , making  $\mathcal{A}$   $G$ -compact. Then there is a nonzero invariant, necessarily lower semicontinuous, trace in  $T_{\text{isc}}(\mathcal{A})$  if and only if there is a nonzero locally bounded trace in  $T_{\text{isc}}(\mathcal{A})$ .*

*If this is the case, then  $T_{\text{isc}}(\mathcal{A} \rtimes G)$  is nonzero; i.e.,  $\mathcal{A} \rtimes G$  admits a nonzero lower semicontinuous densely defined trace.*

*Proof.* The cone  $T_{\text{isc}}(\mathcal{A})$  is weakly complete by Proposition 3.1, and the action of  $G$  on  $T_{\text{isc}}(\mathcal{A})$  is affine and continuous. By the assumption on  $G$  there is a nonzero invariant trace in  $T_{\text{isc}}(\mathcal{A})$  if the action of  $G$  on  $T_{\text{isc}}(\mathcal{A})$  is of cobounded type and locally bounded. The former holds by Proposition 3.6 since  $\mathcal{A}$  is assumed to be  $G$ -compact, and the latter holds by Lemma 3.8 if there is a nonzero locally bounded trace. Conversely, any invariant trace in  $T_{\text{isc}}(\mathcal{A})$  is trivially  $G$ -bounded. The last claim follows from Lemma 2.14.  $\square$

Consider the class of groups  $G$  for which Theorem 3.9 holds. This class contains the class of groups with the fixed-point property for cones, and it is contained in the class of supramenable groups; cf. the proposition below, which paraphrases [11, Theorem 1.1]. We do not know if Theorem 3.9 characterizes groups with the fixed-point property for cones, or if it characterizes the class of supramenable groups, or some intermediate class of groups.

**Proposition 3.10.** *The following conditions are equivalent for each group  $G$ :*

- (i)  $G$  is supramenable.
- (ii) Whenever  $G$  acts on a commutative  $C^*$ -algebra  $\mathcal{A}$  such that  $\mathcal{A}$  is  $G$ -compact, then there is a nonzero invariant trace in  $T_{\text{isc}}(\mathcal{A})$ .
- (iii) Whenever  $G$  acts on a commutative  $C^*$ -algebra  $\mathcal{A}$ , for each projection  $p \in \mathcal{A}$  there is an invariant lower semicontinuous trace  $\tau \in T^+(\mathcal{A})$  with  $\tau(p) = 1$ .
- (iv) Whenever  $G$  acts on a commutative  $C^*$ -algebra  $\mathcal{A}$ , then for each projection  $p \in \mathcal{A}$  there is a lower semicontinuous trace  $\tau \in T^+(\mathcal{A} \rtimes G)$  with  $\tau(p) = 1$ .

*Proof.*

(i)  $\Rightarrow$  (ii) Let  $X$  be the spectrum of  $\mathcal{A}$  such that  $\mathcal{A} = C_0(X)$ . If  $\mathcal{A}$  is  $G$ -compact, then the action of  $G$  on  $X$  is co-compact (see Proposition 3.5), so by [11, Theorem 1.1] there is a nonzero invariant Radon measure on  $X$  since  $G$  is supramenable. Integrating with respect to this measure gives a nonzero invariant trace in  $T_{\text{isc}}(\mathcal{A})$ .

(ii)  $\Rightarrow$  (iii) The  $C^*$ -algebra  $\mathcal{B} = \overline{\mathcal{I}_{\mathcal{A}}^G(p)}$  is  $G$ -compact, being generated by a projection, so (ii) implies that there is a nonzero invariant  $\tau \in T_{\text{isc}}(\mathcal{B})$ . We must show that  $0 < \tau(p) < \infty$ . The latter inequality holds because  $p \in \text{Ped}(\mathcal{B})$ . If  $\tau(p) = 0$ , then  $\tau(x) = 0$  for all  $x \in \mathcal{I}_{\mathcal{B}}^G(p)$ , and  $\mathcal{I}_{\mathcal{B}}^G(p) = \text{Ped}(\mathcal{B})$  (see Lemma 2.6), contradicting the condition that  $\tau \neq 0$ . Finally, we can extend  $\tau$  to a lower semicontinuous trace in  $T^+(\mathcal{A})$  by (2.3).

(iii)  $\Rightarrow$  (iv) This follows from Lemma 2.14.

(iv)  $\Rightarrow$  (i) If  $G$  is nonsupramenable, then, by [11, Theorem 1.1], it admits a minimal, free, purely infinite action on the locally compact noncompact Cantor set  $\mathbf{K}^*$ , making the crossed product  $C_0(\mathbf{K}^*) \rtimes G$  purely infinite (and simple). In particular, there is no nonzero lower semicontinuous trace on  $C_0(\mathbf{K}^*) \rtimes G$ .  $\square$

*Remark 3.11.* If  $\tau$  is a trace on a  $C^*$ -algebra  $\mathcal{B}$  such that  $\tau(p) = 1$ , for some projection  $p \in \mathcal{B}$ , then the restriction of  $\tau$  to the unital corner  $C^*$ -algebra  $p\mathcal{B}p$  is a tracial

state. Conversely, each tracial state  $\tau$  on  $p\mathcal{B}p$  extends to a lower semicontinuous trace in  $T^+(\mathcal{B})$  normalizing  $p$ .

To see this, consider the family  $\{\mathcal{H}_\alpha\}_\alpha$  of all  $\sigma$ -unital hereditary sub- $C^*$ -algebras of  $\overline{\mathcal{I}_\mathcal{B}(p)}$  containing  $p\mathcal{B}p$ , and observe that  $p\mathcal{B}p$  is a full hereditary sub- $C^*$ -algebra of  $\mathcal{H}_\alpha$ , for each  $\alpha$ . By Brown's theorem, the inclusion  $p\mathcal{B}p \rightarrow p\mathcal{B}p \otimes \mathcal{K}$ , where  $\mathcal{K}$  is the  $C^*$ -algebra of compact operators on a separable Hilbert space, extends to an inclusion  $\mathcal{H}_\alpha \rightarrow p\mathcal{B}p \otimes \mathcal{K}$ , for each  $\alpha$ . The tracial state  $\tau$  on  $p\mathcal{B}p$  extends (uniquely) to a lower semicontinuous trace on the positive cone of  $p\mathcal{B}p \otimes \mathcal{K}$ , and thus it restricts to a lower semicontinuous trace  $\tau_\alpha$  on the positive cone of  $\mathcal{H}_\alpha$ . The extension of  $\tau$  to  $\tau_\alpha$  on  $\mathcal{H}_\alpha$  is unique.

The family  $\{\mathcal{H}_\alpha\}_\alpha$  is upward directed with union  $\bigcup_\alpha \mathcal{H}_\alpha = \overline{\mathcal{I}_\mathcal{B}(p)}$ . To see the first claim, recall that a  $C^*$ -algebra is  $\sigma$ -unital precisely if it contains a strictly positive element. Let  $b_\alpha \in \mathcal{H}_\alpha$  be strictly positive. Then  $b_\alpha + b_\beta$  is a strictly positive element in  $\overline{(b_\alpha + b_\beta)\mathcal{B}(b_\alpha + b_\beta)}$ , and the latter is therefore a  $\sigma$ -unital sub- $C^*$ -algebra of  $\overline{\mathcal{I}_\mathcal{B}(p)}$  containing  $\mathcal{H}_\alpha$  and  $\mathcal{H}_\beta$ . For the second claim, if  $a \in \overline{\mathcal{I}_\mathcal{B}(p)}$  is positive, then  $\overline{(a + p)\mathcal{B}(a + p)}$  is a  $\sigma$ -unital hereditary sub- $C^*$ -algebra of  $\overline{\mathcal{I}_\mathcal{B}(p)}$  containing  $p\mathcal{B}p$  and  $a$ . There exists therefore a (unique) lower semicontinuous trace  $\tau$  in  $T^+(\overline{\mathcal{I}_\mathcal{B}(p)})$  that extends each  $\tau_\alpha$ , and hence  $\tau$ .

Extend  $\tau$  further to a lower semicontinuous trace in  $T^+(\mathcal{B})$  using (2.3). Finally, if we wish, we can linearize  $\tau$  using Proposition 2.13.

If we depart from lower semicontinuous densely defined traces and consider possibly singular traces, then we do obtain a  $C^*$ -algebraic characterization of groups with the fixed-point property for cones. The theorem below is a noncommutative analogue of the equivalence between (1) and (4) of [16, Theorem 7].

**Theorem 3.12.** *The following conditions are equivalent for any group  $G$ :*

- (i)  $G$  has the fixed-point property for cones.
- (ii) Whenever  $G$  acts on a  $C^*$ -algebra  $\mathcal{A}$  and whenever  $e$  is a positive element in  $\mathcal{A}$  for which there exists a nonzero trace in  $T(\mathcal{J}_\mathcal{A}^G(e), \mathcal{A})$ , which is bounded on the  $G$ -orbit of  $e$ , then there exists an invariant trace in  $T(\mathcal{J}_\mathcal{A}^G(e), \mathcal{A})$  normalizing  $e$ .
- (iii) Whenever  $G$  acts on a  $C^*$ -algebra  $\mathcal{A}$  and whenever  $e$  is a positive element in  $\mathcal{A}$  for which there exists a trace in  $T^+(\mathcal{A})$  which is nonzero and bounded on the  $G$ -orbit of  $e$ , then there is an invariant trace in  $T^+(\mathcal{A})$  normalizing  $e$ .

*Proof.*

(i)  $\Rightarrow$  (ii) Set  $\mathcal{I} = \mathcal{J}_\mathcal{A}^G(e)$ . Then  $T(\mathcal{I}, \mathcal{A})$  is weakly complete by Proposition 3.1, the action of  $G$  on  $T(\mathcal{I}, \mathcal{A})$  is locally bounded by Lemma 3.8, and of cobounded type by Corollary 3.4. The cone of traces  $T(\mathcal{I}, \mathcal{A})$  therefore has a nonzero fixed-point  $\tau_0$  since  $G$  has the fixed-point property for cones. As  $\tau_0$  is nonzero and invariant, we can use Lemmas 2.9 and 2.5 and the fact that  $e \in \mathcal{I}$  to conclude that  $0 < \tau_0(e) < \infty$ .

(ii)  $\Rightarrow$  (i) Property (ii), applied to  $\mathcal{A} = \ell^\infty(G)$  (see also Section 4), says that condition (4) in [16, Theorem 7] holds, which again, by that theorem, is equivalent to (i).

(ii)  $\Leftrightarrow$  (iii) This follows from Proposition 2.13 and the remarks below that proposition. □

If  $\mathcal{A}$  is commutative, or, more generally, if  $\mathcal{A}$  admits a separating family of bounded traces, then the condition in Theorem 3.12(ii) and (iii), that there exists a nonzero locally bounded traces, is always satisfied independent on the action of  $G$  on  $\mathcal{A}$ .

Below we specialize Theorem 3.12 to the case in which the positive element is a projection.

**Corollary 3.13.** *Let  $G$  be a group with the fixed-point property for cones acting on a  $C^*$ -algebra  $\mathcal{A}$ , and let  $p \in \mathcal{A}$  be a nonzero projection. The following conditions are equivalent:*

- (i) *There is a nonzero lower semicontinuous trace with domain  $\mathcal{J}_{\mathcal{A}}^G(p)$ , which is bounded on the  $G$ -orbit of  $p$ .*
- (ii) *There is an invariant lower semicontinuous trace in  $T^+(\mathcal{A})$  normalized on  $p$ .*
- (iii) *There is a lower semicontinuous trace in  $T^+(\mathcal{A} \rtimes G)$  normalized on  $p$ .*

*If, in addition,  $\mathcal{A} \rtimes G$  is exact, then the conditions above are equivalent to the following:*

- (iv) *The  $C^*$ -algebra  $p(\mathcal{A} \rtimes G)p \otimes M_n$  is not properly infinite, for all  $n \geq 1$ .*

*Proof.*

(i)  $\Rightarrow$  (ii) It follows from Theorem 3.12 that there is an invariant trace  $\tau$  with domain  $\mathcal{J}_{\mathcal{A}}^G(p)$  normalizing  $p$ . As remarked below Theorem 2.11, each trace on  $\mathcal{J}_{\mathcal{A}}^G(p)$  is automatically lower semicontinuous.

(ii)  $\Rightarrow$  (iii) Any invariant lower semicontinuous trace  $\tau$  on  $\mathcal{A}$  with  $\tau(p) = 1$  extends to a lower semicontinuous trace on  $\mathcal{A} \rtimes G$ , by Lemma 2.14.

(iii)  $\Rightarrow$  (i) Take the restriction to  $\mathcal{A}$  of the trace whose existence is claimed in (iii) and linearize as in Proposition 2.13.

(iii)  $\Leftrightarrow$  (iv) It is well known that this equivalence holds when  $\mathcal{A} \rtimes G$  is exact; see Remark 3.11 and the comments above Theorem 2.15.  $\square$

We can rephrase the corollary above as follows. A projection  $p$  in a  $C^*$ -algebra  $\mathcal{A}$  with a  $G$ -action can fail to be normalized by an invariant trace on  $\mathcal{A}$  for two reasons. Either  $\mathcal{A}$  does not have a trace that is nonzero and bounded on the  $G$ -orbit of  $p$  or the group  $G$  possesses some amount of “paradoxicality” materialized in failing to have the fixed-point property for cones. In Lemma 5.7 we give nonobvious examples of projections for which there exist a trace that is nonzero and bounded on the  $G$ -orbit of the projection.

While we do not know that every group without the fixed-point property for cones has the ability of acting on a  $C^*$ -algebra  $\mathcal{A}$  in such a way that it obstructs the existence of an invariant trace normalizing a given projection  $p$  in  $\mathcal{A}$ , for which there is a trace on  $\mathcal{A}$  that is nonzero and bounded on the orbit of  $p$ , it does follow from Proposition 3.10 that all nonsupramenable groups have this quality.

In conclusion, we do not know if Corollary 3.13 characterizes the class of groups with the fixed-point property for cones, the class of supramenable groups, or some intermediate class of groups.

#### 4. INVARIANT INTEGRALS ON $\ell^\infty(G)$

The bounded (complex valued) functions,  $\ell^\infty(G)$ , on a (discrete) group  $G$  is a unital  $C^*$ -algebra equipped with an action of the group  $G$  by left-translation.

Following the notation of Monod [16], for each positive  $f \in \ell^\infty(G)$ , let  $\ell^\infty(G, f)$  denote the set of bounded functions  $g$  on  $G$  whose absolute value is  $G$ -bounded by  $f$ , i.e., for which  $|g| \leq \sum_{j=1}^n t_j \cdot f$ , for some  $n \geq 1$  and some  $t_1, \dots, t_n \in G$ , where  $t \cdot f$  is the left-translate of  $f \in \ell^\infty(G)$  by  $t \in G$ . In the language of  $C^*$ -algebras,  $\ell^\infty(G, f)$  is the smallest (automatically symmetric) hereditary  $G$ -invariant ideal in  $\ell^\infty(G)$  containing  $f$ , denoted by  $\mathcal{I}_{\ell^\infty(G)}^G(f)$  in the previous sections. The Pedersen ideal of the uniform closure of  $\ell^\infty(G, f)$  is denoted by  $\ell_c^\infty(G, f)$ , and we have the following inclusions:

$$\ell_c^\infty(G, f) \subseteq \ell^\infty(G, f) \subseteq \overline{\ell^\infty(G, f)}.$$

An *invariant integral on  $G$  normalized for  $f$*  is a  $G$ -invariant positive linear functional  $\mu$  on  $\ell^\infty(G, f)$  satisfying  $\mu(f) = 1$ . In the language of  $C^*$ -algebras, an (invariant) integral on  $\ell^\infty(G, f)$  is an (invariant) trace on  $\ell^\infty(G)$  with domain  $\ell^\infty(G, f)$ .

Monod observed in [16, Theorem 7] that  $G$  has the fixed-point property for cones if and only if for each positive function  $f$  in  $\ell^\infty(G)$  there is an invariant integral on  $G$  normalized for  $f$ . (This result is extended to general  $C^*$ -algebras in Theorem 3.12.)

Let  $\mu$  be an invariant integral on  $\ell^\infty(G)$  normalized on some positive function  $f \in \ell^\infty(G)$ . We say that  $\mu$  is lower semicontinuous if  $\mu(h) = \lim_{n \rightarrow \infty} \mu(h_n)$  whenever  $\{h_n\}_{n=1}^\infty$  is an increasing sequence of positive functions in  $\ell^\infty(G, f)$  converging uniformly to  $h \in \ell^\infty(G, f)$ ; cf. Lemma 2.10. The restriction of  $\mu$  to  $\ell_c^\infty(G, f)$  is automatically lower semicontinuous by Theorem 2.11 (Pedersen). If the restriction of  $\mu$  to  $\ell_c^\infty(G, f)$  is 0, then  $\mu$  is said to be singular.

We shall in this section find conditions that will ensure that an (invariant) integral is lower semicontinuous and also exhibit situations where such integrals necessarily are singular. We start by rephrasing what Proposition 3.10 says about the existence of lower semicontinuous integrals.

**Proposition 4.1.** *Let  $G$  be a supramenable group, let  $f$  be a positive function in  $\ell^\infty(G)$ , and suppose that  $\overline{\ell^\infty(G, f)}$  is  $G$ -compact (see Proposition 3.6). Then there is a nonzero invariant lower semicontinuous integral on  $\ell_c^\infty(G, f)$ .*

**Example 4.2.** Let  $G$  be a group, and let  $f$  be a positive function in  $\ell^\infty(G)$ . Then  $\overline{\ell^\infty(G, f)}$  is  $G$ -compact if and only if there exists a  $\delta > 0$  such that  $A(f, \varepsilon) \times_G A(f, \delta)$ , for all  $0 < \varepsilon < \delta$ , where  $A(f, \eta) = \{t \in G : f(t) > \eta\}$ , and where  $A \times_G B$ , for subsets  $A, B$  of  $G$ , means that  $A \subseteq \bigcup_{t \in F} tB$ , for some finite subset  $F$  of  $G$ . We will not go into the details of the proof of this, but just mention that to prove the “if” part, one observes that  $A(f, \varepsilon) \times_G A(f, \delta)$  implies that  $(f - \varepsilon)_+ \in \mathcal{I}_{\ell^\infty(G)}^G((f - \delta')_+)$ , for all  $0 < \delta' < \delta$ .

The condition above ensuring  $G$ -compactness of  $\overline{\ell^\infty(G, f)}$  can further be rewritten as follows: Set  $A_n = \{t \in G : \frac{1}{n} < f(t) \leq \frac{1}{n-1}\}$ , for all  $n \geq 1$ . Then  $\overline{\ell^\infty(G, f)}$  is  $G$ -compact if and only if there exists an  $N_0 \geq 1$  such that  $\bigcup_{n=1}^N A_n \times_G \bigcup_{n=1}^{N_0} A_n$ , for all  $N \geq N_0$ . This condition holds if  $A_k \times_G \bigcup_{n=1}^{N_0} A_n = A(f, 1/N_0)$ , for all  $k > N_0$ . In other words,  $\overline{\ell^\infty(G, f)}$  is  $G$ -compact if the set of  $t \in G$  where  $f(t)$  is “very small” can be controlled by the set where  $f(t)$  is “small enough”.

Proposition 4.1 and the example above does not give information about the existence of invariant lower semicontinuous integrals normalizing the given positive function  $f \in \ell^\infty(G)$ . See Example 4.4 for more about this problem. Using an example from [15], we proceed to show that the conclusion of Proposition 4.1 fails without the assumption on  $G$ -compactness.

**Proposition 4.3.** *For each countably infinite group  $G$  there is a positive function  $f \in \ell^\infty(G)$  such that  $\ell_c^\infty(G, f)$  admits no nonzero invariant integral. In particular, if an invariant integral on  $G$  normalized on  $f$  exists, as is the case when  $G$  has the fixed-point property for cones, then it is singular.*

*Proof.* Let  $G$  be a countably infinite group. By [15, Proposition 4.3] and its proof there is an increasing sequence  $\{A_n\}_{n \geq 1}$  of (infinite) subsets of  $G$  such that if  $K_n$  is the (compact-open) closure of (the open set)  $A_n$  in  $\beta G$ , and if  $X_n = \bigcup_{t \in G} t.K_n$ , then  $X := \bigcup_{n \geq 1} X_n$  is an open  $G$ -invariant subset of  $\beta G$  that admits no nonzero invariant Radon measure.

Let  $\varphi: \ell^\infty(G) \rightarrow C(\beta G)$  be the canonical  $*$ -isomorphism of  $C^*$ -algebras, and let  $\mathcal{I}$  be the closed  $G$ -invariant ideal of  $\ell^\infty(G)$  such that  $\varphi(\mathcal{I}) = C_0(X)$ . Observe that  $C_0(X)$  is the closed  $G$ -invariant ideal in  $C(\beta G)$  generated by the indicator functions  $1_{K_n}$ ,  $n \geq 1$ . Since  $\varphi(1_{A_n}) = 1_{K_n}$ , we conclude that  $\mathcal{I}$  is the closed  $G$ -invariant ideal in  $\ell^\infty(G)$  generated by the projections  $1_{A_n}$ ,  $n \geq 1$ , or by the positive function  $f = \sum_{n \geq 1} n^{-2} 1_{A_n} \in \ell^\infty(G)$ . Hence  $\mathcal{I} = \overline{\ell^\infty(G, f)}$ .

Since  $X$  admits no nonzero invariant Radon measure,  $\ell_c^\infty(G, f) \cong C_c(X)$  admits no nonzero invariant integrals. □

**Example 4.4** (On the ideal  $c_0(G)$ ). For a countably infinite group  $G$  the subspace  $c_0(G)$  is a closed  $G$ -simple invariant ideal of  $\ell^\infty(G)$ , and the Pedersen ideal of  $c_0(G)$  is  $c_c(G)$ . It follows that if  $f$  is a positive nonzero function in  $c_0(G)$ , then

$$\ell_c^\infty(G, f) = c_c(G), \quad \overline{\ell^\infty(G, f)} = c_0(G),$$

while  $\ell^\infty(G, f)$  is some invariant hereditary ideal between these two ideals. Being  $G$ -simple, the ideal  $c_0(G)$  is  $G$ -compact.

The only lower semicontinuous invariant integrals defined on  $c_c(G)$  are multiples of the counting measure on  $G$ . Hence if  $f$  is a positive function in  $c_0(G)$ , then there is an invariant lower semicontinuous integral on  $G$  normalized on  $f$  if and only if  $f$  belongs to  $\ell^1(G)$ . If a positive function  $f$  in  $c_0(G) \setminus \ell^1(G)$  is normalized by an invariant integral on  $G$ , which is the case if  $G$  has the fixed-point property for cones, then this integral is necessarily singular.

Consider now a countably infinite group  $G$  with the fixed-point property for cones, and take a nonzero positive function  $f \in c_0(G)$ . The crossed product  $C^*$ -algebra  $c_0(G) \rtimes G$  is isomorphic to the  $C^*$ -algebra  $\mathcal{K}$  of compact operators (on a separable infinite-dimensional Hilbert space). There is an invariant integral on  $G$  normalized for  $f$ , and we can view this integral as a invariant densely defined trace  $\tau$  on  $c_0(G)$  with  $\tau(f) = 1$ . The image of  $f$  in  $c_0(G) \rtimes G$  is a positive compact operator with eigenvalues  $\{f(t)\}_{t \in G}$ . It was shown in [1] that for a positive compact operator  $T$  with eigenvalues  $\{\lambda_n\}_{n=1}^\infty$ , listed in decreasing order (and with multiplicity), there exists a densely defined trace on  $\mathcal{K}$  normalized for  $T$  if and only if

$$(4.1) \quad \liminf_{n \rightarrow \infty} \sigma_{2n} / \sigma_n = 1,$$

where  $\sigma_n = \sum_{j=1}^n \lambda_j$ .

Let  $s > 0$ , and choose  $f \in c_0(G)$  such that the eigenvalues of the positive compact operator  $f \in c_0(G) \rtimes G$  is the sequence  $\{n^{-s}\}_{n=1}^\infty$ . Then  $f$  is trace class if and only if  $s > 1$ , and the limit in (4.1) is 1 if and only if  $s \geq 1$ . If  $s = 1$ , then  $f$  is normalized by the *Dixmier trace* on  $\mathcal{K}$ . If  $0 < s < 1$ , then there is no trace on  $c_0(G) \rtimes G$  (lower semicontinuous or not) that normalizes  $f$ . For such a choice of  $f$ , the invariant

densely defined trace  $\tau$  on  $c_0(G)$  does not extend to a trace on  $c_0(G) \rtimes G$ , thus showing that the conclusion of Lemma 2.14 fails without the assumption that the trace is lower semicontinuous.

By the remark in the example above, that for no positive functions  $f$  in  $c_0(G) \setminus \ell^1(G)$  does there exist an invariant lower semicontinuous integral  $\mu$  on  $\ell^\infty(G, f)$  with  $\mu(f) = 1$ , we get the corollary below, which implies that at least some integrals witnessing the fixed-point property for cones for infinite groups necessarily must be singular.

**Corollary 4.5.** *Let  $G$  be a countable group with the property that for each positive function  $f$  in  $\ell^\infty(G)$  there exists a  $G$ -invariant lower semicontinuous integral  $\mu$  on  $\ell^\infty(G, f)$  normalized for  $f$ . Then  $G$  must be a finite group.*

## 5. THE ROE ALGEBRA

As an application of the results developed in the previous sections, we shall in this last section prove some results about existence (and nonexistence) of traces on the Roe algebra  $\ell^\infty(G, \mathcal{K}) \rtimes G$  associated with a (countably infinite) group  $G$ , where  $\mathcal{K}$  denotes the  $C^*$ -algebra of compact operators on a separable infinite-dimensional Hilbert space. These  $C^*$ -algebras originate from the thesis of Roe, published in [21], where the index of elliptic operators are computed using traces on a (variant of) what is now called the Roe algebra.

The  $C^*$ -algebra  $\ell^\infty(G, \mathcal{K})$  is equipped with the natural action of  $G$  given by left-translation. We shall consider existence (and nonexistence) of invariant traces on this  $C^*$ -algebra. First we give a complete description of the densely defined lower semicontinuous traces on these  $C^*$ -algebras (there are not so many). The group  $G$  plays no role in Lemma 5.1 and Proposition 5.2 other than as a set, and it could be replaced by the set of natural numbers  $\mathbb{N}$ .

**Lemma 5.1.** *The Pedersen ideal of  $\ell^\infty(G, \mathcal{K})$  is equal to  $\ell^\infty(G, \mathcal{F})$ , where  $\mathcal{F} = \text{Ped}(\mathcal{K})$  is the set of finite rank operators.*

*Proof.* The “point evaluation” at  $t \in G$  gives a surjective  $*$ -homomorphism  $\ell^\infty(G, \mathcal{K}) \rightarrow \mathcal{K}$  that maps  $\text{Ped}(\ell^\infty(G, \mathcal{K}))$  onto  $\text{Ped}(\mathcal{K}) = \mathcal{F}$ . This shows that  $\text{Ped}(\ell^\infty(G, \mathcal{K})) \subseteq \ell^\infty(G, \mathcal{F})$ . Conversely, if  $x \in \ell^\infty(G, \mathcal{F})$ , then  $x(t)$  has finite rank, for each  $t \in G$ , so there exists a finite-dimensional projection  $p(t) \in \mathcal{K}$  with  $p(t)x(t) = x(t)$ . The function  $t \mapsto p(t)$  defines a projection  $p$  in  $\ell^\infty(G, \mathcal{K})$  satisfying  $px = x$ . As  $p$  belongs to the Pedersen ideal, being a projection, it follows that also  $x$  belongs to  $\text{Ped}(\ell^\infty(G, \mathcal{K}))$ .  $\square$

It is worth mentioning that the Pedersen ideal of a  $C^*$ -algebra of the form  $C(T, \mathcal{K})$ , where  $T$  is a compact Hausdorff space, may be properly contained in  $C(T, \mathcal{F})$ ; see [9].

For each  $s \in G$ , let  $\tau_s$  be the (lower semicontinuous densely defined) trace on  $\ell^\infty(G, \mathcal{K})$  given by  $\tau_s(f) = \text{Tr}(f(s))$ , for  $f$  either in  $\ell^\infty(G, \mathcal{K})^+$  or in  $\ell^\infty(G, \mathcal{F})$ , where  $\text{Tr}$  is the standard trace on the compact operators  $\mathcal{K}$ .

In the proof of the proposition below, we shall view the  $C^*$ -algebra  $\ell^\infty(G, \mathcal{K})$  as an  $\ell^\infty(G)$ -algebra via the natural (unital) embedding of  $\ell^\infty(G)$  into the center of the multiplier algebra of  $\ell^\infty(G, \mathcal{K})$ .

**Proposition 5.2.** *For each countable group  $G$ , the cone,  $T_{\text{isc}}(\ell^\infty(G, \mathcal{K}))$ , of densely defined lower semicontinuous traces on  $\ell^\infty(G, \mathcal{K})$  is equal to the cone of finite positive linear combinations of the traces  $\tau_s$ ,  $s \in G$ , defined above, i.e., traces of the form  $\sum_{s \in G} c_s \tau_s$ , where  $c_s \geq 0$  and  $c_s \neq 0$  for only finitely many  $s \in G$ .*

*Proof.* Clearly, any trace of the form  $\sum_{s \in G} c_s \tau_s$ , with  $c_s \geq 0$  and  $c_s \neq 0$  only for finitely many  $s \in G$ , is lower semicontinuous and densely defined.

For the converse direction, take  $\tau$  in  $T_{\text{isc}}(\ell^\infty(G, \mathcal{K}))$ . Fix a one-dimensional projection  $e \in \mathcal{K}$ . For each  $s \in G$ , let  $e_s \in \ell^\infty(G, \mathcal{F})$  be given by  $e_s(s) = e$  and  $e_s(t) = 0$  when  $t \neq s$ . Set  $c_s = \tau(e_s) \geq 0$ .

We show first that the set  $F = \{s \in G : c_s \neq 0\}$  is finite. Suppose that it were infinite, and let  $\{s_1, s_2, s_3, \dots\}$  be an enumeration of the elements in  $F$ . Let  $p \in \ell^\infty(G, \mathcal{K})$  be a projection such that  $\text{Tr}(p(s_n)) \geq nc_{s_n}^{-1}$ , for all  $n \geq 1$ . As  $p \geq p \cdot 1_{\{s\}}$ , for all  $s \in G$ , we get  $\tau(p) \geq \tau(p \cdot 1_{\{s_n\}}) \geq c_{s_n} \text{Tr}(p(s_n)) \geq n$ . As this cannot be true for all  $n \geq 1$ , we conclude that  $F$  is finite.

Set  $\tau_0 = \sum_{s \in F} c_s \tau_s$ , and observe that  $\tau_0(f) = \tau(f \cdot 1_F)$ , for all  $f \in \ell^\infty(G, \mathcal{K})$ . It follows that  $\tau' := \tau - \tau_0$  is a positive trace on  $\ell^\infty(G, \mathcal{K})$ , satisfying  $\tau'(f) = \tau(f \cdot 1_{F^c})$ , for all  $f \in \ell^\infty(G, \mathcal{K})$ . We show that  $\tau' = 0$ . Assume, to reach a contradiction, that  $\tau' \neq 0$ . Notice that  $\tau'$  vanishes on  $c_c(G, \mathcal{K})$  by construction of  $\tau_0$ . Since the Pedersen ideal of  $\ell^\infty(G, \mathcal{K})$  is generated (as a hereditary ideal) by its projections (see the proof of Lemma 5.1), there is a projection  $p \in \ell^\infty(G, \mathcal{K})$  such that  $\tau'(p) > 0$ . Write  $G = \bigcup_{n=1}^\infty F_n$ , where  $\{F_n\}_{n \geq 1}$  is a strictly increasing sequence of finite subsets of  $G$ . Find a sequence  $\{p_k\}_{k \geq 1}$  of pairwise orthogonal projections in  $\ell^\infty(G, \mathcal{K})$  such that each  $p_k$  is equivalent to  $p$ . Let  $q \in \ell^\infty(G, \mathcal{K})$  be the projection given by

$$q(s) = p_1(s) + p_2(s) + \dots + p_n(s), \quad s \in F_{n+1} \setminus F_n,$$

for  $n \geq 0$  (with the convention  $F_0 = \emptyset$ ). Then

$$q \cdot 1_{F_n^c} \geq (p_1 + p_2 + \dots + p_n) \cdot 1_{F_n^c},$$

for all  $n \geq 1$ , and as  $\tau'(g \cdot 1_{E^c}) = \tau'(g)$ , for all  $g \in \ell^\infty(G, \mathcal{K})$  and all finite subsets  $E \subseteq G$ , we conclude that

$$\tau'(q) = \tau'(q \cdot 1_{F_n^c}) \geq \tau'((p_1 + p_2 + \dots + p_n) \cdot 1_{F_n^c}) = \tau'(p_1 + p_2 + \dots + p_n) = n\tau'(p),$$

for all  $n \geq 1$ , which is impossible. □

A nonzero trace of the form as in Proposition 5.2 can clearly not be  $G$ -invariant when  $G$  is infinite, so we obtain the following.

**Corollary 5.3.** *The  $C^*$ -algebra  $\ell^\infty(G, \mathcal{K})$  admits no nonzero lower semicontinuous invariant densely defined traces when  $G$  is a countably infinite group, and, consequently, the Roe algebra  $\ell^\infty(G, \mathcal{K}) \rtimes G$  admits no nonzero densely defined lower semicontinuous trace.*

When combining the conclusion of the corollary above with Theorem 3.9, we see that the action of  $G$  on  $T_{\text{isc}}(\ell^\infty(G, \mathcal{K}))$  either must fail to be of cobounded type or there is no nonzero locally bounded trace in  $T_{\text{isc}}(\ell^\infty(G, \mathcal{K}))$ . In fact, both fail. That the latter fails follows easily from the description of  $T_{\text{isc}}(\ell^\infty(G, \mathcal{K}))$  in Proposition 5.2.

**Lemma 5.4.** *If  $G$  is a countably infinite group, then the action of  $G$  on  $T_{\text{isc}}(\ell^\infty(G, \mathcal{K}))$  is not of cobounded type.*

*Proof.* Let  $e$  be a positive contraction in  $\text{Ped}(\ell^\infty(G, \mathcal{K}))$ . We must find another positive contraction  $a$  in  $\text{Ped}(\ell^\infty(G, \mathcal{K}))$  that is not tracially  $G$ -dominated by  $e$ ; cf. Lemma 3.2. In other words, for all finite sets  $t_1, t_2, \dots, t_n \in G$ , the inequality  $\tau(a) \leq \sum_{j=1}^n \tau(t_j \cdot e)$  will fail for at least one  $\tau \in T_{\text{isc}}(\ell^\infty(G, \mathcal{K}))$ , or, taking Proposition 5.2 into account,

$$\text{Tr}(a(s)) \leq \sum_{j=1}^n \text{Tr}(e(t_j^{-1}s)),$$

will fail for at least one  $s \in G$ . It follows from Lemma 5.1 and its proof that  $e$  is dominated by a projection in  $\ell^\infty(G, \mathcal{K})$ , so we may without loss of generality assume that  $e$  itself is a projection. Set  $f(s) = \text{Tr}(e(s))$ , for  $s \in G$ .

Let  $G = \{s_1, s_2, s_3, \dots\}$  be an enumeration of the elements in  $G$ , and let  $\{u_j\}_{j=1}^\infty$  be a sequence in which each element of  $G$  is repeated infinitely often. Set  $f_N = \sum_{j=1}^N u_j \cdot f$ , for each  $N \geq 1$ , and let  $g: G \rightarrow \mathbb{N}_0$  be given by  $g(s_N) = f_N(s_N) + 1$ , for  $N \geq 1$ . For each finite set  $t_1, t_2, \dots, t_n \in G$  there exists an  $N \geq 1$  such that  $\sum_{j=1}^n t_j \cdot f \leq f_N$ , which entails that  $g(s_N) \not\leq \sum_{j=1}^n t_j \cdot f(s_N)$ , for each  $N \geq 1$ .

Let now  $a \in \ell^\infty(G, \mathcal{K})$  be a projection such that  $\text{Tr}(a(t)) = g(t)$ , for all  $t \in G$ . Then  $a$  is not tracially  $G$ -dominated by  $e$ . □

Although there are no *densely defined* lower semicontinuous invariant traces on  $\ell^\infty(G, \mathcal{K})$ , when  $G$  is infinite, there are still interesting lower semicontinuous traces on the Roe algebra. In the results to follow, we describe the class of projections  $p \in \ell^\infty(G, \mathcal{K})$  for which there exists a lower semicontinuous trace on  $\ell^\infty(G, \mathcal{K})$  which is bounded and nonzero on the orbit of  $p$  (resp., which is invariant and normalizes  $p$ ). These two classes of projections agree when  $G$  has the fixed-point property for cones; cf. Corollary 3.13. First we note the following general result that holds for locally finite groups.

**Proposition 5.5.** *The Roe algebra  $\ell^\infty(G, \mathcal{K}) \rtimes G$  of any locally finite group  $G$  is stably finite. Moreover, for each nonzero projection  $p \in \ell^\infty(G, \mathcal{K})$  there exists a lower semicontinuous trace  $\tau \in T^+(\ell^\infty(G, \mathcal{K}) \rtimes G)$  with  $\tau(p) = 1$ , and hence an invariant lower semicontinuous trace on  $\ell^\infty(G, \mathcal{K})$  normalizing  $p$ .*

*Proof.* Write  $G = \bigcup_{n=1}^\infty G_n$  as an increasing union of finite groups. The  $C^*$ -algebra  $\ell^\infty(G, \mathcal{K})$  is stably finite, as can be witnessed by the separating family of densely defined traces from Proposition 5.2. The crossed product  $\ell^\infty(G, \mathcal{K}) \rtimes G_n$  is stably finite because it embeds in the stably finite  $C^*$ -algebra  $\ell^\infty(G, \mathcal{K}) \otimes B(\ell^2(G_n))$ . It follows that

$$\ell^\infty(G, \mathcal{K}) \rtimes G = \varinjlim \ell^\infty(G, \mathcal{K}) \rtimes G_n$$

is stably finite, being an inductive limit of stably finite  $C^*$ -algebras.

Let next  $p \in \ell^\infty(G, \mathcal{K})$  be a nonzero projection. Then

$$(5.1) \quad p(\ell^\infty(G, \mathcal{K}) \rtimes G)p = \varinjlim p(\ell^\infty(G, \mathcal{K}) \rtimes G_n)p.$$

The unital  $C^*$ -algebra  $p(\ell^\infty(G, \mathcal{K}) \rtimes G_n)p$  admits a tracial state, for each  $n \geq 1$ . To see this, choose  $m \geq n$  such that the restriction  $p'$  of  $p$  to  $\ell^\infty(G_m, \mathcal{K})$  is nonzero. The restriction mapping  $\ell^\infty(G, \mathcal{K}) \rightarrow \ell^\infty(G_m, \mathcal{K})$  is  $G_n$ -equivariant and therefore extends to a  $*$ -homomorphism  $\ell^\infty(G, \mathcal{K}) \rtimes G_n \rightarrow \ell^\infty(G_m, \mathcal{K}) \rtimes G_n$ , and in turn to a unital  $*$ -homomorphism  $p(\ell^\infty(G, \mathcal{K}) \rtimes G_n)p \rightarrow p'(\ell^\infty(G_m, \mathcal{K}) \rtimes G_n)p'$ . Composing this  $*$ -homomorphism with any tracial state on the (finite-dimensional)  $C^*$ -algebra  $p'(\ell^\infty(G_m, \mathcal{K}) \rtimes G_n)p'$  gives a tracial state on  $p(\ell^\infty(G, \mathcal{K}) \rtimes G_n)p$ . As the inductive

limit of a sequence of unital  $C^*$ -algebras with unital connecting mappings (e.g., as in (5.1)) admits a tracial state if and only if each  $C^*$ -algebra in the sequence does, we conclude that  $p(\ell^\infty(G, \mathcal{K}) \rtimes G)p$  admits a tracial state. By Remark 3.11 there is a lower semicontinuous trace on  $\ell^\infty(G, \mathcal{K}) \rtimes G$  normalizing the projection  $p$ . The restriction of this trace to  $\ell^\infty(G, \mathcal{K})$  is an invariant lower semicontinuous trace still normalizing  $p$ .  $\square$

The Roe algebra of any infinite, locally finite group provides yet another example of a stably finite  $C^*$ -algebra with an approximate unit consisting of projections which has no densely defined lower semicontinuous trace; see Theorem 2.15 and Proposition 2.16.

For the more general class of groups  $G$  with the fixed-point property for cones, it follows from Corollary 3.13 that if  $p$  is a projection in  $\ell^\infty(G, \mathcal{K})$ , then there is a lower semicontinuous invariant trace on  $\ell^\infty(G, \mathcal{K})$  that normalizes  $p$  (and hence there is a trace on the Roe algebra  $\ell^\infty(G, \mathcal{K}) \rtimes G$  normalizing  $p$ ) if and only if there is a trace on  $\ell^\infty(G, \mathcal{K})$  which is nonzero and bounded on the orbit  $\{t.p\}_{t \in G}$ . If  $p$  has *uniformly bounded dimension*; i.e., if  $\sup_{t \in G} \text{Tr}(p(t)) < \infty$ , then such a trace clearly exists—take, for example,  $\tau_s$  (defined above Proposition 5.2), for any fixed  $s \in G$ . One can do a bit better: In Proposition 5.10 it is shown that for each projection in  $\ell^\infty(G, \mathcal{K})$  with uniformly bounded dimension there is a trace on the Roe algebra normalizing this projection provided that the group  $G$  is *supramenable* (a formally weaker condition than having the fixed-point property for cones).

Let  $G$  be a countably infinite group, and let  $\ell: G \rightarrow \mathbb{N}_0$  be a proper length function on  $G$ ; i.e.,  $\ell(t) = 0$  if and only if  $t = e$ ,  $\ell(s + t) \leq \ell(s) + \ell(t)$ , for all  $s, t \in G$ , and  $W_n := \{t \in G : \ell(t) \leq n\}$  is finite, for all  $n \geq 0$ . Such proper length functions always exist, and if  $G$  is finitely generated, then we can take  $\ell$  to be the word length function with respect to some finite generating set for  $G$ . Set

$$(5.2) \quad \alpha_n = \max_{t \in W_n} \text{Tr}(p(t)), \quad Z_n = \{t \in W_n : \text{Tr}(p(t)) = \alpha_n\}.$$

Let  $\tau_n$  be the (lower semicontinuous densely defined) trace on  $\ell^\infty(G, \mathcal{K})$  given by

$$\tau_n(f) = \frac{1}{|Z_n|} \sum_{t \in Z_n} \alpha_n^{-1} \text{Tr}(f(t)),$$

for  $f$  in  $\ell^\infty(G, \mathcal{K})^+$  or in  $\ell^\infty(G, \mathcal{F})$ . Observe that  $\tau_n(p) = 1$ , for all  $n \geq 1$ . Let  $\omega$  be a free ultrafilter on  $\mathbb{N}_0$ , and define a trace  $\tau_{\omega,p}$  on the positive cone  $\ell^\infty(G, \mathcal{K})^+$  by

$$\tau_{\omega,p}(f) = \lim_{\omega} \tau_n(f), \quad f \in \ell^\infty(G, \mathcal{K})^+$$

(where the limit along the ultrafilter is taken in the compact set  $[0, \infty]$ ). Let  $C_{\omega,p}$  be the cone of positive functions  $f$  in  $\ell^\infty(G, \mathcal{K})$ , for which  $\tau_{\omega,p}(f) < \infty$ , and let  $\mathcal{I}_{\omega,p}$  be the linear span of  $C_{\omega,p}$ . Then  $\mathcal{I}_{\omega,p}$  is a hereditary symmetric ideal in  $\ell^\infty(G, \mathcal{K})$ , and  $\tau_{\omega,p}$  defines a linear trace on  $\mathcal{I}_{\omega,p}$ ; cf. Proposition 2.13.

We say that the projection  $p \in \ell^\infty(G, \mathcal{K})$  has *subexponentially growing dimension* if

$$(5.3) \quad \liminf_{n \rightarrow \infty} \frac{\alpha_{n+m}}{\alpha_n} = 1,$$

for all  $m \geq 0$ . (This definition may depend on the choice of proper length function  $\ell$  and should be understood to be with respect to *some* proper length function.)

**Lemma 5.6.** *Let  $\{\alpha_n\}_{n=0}^\infty$  be an increasing sequence of strictly positive real numbers satisfying (5.3). Then there is a free ultrafilter  $\omega$  on  $\mathbb{N}_0$  such that*

$$\lim_{\omega} \frac{\alpha_{n+m}}{\alpha_n} = 1,$$

for all  $m \geq 1$ .

*Proof.* For each  $m \geq 0$  and  $\varepsilon > 0$  set  $A_{m,\varepsilon} = \{n \geq 0 : \alpha_{n+m}/\alpha_n \leq 1 + \varepsilon\}$ . By the assumption that (5.3) holds, each of the sets  $A_{m,\varepsilon}$  is infinite. The collection of sets  $A_{m,\varepsilon}$  is downward directed, since  $A_{m_1,\varepsilon_1} \subseteq A_{m_2,\varepsilon_2}$ , when  $m_1 \geq m_2$  and  $\varepsilon_1 \leq \varepsilon_2$ . It follows that the intersection of any finite collection of these sets is infinite. We can therefore find a free ultrafilter  $\omega$  which contains all of the sets  $A_{m,\varepsilon}$ , and any such ultrafilter will satisfy the conclusion of the lemma.  $\square$

**Lemma 5.7.** *Let  $G$  be a countably infinite group, let  $p \in \ell^\infty(G, \mathcal{K})$  be a projection of subexponentially growing dimension, and let  $\omega$  be a free ultrafilter as in Lemma 5.6 for the sequence  $\{\alpha_n\}_{n=0}^\infty$  associated with the projection  $p$  as in (5.2). Then  $\tau_{\omega,p}(p) = 1$  and  $\tau_{\omega,p}(t.p) \leq 1$ , for all  $t \in G$ .*

*Proof.* We have already noted that  $\tau_n(p) = 1$ , for all  $n \geq 1$ , which implies that  $\tau_{\omega,p}(p) = 1$ . Let  $m \geq 0$ , and let  $s \in W_m$ . For  $n \geq 0$  and  $t \in Z_n \subseteq W_n$  we have  $st \in W_{n+m}$ , so  $\text{Tr}(p(st)) \leq \alpha_{n+m}$ , which shows that

$$\tau_n(s^{-1}.p) = \frac{1}{|Z_n|} \sum_{t \in Z_n} \alpha_n^{-1} \text{Tr}(p(st)) \leq \frac{\alpha_{n+m}}{\alpha_n}.$$

By the assumption that  $p$  has subexponentially growing dimension, by Lemma 5.6, and by the choice of  $\omega$ , we conclude that  $\tau_{\omega,p}(s^{-1}.p) \leq 1$ .  $\square$

**Theorem 5.8.** *Let  $G$  be a countably infinite group with the fixed-point property for cones, and let  $p \in \ell^\infty(G, \mathcal{K})$  be a projection of subexponentially growing dimension. Then there is an invariant lower semicontinuous trace on  $\ell^\infty(G, \mathcal{K})$  normalized on  $p$ , and there is a lower semicontinuous trace on the Roe algebra  $\ell^\infty(G, \mathcal{K}) \rtimes G$  also normalized on  $p$ .*

*Proof.* This follows from Corollary 3.13, where condition (i) is satisfied with  $\tau = \tau_{p,\omega}$ ; cf. Lemma 5.7.  $\square$

We proceed to examine the case of projections in  $\ell^\infty(G, \mathcal{K})$  of uniformly bounded dimension. In the lemma below we embed  $\ell^\infty(G)$  in  $\ell^\infty(G) \rtimes G$ , which again embeds in (the upper left corner of)  $(\ell^\infty(G) \rtimes G) \otimes M_n$ , for each  $n \geq 1$ . Note that projections in  $\ell^\infty(G)$  are indicator functions  $1_E$ , for some subset  $E$  of  $G$ .

**Lemma 5.9.** *Let  $G$  be an exact group, and let  $n \geq 1$  be an integer. Then for each projection  $p$  in  $(\ell^\infty(G) \rtimes G) \otimes M_n$ , there is a projection  $r$  in  $\ell^\infty(G)$  such that  $p$  and  $r$  generate the same closed two-sided ideal in  $(\ell^\infty(G) \rtimes G) \otimes M_n$ .*

*Proof.* Let  $n \geq 1$ , let  $p \in (\ell^\infty(G) \rtimes G) \otimes M_n$  be a projection, and let  $\mathcal{I}$  be the closed two-sided ideal in  $(\ell^\infty(G) \rtimes G) \otimes M_n$  generated by  $p$ . It follows from [26, Theorem 1.16] that  $\mathcal{I}$  is the closed two-sided ideal in  $(\ell^\infty(G) \rtimes G) \otimes M_n$  generated by  $\mathcal{J} := \mathcal{I} \cap (\ell^\infty(G) \otimes M_n)$ . Arguing as in [11, proof of Proposition 5.3], we find a projection  $q \in \mathcal{J}$  which generates the ideal  $\mathcal{I}$  in  $(\ell^\infty(G) \rtimes G) \otimes M_n$ . By [28], since  $\ell^\infty(G)$  is of real rank 0,  $q$  is equivalent to a diagonal projection  $\text{diag}(q_1, q_2, \dots, q_n)$  in  $\ell^\infty(G) \otimes M_n$ . Let  $r \in \ell^\infty(G)$  be the supremum of the projections  $q_1, q_2, \dots, q_n$ .

Then  $q$  and  $r$  generate the same ideal of  $\ell^\infty(G) \otimes M_n$ , so  $r$  and  $p$  generate the same ideal in  $(\ell^\infty(G) \rtimes G) \otimes M_n$ .  $\square$

The result below extends the characterization in [11] of supramenable groups in terms of nonexistence of properly infinite projections in the uniform Roe algebra.

**Proposition 5.10.** *Let  $G$  be a group. The following conditions are equivalent:*

- (i)  $G$  is supramenable.
- (ii) Each nonzero projection in the stabilized uniform Roe algebra  $(\ell^\infty(G) \rtimes G) \otimes \mathcal{K}$  is normalized by a lower semicontinuous trace in  $T^+(\ell^\infty(G) \rtimes G) \otimes \mathcal{K}$ .
- (iii) Each projection in  $\ell^\infty(G, \mathcal{K})$  of uniformly bounded dimension is normalized by an invariant lower semicontinuous trace in  $T^+(\ell^\infty(G, \mathcal{K}))$ , and by a lower semicontinuous trace in  $T^+(\ell^\infty(G, \mathcal{K}) \rtimes G)$ .
- (iv) The stabilized uniform Roe algebra  $(\ell^\infty(G) \rtimes G) \otimes \mathcal{K}$  contains no properly infinite projections.

*Proof.*

(i)  $\Rightarrow$  (ii) Since  $(\ell^\infty(G) \rtimes G) \otimes \mathcal{K}$  is the inductive limit of  $C^*$ -algebras of the form  $(\ell^\infty(G) \rtimes G) \otimes M_n$ , for  $n \geq 1$ , it suffices to show that each projection  $p$  in  $(\ell^\infty(G) \rtimes G) \otimes M_n$  is normalized by a lower semicontinuous trace on this  $C^*$ -algebra.

By Lemma 5.9 there is a projection  $q = 1_E \in \ell^\infty(G)$  that generates the same closed two-sided ideal in  $(\ell^\infty(G) \rtimes G) \otimes M_n$  as  $p$ . Since  $G$  is supramenable, the set  $E$  must be nonparadoxical, so by Tarski’s theorem there is an invariant trace  $\tau_0$  on  $\ell^\infty(G, q)$  which normalizes  $q$  (see [11, Proposition 5.3]). Extend  $\tau_0$  to a trace  $\tau$  on  $\mathcal{J} := \mathcal{J}_{(\ell^\infty(G) \rtimes G) \otimes M_n}(q)$  satisfying  $\tau(q) = 1$ . Now,  $p \in \mathcal{J}$  and  $0 < \tau(p) < \infty$ , by the assumption on  $p$  and  $q$ , so upon rescaling we obtain a trace  $\tau$  on  $\mathcal{J}$  satisfying  $\tau(p) = 1$ , and  $\tau$  is lower semicontinuous by the remarks below Theorem 2.11. Finally, we can extend  $\tau$  to a trace in  $T^+(\ell^\infty(G) \rtimes G) \otimes M_n$  using (2.3).

(ii)  $\Rightarrow$  (iii) Let  $p$  be a projection in  $\ell^\infty(G, \mathcal{K})$  such that  $n := \sup_{t \in G} \text{Tr}(p(t)) < \infty$ . Let  $e \in \mathcal{K}$  be a projection of dimension  $n$ , and let  $\bar{e} \in \ell^\infty(G, \mathcal{K})$  be the projection given by  $\bar{e}(t) = e$ , for all  $t \in G$ . Then  $\bar{e}$  is fixed by the action of  $G$ , and we have the isomorphisms

$$\bar{e}(\ell^\infty(G, \mathcal{K}))\bar{e} \cong \ell^\infty(G) \otimes M_n, \quad \bar{e}(\ell^\infty(G, \mathcal{K}) \rtimes G)\bar{e} \cong (\ell^\infty(G) \rtimes G) \otimes M_n.$$

Moreover,  $p$  is equivalent to a projection  $p_0$  in  $\bar{e}(\ell^\infty(G, \mathcal{K}))\bar{e}$ , which, under the isomorphism above, corresponds to a projection  $p_1 \in \ell^\infty(G) \otimes M_n$ . By (ii) there is a lower semicontinuous trace on  $(\ell^\infty(G) \rtimes G) \otimes M_n$  normalizing  $p_1$ . Hence there is a lower semicontinuous trace on  $\bar{e}(\ell^\infty(G, \mathcal{K}) \rtimes G)\bar{e}$  normalizing  $p_0$ . Arguing as in Remark 3.11, we can extend this trace to a lower semicontinuous trace in  $T^+(\ell^\infty(G, \mathcal{K}) \rtimes G)$ . The restriction of  $\tau$  to  $\ell^\infty(G, \mathcal{K})$  becomes an invariant lower semicontinuous trace.

(iii)  $\Rightarrow$  (iv) This is clear (no properly infinite projection can be normalized by a trace). If  $G$  is nonsupramenable, then  $\ell^\infty(G) \rtimes G$  contains a properly infinite projection (see [20, Proposition 5.5]), in which case (iv) cannot be true, which proves (iv)  $\Rightarrow$  (i).  $\square$

It was shown in [20] that the uniform Roe algebra  $\ell^\infty(G) \rtimes G$  is properly infinite, i.e., its unit is a properly infinite projection, if and only if  $G$  is nonamenable, and in [11] it was shown that the uniform Roe algebra contains a properly infinite projection if and only if  $G$  is nonsupramenable. The proposition above allows us

to conclude that no matrix algebras over the uniform Roe algebra contain properly infinite projections when  $G$  is supramenable.

It was shown by Wei in [27] and Scarparo in [25], answering a question in [11], that  $\ell^\infty(G) \rtimes G$  is finite if and only if  $G$  is a locally finite group. Using a similar idea as in [25], we show below that the Roe algebra  $\ell^\infty(G, \mathcal{K}) \rtimes G$  contains a properly infinite projection whenever  $G$  is not locally finite. We may therefore strengthen Proposition 5.5 as follows: The Roe algebra  $\ell^\infty(G, \mathcal{K}) \rtimes G$  is stably finite if and only if  $G$  is locally finite, and if  $\ell^\infty(G, \mathcal{K}) \rtimes G$  is not stably finite, then it contains a properly infinite projection.

**Lemma 5.11.** *Let  $G$  be a nonlocally finite group. Then there is a finite subset  $S$  of  $G$  such that for each integer  $N \geq 1$  there exists a nonzero function  $\varrho: G \rightarrow \mathbb{N}_0$  satisfying*

$$(5.4) \quad N\varrho \leq \sum_{s \in S} s.\varrho.$$

*Proof.* Let  $G_0$  be an infinite, finitely generated subgroup of  $G$ , and let  $S$  be a finite symmetric generating set for  $G_0$ . By [29, Lemma 1] there is a one-sided geodesic  $\{t_n\}_{n=0}^\infty$  in  $G_0$ , where  $t_n t_{n+1}^{-1} \in S$ , for all  $n \geq 0$ , and  $n \mapsto t_n$  is injective. Fix  $N \geq 1$ , define  $\varrho: G \rightarrow \mathbb{N}_0$  by  $\varrho(t_n) = N^n$ , for all  $n \geq 0$ , and set  $\varrho(t) = 0$  if  $t \notin \{t_n : n \geq 0\}$ .

Fix  $n \geq 0$ , and set  $s = t_n t_{n+1}^{-1} \in S$ . Then  $s.\varrho(t_n) = \varrho(t_{n+1}) = N\varrho(t_n)$ . As  $\varrho(t) = 0$  when  $t \notin \{t_n : n \geq 0\}$ , we see that (5.4) holds.  $\square$

**Proposition 5.12.** *For each countable nonlocally finite group  $G$  there is a projection  $p \in \ell^\infty(G, \mathcal{K})$  satisfying the following:*

- (i)  $p$  is properly infinite in the Roe algebra  $\ell^\infty(G, \mathcal{K}) \rtimes G$ .
- (ii) There is no trace on the Roe algebra  $\ell^\infty(G, \mathcal{K}) \rtimes G$  normalizing the projection  $p$ .
- (iii) Each trace on  $\ell^\infty(G, \mathcal{K})$  is either unbounded or 0 on the orbit  $\{t.p\}_{t \in G}$ .

*Proof.* Note first that if  $e$  and  $f$  are projections in  $\ell^\infty(G, \mathcal{K})$ , then  $e \sim f$  (resp.,  $e \precsim f$ ) in  $\ell^\infty(G, \mathcal{K})$  if and only if  $\text{Tr}(e(t)) = \text{Tr}(f(t))$  (resp.,  $\text{Tr}(e(t)) \leq \text{Tr}(f(t))$ ), for all  $t \in G$ . Let  $\varrho: G \rightarrow \mathbb{N}_0$  be as in Lemma 5.11 with respect to a finite subset  $S$  of  $G$  and with  $N \geq 2|S|$ . Find a projection  $q \in \ell^\infty(G, \mathcal{K})$  with  $\text{Tr}(q(t)) = \varrho(t)$ , for all  $t \in G$ .

Let  $p$  (resp.,  $p'$ ) be the  $|S|$ -fold (resp., the  $N$ -fold) direct sum of  $q$  with itself in  $\ell^\infty(G, \mathcal{K})$ , and set  $e = \bigoplus_{s \in S} s.q$ . Then  $\text{Tr}(p'(t)) = N\varrho(t)$  and  $\text{Tr}(e(t)) = \sum_{s \in S} s.\varrho(t)$ , for all  $t \in G$ , so  $p' \precsim e$  and  $p \oplus p \precsim p'$  in  $\ell^\infty(G, \mathcal{K})$ . Moreover,  $p \sim e$  in  $\ell^\infty(G, \mathcal{K}) \rtimes G$  (since  $q \sim t.q$  in  $\ell^\infty(G, \mathcal{K}) \rtimes G$ , for all  $t \in G$ ). Hence

$$p \oplus p \precsim p' \precsim e \sim p$$

in  $\ell^\infty(G, \mathcal{K}) \rtimes G$ , which shows that (i) and (ii) hold.

(iii) follows from (ii) when  $G$  has the fixed-point property for cones (cf. Corollary 3.13) but requires a separate argument for general groups. Applying the operator  $\sigma \mapsto \sum_{s \in S} s.\sigma$ , on functions  $\sigma: G \rightarrow \mathbb{N}_0$ , to the left- and right-hand sides of (5.4)  $k - 1$  times, we get  $N^k \varrho \leq \sum_{s \in S^k} \bar{s}.\varrho$ , for all  $k \geq 1$ , where  $S^k$  is the set of  $k$ -tuples of elements from  $S$ , and  $\bar{s} \in G$  is the product of the  $k$  elements in the  $k$ -tuple  $s \in S^k$ . For  $k \geq 1$ , set  $e_k = \bigoplus_{s \in S^k} \bar{s}.q$ , and let  $p'_k$  be the  $N^k$ -fold direct sum of  $q$  with itself. Then  $\text{Tr}(p'_k(t)) = N^k \varrho(t)$  and  $\text{Tr}(e_k(t)) = \sum_{s \in S^k} \bar{s}.\varrho(t)$ , for all  $t \in G$ , so  $p'_k \precsim e_k$ .

Let  $\tau$  be any trace in  $T^+(\ell^\infty(G, \mathcal{K}))$ . Then

$$N^k \tau(q) = \tau(p'_k) \leq \tau(e_k) = \sum_{s \in S^k} \tau(\bar{s}.q) \leq |S|^k \sup_{t \in G} \tau(t.q).$$

As this holds for all  $k \geq 1$ , either  $\tau(q) = 0$  or  $\sup_{t \in G} \tau(t.q) = \infty$ . Suppose that  $\{\tau(t.p)\}_{t \in G}$  is bounded. Then  $\{\tau(t.q)\}_{t \in G}$  is also bounded. Fix  $t \in G$ , and let  $\tau'$  be the trace on  $\ell^\infty(G, \mathcal{K})$  given by  $\tau'(f) = \tau(t.f)$ , for  $f \in \ell^\infty(G, \mathcal{K})^+$ . As  $\tau'$  also is bounded on the orbit  $\{s.q\}_{s \in G}$ , the argument above implies that  $0 = \tau'(q) = \tau(t.q)$ , so  $\tau(t.p) = |S|\tau(t.q) = 0$ . This proves that  $\tau$  is 0 on the orbit  $\{t.p\}_{t \in G}$ .  $\square$

Any projection satisfying the conclusion of part (iii) of the proposition above must have exponentially growing dimension with respect to any proper length function on the group  $G$  (cf. Lemma 5.7), and when  $G$  is a group with the fixed-point property for cones, then the same holds for any projection satisfying part (ii), cf. Theorem 5.8.

#### ACKNOWLEDGMENTS

I thank Nicolas Monod, Nigel Higson, Guoliang Yu, and Eduardo Scarparo for useful discussions related to this paper. I also thank the referee for suggesting improvements of the exposition and for pointing out a couple of mistakes in an earlier version of this paper.

#### REFERENCES

- [1] S. Albeverio, D. Guido, A. Ponosov, and S. Scarlatti, *Singular traces and compact operators*, J. Funct. Anal. **137** (1996), no. 2, 281–302, DOI 10.1006/jfan.1996.0047. MR1387512
- [2] B. E. Blackadar and J. Cuntz, *The structure of stable algebraically simple  $C^*$ -algebras*, Amer. J. Math. **104** (1982), no. 4, 813–822, DOI 10.2307/2374206. MR667536
- [3] B. Blackadar and D. Handelman, *Dimension functions and traces on  $C^*$ -algebras*, J. Funct. Anal. **45** (1982), no. 3, 297–340, DOI 10.1016/0022-1236(82)90009-X. MR650185
- [4] B. Blackadar and M. Rørdam, *Extending states on preordered semigroups and the existence of quasitraces on  $C^*$ -algebras*, J. Algebra **152** (1992), no. 1, 240–247, DOI 10.1016/0021-8693(92)90098-7. MR1190414
- [5] K. T. Coward, G. A. Elliott, and C. Ivanescu, *The Cuntz semigroup as an invariant for  $C^*$ -algebras*, J. Reine Angew. Math. **623** (2008), 161–193, DOI 10.1515/CRELLE.2008.075. MR2458043
- [6] J. Cuntz, *Dimension functions on simple  $C^*$ -algebras*, Math. Ann. **233** (1978), no. 2, 145–153, DOI 10.1007/BF01421922. MR0467332
- [7] J. Dixmier, *Traces sur les  $C^*$ -algèbres* (French), Ann. Inst. Fourier (Grenoble) **13** (1963), no. fasc. 1, 219–262. MR0149317
- [8] G. A. Elliott, L. Robert, and L. Santiago, *The cone of lower semicontinuous traces on a  $C^*$ -algebra*, Amer. J. Math. **133** (2011), no. 4, 969–1005, DOI 10.1353/ajm.2011.0027. MR2823868
- [9] R. M. Gillette and D. C. Taylor, *A characterization of the Pedersen ideal of  $CD0(T, B_0(H))$  and a counter-example*, Proc. Amer. Math. Soc. **68** (1978), no. 1, 59–63, DOI 10.2307/2040908. MR0461156
- [10] U. Haagerup, *Quasitraces on exact  $C^*$ -algebras are traces* (English, with English and French summaries), C. R. Math. Acad. Sci. Soc. R. Can. **36** (2014), no. 2-3, 67–92. MR3241179
- [11] J. Kellerhals, N. Monod, and M. Rørdam, *Non-supramenable groups acting on locally compact spaces*, Doc. Math. **18** (2013), 1597–1626. MR3158244
- [12] E. Kirchberg, *On the existence of traces on exact stably projectionless simple  $C^*$ -algebras*, Operator algebras and their applications (Waterloo, ON, 1994/1995), Fields Inst. Commun., vol. 13, Amer. Math. Soc., Providence, RI, 1997, pp. 171–172. MR1424961
- [13] E. Kirchberg and M. Rørdam, *Non-simple purely infinite  $C^*$ -algebras*, Amer. J. Math. **122** (2000), no. 3, 637–666. MR1759891

- [14] E. Kirchberg and M. Rørdam, *Infinite non-simple  $C^*$ -algebras: Absorbing the Cuntz algebras  $\mathcal{O}_\infty$* , Adv. Math. **167** (2002), no. 2, 195–264, DOI 10.1006/aima.2001.2041. MR1906257
- [15] H. Matui and M. Rørdam, *Universal properties of group actions on locally compact spaces*, J. Funct. Anal. **268** (2015), no. 12, 3601–3648, DOI 10.1016/j.jfa.2015.01.022. MR3341960
- [16] N. Monod, *Fixed points in convex cones*, Trans. Amer. Math. Soc. Ser. B **4** (2017), 68–93, DOI 10.1090/btran/15. MR3672898
- [17] C. Pasnicu and M. Rørdam, *Purely infinite  $C^*$ -algebras of real rank zero*, J. Reine Angew. Math. **613** (2007), 51–73, DOI 10.1515/CRELLE.2007.091. MR2377129
- [18] G. K. Pedersen,  *$C^*$ -algebras and their automorphism groups*, London Mathematical Society Monographs, vol. 14, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London–New York, 1979. MR548006
- [19] G. K. Pedersen, *Measure theory for  $C^*$  algebras*, Math. Scand. **19** (1966), 131–145, DOI 10.7146/math.scand.a-10802. MR0212582
- [20] M. Rørdam and A. Sierakowski, *Purely infinite  $C^*$ -algebras arising from crossed products*, Ergodic Theory Dynam. Systems **32** (2012), no. 1, 273–293, DOI 10.1017/S0143385710000829. MR2873171
- [21] J. Roe, *An index theorem on open manifolds. I, II*, J. Differential Geom. **27** (1988), no. 1, 87–113, 115–136. MR918459
- [22] M. Rørdam, *On the structure of simple  $C^*$ -algebras tensored with a UHF-algebra. II*, J. Funct. Anal. **107** (1992), no. 2, 255–269, DOI 10.1016/0022-1236(92)90106-S. MR1172023
- [23] J. M. Rosenblatt, *Invariant measures and growth conditions*, Trans. Amer. Math. Soc. **193** (1974), 33–53, DOI 10.2307/1996899. MR0342955
- [24] W. Rudin, *Functional analysis*, 2nd ed., International Series in Pure and Applied Mathematics, McGraw-Hill, Inc., New York, 1991. MR1157815
- [25] E. Scarparo, *Characterizations of locally finite actions of groups on sets*, Glasg. Math. J. **60** (2018), no. 2, 285–288, DOI 10.1017/S001708951700009X. MR3784048
- [26] A. Sierakowski, *The ideal structure of reduced crossed products*, Münster J. Math. **3** (2010), 237–261. MR2775364
- [27] S. Wei, *On the quasidiagonality of Roe algebras*, Sci. China Math. **54** (2011), no. 5, 1011–1018, DOI 10.1007/s11425-011-4205-x. MR2800923
- [28] S. Zhang, *Diagonalizing projections in multiplier algebras and in matrices over a  $C^*$ -algebra*, Pacific J. Math. **145** (1990), no. 1, 181–200. MR1066403
- [29] A. Żuk, *On an isoperimetric inequality for infinite finitely generated groups*, Topology **39** (2000), no. 5, 947–956, DOI 10.1016/S0040-9383(99)00043-9. MR1763958

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF COPENHAGEN, UNIVERSITETSPARKEN  
5, DK-2100, COPENHAGEN Ø, DENMARK  
Email address: rordam@math.ku.dk