

pression is $4n^2/(\log_2 n - \log_2 \log_2 n - 1)$. The result stated at the beginning of this section now follows immediately.

4. Generalizations. The corresponding problem may be considered for n by n matrices whose entries are taken from the integers $0, 1, \dots, k-1$. If an operation on such a matrix consists of adding a multiple of some row to some other row modulo k , then it can be shown that the foregoing theorem remains valid in this more general situation for any fixed value of k . In fact, the bounds in Sections 2 and 3 will still hold if $\log_2 n$ is replaced by $\log_k n$.

1. N. J. FINE & I. NIVEN, "The probability that a determinant be congruent to a (mod m)," *Bull. Amer. Math. Soc.*, v. 50, 1944, pp. 89-93.

Evaluation of $I_n(b) = 2\pi^{-1} \int_0^\infty \left(\frac{\sin x}{x}\right)^n \cos(bx) dx$ and of Similar Integrals

By Rory Thompson

Medhurst and Roberts [1] suggest the problem of evaluating $I_n(b)$ for non-integral values of b . There will be developed in this note an effective recursion scheme for such a calculation. In particular, it can be used to evaluate $I_n(0)$ for moderate values of n .

Following a suggestion by Hamming [2, p. 164], we differentiate $I_n(b)$ with respect to the parameter b , which is permissible by virtue of uniform convergence of the resulting integral for $n > 2$ and continuity of the corresponding integrand with respect to both x and b .

Thus we obtain

$$\begin{aligned} I_n'(b) &= -2\pi^{-1} \int_0^\infty \left(\frac{\sin x}{x}\right)^{n-1} \sin x \sin(bx) dx \\ &= \pi^{-1} \int_0^\infty \left(\frac{\sin x}{x}\right)^{n-1} [\cos(b+1)x - \cos(b-1)x] dx \\ &= \frac{1}{2} [I_{n-1}(b+1) - I_{n-1}(b-1)]. \end{aligned}$$

If the first expression for $I_n'(b)$ is integrated by parts there results the relation

$$\begin{aligned} I_n'(b) &= (n-1)b^{-1} 2\pi^{-1} \int_0^\infty \left(\frac{\sin x}{x}\right)^n \cos(bx) dx \\ &\quad - nb^{-1} 2\pi^{-1} \int_0^\infty \left(\frac{\sin x}{x}\right)^{n-1} \cos x \cos(bx) dx \\ &= \frac{n-1}{b} I_n(b) - \frac{n}{2b} [I_{n-1}(b+1) + I_{n-1}(b-1)]. \end{aligned}$$

Elimination of $I_n'(b)$ between these expressions yields the recurrence relation

$$I_n(b) = \frac{1}{2(n-1)} \left[(n+b)I_{n-1}(b+1) + (n-b)I_{n-1}(b-1) \right].$$

This recursion scheme for calculating $I_n(b)$ is reasonably stable, owing to the addition of comparable positive numbers and to the fact that an error in either $I_{n-1}(b+1)$ or $I_{n-1}(b-1)$ is multiplied by at most $(n+b)/2(n-1)$, which is less than unity, since $b < n$ for nonzero values of $I_n(b)$.

Starting values are readily given by the relations

$$\begin{aligned} I_3(b) &= \frac{1}{8}[(b+3)^2 - 3(b+1)^2], & 0 \leq b \leq 1 \\ &= \frac{1}{8}[(b+3)^2 - 3(b+1)^2 + 3(b-1)^2], & 1 \leq b \leq 3 \\ &= 0, & b \geq 3. \end{aligned}$$

Furthermore, we observe that negative values of b can be taken into account by use of the relation

$$I_n(-b) = I_n(b).$$

The iterative procedure just described was used to produce an 8D table of $I_n(b)$ for $n = 3(1)100$, $b = 0(0.1)9$ in approximately 0.9 minute on an IBM 7094 system at the MIT Computation Center. This table has been deposited in the UMT file of this journal.

For integral values of b the table was checked by using the recurrence formula in the form

$$I_{n-1}(b+1) = \frac{2(n-1)}{n+b} I_n(b) - \frac{n-b}{n+b} I_{n-1}(b-1).$$

This implies

$$I_{n-1}(1) = \frac{n-1}{n} I_n(0).$$

Values of $I_n(0)$ may be obtained from 10D tables in [1] and [3], so that $I_n(b)$ can be evaluated for integral values of b in a more compact form than the formulas in [1].

The use of recurrence formulas is applicable to the numerical evaluation of other integrals, including indefinite ones. One such example is the calculation of the chi-square distribution, which was accomplished by Harter [4] essentially by direct integration. In this case, integrating

$$F_n(u) = \left[2^{n/2} \Gamma\left(\frac{n}{2}\right) \right]^{-1} \int_0^u \exp\left(-\frac{x}{2}\right) x^{n/2-1} dx$$

by parts yields the recurrence relation

$$F_n(u) = F_{n-2}(u) - H_n(u),$$

where

$$H_n(u) = u^{n/2-1} / \left[2^{n/2-1} \exp\left(\frac{u}{2}\right) \Gamma\left(\frac{n}{2}\right) \right] = \frac{u}{n-2} H_{n-1}(u)$$

and

$$H_1(u) = \exp\left(-\frac{u}{2}\right),$$

$$F_2(u) = 1 - \exp\left(-\frac{u}{2}\right).$$

This procedure for evaluating $F_n(u)$ is sufficiently fast to permit a direct search for percentage points, in lieu of interpolation. Thus eleven critical levels were calculated to 5D for $n = 2(2)100$ in 1.8 minutes on an IBM 7094.

Many other types of integrals exist for which this recursion scheme is feasible, in particular, Fourier (and other) transforms similar to $I_n(b)$.

Massachusetts Institute of Technology
Cambridge, Massachusetts

1. R. G. MEDHURST & J. H. ROBERTS, "Evaluation of the integral $I_n(b) = 2/\pi \int_0^\infty ((\sin x)/x)^n \cos(bx) dx$," *Math. Comp.*, v. 19, 1965, pp. 113-117.

2. R. W. HAMMING, *Numerical Methods for Scientists and Engineers*, International Series in Pure and Applied Mathematics, McGraw-Hill, New York, 1962. MR 25 #735.

3. K. HARUMI, S. KATSURA & J. W. WRENCH, JR., "Values of $2/\pi \int_0^\infty ((\sin t)/t)^n dt$," *Math. Comp.*, v. 14, 1960, p. 379. MR 22 #12737.

4. H. L. HARTER, *New Tables of the Incomplete Gamma-Function Ratio and of Percentage Points of the Chi-Square and Beta Distributions*, U. S. Government Printing Office, Washington, D. C., 1964. MR 30 #1562.

Evaluation of Some Integrals Involving the ψ -Function

By M. L. Glasser

In the Bateman manuscript project tables, Erdelyi et al. [1] list five integrals over the unit interval involving the ψ -function (logarithmic derivative of the gamma function). The first of these is trivial, the second is easily derived by integrating by parts to derive a differential equation in terms of the parameter a . The fourth and fifth formulas are obtained by equating the imaginary and real parts of the second and the third is simply the case $a = 0$ of the fourth. The purpose of this note is to point out that this table can be easily extended by simple use of the properties of the ψ -function. For example, many convergent integrals of the form

$$I = \int_m^n f(x)\psi(x) dx,$$

where m and n are integers and $f(x)$ is a function such that $f(x) = -f(m+n-x)$, can be evaluated exactly. Thus, by symmetry

$$I = \frac{1}{2} \int_m^n f(x) \{\psi(x) - \psi(m+n-x)\} dx.$$

Now use of the relations $\psi(y+1) = \psi(y) + y^{-1}$ and $\psi(y) - \psi(1-y) = -\pi \cot \pi y$ gives immediately