On the Existence of Regions with Minimal Third Degree Integration Formulas*

By F. N. Fritsch

Abstract. A. H. Stroud has shown that $n + 1$ is the minimum possible number of nodes in an integration formula of degree three for any region in E_n . In this paper, in answer to the question of the attainability of this minimal number, we exhibit for each n a region that possesses a third degree formula with $n + 1$ nodes. This is accomplished by first deriving an $(n + 2)$ -point formula of degree three for an arbitrary region that is invariant under the group of affine transformations that leave an n -simplex fixed. The formula is then applied to a one-parameter family of such regions, and a value of the parameter is determined for which the weight at the centroid vanishes.

1. Introduction. An approximate N -point integration formula of the form

$$I(f) \equiv \int_R f d\mu = \sum_{k=1}^N A_k f(\mathbf{x}_k) + E(f),$$

with nodes $\mathbf{x}_k$ and weights A_k for a region R in n -dimensional Euclidean space E_n , where $d\mu$ is ordinary n -dimensional Lebesgue measure, is said to be of *degree m* if $E(f) = 0$ whenever f is a polynomial of degree at most m in the n variables $\mathbf{x} = (x_1, \dots, x_n)$. Such a formula is said to be *positive* if $A_k > 0$ for $k = 1, \dots, N$; *self-contained* if $\mathbf{x}_k \in R$ for $k = 1, \dots, N$.

Let S_n be the n -simplex with vertices $\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_n$ and let $\mathbf{c}$ be the centroid of S_n . Let R be any subset of E_n with positive Lebesgue measure which is invariant under the group of affine transformations that map S_n onto itself. A region that possesses this property for some S_n will be called *simplicially-symmetric*. We assume that all polynomials of degree at most three in the n variables are integrable over R . We shall consider third degree $(n + 2)$ -point formulas of the form

$$(1) \quad \int_R f d\mu = A \sum_{k=0}^n f(\mathbf{x}_k) + Bf(\mathbf{c}) + E(f),$$

where

$$\mathbf{x}_k = r\mathbf{v}_k + (1 - r)\mathbf{c} \quad (k = 0, 1, \dots, n).$$

In this paper we obtain a condition on the simplicially-symmetric region R for the existence of a formula of form (1) which is of degree three. We derive general expressions for the unknowns A , B and r , and show that the weight A must be positive, while B is unrestricted in sign. We exhibit a region for which $B = 0$, so that the formula actually involves only $n + 1$ points. Stroud [6] has shown that $n + 1$ is the smallest number of nodes possible in an integration formula of degree three

Received May 1, 1969, revised February 4, 1970.

AMS 1968 subject classifications. Primary 6555, 4144.

Key words and phrases. Approximate integration formulas, numerical integration formulas, minimal integration formulas, approximation of linear functionals, symmetric regions.

* Work performed under the auspices of the U. S. Atomic Energy Commission.

Copyright © 1971, American Mathematical Society

for any region, but it was not known in general prior to the publication of this example whether this minimal number was attained by any region. In fact a conjecture to the contrary was made by Hammer and Stroud [3].

Specifically, a one-parameter family of star-shaped simplicially-symmetric regions $S_n(d)$ is studied. There is a unique value $d = d_n^*$ for which the formula fails to exist. A result due to Mysovskikh [5] enables us to explain the failure for $n = 2$ and 3, but the significance of $S_n(d_n^*)$ is still unexplained for $n > 3$. A unique value $d = d_n \neq d_n^*$ is determined for which the number of nodes reduces to $n + 1$. The formula is positive for $d \geq d_n$. It is further shown that the formula is self-contained for all d . We have proved [2, pp. 96-104] that $S_2(d_2)$ is an isolated example, in the sense that any three-point third degree formula for a member $S_2(d)$ of the family must be of form (1) with $B = 0$. It is not known whether the same is true for $n > 2$.

The results reported here provide further evidence in support of the concluding remark of Stroud [7]: "... the minimal point formulas of degree 3 for a region are related to the group of symmetries of the region."

2. Derivation of Formulas for the Unknowns. As discussed by Hammer and Wymore [4], we may identify any two regions that are affine-equivalent for the purpose of deriving approximate integration formulas. We may accordingly take S_n to be the n -simplex

$$(2) \quad \{x \in E_n: x_i \geq 0, x_1 + x_2 + \cdots + x_n \leq 1\}.$$

THEOREM. *If we determine A , B , and r so that formula (1) is exact for the three monomials 1 , x_1^2 , x_1^3 , then the formula is of degree three.*

This theorem is proved in Stroud [8]. His proof is based on a special case of the following lemma.

LEMMA. *Let L be a linear functional which is invariant under the symmetries of S_n ; that is, if T is an affine transformation for which $TS_n = S_n$ and if $g(\mathbf{x}) = f(T\mathbf{x})$, then $L(g) = L(f)$. If $0 \leq k < n$ and $\{\alpha_i\}_{i=1}^k$ is a sequence of positive integers, then $L(x_1^{\alpha_1} \cdots x_k^{\alpha_k} x_{k+1})$ can be expressed as a linear combination of the values of L on the $k + 1$ monomials*

$$\prod_{i \leq k} x_i^{\alpha_i}, \quad \prod_{i \leq k} x_i^{\alpha_i + \delta_{ij}} \quad (j = 1, 2, \dots, k),$$

where δ_{ij} is the Kronecker delta symbol, and the empty product

$$\prod_{i \leq 0} x_i^{\alpha_i} = 1.$$

The proof depends mainly on the invariance of L under the affine transformation which interchanges vertices $\mathbf{v}_0 (= 0)$ and $\mathbf{v}_{k+1} (= (k + 1)\text{th unit vector})$ of S_n . Use of the Lemma allows us to express $L(f)$, where f is any polynomial of degree at most three, as a linear combination of $L(1)$, $L(x_1^2)$, $L(x_1^3)$. Noting that both sides of (1) have the required invariance proves the Theorem.

By application of the Theorem, a necessary and sufficient condition for (1) to be a formula of degree three for the simplicially-symmetric region R is that A , B , and r be solutions of the following system of nonlinear equations:

$$(3) \quad (n + 1)A + B = I_0 \equiv I(1);$$

$$(4) \quad \frac{1 + nr^2}{n + 1} A + \frac{1}{(n + 1)^2} B = I_2 \equiv I(x_1^2);$$

$$(5) \quad \frac{1 + 3nr^2 + n(n - 1)r^3}{(n + 1)^2} A + \frac{1}{(n + 1)^3} B = I_3 \equiv I(x_1^3).$$

We may use Eq. (3) to eliminate B from the other two equations

$$(4') \quad n(n + 1)Ar^2 = J_2 \equiv (n + 1)^2 I_2 - I_0;$$

$$(5') \quad n(n + 1)[(n - 1)r + 3]Ar^2 = J_3 \equiv (n + 1)^3 I_3 - I_0.$$

Since

$$J_2 = \int_R [(n + 1)x_1 - 1]^2 d\mu > 0,$$

necessary conditions for the existence of a solution are $A > 0$ and $r \neq 0$. Substituting Eq. (4') into (5') and solving for r , we obtain

$$(6) \quad r = \frac{J_3 - 3J_2}{(n - 1)J_2} \equiv \frac{D}{(n - 1)J_2}.$$

Thus a necessary and sufficient condition on the region R for the existence of a third degree formula of form (1) is that

$$(7) \quad D = \int_R [(n + 1)x_1 - 1]^3 d\mu \neq 0.$$

If R satisfies Eq. (7), then there exists a unique formula (1) given by Eq. (6),

$$(8) \quad A = \frac{J_2}{n(n + 1)r^2} = \frac{(n - 1)^2 J_2^3}{n(n + 1)D^2},$$

and

$$(9) \quad B = I_0 - (n + 1)A = \frac{nI_0 D^2 - (n - 1)^2 J_2^3}{nD^2} \equiv \frac{P}{nD^2}.$$

While A is always positive, the sign of B is determined by that of P . We observe that, since $J_2 > 0$, $P = 0$ implies $D \neq 0$. Thus if we can find a region R for which $P = 0$ we will have an example of a region that possesses a positive formula of degree three with the minimum possible number of nodes.

3. A Family of Star-Shaped Simplicially-Symmetric Regions. In the remainder of this paper we shall apply these formulas to a family of star-shaped simplicially-symmetric regions $S_n(d)$ over which the necessary integrals can be computed, in order to prove the existence of a region for which this minimal number of nodes is attained. For notational clarity, if Q is one of the quantities introduced in the preceding section, then $Q_n(d)$ will denote the value of this quantity for $S_n(d)$. The order of presentation will be as follows:

1. Derive formulas for $A_n(d)$, $B_n(d)$, and $r_n(d)$ as rational functions of the parameter d , with coefficients which are polynomials in the dimension n .

2. Prove the existence of a unique value d_n of d for which $P_n(d_n) = 0$, so that $S_n(d_n)$ possess a formula of form (1) with $B = 0$.

3. Prove that there exists a unique value d_n^* of d for which $D_n(d_n^*) = 0$, and consequently formula (1) fails to exist for $S_n(d_n^*)$.

4. Finally, show that the formula is self-contained for all $d > 0$, $d \neq d_n^*$. (Note that formula (1) need not be self-contained, in general.)

The regions $S_n(d)$ are defined as follows. Let S_n be the n -simplex with vertices $v_0, v_1, \dots, v_n$. Let F_k be the face of S_n that does not contain the vertex v_k and let c_k be the centroid of F_k . Let $d > 0$ and define the points $u_k(d)$ by:

$$u_k(d) = dc_k + (1 - d)c \quad (k = 0, 1, \dots, n).$$

Let $S_{nk}(d)$ be the pyramid (simplex) with base F_k and vertex $u_k(d)$. ($S_{nk}(d)$ degenerates to the face F_k when $d = 1$.) Define

$$\begin{aligned} S_n(d) &= S_n \cup \left(\bigcup_{k=0}^n S_{nk}(d) \right) \quad \text{if } d \geq 1; \\ &= S_n - \left(\bigcup_{k=0}^n S_{nk}(d) \right) \quad \text{if } 0 < d < 1. \end{aligned}$$

Clearly, $S_n(1) = S_n$. For $d > 1$, this star-shaped polyhedral region can be visualized as the result of "pushing out" the center of each face of S_n . This family of regions was previously studied for $n = 2$ by De Vogelaere [1], who knew of the existence of $S_2(d_2)$. Since $u_k(d)$ lies on the line determined by v_k and c , the set $\{u_0(d), u_1(d), \dots, u_n(d)\}$ is invariant under any affine transformation that leaves S_n invariant, and $S_n(d)$ is simplicially-symmetric.

4. Application of the Formulas to $S_n(d)$. Let us now specialize S_n to the simplex (2). If we use the natural simplicial decomposition of $S_n(d)$, a straightforward but extremely tedious computation [2, pp. 72-75, 151-159] yields the following values for the needed monomial integrals:

$$I_{0n}(d) = \frac{d}{n!}.$$

$$I_{2n}(d) = \frac{2d}{n(n+1)^2(n+2)!} [d^2 + (n-1)d + n^2(n+2)],$$

$$\begin{aligned} I_{3n}(d) &= \frac{6d}{n^2(n+1)^3(n+3)!} [(1-n)d^3 + (5n-1)d^2 \\ &\quad + n(n-1)(n+4)d + n^2(n^3 + 3n^2 + 2n - 2)]. \end{aligned}$$

We may now compute $A_n(d)$, $B_n(d)$, and $r_n(d)$.

$$J_{2n}(d) = (n+1)^2 I_{2n}(d) - I_{0n}(d) = \frac{d}{n(n+2)!} \psi_n(d),$$

where

$$(10) \quad \psi_n(d) = 2d^2 + 2(n-1)d + n(n-1)(n+2).$$

$$J_{3n}(d) = (n+1)^3 I_{3n}(d) - I_{0n}(d) = \frac{d}{n^2(n+3)!} \chi_n(d),$$

where

$$\chi_n(d) = 6(1 - n)d^3 + 6(5n - 1)d^2 + 6n(n - 1)(n + 4)d + n^2(n - 1)(5n^2 + 17n + 18).$$

$$(11) \quad D_n(d) = J_{3n}(d) - 3J_{2n}(d) = \frac{2(n - 1)d}{n(n^2 + 3)!} \phi_n(d),$$

where

$$(12) \quad \phi_n(d) = 3d^3 + 3(n - 1)d^2 - 3nd - n^3(n + 1).$$

From Eq. (6) we obtain:

$$(13) \quad r_n(d) = \frac{D_n(d)}{(n - 1)J_{2n}(d)} = \frac{-2}{n(n + 3)} \frac{\phi_n(d)}{\psi_n(d)}.$$

Provided $D_n(d) \neq 0$, we similarly obtain from Eqs. (8) and (9) the following:

$$(14) \quad A_n(d) = \frac{(n + 3)^2 d}{4(n + 1)(n + 2)!} \frac{[\psi_n(d)]^3}{[\phi_n(d)]^2}.$$

$$(15) \quad B_n(d) = \frac{P_n(d)}{n[D_n(d)]^2} = \frac{d}{4(n + 2)!} \frac{F_n(d)}{[\phi_n(d)]^2},$$

where

$$(16) \quad F_n(d) = 4(n + 1)(n + 2)[\phi_n(d)]^2 - (n + 3)^2[\psi_n(d)]^3.$$

5. A Region that Possesses an $(n + 1)$ -Point Formula of Degree Three. We have seen that, whenever $D_n(d) \neq 0$, $S_n(d)$ possesses a third degree formula of the form (1) with $r = r_n(d)$, $A = A_n(d)$, and $B = B_n(d)$ given by Eqs. (13), (14), and (15), respectively. From Eq. (15) we see that $B_n(d)$, the weight at the centroid, is negative, zero, or positive according as the polynomial $F_n(d)$ is negative, zero, or positive. Hammer and Stroud [3] showed that $B_n(1)$ is negative, while the leading coefficient of $F_n(d)$ is $4n(7n + 15) > 0$. Hence there exists a $d_n > 1$ such that $F_n(d_n) = 0$. As remarked above, $D_n(d_n) \neq 0$. Thus, $S_n(d_n)$ possesses a positive $(n + 1)$ -point formula of degree three, providing the first known example of a region with the minimal number of nodes.

A consideration based on the Descartes rule of signs and the use of a polynomial root finder to compute the real and complex roots of the polynomials in n that appear in the coefficients of $F_n(d)$ in Eq. (16) enables us to conclude [2, pp. 77-80] that $F_n(d)$ possesses a single positive root and d_n is uniquely determined for $n \geq 3$. The case $n = 2$ deserves special attention. $F_2(d)$ has two positive roots, one between 0 and 1 and the other greater than 1. One may easily verify that $F_2(4/d) = (2/d)^6 F_2(d)$, so that F_2 is "reciprocal" in the sense that if d is a root, so is $4/d$. The geometrical significance of this property of F_2 is that $S_2(4/d)$ is similar to $S_2(d)$. Thus, the two positive roots of F_2 must result in similar figures, and d_2 is essentially unique. The two reciprocal figures $S_2(d_2)$ and $S_2(4/d_2)$, based on an equilateral triangle, are depicted in Fig. 1.

6. The Exceptional Member of the Family. From Eq. (12) we see by the Descartes rule of signs that ϕ_n has exactly one positive root, which we shall call d_n^* .

FIGURE 1. $S_2(d_2)$ and (shaded) $S_2(4/d_2)$, two planar regions which possess three-point formulas of degree three. The x 's locate the nodes for $S_2(d_2)$.

From Eq. (11) we see that $D_n(d_n^*) = 0$ and $D_n(d) \neq 0$ for $d \neq d_n^*$. In other words, there is only one member $S_n(d_n^*)$ of the family for which a formula of the desired form fails to exist. Since $J_{2n}(d) > 0$ for all $d > 0$, Eq. (13) shows that the sign of $r_n(d)$ is the same as that of $D_n(d)$, and that $r_n(d_n^*) = 0$. Thus, the formula fails to exist because all $n + 2$ nodes coincide, and their associated weights become infinite.

For $n = 2$, $d_2^* = 2$ is the unique positive value of d for which $4/d = d$. When based on an equilateral triangle, $S_2(2)$ is a regular hexagon. It is the only centrally-symmetric member of the family. As Mysovskikh [5] shows, there exist infinitely many third degree four-point formulas for $S_2(2)$, but none of them has c as one of its nodes. A formula of form (1) *must* fail to exist, because the four nodes are not centrally-symmetric with respect to c . For $n = 3$, $d_3^* = 3$. Again, $S_3(3)$ is the only centrally-symmetric member of the family. (When based on a regular tetrahedron, it is a cube.) In this case, since the Mysovskikh result shows that six is the minimal number of nodes for third degree formulas on $S_3(3)$, a 5-point formula *must* fail to exist. For $n > 3$, $d_n^* > n$. We remark that in this case there is no centrally-symmetric member of the family, and the significance of the nonexistence of such a formula for $S_n(d_n^*)$ is still not known for $n > 3$.

We know from the general theory of Section 2 that $d_n \neq d_n^*$. In fact, one can show [2, pp. 82–85] that for all $n \geq 2$,

$$d_n^* < n^{4/3} < d_n.$$

This separation property is illustrated in Fig. 5-1 of [2, p. 95].

7. Self-Containment of the Formulas. We show further that *the formula is self-contained for all $d > 0$ ($d \neq d_n^*$) and all $n \geq 2$* . Note that this does not follow from the general theory. It is evident that $u_k(d)$ and x_k both lie on the line determined by v_k and c . $u_k(d)$ is on the opposite side of c from v_k . If $r_n(d) > 0$, x_k is on the same side as v_k , and the condition for self-containment in this case is

$$(17) \quad r_n(d) \leq 1 \quad \text{for } 0 < d < d_n^*.$$

If $r_n(d) < 0$, x_k is on the same side as $u_k(d)$, and the condition for self-containment is

$$(18) \quad \rho_n(d) \equiv -\frac{n}{d} r_n(d) \leq 1 \quad \text{for } d > d_n^*.$$

Condition (17) can be verified by differentiating Eq. (13) and observing that $r'_n(d) < 0$ for all $d > 0$, $n \geq 2$. Since one can easily see that $r_n(0) < 1$, (17) holds with strict inequality. From Eqs. (10), (12), (13), condition (18) can be shown to hold with strict inequality by observing that $2\phi_n(d) < (n+3)d\psi_n(d)$ for all $d > d_n^*$, $n \geq 2$. We remark that numerical computations have shown that $\rho_n(d_n) < .5$ for $n = 2(1)50$, and indicate that $\rho_n(d_n)$ decreases monotonically to zero as $n \rightarrow \infty$.

Acknowledgments. The author wishes to express his deep gratitude to Professor R. Sherman Lehman for the direction of his thesis research and to A. C. Hindmarsh and R. E. von Holdt for their suggestions concerning the organization of this paper.

Numerical Analysis Group
Lawrence Radiation Laboratory
University of California
Livermore, California 94550

1. R. J. DE VOGELAERE, Private communication.
2. F. N. FRITSCH, *On Minimal Positive and Self-Contained Multi-Dimensional Integration Formulas*, Ph.D. Thesis, University of California, Berkeley, Calif., 1969. (Available as UCRL-50600, Lawrence Radiation Laboratory, Livermore, Calif.)
3. P. C. HAMMER & A. H. STROUD, "Numerical integration over simplexes," *Math. Tables Aids Comput.*, v. 10, 1956, pp. 137-139. MR 19, 177.
4. P. C. HAMMER & A. W. WYMORE, "Numerical evaluation of multiple integrals. I," *Math. Tables Aids Comput.*, v. 11, 1957, pp. 59-67. MR 19, 323.
5. I. P. MYSOVSKIKH, "Proof of the minimality of the number of nodes in the cubature formula for a hypersphere," *Ž. Vyčisl. Mat. i Mat. Fiz.*, v. 6, 1966, pp. 621-630 = *U.S.S.R. Comput. Math. and Math. Phys.*, v. 6, no. 4, 1966, pp. 15-27. MR 33 #8104.
6. A. H. STROUD, "Quadrature methods for functions of more than one variable," *Ann. New York Acad. Sci.*, v. 86, 1960, pp. 776-791. MR 22 #10179.
7. A. H. STROUD, "Numerical integration formulas of degree 3 for product regions and cones," *Math. Comp.*, v. 15, 1961, pp. 143-150. MR 22 #12717.
8. A. H. STROUD, "A fifth degree integration formula for the n -simplex," *SIAM J. Numer. Anal.*, v. 6, 1969, pp. 90-98.