

Calculation of the Ramanujan τ -Dirichlet Series

By Robert Spira

Abstract. A method is found for calculating the Ramanujan τ -Dirichlet series $F(s)$. An inequality connecting points symmetric with the critical line, $\sigma = 6$, is proved, and a table is given for $\Gamma(s)F(s)$ for $\sigma = 6.0, 6.5, t = 0(.25)16$. Two zeros are found in $0 < t \leq 16$; they appear to be simple and on the critical line.

The Ramanujan τ -function (Hardy [1]) is defined by

$$(1) \quad g(x) = x \prod_{k=1}^{\infty} (1 - x^k)^{24} = \sum_{n=1}^{\infty} \tau(n)x^n,$$

and we consider the series

$$(2) \quad F(s) = \sum_{n=1}^{\infty} \tau(n)/n^s.$$

We have

$$(3) \quad \Gamma(s)F(s) = \int_0^{\infty} y^{s-1}g(e^{-y}) dy$$

and

$$(4) \quad \Gamma(12 - s)F(12 - s) = (2\pi)^{2(6-s)}\Gamma(s)F(s).$$

Further,

$$(5) \quad g(e^{-y}) = (2\pi/y)^{12}g(e^{-4\pi^2/y}).$$

To calculate (3), we break the integral at 1. We have

$$(6) \quad \begin{aligned} \int_1^{\infty} y^{s-1}g(e^{-y}) dy &= \int_1^{\infty} y^{s-1} \sum_{n=1}^{\infty} \tau(n)e^{-ny} dy \\ &= \sum_{n=1}^{\infty} \tau(n) \int_1^{\infty} y^{s-1}e^{-ny} dy \\ &= \sum_{n=1}^{\infty} \tau(n)n^{-s} \int_n^{\infty} u^{s-1}e^{-u} du \\ &= \sum_{n=1}^{\infty} \tau(n)n^{-s}\Gamma(s, n) \end{aligned}$$

and, using (5),

Received June 29, 1972.

AMS (MOS) subject classifications (1970). Primary 10H10, 65A05.

Copyright © 1973, American Mathematical Society

$$\begin{aligned}
 \int_0^1 y^{s-1} g(e^{-y}) dy &= \int_0^1 y^{s-1} (2\pi/y)^{12} \left(\sum_{n=1}^{\infty} \tau(n) e^{-4\pi^2 n/y} \right) dy \\
 (7) \qquad \qquad \qquad &= (2\pi)^{12} \sum_{n=1}^{\infty} \tau(n) \int_1^{\infty} u^{11-s} e^{-4\pi^2 nu} du \\
 &= (2\pi)^{2(s-6)} \sum_{n=1}^{\infty} \tau(n) n^{s-12} \Gamma(12-s, 4\pi^2 n).
 \end{aligned}$$

In order to truncate the series in (6) and (7), we need estimates for $\tau(n)$ and $\Gamma(s, y)$, where $|\operatorname{Re} s - 6| \leq \frac{1}{2}$. It has been proved by Rankin that $\tau(n) = O(n^{29/5})$, but that, for an infinity of n , $\tau(n) \geq n^{11/2}$. We content ourselves here with

LEMMA 1. For $n \geq 1$, $|\tau(n)| \leq 51,000n^8$.

Proof. For $0 < x < 1$, using Hardy [1, p. 156],

$$\sum \tau(n)x^n = x \left\{ \prod_{n=1}^{\infty} (1 - x^n)^{24} \right\} = x(1 - 3x + 5x^3 - 7x^6 + \dots)^8$$

and defining $\epsilon(n)$ by

$$\sum \epsilon(n)x^n = x(1 + 3x + 5x^3 + 7x^6 + \dots)^8,$$

we have $0 \leq |\tau(n)| \leq \epsilon(n)$. Thus,

$$|\tau(n)| x^n \leq \sum |\tau(n)| x^n \leq x \left\{ \sum_{j=0}^{\infty} (2j+1)x^{j(j+1)/2} \right\}^8.$$

Now, write $x = e^{-y}$ and take $x = 1 - 1/n$. Then, for $n \geq 25$, we have $0 < y < .1$, and

$$\sum_{j=0}^{\infty} (2j+1)x^{j(j+1)/2} \leq 1 + 2 \sum_{j=1}^{\infty} (j + \frac{1}{2})x^{j^2/2} \leq 1 + 3 \sum_{j=1}^{\infty} je^{-j^2 y/2}.$$

Now,

$$(d/dt)(te^{-vt^2/2}) = e^{-vt^2/2}(1 - yt^2),$$

which is zero for $t^2 = 1/y < 1$. Hence,

$$\begin{aligned}
 1 + 3 \sum_{j=1}^{\infty} je^{-j^2 y/2} &\leq 1 + 3e^{-y/2} + 3 \int_1^{\infty} te^{-t^2 y/2} dt \\
 &= \frac{1}{y} \{ 3e^{-y/2} + y + 3ye^{-y/2} \} \leq \frac{1}{y} \{ 3 + 4y \} \leq \frac{3.4}{y}.
 \end{aligned}$$

Thus, for $n \geq 25$,

$$|\tau(n)| x^n \leq x(3.4/y)^8,$$

or

$$\begin{aligned}
 |\tau(n)| &\leq (1 - 1/n)^{-n} \cdot (1 - 1/n)(3.4)^8 \{ -\log(1 - 1/n) \}^{-8} \\
 &\leq 2.84(3.4)^8 \{ 1/n + 1/2n^2 + 1/3n^3 + \dots \}^{-8} \leq 51,000n^8,
 \end{aligned}$$

so we have the result for $n \geq 25$. For $1 \leq n < 25$, the result follows from the tables of $\tau(n)$. We have used the fact that $(1 - 1/n)^n$ is increasing. We next have

TABLE I. $\Gamma(s) \cdot \sum \tau(n)n^{-s}$

t	$\sigma = 6.0$		$\sigma = 6.5$	
	Re ΓF	Im ΓF	Re ΓF	Im ΓF
0.00	95.054741	0.0	241.635216	0.0
0.25	84.897790	42.006402	214.297736	109.802252
0.50	56.861760	74.510262	139.175001	193.401836
0.75	17.608952	90.396027	35.036781	231.495179
1.00	-23.713754	86.667437	- 72.600156	216.603934
1.25	-57.823930	65.064019	- 158.323399	154.660635
1.50	-77.603739	31.426603	- 203.406452	62.899797
1.75	-79.731717	- 5.966438	- 200.276555	- 35.205315
2.00	-65.241565	- 38.591744	- 153.721738	- 116.389995
2.25	-38.925282	- 59.781593	- 78.700101	- 163.541602
2.50	- 7.825106	- 66.178287	4.392353	- 169.519207
2.75	20.707071	- 58.209598	75.468918	- 137.964276
3.00	40.826033	- 39.552406	119.825714	- 81.159823
3.25	49.417013	- 15.834251	131.207861	- 15.838631
3.50	46.472398	6.994358	112.251787	41.665388
3.75	34.630023	24.144535	72.510735	79.501835
4.00	18.111104	32.958140	24.936651	92.601304
4.25	1.427858	33.182355	- 17.985120	82.813237
4.50	- 11.760975	26.588121	- 47.326988	57.184320
4.75	- 19.369256	16.145104	- 59.311148	25.163485
5.00	- 21.063762	5.051909	- 55.227435	- 4.302722
5.25	- 17.961022	- 4.093386	- 40.013489	- 24.958162
5.50	- 12.019054	- 9.790036	- 20.156443	- 34.241392
5.75	- 5.351287	- 11.736869	- 1.642814	- 33.071710
6.00	0.336026	- 10.600628	11.479144	- 24.742209

LEMMA 2. For $|\sigma - 6| \leq \frac{1}{2}$, $s = \sigma + it$ and $x \geq 10$,

$$|\Gamma(s, x)| \leq 1.87x^{\sigma-1}e^{-x}.$$

Proof. From [1, 6.5.32], we have

$$\begin{aligned}
 |\Gamma(s, x)| &\leq \int_x^\infty e^{-t} t^{\sigma-1} dt \leq \Gamma(\sigma, x) \\
 &= x^{\sigma-1} e^{-x} \left[1 + \frac{5.5}{x} + \frac{(5.5)(4.5)}{x^2} + \frac{(5.5)(4.5)(3.5)}{x^3} \right. \\
 &\quad \left. + \frac{(5.5)(4.5)(3.5)(2.5)}{x^4} + \frac{(5.5)(4.5)(3.5)(2.5)(1.5)}{x^5} + R_5 \right]
 \end{aligned}$$

where $|R_5| \leq (5.5)(4.5)(3.5)(2.5)(1.5)(.5)/x^6$.

t	$\sigma = 6.0$		$\sigma = 6.5$	
	Re ΓF	Im ΓF	Re ΓF	Im ΓF
6.25	4.060183	- 7.590891	17.659110	- 13.435161
6.50	5.606613	- 4.004048	17.576528	- 2.870248
6.75	5.358784	- 0.868649	13.326154	4.561652
7.00	4.017186	1.23 7244	7.432091	8.055152
7.25	2.315988	2.193480	2.021947	8.125970
7.50	0.817445	2.218105	- 1.641478	6.047054
7.75	- 0.180696	1.692174	- 3.234755	3.237683
8.00	- 0.632869	0.996442	- 3.148548	0.802248
8.25	- 0.665268	0.403751	- 2.121957	- 0.686960
8.50	- 0.473370	0.040814	- 0.892740	- 1.172424
8.75	- 0.235696	- 0.092358	0.032502	- 0.940076
9.00	- 0.064737	- 0.071203	0.454587	- 0.398768
9.25	0.002628	0.009194	0.431195	0.101857
9.50	- 0.013714	0.074908	0.159194	0.368353
9.75	- 0.067839	0.091012	- 0.147665	0.372226
10.00	- 0.115695	0.058884	- 0.342453	0.198228
10.25	- 0.131613	0.001489	- 0.374718	- 0.028725
10.50	- 0.111796	- 0.053749	- 0.274697	- 0.204745
10.75	- 0.068324	- 0.087460	- 0.113729	- 0.278673
11.00	- 0.019278	- 0.093322	0.036463	- 0.253193
11.25	0.019980	- 0.076408	0.130413	- 0.165736
11.50	0.041557	- 0.047840	0.156643	- 0.063219
11.75	0.045459	- 0.019011	0.130285	0.017928
12.00	0.037030	0.002500	0.078600	0.060969

Hence, for $x \geq 10$,

$$|\Gamma(s, x)| \leq x^{\sigma-1} e^{-x} \{1 + .550 + .222 + .071 + .018 + .003 + .002\},$$

and the lemma follows. We next estimate the truncation error for the series (6) and (7).

We have, for $N \geq 8$,

$$\begin{aligned} \left| \sum_{n=N+1}^{\infty} \tau(n) n^{-s} \Gamma(s, n) \right| &\leq \sum_{n=N+1}^{\infty} 51,000 n^8 n^{-\sigma} n^{\sigma-1} e^{-n} (1.87) \\ &\leq 9.6 \times 10^4 \sum_{n=N+1}^{\infty} n^7 e^{-n} \\ &\leq 9.6 \times 10^4 \int_N^{\infty} x^7 e^{-x} dx \\ &= 9.6 \times 10^4 e^{-N} \{N^7 + 7N^6 + 7 \cdot 6N^5 + \dots + 7!\} \\ &\leq 3.2 \times 10^5 N^7 e^{-N}, \end{aligned}$$

t	$\sigma = 6.0$		$\sigma = 6.5$	
	Re ΓF	Im ΓF	Re ΓF	Im ΓF
12.25	0.023387	0.013479	0.027219	0.067722
12.50	0.010481	0.015706	- 0.008103	0.051173
12.75	0.001620	0.012488	- 0.023197	0.026800
13.00	- 0.002558	0.007458	- 0.022402	0.006247
13.25	- 0.003192	0.003163	- 0.013631	- 0.005196
13.50	- 0.002047	0.000682	- 0.004077	- 0.007832
13.75	- 0.000630	- 0.000088	0.002079	- 0.005069
14.00	0.000232	0.000158	0.003914	- 0.000801
14.25	0.000377	0.000668	0.002673	0.002293
14.50	0.000052	0.000961	0.000318	0.003261
14.75	- 0.000384	0.000896	- 0.001586	0.002497
15.00	- 0.000671	0.000572	- 0.002344	0.000984
15.25	- 0.000717	0.000181	- 0.002045	- 0.000375
15.50	- 0.000570	- 0.000123	- 0.001193	- 0.001102
15.75	- 0.000341	- 0.000272	- 0.000318	- 0.001170
16.00	- 0.000131	- 0.000279	0.000254	- 0.000828

so $N \geq 70$ will give a truncation estimate of $< 10^{-10}$. Using $|\tau(n)n^{-\sigma}| \leq 2$ in this range, we can take $N \geq 51$. For (7), we estimate

$$\begin{aligned}
 \left| \sum_{n=N+1}^{\infty} \tau(n)n^{s-12}\Gamma(12-s, 4\pi^2 n) \right| &\leq \sum_{n=N+1}^{\infty} 51,000n^8 n^{\sigma-12}(1.87)(4\pi^2 n)^{11-\sigma} e^{-4\pi^2 n} \\
 &\leq 9.6 \times 10^4 \sum_{n=N+1}^{\infty} n^7 e^{-4\pi^2 n} (4\pi^2)^{11-\sigma} \\
 &\leq 9.6 \times 10^4 (4\pi^2)^{5.5} \int_N^{\infty} x^7 e^{-4\pi^2 x} dx \\
 &= 9.6 \times 10^4 (4\pi^2)^{5.5-8} \int_{4\pi^2 N}^{\infty} u^7 e^{-u} du \\
 &\leq 3.2 \times 10^5 (4\pi^2)^{5.5-8+7} N^7 e^{-4\pi^2 N} \\
 &\leq 5 \times 10^{12} N^7 e^{-4\pi^2 N}
 \end{aligned}$$

which is $\leq 10^{-10}$ for $N \geq 4$. Thus, for this series, we need only a very few terms.

For convenience in the calculation, one calculates the real and imaginary parts of

$$\int_n^{n+1} u^{s-1} e^{-u} du, \quad n = 1, 2, \dots,$$

using Simpson's rule, and sums the results to get $\Gamma(s, n)$, $n = 1, 2, \dots$. The number of steps for each integration varies directly as t , and also the Γ factor shrinks the size of the answer as t increases, so the method is only suitable for a restricted range of t .

Table I gives the rounded real and imaginary parts of $\Gamma(s)F(s)$ for $\sigma = 6.0, 6.5$

and $t = 0(.25)16$. Two zeros $6 + iz_j$ were found in this range

$$z_1 = 9.22238, \quad z_2 = 13.90755,$$

and the change of argument around the rectangle with vertices $5.5, 6.5, 6.5 + 16i, 5.5 + 16i$ indicates that these are the only zeros in this region. A rigorous machine proof could be given by taking a mesh sufficiently fine. The third zero appears to be

$$z_3 = 17.443.$$

In principle, the method can be used as high up in the critical strip as one desires, but in practice, the problem is that the calculation of the incomplete Γ -function starts to take too much time. J. B. Rosser [5] studied this problem. Several other possible methods of calculation were studied. What one would like is to take a number of terms of $\sum \tau(n)n^{-s}$ and find a correction term, or to use some forms of the approximate functional equation (Apostol and Sklar [3]). The present author could not see how to do it.

We close this paper with the following

THEOREM. For $6 < \sigma < 6.5$, $|t| \geq 4.35$, if $F(s) \neq 0$, then

$$|F(12 - s)| > |F(s)|.$$

Proof. We proceed as in Schoenfeld and Dixon [4]. Write, from (2) and (4),

$$F(12 - s) = f(s)F(s),$$

$$f(s) = (2\pi)^{2(6-s)} \Gamma(s)/\Gamma(12 - s).$$

Taking $s_0 = 6 + it$, we have $|f(s_0)| = 1$ for all t . Setting $h(s) = \log |g(s)/g(s_0)|$, it suffices to prove $h(s) > 0$ for $6 < \sigma < 6.5$ and $t \geq 4.35$. We have

$$\begin{aligned} h(s) &= \log \left| \frac{(2\pi)^{2(6-s)}}{(2\pi)^{2it}} \frac{\Gamma(s)}{\Gamma(12-s)} \frac{\Gamma(6+it)}{\Gamma(6-it)} \right| \\ &= -2(\sigma - 6) \log 2\pi + \log |\Gamma(s)| - \log |\Gamma(12 - s)| \\ &= -2(\sigma - 6) \log 2\pi + 2(\sigma - 6)(\partial/\partial\sigma) \log |\Gamma(\sigma + it)|_{\sigma=\sigma_1} \end{aligned}$$

where $5.5 \leq \sigma_1 \leq 6.5$. Thus

$$\frac{h(s)}{2(\sigma - 6)} = \frac{\partial}{\partial\sigma} \log |\Gamma(\sigma + it)|_{\sigma=\sigma_1} - \log 2\pi,$$

and, as in [4],

$$\begin{aligned} \frac{\partial}{\partial\sigma} \log |\Gamma(\sigma + it)| &\geq \log |s| - \frac{1}{2|s|} - \frac{1}{12|s|^2} - \frac{1}{8} \int_0^\infty \frac{dx}{\{(\sigma + x)^2 + t^2\}^2} \\ &\geq \log |5.5 + it| - \frac{1}{14} - \frac{1}{12 \cdot 7^2} - \frac{1}{8} \int_0^\infty \frac{dx}{(5.5 + x)^2} \\ &= \log |5.5 + it| - \frac{1}{12} - \frac{1}{588} - \frac{1}{44}. \end{aligned}$$

Thus, the result holds if

$$\log |5.5 + it| > \log 2\pi + .108$$

or if $\log |5.5 + it| > \log 7$, or $t^2 > 18.75$, or $t \geq 4.35$, and the proof is complete. Berndt [6] proved the result for $|t| \geq 6.8$, $\sigma > 6$. Empirically, for $\sigma = 6.5$, the inequality appears to fail for $0 \leq t \leq 3$ and appears to hold for $t \geq 3.25$.

Department of Mathematics
Michigan State University
East Lansing, Michigan 48823

1. G. H. HARDY, *Ramanujan: Twelve Lectures on Subjects Suggested by His Life and Work*, Chelsea, New York, 1959. MR 21 #4881.
2. M. ABRAMOWITZ & I. A. STEGUN (Editors), *Handbook of Mathematical Functions, With Formulas, Graphs and Mathematical Tables*, Nat. Bur. Standards Appl. Math. Series, 55, Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964. MR 29 #4914.
3. T. M. APOSTOL & A. SKLAR, "The approximate functional equation of Hecke's Dirichlet series," *Trans. Amer. Math. Soc.*, v. 86, 1957, pp. 446–462. MR 20 #838.
4. R. D. DIXON & L. SCHOENFELD, "The size of the Riemann zeta-function at places symmetric with respect to the point $\frac{1}{2}$," *Duke Math. J.*, v. 33, 1966, pp. 291–292. MR 32 #7517.
5. J. B. ROSSER, "Explicit remainder terms for some asymptotic series," *J. Rational Mech. Anal.*, v. 4, 1955, pp. 595–626. MR 17, 360.
6. B. C. BERNDT, "On the zeros of a class of Dirichlet series. I," *Illinois J. Math.*, v. 14, 1970, pp. 244–258. MR 42 #3261.