On the Vanishing of the Iwasawa Invariant μ_p for $p < 8000$

By Wells Johnson

Abstract. The irregular primes less than 8000 are computed, and it is shown that the Iwasawa invariant $\mu_p = 0$ for all primes $p < 8000$.

1. Introduction. Let $p = 2m + 1$ be an odd prime number, and let F_n ($n \geq 0$) be the cyclotomic field of p^{n+1} th roots of unity over the rational field $\mathbb{Q}$. Let $p^{e(n)}$ be the exact power of p which divides the class number h_n of F_n . Iwasawa [4] has shown that there exist integers $\mu_p \geq 0$, $\lambda_p \geq 0$ and ν_p such that

$$e(n) = \mu_p p^n + \lambda_p n + \nu_p$$

for all n sufficiently large. Iwasawa and Sims [7] have computed the cyclotomic invariants μ_p , λ_p , and ν_p for all primes $p \leq 4001$. In particular, they showed that $\mu_p = 0$ for every $p \leq 4001$.

In this paper, we derive some conditions on p which are necessary if $\mu_p > 0$. Computations have been performed which show that these conditions are not satisfied for any prime p , $p < 8000$, so that $\mu_p = 0$ for all such primes. In particular, the computations of μ_p in [7] have been verified, although these appear to have been incomplete, since they were based upon the incomplete tables in the first paper of [8].

The author wishes to acknowledge the assistance of his colleagues, R. B. S. Brooks and M. W. Curtis, in the preparation of the computer programs.

2. Notation. Let $\mathbb{Z}_p$ denote the ring of p -adic integers. Let U be the group of units of $\mathbb{Z}_p$ and let V denote the cyclic subgroup of U consisting of the $(p - 1)$ st roots of unity.

Any x in $\mathbb{Z}_p$ has the p -adic representation

$$x = \sum_{k=0}^{\infty} x_k p^k,$$

where the x_k are rational integers satisfying $0 \leq x_k < p$ for $k \geq 0$. In the following, the subscript notation x_k will always denote the coefficient of p^k in the p -adic expansion of the p -adic integer x . If x is given as above, we define the truncated sum $s_n(x)$ by

$$s_n(x) = \sum_{k=0}^n x_k p^k.$$

Received June 15, 1972.

AMS (MOS) subject classifications (1970). Primary 12A35, 12A50; Secondary 10A40.

Key words and phrases. Cyclotomic fields, class numbers, irregular primes, Γ -extensions, cyclotomic invariants, Fermat's Last Theorem.

Copyright © 1973, American Mathematical Society

Thus, $x \equiv s_n(x) \pmod{p^{n+1}}$ and $0 \leq s_n(x) < p^{n+1}$ for all $n \geq 0$.

For any rational integer a , $1 \leq a \leq p-1$, we let $v(a)$ denote the unique member of V satisfying $v(a) \equiv a \pmod{p}$. In particular, we always have $v(a)_0 = a$.

We use the so-called "even-index" notation for the sequence of Bernoulli numbers, B_n . This notation and the basic results on Bernoulli numbers used here are given in [1].

3. Results of Iwasawa. Iwasawa ([3], [5], and [6]) has proved the following fundamental theorem on the cyclotomic invariant μ_p :

THEOREM 1. $\mu_p > 0$ if and only if there exists an odd index i , $1 \leq i \leq p-4$, such that

$$(1) \quad \sum_{v \in V} s_n(uv)v^i \equiv 0 \pmod{p^{n+2}}$$

for all units $u \in U$ and for all $n \geq 0$.

In [3], Iwasawa proves the equivalence of (1) with another set of congruences modulo p , using the relation $(uv)_n p^n = s_n(uv) - s_{n-1}(uv)$ for $n \geq 1$:

THEOREM 2. $\mu_p > 0$ if and only if there exists an odd index i , $1 \leq i \leq p-4$, such that

$$\sum_{v \in V} (uv)_n v^i \equiv 0 \pmod{p}$$

for all $u \in U$ and for all $n \geq 1$.

By choosing $u = 1$ in Theorems 1 and 2, we obtain

THEOREM 3. If $\mu_p > 0$, then there exists an odd index i , $1 \leq i \leq p-4$, such that

$$(1) \quad \sum_{v \in V} v_0 v^i \equiv 0 \pmod{p^2}, \quad \text{and}$$

$$(2) \quad \sum_{v \in V} v_n v_0^i \equiv 0 \pmod{p} \quad \text{for all } n \geq 1.$$

It is known from the general theory of Γ -extensions that $\mu_p = 0$ for all regular primes p (see [11] for a nice proof). We show next how this follows at once from Theorem 3.

COROLLARY. If $\mu_p > 0$, then there exists an odd index i , $1 \leq i \leq p-4$, such that $B_{i+1} \equiv 0 \pmod{p}$. Hence p is an irregular prime.

Proof. Part (1) of Theorem 3 implies that

$$\sum_{v \in V} v_0(v_0 + v_1 p)^i \equiv \sum_{v \in V} v_0^{i+1} + p i \sum_{v \in V} v_1 v_0^i \equiv 0 \pmod{p^2}.$$

But the second term is $0 \pmod{p^2}$ by part (2) of Theorem 3. Hence

$$B_{i+1} p \equiv \sum_{a=1}^{p-1} a^{i+1} = \sum_{v \in V} v_0^{i+1} \equiv 0 \pmod{p^2},$$

as desired.

4. Additional Conditions for Positive μ_p . In this section, we investigate the implications of Theorem 2 for different choices of the units $u \in U$, obtaining additional necessary conditions that μ_p be positive.

THEOREM 4. If $\mu_p > 0$, then there exists an odd index i , $1 \leq i \leq p-4$, such that

$$(1) \quad B_{i+1} \equiv 0 \pmod{p},$$

$$(2) \quad \sum_{a=1; a+v(a)_n < p}^{p-1} a^i \equiv \sum_{a=1; a+v(a)_n \geq p}^{p-1} a^i \equiv 0 \pmod{p} \quad \text{for all } n \geq 1.$$

Proof. We have already seen that (1) is true. If $n \geq 1$ and $v \in V$, we can write

$$v \equiv v_0 + v_1 p + \cdots + v_{n+1} p^{n+1} \pmod{p^{n+2}}.$$

For $u = 1 + p^n$, we have

$$uv \equiv v_0 + \cdots + v_{n-1} p^{n-1} + (v_0 + v_n) p^n + (v_1 + v_{n+1}) p^{n+1} \pmod{p^{n+2}}.$$

Thus, if $v_0 + v_n < p$, we have that $(uv)_n = v_0 + v_n$ and $(uv)_{n+1} \equiv v_1 + v_{n+1} \pmod{p}$. However, if $v_0 + v_n \geq p$, then $(uv)_n = v_0 + v_n - p$ and $(uv)_{n+1} \equiv v_1 + v_{n+1} + 1 \pmod{p}$.

By Theorem 2, $\sum_{v \in V} (uv)_{n+1} v^i \equiv 0 \pmod{p}$, or

$$\sum_{v \in V} (v_1 + v_{n+1}) v_0^i + \sum_{v \in V; v_0 + v_n \geq p} v_0^i \equiv 0 \pmod{p}.$$

But by Theorem 3, the first sum is $0 \pmod{p}$, and therefore, so is the second. Since $\sum_{v \in V} v_0^i \equiv 0 \pmod{p}$, we have

$$\sum_{v \in V; v_0 + v_n < p} v_0^i \equiv \sum_{v \in V; v_0 + v_n \geq p} v_0^i \equiv 0 \pmod{p},$$

which is the same as (2).

We remark that for several other choices of the unit u in Theorem 2 (e.g., $u = p - 1$ and $u = 1 + 2p^n$) we have derived additional congruences which are also necessary conditions for $\mu_p > 0$. These are omitted here since they are not required for any of our computations. We have selected the congruences of Theorem 4 since they lead (in the next section) to a sum with relatively few terms, thus providing for the greatest computational efficiency.

5. Main Theorem. For the actual computation of μ_p for $p < 8000$, it was necessary to use Theorem 4 only in the case that $n = 1$. In this section, we derive some simplifications of Theorem 4 when $n = 1$.

By expanding the congruence

$$1 = v(a)^{p-1} \equiv (a + v(a)_1 p)^{p-1} \pmod{p^2},$$

we see that $v(a)_1$ is completely determined by the conditions

$$(2) \quad v(a)_1 \equiv (a^p - a)/p \pmod{p} \quad \text{and} \quad 0 \leq v(a)_1 < p.$$

It is easy to see by (2) that

$$(3) \quad v(a)_1 + v(p - a)_1 = p - 1, \quad 1 \leq a \leq p - 1.$$

It follows immediately that $a + v(a)_1 < p$ if and only if $(p - a) + v(p - a)_1 \geq p$, so that, letting $b = p - a$ and recalling that i is odd, we obtain

$$\sum_{a=1; a+v(a)_1 < p}^m a^i \equiv - \sum_{b=m+1; b+v(b)_1 \geq p}^{p-1} b^i \pmod{p}.$$

By the Mirimanoff congruence [9]

$$2^i(i+1) \sum_{a=1}^m a^i \equiv (1 - 2^{i+1}) B_{i+1} \pmod{p},$$

we see that $B_{i+1} \equiv 0 \pmod{p}$ implies that

$$\sum_{a=1}^m a^i \equiv \sum_{a=m+1}^{p-1} a^i \equiv 0 \pmod{p}.$$

Combining the above congruences with the results of Theorem 4, we arrive at our main result:

THEOREM 5. *If $\mu_p > 0$, then there exists an odd index i , $1 \leq i \leq p-4$, such that*

$$(1) \quad B_{i+1} \equiv 0 \pmod{p}, \quad \text{and}$$

$$(2) \quad \sum_{a=1; a+v(a) \geq p}^m a^i \equiv 0 \pmod{p}.$$

The computations of Theorem 5 were carried out on the PDP-10 computer at Bowdoin College for all primes $p < 8000$, and the results are included in the Table accompanying this paper. About 60 hours of computing time were required for all the computations, the bulk of it in the search for the irregular primes and for the Bernoulli numbers satisfying (1). For no prime $p < 8000$ is the conclusion of Theorem 5 satisfied, so that we have

THEOREM 6. *The Iwasawa invariant $\mu_p = 0$ for all primes $p < 8000$.*

A more detailed explanation of how the computations of Theorem 5 were carried out is given in the following sections.

6. The Irregular Primes. The first condition of Theorem 5, that involving the Bernoulli numbers, has been of interest since Kummer's fundamental work on Fermat's Last Theorem in the nineteenth century. In a series of papers, Vandiver and others [8] claimed to have found all ordered pairs $(p, i+1)$ satisfying $B_{i+1} \equiv 0 \pmod{p}$, for $p \leq 4001$. They then verified that Fermat's Last Theorem is true for all exponents in this range. These pairs were used by Iwasawa and Sims [7] for their computation of the cyclotomic invariants μ_p , λ_p , and ν_p for primes $p \leq 4001$.

Our approach to finding these pairs was somewhat different from that used in [8]. The Bernoulli numbers satisfy the recursion relation

$$(4) \quad \sum_{j=0}^k \binom{k+1}{j} B_j = 0,$$

with $B_0 = 1$. Computing the binomial coefficients $\pmod{p}$, we can use the above to compute $B_k \pmod{p}$ recursively. This requires, of course, that we store the B_j 's as we go along. Since $B_j = 0$ for j odd, $j \geq 3$, there are really only approximately $k/2$ terms in the sum defining B_k .

Carlitz posed the following identity for the Bernoulli numbers as a problem in [2]:

$$(5) \quad (-1)^m \sum_{r=0}^m \binom{m}{r} B_{n+r} = (-1)^n \sum_{s=0}^n \binom{n}{s} B_{m+s}, \quad m, n \geq 0.$$

If we let $f(m, n)$ be the left-hand side of this equation, the problem is to show that $f(m, n) = f(n, m)$ for all $m, n \geq 0$. This is easily done by induction on m , using the

identity $f(m+1, n) = -f(m, n) - f(m, n+1)$.

If we now consider the special case $m = n+1$ in (5), we obtain

$$(6) \quad \sum_{r=0}^n C(n, r) B_{n+r} = 0, \quad n \geq 1,$$

where $C(n, 0) = 1$ and

$$C(n, r) = \binom{n+1}{r} + \binom{n}{r-1} \quad \text{for } 1 \leq r \leq n+1.$$

Equation (6) defines B_{2n} recursively in terms of $B_n, B_{n+1}, \dots, B_{2n-2}$, a considerable saving in computation time over the recursive relation (4). The coefficients $C(n, r)$ are easily computed modulo p , since they form a Pascal triangle whose first row is 1 2.

Using Eq. (6) modulo p , we found all pairs $(p, i+1)$ with $B_{i+1} \equiv 0 \pmod{p}$ for all primes $p < 8000$, at which time the program was terminated, since each additional prime took an excessively long time to run. Four additions were found to the tables in the first paper of [8], and these are marked with an asterisk in the accompanying Table. These omissions also occur in the table on p. 430–431 of [1], and, presumably, in the table (not completely published) of [7].

In [8], a prime p was first tested for irregularity by means of the congruence

$$2(i+1) \sum_{p/8 < a < p/4} a^i \equiv (4^{p-i-1} + 3^{p-i-1} - 6^{p-i-1} - 1) B_{i+1} \pmod{p}$$

which holds for $p > 7$. As a check on our computations, we ran another program for the four pairs $(p, i+1)$ omitted from [8] as well as for those pairs for which $4002 < p < 8000$. For each of these pairs, it was found that the sum on the left-hand side of the congruence above is 0 modulo p , while the coefficient of B_{i+1} is not, so that indeed, $B_{i+1} \equiv 0 \pmod{p}$.

Selfridge and Pollack [10] have found all pairs $(p, i+1)$ with $B_{i+1} \equiv 0 \pmod{p}$ for primes $p < 25,000$, and they have verified that Fermat's Last Theorem is true for all exponents less than or equal to 25,000 using the methods of [8]. A complete table of their calculations has not yet been published, but when it appears, we intend to use it to make further computations of the Iwasawa invariant μ_p . The validity of Fermat's Last Theorem for exponents less than or equal to 8000 was also verified by us in still another machine computation, using the criteria developed in [8].

7. Computation of the Sum in Theorem 5. In this section, we make some remarks on the algorithm that we devised for computing the sum in Theorem 5. The real problem lies in computing $v(a)_1$ for $a = 1, 2, \dots, m$. This can be done, of course, by Eq. (2), but those computations really have to be done modulo p^2 . Below, we indicate how certain of the $v(a)_1$'s can be found from others by means of a linear congruence modulo p .

Clearly, $v(1)_1 = 0$, so that the index $a = 1$ is never included in the sum. We first computed $v(2)_1$, using Eq. (2). The following identities can be derived from (2) and (3):

$$v(a)_1 - 2v(a/2)_1 - (a/2)v(2)_1 \equiv 0 \pmod{p} \quad (a \text{ even}),$$

$$v(a)_1 + 2v((p-a)/2)_1 + ((p-a)/2)v(2)_1 + 1 \equiv 0 \pmod{p} \quad (a \text{ odd}).$$

Hence, given $v(a)_1$ for some a , $1 \leq a \leq m$, these identities may be used to compute either $v(a/2)_1$ if a is even or $v((p-a)/2)_1$ if a is odd, without resorting to Eq. (2).

<i>p</i>	<i>i+1</i>	<i>terms</i>	<i>sum</i>	<i>p</i>	<i>i+1</i>	<i>terms</i>	<i>sum</i>	<i>p</i>	<i>i+1</i>	<i>terms</i>	<i>sum</i>
37	32	3	7	617	20	66	303	1201	676	159	1048
59	44	6	25	617	174	68	385	1217	784	174	57
67	58	7	43	617	338	66	191	1217	866	19	118
101	68	11	91	619	428	83	334	1217	1118	174	739
103	24	9	88	631	80	81	551	1229	784	139	
131	22	16	49	631	226	81	361	1237	874	148	1227
149	130	17	79	647	236	84	221	1279	518	146	861
157	62	21	108	647	242	84	320	1283	510	168	1233
157	110	21	69	647	554	84	219	1291	206	173	702
233	84	24	140	653	48	76	613	1291	824	173	1234
257	164	36	81	659	224	88	277	1297	202	151	1271
263	100	32	99	673	408	91	8	1297	220	151	250
271	84	30	179	673	502	91	571	1301	176	163	165
283	20	39	174	677	628	85	624	1307	382	184	48
293	156	34	217	683	32	87	2	1307	852	184	896
307	88	40	220	691	12	87	555	1319	304	165	1289
311	292	34	200	691	200	87	598	1327	466	163	332
347	280	49	198	727	378	88	628	1367	234	161	755
353	186	43	49	751	290	87	16	*1381	266	182	836
353	300	43	232	757	514	90	719	1409	358	171	1193
379	100	47	362	761	260	80	62	1429	936	178	94
379	174	47	164	773	732	106	184	1439	574	161	1230
389	200	50	159	797	220	95	38	1483	224	200	252
401	382	45	209	809	330	86	673	1499	94	191	1082
409	126	46	282	809	628	88	265	1523	1310	202	427
421	240	57	287	811	544	102	39	1559	862	186	1409
433	366	55	18	821	744	100	83	*1597	842	184	1579
461	196	53	270	827	102	97	691	1609	1356	191	1594
463	130	59	456	839	66	96	150	1613	172	207	221
467	94	67	169	877	868	114	455	1619	560	188	1199
467	194	67	400	881	162	117	610	1621	980	199	247
491	292	68	397	887	418	119	174	1637	718	205	434
491	336	68	139	929	520	117	679	1663	209	1599	1599
491	388	68	25	929	820	117	907	*1663	1503	209	1155
523	400	63	142	953	156	119	510	1669	308	200	1231
541	86	65	327	971	166	114	759	1669	1086	200	728
547	270	67	174	1061	474	123	488	1721	30	220	164
547	486	67	172	1091	888	150	1042	1733	810	218	1634
557	222	61	238	1117	794	131	167	1733	942	218	1073
577	52	68	558	1129	348	143	68	1753	712	233	1088
587	90	78	373	1151	534	140	751	1759	1520	218	1626
587	92	78	476	1151	784	140	74	1777	1192	240	1553
593	22	84	102	1151	968	140	968	1787	1606	217	1591
607	592	75	515	1153	802	156	1112	1789	848	233	733
613	522	72	428	1193	262	149	344	1799	1442	223	1468

p	$i+1$	$terms$	sum	p	$i+1$	$terms$	sum	p	$i+1$	$terms$	sum
1811	550	228	900	2383	842	327	1786	3089	1706	402	2832
1811	698	228	1044	2383	2278	327	662	3119	1704	374	1185
1811	1520	228	962	2389	776	299	2077	3181	3142	382	2247
1831	1274	232	575	2411	2126	319	1033	3203	2368	411	1303
1847	954	224	866	2423	290	315	1243	3221	98	413	1129
1847	1016	224	1511	2423	884	315	507	3229	1634	392	3059
1847	1558	224	1528	2441	366	307	2353	3257	922	436	629
1871	1794	233	370	2441	1750	307	1323	3313	2222	399	1954
*1877	1026	237	1539	2503	1044	311	90	3323	3292	412	1176
1879	1260	241	1739	2543	2374	318	1915	3329	1378	428	2749
1889	242	251	1849	2557	1464	313	518	3391	2232	429	2415
1901	1722	228	1365	2579	1730	324	624	3391	2534	429	673
1933	1058	237	1489	2591	854	314	1722	3407	2076	444	2201
1933	1320	237	162	2591	2574	314	710	3407	2858	444	2335
1951	1656	252	302	2621	1772	322	1473	3433	1300	436	2120
1979	148	257	1462	2633	1416	316	2054	3469	1174	432	2415
1987	510	248	286	2647	1172	340	2080	3491	2544	411	2053
1993	912	233	1117	2657	710	340	408	3511	1816	438	2441
1997	772	246	1959	2663	1244	340	1260	3511	1724	438	1666
1997	1888	246	1026	2671	404	336	400	3517	1836	419	1157
2003	60	255	1850	2671	2394	336	1344	3517	2586	419	943
2003	600	255	1079	2689	926	346	2243	3529	3490	451	2453
2017	1204	266	1423	2753	482	328	2611	3533	2314	422	1462
2039	1300	256	764	2767	2528	328	13	3533	3136	422	478
2053	1932	252	1067	2777	1600	362	2014	3539	2082	422	1902
2087	376	263	1109	2789	1984	356	164	3539	2130	422	2676
2087	1238	263	1324	2789	2154	356	312	3559	344	438	116
2099	1230	277	492	2791	2554	308	1507	3559	1592	439	3289
2111	1038	243	557	2833	1832	326	1693	3591	1466	417	275
2137	1624	270	1775	2857	98	331	1828	3593	1922	434	611
2143	1916	277	2036	2861	352	349	1251	3593	360	464	565
2153	1832	266	1585	2909	400	396	930	3593	642	464	3097
2213	154	295	461	2909	950	396	1239	3607	1976	446	546
2239	1826	258	1611	2927	242	383	2702	3613	2092	449	1491
2267	2234	289	1182	2939	332	366	334	3617	16	446	2869
2273	876	275	1210	2939	1102	366	280	3617	2056	446	1264
2273	2166	275	1580	2939	2748	366	465	3631	1104	464	1447
2293	2040	257	669	2957	138	361	31	3637	2526	473	1863
2309	1680	208	2018	2957	788	361	1776	3637	3202	473	28
2309	1772	288	1616	2999	776	372	371	3671	1580	468	2742
2357	2204	289	863	3011	1496	395	2526	3677	2238	436	2728
2371	242	298	1139	3023	2020	347	2295	3697	1884	464	1137
2371	2274	298	134	3049	700	397	2693	3779	2362	456	2243
2377	1226	285	1496	3061	2522	399	399	3797	1256	487	2179
2381	2060	271	1106	3083	1450	370	1734	3821	3296	477	2655

<i>p</i>	<i>i+1</i>	<i>terms</i>	<i>sum</i>	<i>p</i>	<i>i+1</i>	<i>terms</i>	<i>sum</i>	<i>p</i>	<i>i+1</i>	<i>terms</i>	<i>sum</i>
3833	1840	476	990	4657	1578	579	2183	5443	1710	652	3387
3833	1999	476	1089	4657	2416	579	3208	5477	1150	631	630
3236	3236	476	2314	4657	4110	579	995	5479	1826	650	5457
3851	216	471	736	4663	216	603	3611	5479	4802	650	5245
3851	404	471	3380	4663	4278	603	3314	5501	666	719	1935
3853	748	435	1422	4679	3592	576	4079	5527	5206	706	4501
3881	1686	462	3356	4651	3450	571	3948	5531	3438	701	5344
3881	2138	462	798	4751	3768	601	4352	5557	3196	714	2146
3917	1490	467	3067	4793	252	604	3786	5569	938	710	2766
3957	106	405	305	4793	2636	626	3762	5573	2032	689	4879
3989	1936	462	2037	4813	2620	585	1692	5639	2672	744	5177
4001	534	518	3299	4861	4678	608	2789	5641	4580	731	3806
4003	82	493	895	4889	2924	640	4440	5641	5258	731	3155
4003	142	493	1459	4903	3106	601	3481	5669	2218	701	3879
4003	2610	493	230	4909	1462	630	2816	5669	2680	701	1311
4021	3228	528	3783	4943	492	624	315	5689	348	711	1190
4027	2332	497	329	4951	1914	609	586	5701	2450	680	712
4049	1854	486	1700	4951	2468	609	4156	5783	2200	757	1074
4051	3548	535	136	4951	3830	609	2946	5791	1258	737	4450
4073	3620	509	1621	4957	3812	619	4069	5813	4284	759	5487
4129	1784	535	581	4969	1940	624	4109	5921	1150	717	2590
4157	658	538	1029	4973	4208	645	3067	5939	2308	709	4879
4157	2322	538	3855	5009	1544	621	22	5861	3554	744	580
4219	4190	556	2220	5039	4956	611	322	5897	2996	729	2862
4243	2712	543	381	5039	594	651	4135	5903	3970	729	2071
4243	4146	543	2191	5077	3092	599	2770	5903	5000	729	5530
4259	3530	558	3355	5081	3016	630	3762	5923	4240	748	1466
4259	3726	558	1208	5099	1378	669	2497	5927	3642	739	4274
4261	2068	482	3725	5101	190	627	3455	5939	342	735	5157
4339	214	547	1243	5107	4872	639	1776	5939	5014	735	5527
4349	2052	511	428	5119	4086	632	4701	5953	3274	763	4360
4409	636	581	3991	5167	4112	626	2294	6007	912	714	118
4409	672	591	3237	5179	4732	678	2277	6011	5870	722	624
4421	3765	560	2845	5189	1102	677	2943	6037	3396	752	4626
4451	2896	551	3365	5209	644	640	3451	6043	1226	770	5010
4451	2978	551	3346	5209	2928	640	528	6091	702	761	5812
4457	444	552	209	5227	308	633	5017	6101	2008	741	2794
4493	746	534	541	5231	3466	642	1224	6173	5008	770	131
4519	848	553	2326	5297	4810	670	4260	6173	5894	770	5093
4523	456	551	3776	5303	4156	660	3140	6217	4186	777	3294
4561	436	567	1428	5309	158	652	1623	6247	1492	799	1767
4591	2292	592	2652	5351	1948	687	872	6247	3474	799	4635
4591	3596	592	1776	5399	1482	674	3705	6257	4272	775	3473
4637	3618	567	1866	5413	1702	685	5133	6263	3286	793	6130
4639	3226	567	958	5441	4726	603	2051	6263	4226	793	6010

p	$i+1$	terms	sum	p	$i+1$	terms	sum	p	$i+1$	terms	sum
6287	4452	803	1358	7039	1454	884	4693	7919	3888	971	1498
6287	5034	603	202	7057	4154	903	669	7927	6448	980	621
6317	2354	793	1683	7057	4972	903	3496	7937	3980	935	601
6329	3102	816	2169	7069	1478	850	3103	7949	2506	1009	1807
6337	1956	772	633	7069	2570	850	3586	7949	3436	1009	926
6343	750	773	555	7109	290	887	3838	7951	4328	961	2743
6343	5820	773	3693	7121	1502	885	2851	7963	4748	1033	74
6367	1190	780	2528	7127	6758	849	4876				
6373	2838	803	4876	7177	962	884	1391				
6373	4226	803	3447	7187	3906	945	6847				
6379	218	779	342	7207	1670	972	1891				
6421	438	793	1151	7207	5774	972	1800				
6449	4884	775	3003	7211	898	906	5989				
6449	5830	775	3431	7213	1436	893	2097				
6451	3236	817	4290	7213	6930	893	2163				
6491	346	807	2178	7229	6236	926	738				
6521	236	816	3448	7309	324	890	1510				
6529	1564	810	3654	7321	348	892	2405				
6547	734	871	6183	7351	1466	928	2113				
6569	1692	813	2814	7411	4712	898	4054				
6569	1776	813	3851	7459	5286	947	6940				
6571	1744	774	3308	7487	2500	924	2778				
6577	1312	848	5700	7489	4250	951	1314				
6619	1952	829	6088	7499	3642	930	827				
6619	3170	829	1719	7507	6924	921	5601				
6659	2950	800	1532	7537	2264	945	7112				
6659	4014	800	2968	7547	5644	914	3305				
6689	5252	858	3333	7559	116	906	3207				
6701	5484	833	2400	7591	2520	965	460				
6733	1690	878	4695	7607	3594	934	1945				
6763	4144	835	6633	7643	5026	960	6969				
6763	6218	835	4046	7681	368	926	5662				
6763	6230	835	2419	7687	1246	969	60				
6779	3934	831	2091	7687	3216	969	192				
6793	2686	852	2303	7687	6516	969	6570				
6823	4952	865	6253	7691	2218	935	4092				
6827	4108	854	6063	7727	950	929	4841				
6833	2254	873	1146	7727	3756	929	1487				
6833	5144	873	369	7817	7346	941	6876				
6857	6676	820	2447	7823	3298	987	2519				
6863	6406	859	6442	7829	1392	988	5036				
6949	2432	854	2726	7853	3494	1014	807				
6971	2010	834	5888	7901	2472	921	1056				
6973	1746	873	240	7901	4286	921	4182				
7001	4642	862	2394	7907	584	963	7541				

Starting with $a = 1$, for example, we next computed $v(m)_1 = v((p - 1)/2)_1$, then $v(m/2)_1$ or $v((p - m)/2)_1 = v((p + 1)/4)_1$ (depending upon whether m was even or odd), and so forth, until a full cycle was completed. We then searched for the first a , $1 \leq a \leq m$, for which $v(a)_1$ had not yet been computed, found the value of $v(a)_1$ by using Eq. (2) again, and then began another cycle using the identities above. This procedure was continued until $v(a)_1$ had been computed for all $a = 1, 2, \dots, m$. It can be shown that the cycles arising in this way all have the same length and that, in the particular case that m is also a prime number, there is but one cycle, indicating the efficiency of the algorithm thus devised.

If we assume that, for fixed a , it is equally likely that $v(a)_1$ assumes any one of the values $0, 1, 2, \dots, p - 1$, then the probability that the term a^i appears in the sum of Theorem 5 (i.e., the probability that $a + v(a)_1 \geq p$) is just a/p . Thus, the expected number of terms in the sum is $\sum_{a=1}^m a/p = p/8 - (8p)^{-1}$, or approximately $p/8$ for large primes p . It is interesting to compare the value $p/8$ with the entries in the third column of the accompanying Table.

8. The Table. In the first two columns of the accompanying Table, we have listed all pairs $(p, i + 1)$, where p is a prime, $p < 8000$, and where the Bernoulli number $B_{i+1} \equiv 0 \pmod{p}$. The four additions to the tables of [8] are marked with an asterisk. The third column contains, for each of these pairs, the number of integers a , $1 \leq a \leq m$, satisfying the condition $a + v(a)_1 \geq p$. This is the same as the number of terms in the sum $\sum_{a=1}^m : a + v(a)_1 \geq p a^i$ of Theorem 5. The value of this sum modulo p is given in the final column of the Table. Since a zero never appears in this final column, we can conclude by Theorem 5 that Theorem 6 must be true.

Department of Mathematics
Bowdoin College
Brunswick, Maine 04011

1. Z. I. BOREVIČ & I. R. ŠAFAREVIČ, *Number Theory*, "Nauka," Moscow, 1964; English transl., Pure and Appl. Math., vol. 20, Academic Press, New York, 1966. MR 30 #1080; MR 33 #4001.
2. L. CARLITZ, "Problem 795," *Math. Mag.*, v. 44, 1971, p. 106.
3. K. IWASAWA, "On some invariants of cyclotomic fields," *Amer. J. Math.*, v. 80, 1958, pp. 773–783; erratum, *ibid.*, v. 81, 1959, p. 280. MR 23 #A1631.
4. K. IWASAWA, "On Γ -extensions of algebraic number fields," *Bull. Amer. Math. Soc.*, v. 65, 1959, pp. 183–226. MR 23 #A1630.
5. K. IWASAWA, "A class number formula for cyclotomic fields," *Ann. of Math.*, (2), v. 76, 1962, pp. 171–179. MR 27 #4806.
6. K. IWASAWA, "On some modules in the theory of cyclotomic fields," *J. Math. Soc. Japan*, v. 16, 1964, pp. 42–82. MR 35 #6646.
7. K. IWASAWA & C. SIMS, "Computation of invariants in the theory of cyclotomic fields," *J. Math. Soc. Japan*, v. 18, 1966, pp. 86–96. MR 34 #2560.
- 8a. D. H. LEHMER, E. LEHMER & H. S. VANDIVER, "An application of high-speed computing to Fermat's last theorem," *Proc. Nat. Acad. Sci. U.S.A.*, v. 40, 1954, pp. 25–33. MR 15, 778.
- 8b. H. S. VANDIVER, "Examination of methods of attack on the second case of Fermat's last theorem," *Proc. Nat. Acad. Sci. U.S.A.*, v. 40, 1954, pp. 732–735. MR 16, 13.
- 8c. J. L. SELFIDGE, C. A. NICOL & H. S. VANDIVER, "Proof of Fermat's last theorem for all prime exponents less than 4002," *Proc. Nat. Acad. Sci. U.S.A.*, v. 41, 1955, pp. 970–973. MR 17, 348.
9. D. MIRIMANOFF, "Sur la congruence $(r^{p-1} - 1) : p \equiv q_r \pmod{p}$," *J. Reine Angew. Math.*, v. 115, 1895, pp. 295–300.
10. J. L. SELFIDGE & B. W. POLLACK, "Fermat's last theorem is true for any exponent up to 25,000," *Notices Amer. Math. Soc.*, v. 11, 1964, p. 97. Abstract #608-138.
11. J.-P. SERRE, *Classes des corps cyclotomiques (d'après K. Iwasawa)*, Séminaire Bourbaki 1958/59, Exposé 174, fasc. 1, Secrétariat mathématique, Paris, 1959. MR 28 #1091.