An Elliptic Integral Identity

By H. S. Wrigge

Abstract. The identity

$$K(\tau) = \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \exp \left[-\frac{1}{2}(x^4 - 2(2\tau^2 - 1)x^2y^2 + y^4) \right] dx dy,$$

where $K(\tau)$ is the complete elliptic integral of the first kind, is used to prove that $K(\sqrt{2} - 1) = \pi^{3/2}(2 + \sqrt{2})^{1/2}/4\Gamma(\frac{3}{8})\Gamma(\frac{5}{8})$.

The identity

$$(1) \quad K(\tau) = \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} \exp \left[-\frac{1}{2}(x^4 - 2(2\tau^2 - 1)x^2y^2 + y^4) \right] dx dy, \quad |\tau| < 1,$$

is easily proved by using the standard transformation $x = r \cos v$, $y = r \sin v$. Before proceeding, we will list a few identities for the benefit of the reader.

$$(2) \quad \int_0^{\infty} e^{-pt} t^{-1/2} e^{-t^{3/4}\alpha} dt = \alpha^{1/2} p^{1/2} e^{\alpha p^{3/2}} K_{1/4}(\frac{1}{2}\alpha p^2), \quad \operatorname{Re} \alpha > 0 \quad [1, \text{p. 146}],$$

$$(3) \quad \int_0^{\infty} e^{-pt} t^{\mu-1/2} K_{\nu+1/2}(\alpha t) dt = 2^{-1/2} \alpha^{-1/2} \pi^{1/2} \Gamma(\mu - \nu) \Gamma(\mu + \nu + 1) s^{-\mu} P_{\nu}^{-\mu}(p/\alpha),$$

$$s = (p^2 - \alpha^2)^{1/2}, \quad \operatorname{Re}(\mu + \nu) > -1, \quad \operatorname{Re}(\mu - \nu) > 0,$$

$$\operatorname{Re} p > -\operatorname{Re} \alpha, \quad [1, \text{p. 198}],$$

$$(4) \quad P_p^q(0) = \frac{2^q \pi^{1/2}}{\Gamma(\frac{1}{2}(p - q) + 1) \Gamma(\frac{1}{2}(-p - q + 1))} \quad [2, \text{p. 354}].$$

We will now calculate the integral (1) in a different way.

Integration with respect to x yields

$$\begin{aligned} \int_{-\infty}^{\infty} \exp \left[-\frac{1}{2}(x^4 - 2(2\tau^2 - 1)x^2y^2) \right] dx &= \int_0^{\infty} t^{-1/2} e^{-t^{3/2}} e^{(2\tau^2-1)y^2 t} dt \\ &= \frac{(1 - 2\tau^2)^{1/2}}{2^{1/2}} |y| \exp \left[\frac{(2\tau^2 - 1)^2 y^4}{4} \right] K_{1/4} \left(\frac{(2\tau^2 - 1)^2 y^4}{4} \right), \quad 1 - 2\tau^2 > 0. \end{aligned}$$

Integration with respect to y now gives

$$K(\tau) = \frac{(1 - 2\tau^2)^{1/2}}{2\pi^{1/2}} \int_{-\infty}^{+\infty} |y| e^{-y^{4/2}} \exp \left[\frac{(2\tau^2 - 1)^2 y^4}{4} \right] K_{1/4} \left(\frac{(2\tau^2 - 1)^2 y^4}{4} \right) dy.$$

A few manipulations finally give

Received October 16, 1972.

AMS (MOS) subject classifications (1970). Primary 33A25.

Copyright © 1973, American Mathematical Society

$$(5) \quad K(\tau) = \frac{(1 - 2\tau^2)^{1/2}}{2\pi^{1/2}} \int_0^\infty t^{-1/2} e^{-t(1+4\tau^2-4\tau^4)} K_{1/4}((1 - 2\tau^2)^2 t) dt.$$

Using Eq. (3) we can write

$$(6) \quad K(\tau) = \frac{\pi}{2(1 - 2\tau^2)^{1/2}} P_{-1/4} \left(\frac{1 + 4\tau^2 - 4\tau^4}{(1 - 2\tau^2)^2} \right).$$

To get (6) in a more convenient form, we use the well-known formula

$$K(i\tau) = \frac{1}{(1 + \tau^2)^{1/2}} K \left(\frac{\tau}{(1 + \tau^2)^{1/2}} \right).$$

Then

$$(7) \quad K \left(\frac{\tau}{(1 + \tau^2)^{1/2}} \right) = \frac{\pi(1 + \tau^2)^{1/2}}{2(1 + 2\tau^2)^{1/2}} P_{-1/4} \left(\frac{1 - 4\tau^2 - 4\tau^4}{(1 + 2\tau^2)^2} \right).$$

Now put $1 - 4\tau^2 - 4\tau^4 = 0$ which yields $\tau^2 = -\frac{1}{2} \pm \frac{1}{2}\sqrt{2}$. A little calculation and Eq. (4) give

$$(8) \quad K(\sqrt{2} - 1) = \frac{\pi^{3/2}(2 + \sqrt{2})^{1/2}}{4\Gamma(\frac{5}{8})\Gamma(\frac{7}{8})}. \quad \text{Q.E.D.}$$

The identity (8) has been proved previously; see for instance [3, pp. 535–536]. The present derivation seems, however, to give a better insight into the problem.

Research Institute of the Swedish National Defense
104 50 Stockholm 80,
Sweden

1. A. ERDÉLYI ET AL., *Tables of Integral Transforms*. Vol. 1, McGraw-Hill, New York, 1954. MR 15, 868.

2. I. M. RYŽIK & I. S. GRADŠTEĬN, *Summen-, Produkt- und Integral-Tafeln*, 2nd rev. ed., VEB Deutscher Verlag, Berlin; Plenum Press, New York, 1963. MR 28 #1326.

3. M. L. GLASSER & V. E. WOOD, "A closed form evaluation of the elliptic integral," *Math. Comp.*, v. 25, 1971, pp. 535–536.