Negative Integral Powers of a Bidiagonal Matrix

By Gurudas Chatterjee

Abstract. The elements of the inverse of a bidiagonal matrix have been expressed in a convenient form. The higher negative integral powers of the bidiagonal matrix exhibit an interesting property: the (ij) th element of the $(-m)$ th power is equal to the product of the corresponding element of the inverse by a Wronski polynomial, viz., the complete symmetric function of degree $(m - 1)$ of the diagonal elements, $d_i, d_{i+1}, \dots, d_j$, of the inverse matrix.

1. Introduction. Positive integral powers of a bidiagonal matrix with a fixed diagonal element b and superdiagonal element 1, have been reported by Varga [1]. In the present note, we shall find the negative integral powers of a general $n \times n$ bidiagonal matrix B , having diagonal elements $b_i, i = 1, 2, \dots, n$, and superdiagonal elements $c_j, j = 1, 2, \dots, n - 1$.

One may express $B = (I - \Gamma)D^{-1}$, where I is the identity matrix and D^{-1} a diagonal matrix composed of the diagonal elements of B . Γ is null, except for the elements $\Gamma_{i,i+1} = -c_i/b_{i+1}$, for $i = 1, 2, \dots, n - 1$, on its first superdiagonal. The powers of Γ can easily be evaluated. In fact, the nonzero elements of Γ^m are given by

$$(\Gamma^m)_{i,i+m} = \prod_{k=i}^{i+m-1} (-c_k/b_{k+1}), \quad \text{for } i = 1, 2, \dots, n - m,$$

occurring only on the m th superdiagonal.

The inverse E_1 of B may be calculated either by the usual method of cofactors, or from the following expansions:

$$\begin{aligned} E_1 &= B^{-1} = D[I - \Gamma]^{-1} \\ &= D[I + \Gamma + \Gamma^2 + \Gamma^3 + \dots + \Gamma^{n-1}]. \end{aligned}$$

The elements of E_1 may be written in a convenient form as:

$$(1a) \quad e_1(i, j) = 0 \quad \text{for } i > j,$$

$$(1b) \quad = 1/b_j = d_j \quad \text{for } i = j,$$

$$(1c) \quad = d_i \prod_{k=i}^{j-1} (-c_k/b_{k+1}) \quad \text{for } i < j.$$

The inverse is upper triangular but is not necessarily bidiagonal.

2. Powers of the Inverse. The product of E_1 with itself is a matrix E_2 , which is also upper triangular. Elements of E_2 are given by

$$\begin{aligned} (2) \quad e_2(i, j) &= e_1(i, j) \sum_{k=i}^j [d_k] \quad \text{for } i \leq j, \\ &= 0 \quad \text{for } i > j. \end{aligned}$$

Received August 6, 1973.

AMS (MOS) subject classifications (1970). Primary 15A09.

Copyright © 1974, American Mathematical Society

Result (2) may be generalized. In fact, the n th power of E_1 is an upper triangular matrix E_n where the (i, j) th element, for $i \leq j$, is given by

$$(3) \quad e_n(i, j) = e_1(i, j) \sum_{k_1=i}^j \sum_{k_2=i}^{k_1} \sum_{k_3=i}^{k_2} \cdots \sum_{k_{n-2}=i}^{k_{n-3}} \sum_{k_{n-1}=i}^{k_{n-2}} [d_{k_1} d_{k_2} d_{k_3} \cdots d_{k_{n-2}} d_{k_{n-1}}].$$

Proof. Let us assume that result (3) is true for $n = m$.

$$e_{m+1}(i, j) = \sum_{k_0=i}^j [e_m(i, k_0) e_1(k_0, j)],$$

the other terms in the summation for $1 \leq k_0 \leq i - 1$ and $j + 1 \leq k_0 \leq n$, are zero, as both $e_m(p, q)$ and $e_1(p, q)$ are zero for $p > q$.

By writing the expression for $e_m(i, k_0)$ from result (3), which is assumed to be valid for $n = m$, we have

$$e_{m+1}(i, j) = \sum_{k_0=i}^j [e_1(i, k_0) e_1(k_0, j)] \\ \cdot \sum_{k_1=i}^j \sum_{k_2=i}^{k_1} \sum_{k_3=i}^{k_2} \cdots \sum_{k_{m-2}=i}^{k_{m-3}} \sum_{k_{m-1}=i}^{k_{m-2}} [d_{k_1} d_{k_2} d_{k_3} \cdots d_{k_{m-2}} d_{k_{m-1}}].$$

The first summation is done by Eq. (2) and the expression reduces to

$$e_{m+1}(i, j) = e_1(i, j) \sum_{k_0=i}^j [d_{k_0}] \cdot \sum_{k_1=i}^j \sum_{k_2=i}^{k_1} \sum_{k_3=i}^{k_2} \cdots \sum_{k_{m-2}=i}^{k_{m-3}} \sum_{k_{m-1}=i}^{k_{m-2}} [d_{k_1} d_{k_2} d_{k_3} \cdots d_{k_{m-2}} d_{k_{m-1}}].$$

After grouping the summations together, we find that the result is true for $n = m + 1$.

It has already been found true for $n = 2$ in Eq. (2), and therefore, by mathematical induction, we have the proof.

3. Thanks are due to the Director of this Institute for his kind permission to publish this paper.

Computer Centre
Central Mechanical Engineering Research Institute
Durgapur-9, West Bengal, India

1. R. S. VARGA, *Matrix Iterative Analysis*, Prentice-Hall, Englewood Cliffs, N.J., 1962, p. 14. MR 28 #1725.