The Design of Slowly Shrinking Labelled Squares*

By Elwyn R. Berlekamp

Abstract. This paper considers the nonlinear operator T , which acts on 4-dimensional vectors with nonnegative real components. T transforms such a vector into a new vector whose components are the magnitudes of the differences of cyclically adjacent pairs of components of the previous vector. We show that T is not nilpotent.

This short paper is devoted to a problem which I first heard from the Lehmers, Professor D. Gale, and Professor A. P. Morse on May 6, 1973. It consumed the greater part of that evening and it was not solved until the next day.

If the four vertices of a square are labelled with a given set of four nonnegative real numbers, then one may label the midpoint of each side as the absolute value of the difference between the labels on the corresponding vertices. Taking these labels on the midpoints of the original square as the labels on the vertices of a new square, the operation may be repeated again and again. For example, if the vertices of the original square are labelled, in clockwise order, as $x = [1, 5, 10, 8]$, then the labels after successive shrinkings are

$$xT = [4, 5, 2, 7],$$

$$xT^2 = [1, 3, 5, 3],$$

$$xT^3 = [2, 2, 2, 2],$$

$$xT^4 = [0, 0, 0, 0].$$

If one knows the ordering of the original labels, then the shrinking operation can be represented as a linear transformation. For example, if $x_1 \leq x_2 \leq x_3 \leq x_4$, then

$$xT = x \begin{bmatrix} -1 & 0 & 0 & -1 \\ +1 & -1 & 0 & 0 \\ 0 & +1 & -1 & 0 \\ 0 & 0 & +1 & +1 \end{bmatrix}.$$

Received July 13, 1974.

AMS (MOS) subject classifications (1970). Primary 10A35.

*Supported in part by NSF Grant GK-36172.

On the other hand, if $x_1 \geq x_2 \geq x_4 \geq x_3$, then

$$xT = x \begin{bmatrix} +1 & 0 & 0 & +1 \\ -1 & +1 & 0 & 0 \\ 0 & -1 & -1 & 0 \\ 0 & 0 & +1 & -1 \end{bmatrix}.$$

Almost any square labelled with positive integers will shrink to zero after only a few applications of the T operator, as the reader is invited to verify. One might even be tempted to conjecture that the nonlinear operator T is nilpotent, i.e., some finite power of it will shrink any square to zero. This conjecture turns out to be false, but the following weakened form of it is true:

LEMMA. *If the components of x with maximum and minimum labels are not cyclically adjacent, then $xT^6 = 0$.*

Proof. If the extreme labels are not adjacent they must be opposite, and without loss of generality we may assume that $x_1 \geq x_2 \geq x_4 \geq x_3$. Setting $y_i = x_i - x_3$, we have

$$y_1 \geq y_2 \geq y_4 \geq y_3 = 0,$$

$$xT = yT = [y_1 - y_2, y_2, y_4, y_1 - y_4],$$

$$xT^2 = [|y_1 - 2y_2|, y_2 - y_4, |y_1 - 2y_4|, y_2 - y_4].$$

If we then set $z_1 = ||y_1 - 2y_2| + y_4 - y_2|$ and $z_3 = ||y_1 - 2y_4| + y_4 - y_2|$ we have

$$xT^3 = [z_1, z_1, z_3, z_3].$$

Setting $w = |z_1 - z_3|$ yields

$$xT^4 = [0, w, 0, w], \quad xT^5 = [w, w, w, w], \quad xT^6 = 0. \quad \text{Q.E.D.}$$

COROLLARY. *If $xT^7 \neq 0$, then the components of x are cyclically ordered.*

Proof. By appropriate choice of coordinates, we may assume that the minimum label occurs at component x_1 . If $xT^6 \neq 0$, then the lemma asserts that the maximum label is not x_3 , so by appropriate choice of direction we may assume that x_4 is the maximum label. So there is no loss of generality in assuming that $x_1 \leq x_2 \leq x_4$ and $x_1 \leq x_3 \leq x_4$. If $x_3 \leq x_2$, then

$$xT = [x_2 - x_1, x_2 - x_3, x_4 - x_3, x_4 - x_1].$$

The maximum component of xT is $x_4 - x_1$, and the minimum component of xT is $x_2 - x_3$, whence by the Lemma, $xT \cdot T^6 = 0$, which contradicts the hypothesis of

this Corollary. The assumption that $x_3 \leq x_2$ is therefore untenable, and we can therefore conclude that $x_1 \leq x_2 < x_3 \leq x_4$. Q.E.D.

If we wish to find labellings which will not shrink to zero in n or fewer operations, then the Corollary tells us that for any $m \leq n - 7$, the components of $y = xT^m$ must be cyclically ordered. Furthermore, for $1 \leq m \leq n - 7$, the biggest component of xT^m must be equal to the sum of the other three components. So, without loss of generality, we may assume that $y_1 \leq y_2 < y_3 \leq y_4 = y_1 + y_2 + y_3$, and that $yT = z = [z_1, z_2, z_3, z_4] = yM$, where M is this 4×4 matrix

$$M = \begin{bmatrix} -1 & 0 & 0 & -1 \\ +1 & -1 & 0 & 0 \\ 0 & +1 & -1 & 0 \\ 0 & 0 & +1 & +1 \end{bmatrix}.$$

The biggest component of z is obviously z_4 , and if $zT^6 \neq 0$, then the components of z must also be cyclically ordered. So either $z_1 \leq z_2 \leq z_3 \leq z_4$ or $z_3 \leq z_2 \leq z_1 \leq z_4$. To see that the latter ordering is not possible, we notice that since $y_2 = y_1 + z_1$ and $y_3 = y_1 + z_1 + z_2$ and $y_4 = y_1 + z_1 + z_2 + z_3$, the equation $y_4 = y_1 + y_2 + y_3$ implies $z_3 = 2y_1 + z_1$, whence $z_1 \leq z_3$. Thus, the components of z are cyclically ordered in the same direction as the components of y . It follows that for $n \geq 9$, $xT^n = (xT)M^{n-8}T^7$.

The eigenvalues of M are the roots of the polynomial

$$|M - \lambda I| = (\lambda + 1)^3(\lambda - 1) + 1 = 0 = \lambda(\lambda^3 + 2\lambda^2 - 2)$$

and the corresponding eigenvectors are

$$x = x_1 [1, (\lambda + 1), (\lambda + 1)^2, (\lambda + 1)^3].$$

One of the eigenvalues of M is 0, corresponding to the trivial eigenvector $x = 1, 1, 1, 1$. There is also one positive real eigenvalue, $= .83928675521 \dots$ and a pair of complex eigenvalues of greater magnitude. It is possible to keep $xT^n \neq 0$ for large n only by choosing an initial vector x for which xT is very close to y , the non-trivial real eigenvector of M . This may be accomplished either by taking x to be a linear combination of y and $[1, 1, 1, 1]$, or by taking x to be a linear combination of $[1, 1, 1, 1]$ and $[y_4, y_3 + y_2, y_3, 0]$. In the latter case the first application of T reverses the ordering of the labels. $\square$

Acknowledgement. I am indebted to Dean Hickerson for pointing out certain omissions in an earlier draft of this paper.