

## Five-Diagonal Sixth Order Methods for Two-Point Boundary Value Problems Involving Fourth Order Differential Equations

By C. P. Katti

**Abstract.** We present a sixth order finite difference method for the two-point boundary value problem  $y^{(4)} + f(x, y) = 0$ ,  $y(a) = A_0$ ,  $y(b) = B_0$ ,  $y'(a) = A_1$ ,  $y'(b) = B_1$ . In the case of linear differential equations, our difference scheme leads to five-diagonal linear systems.

Consider the two-point boundary value problem

$$(1) \quad y^{(4)} + f(x, y) = 0, \quad y(a) = A_0, \quad y(b) = B_0, \quad y'(a) = A_1, \quad y'(b) = B_1.$$

In a recent paper Usmani [1] has given finite difference methods of orders two, four and six for the boundary value problem (1) in the case when  $f(x, y)$  is linear. While his methods of orders two and four can be easily adapted for nonlinear  $f(x, y)$  and lead to five-diagonal linear systems when  $f(x, y)$  is linear, the sixth order method given by Usmani leads to a nine-diagonal linear system. In the following we present a sixth order method for the nonlinear boundary value problem (1) which, in the case of linear  $f(x, y)$ , leads to five-diagonal linear systems.

At the grid points  $x_k$ ,  $k = 2(1)N-1$ , where  $x_k = a + kh$ ,  $k = 0(1)N+1$ ,  $N \geq 5$ , the differential equation in (1) can be discretized by

$$(2) \quad \delta^4 y_k + h^4 [2a_0 f_k + a_1(f_{k+1} + f_{k-1}) + a_2(f_{k+2} + f_{k-2})] + T_k(h) = 0,$$

where we have set  $y_k = y(x_k)$  and  $f_k = f(x_k, y_k)$ .

In order that  $T_k(h) = O(h^{10})$ , we find that

$$(a_0, a_1, a_2) = (1/720) (237, 124, -1).$$

Note that  $y_0 = A_0$ ,  $y_{N+1} = B_0$ . Let  $y'_k = y'(x_k)$ ,  $k = 0, N+1$ . The discretizations for the boundary conditions  $y'_0 = A_1$ ,  $y'_{N+1} = B_1$  can be obtained following Chawla and Katti [2]. Now, for the boundary conditions  $y'_0 = A_1$  and  $y'_{N+1} = B_1$ , consider the discretizations

$$(3a) \quad \sum_{k=0}^3 b_k y_k + ch y'_0 + h^4 \left( \sum_{k=0}^3 d_k f_k + \sum_{k=0}^1 d_k^* \bar{f}_{k+1/2} \right) + T_1(h) = 0,$$

---

Received August 20, 1979.

1980 *Mathematics Subject Classification.* Primary 65L10.

© 1980 American Mathematical Society  
0025-5718/80/0000-0158/\$01.75

and

$$(3b) \quad \sum_{k=0}^3 b_k y_{N+1-k} - ch y'_{N+1} + h^4 \left( \sum_{k=0}^3 d_k f_{N+1-k} + \sum_{k=0}^1 d_k^* \bar{f}_{N-k+1/2} \right) + T_N(h) = 0,$$

where we have set  $\bar{f}_{k+1/2} = f(x_{k+1/2}, \bar{y}_{k+1/2})$ ,  $x_{k+1/2} = x_k + h/2$ ,  $k = O(1)N$ , and where

$$(4a) \quad \bar{y}_{k+1/2} = \sum_{m=0}^3 u_{k,m} y_m + h^4 \sum_{m=0}^3 w_{k,m} f_m, \quad k = 0, 1,$$

and

$$(4b) \quad \bar{y}_{N-k+1/2} = \sum_{m=0}^3 u_{k,m} y_{N+1-m} + h^4 \sum_{m=0}^3 w_{k,m} f_{N+1-m}, \quad k = 0, 1.$$

In order that  $T_1(h)$  and  $T_N(h) = O(h^{10})$ , we find the following values for the parameters in (3) and (4):

$$(b_0, b_1, b_2, b_3) = \left( -\frac{11}{2}, 9, -\frac{9}{2}, 1 \right), \quad c = -3,$$

$$(d_0, d_1, d_2, d_3) = \left( \frac{1}{4200} \right) (20, 1335, 460, 7),$$

$$(d_0^*, d_1^*) = \left( \frac{1}{4200} \right) (488, 840),$$

$$(u_{0,0}, u_{0,1}, u_{0,2}, u_{0,3}) = \left( \frac{1}{16} \right) (5, 15, -5, 1),$$

$$(w_{0,0}, w_{0,1}, w_{0,2}, w_{0,3}) = \left( \frac{1}{256} \right) (-3, 13, 0, 0),$$

$$(u_{1,0}, u_{1,1}, u_{1,2}, u_{1,3}) = \left( \frac{1}{16} \right) (-1, 9, 9, -1),$$

$$(w_{1,0}, w_{1,1}, w_{1,2}, w_{1,3}) = \left( \frac{3}{256} \right) (0, -1, -1, 0),$$

While determining these parameters, we have set the free parameters  $w_{0,2}, w_{0,3}, w_{1,0}, w_{1,3} = 0$  for simplicity, and we have fixed  $b_3 = 1$  for the reason that when the discretization is written in a matrix form the inverse of the coefficient matrix ( $D$ ) would be available in [1]. We also note that then

$$T_k(h) = -\frac{h^{10}}{3024} y_k^{(10)} + O(h^{12}), \quad k = 2(1)N - 1,$$

and

$$T_1(h) = h^{10} \left[ \frac{1}{38400} y_0^{(10)} - \left\{ \frac{(61F_{1/2} + 9F_{3/2})}{46080} \right\} y_0^{(6)} \right] + O(h^{11}),$$

$$T_N(h) = h^{10} \left[ \frac{1}{38400} y_{N+1}^{(10)} - \left\{ \frac{(61F_{N+1/2} + 9F_{N-1/2})}{46080} \right\} y_{N+1}^{(6)} \right] + O(h^{11}),$$

where  $F = \partial f / \partial y$ .

Now, a method based on the discretizations (3a), (2) and (3b) can be expressed in the matrix form as

$$(5) \quad D\tilde{Y} + G(\tilde{Y}) = 0.$$

For the derivation of the above difference scheme, we have assumed that  $y \in C^{10}[a, b]$ , and for  $x \in [a, b]$ ,  $-\infty < y < \infty$ ,  $f$  is six times continuously differentiable and that  $\partial f / \partial y$  exists and is continuous.

Following arguments given in Usmani [1], we can show that if  $E = \tilde{Y} - Y$ , then in the uniform norm, for sufficiently small  $h$ ,

$$\|E\| = O(h^6),$$

provided  $U^* < 2592/(7K(b-a)^4)$ , where

$$K = \frac{181}{180} \quad \text{and} \quad U^* = \max \left| \frac{\partial f}{\partial y} \right|.$$

**Acknowledgement.** I would like to thank Professor M. M. Chawla for his helpful comments.

Department of Mathematics  
Indian Institute of Technology  
Hauz Khas, New Delhi-29, India

1. RIAZ A. USMANI, "Discrete variable methods for a boundary value problem with engineering applications," *Math. Comp.*, v. 32, 1978, pp. 1087-1096.
2. M. M. CHAWLA & C. P. KATTI, "Finite difference methods for two-point boundary value problems involving high order differential equations," *BIT*, v. 19, 1979, pp. 27-33.