Odd Perfect Numbers Not Divisible By 3. II

By Masao Kishore*

Abstract. We prove that odd perfect numbers not divisible by 3 have at least eleven distinct prime factors.

1. N is called a perfect number if $\sigma(N) = 2N$, where $\sigma(N)$ is the sum of positive divisors of N . Twenty-seven even perfect numbers are known; however, no odd perfect (OP) numbers have been found.

Suppose N is OP and $\omega(N)$ is the number of distinct prime factors of N . Gradstein (1925), Kühnel (1949), Weber (1951), and the author (1978, [5]) proved that $\omega(N) \geq 6$. Pomerance (1972, [7]) and Robbins (1972) proved that $\omega(N) \geq 7$. Hagis (1975, [2]) and Chein (1978, [1]) proved that $\omega(N) \geq 8$.

Hagis and McDaniel [3] proved that the largest prime factor of $N \geq 100129$, and Pomerance [8] proved that the second largest prime factor of $N \geq 139$.

If $3 \nmid N$, then Kanold (1949) proved that $\omega(N) \geq 9$, and the author (1977, [4]) proved that $\omega(N) \geq 10$.

In this paper we prove

THEOREM. *If N is OP and $3 \nmid N$, then $\omega(N) \geq 11$.*

2. In the remainder of this paper we assume that N is OP and

$$N = \prod_{i=1}^{10} p_i^{a_i},$$

where p_i 's are primes, $5 \leq p_1 < \cdots < p_{10}$ and a_i 's are positive integers, and we will get a contradiction. We write $p_i^{a_i} \parallel N$ and $a_i = V_{p_i}(N)$.

The following lemmas were proved in [4] and [7]:

LEMMA 1. *Suppose $p^a \parallel N$. Then*

(a) *All a 's are even except for one a in which case $a \equiv p \equiv 1(4)$. We write π for p .*

(b) *If $p \equiv 1(3)$, $a \not\equiv 2(3)$.*

(c) *If $p \equiv 1(4)$ and $p \equiv 2(3)$, then $p \neq \pi$.*

LEMMA 2. *Suppose $q = 5$ or 17 and $p^a \parallel N$. Then*

$$V_q(\sigma(p^a)) = \begin{cases} V_q(a+1) & \text{if } p \equiv 1(q), \\ V_q(p+1) + V_q(a+1) & \text{if } p \equiv -1(q) \text{ and } p = \pi, \\ 0 & \text{otherwise.} \end{cases}$$

Received November 2, 1981; revised June 21, 1982.

1980 *Mathematics Subject Classification.* Primary 10A20.

*Current address: Department of Mathematics, East Carolina University, Greenville, North Carolina 27834.

LEMMA 3. Suppose $p^a \parallel N$, q is a prime and $q^b \parallel a + 1$. Then N is divisible by at least c distinct primes $\equiv 1(q)$ other than p , where $c = b$ if $q^b = a + 1$, and $c = 2b$ if $q^b \neq a + 1$.

The proof of the next lemma is similar to that of Lemma 6 in [4].

LEMMA 4. $p_1 = 5, p_2 = 7, p_3 = 11, p_4 \leq 17, p_5 \leq 23, p_6 \leq 37, p_7 \leq 107, p_9 \geq 139, p_{10} \geq 100129$. If $p_7 \geq 103$, then $p_8 \leq 113$.

LEMMA 5. Suppose p, q are odd primes, a, b are positive integers, $p^b \mid q + 1, p \geq 5$ and $2b \geq a$. Then $q \nmid \sigma(p^a)$.

Proof. Since p and q are odd primes, $q \geq 2p^b - 1$. Suppose $\sigma(p^a) = mq$ for some integer m . Then $a \geq b$ and

$$\sigma(p^{b-1}) + m \equiv \sigma(p^a) + m = m(q + 1) \equiv o(p^b).$$

Hence

$$m \geq p^b - \sigma(p^{b-1}) = (p^{b+1} - 2p^b + 1)/(p - 1),$$

and

$$\begin{aligned} \sigma(p^a) = mq &\geq (2p^b - 1)(p^{b+1} - 2p^b + 1)/(p - 1) \\ &= (2p^{2b+1} - 4p^{2b} - p^{b+1} + 4p^b - 1)/(p - 1) \\ &> (p^{2b+1} - 1)/(p - 1) = \sigma(p^{2b}) \geq \sigma(p^a), \end{aligned}$$

because $p \geq 5$ and $2b \geq a$, a contradiction. Q.E.D.

Remark. Lemma 5 also holds if $p = 3$ and $2b > a$.

The next lemma is due to Hagis.

LEMMA 6. Suppose $p = 5$ or 17 and p^a is a component of an OP number. Then $\sigma(p^a)$ has at least one prime factor ≥ 100129 except

- (a) if $p = 5, a = 1, 2, 4, 5, 6, 8, 9, 13, 14, 17, 26, 29$.
- (b) If $p = 17, a = 1, 2, 4, 5, 9$.

COROLLARY 6. Suppose $p = 5$ or 17 and $p^a \parallel N$. Then $\sigma(p^a)$ has at least one prime factor ≥ 100129 except

- (a) if $p = 5, a = 2, 4, 6, 8$,
- (b) if $p = 17, a = 2, 4$.

Proof. We can easily show that $5^{14} \nmid N$ and $5^{26} \nmid N$ because $\sigma(5^{14}) = 11 \cdot 13 \cdot 71 \cdot 181 \cdot 1741$ and $\sigma(5^{26}) = 19 \cdot 31 \cdot 109 \cdot 271 \cdot 829 \cdot 4159 \cdot 31051$. Then Corollary 6 follows from Lemmas 1 and 6. Q.E.D.

LEMMA 7. If $17^a \parallel N$ and $a \geq 8$, then $p_9 \geq 100129, p_{10} \geq 2 \cdot 17^{a-3} - 1 > 2 \cdot 10^6$, and $17^{a-3} \mid \pi + 1$.

Proof. If p is a prime and $p \leq 113$, then $p \not\equiv \pm 1 (17)$ except for $p = 103$. Hence by Lemmas 1, 2 and 4 if $17 \mid \sigma(p_i^{a_i})$ for $1 \leq i \leq 7$, then $i = 7, p_7 = 103$ and $17 \nmid \sigma(p_8^{a_8})$.

Suppose $p_7 \neq 103$. Then $17^a \mid \sigma(p_8^{a_8} p_9^{a_9} p_{10}^{a_{10}})$. Since $a \geq 8, 17^3 \mid \sigma(p_i^{a_i})$ for some $8 \leq i \leq 10$. If $p_i \equiv 1 (17)$, then $17^3 \mid a_i + 1$, and by Lemma 3 N would have at least three more primes $\equiv 1 (17)$, a contradiction. Hence $p_i \equiv -1 (17)$ and $p_i = \pi$. Then by the same lemma $17^2 \nmid \sigma(p_j^{a_j})$ for $j \neq i, 8 \leq j \leq 10, 17^{a-2} \mid \sigma(p_i^{a_i}), 17^2 \nmid a_i + 1$,

and $17^{a-3} \mid p_i + 1$ by Lemma 2. By Lemma 5 $p_i \nmid \sigma(17^a)$, and by Corollary 6 $\sigma(17^a)$ has at least one prime factor ≥ 100129 .

The same arguments hold if $p_7 = 103$ because $17 \nmid \sigma(p_8^a)$. Q.E.D.

LEMMA 8. *If $5^a \parallel N$ and $a \geq 14$, then $a \geq 16$, $p_9 \geq 100129$ and $p_{10} \geq 579281$.*

Proof. We showed that $a \neq 14$ in the proof of Corollary 6. Suppose $p^b \parallel N$, $p \equiv 1$ (5) and $p \leq 107$. Then $p = 11, 31, 41, 61, 71$, or 101 . If $5 \mid \sigma(p^b)$, then by Lemma 2 $\sigma(p^4) \mid \sigma(p^b)$. Since $131 \cdot 21491 \mid \sigma(61^4)$, $211 \cdot 2221 \mid \sigma(71^4)$ and $31 \cdot 391 \cdot 1381 \mid \sigma(101^4)$, it is easy to show that if $5 \mid \sigma(p^b)$, $p \neq 61, 71$, or 101 .

Suppose $5 \nmid \sigma(41^b)$. Then $579281 \mid \sigma(41^4)$. Since the order of 5 mod 579281 is 72410 and a is even, $579281 \nmid \sigma(5^a)$, and $\sigma(5^a)$ has at least one prime factor ≥ 100129 by Corollary 6. Hence we may assume that $5 \nmid \sigma(41^b)$.

Since $3001 \cdot 3221 \cdot 24151 \mid \sigma(11^{24})$ and $101 \cdot 4951 \cdot 17351 \mid \sigma(31^{24})$, $5^2 \nmid \sigma(11^b)$ and $5^2 \nmid \sigma(31^b)$.

Suppose $5 \mid \sigma(11^b)$. Then $3221 \mid \sigma(11^4)$, and if $3221^c \parallel N$, $5^2 \nmid \sigma(3221^c)$ because $151 \cdot 601 \cdot 1301 \cdot 1601 \mid \sigma(3221^{24})$. Similarly, if $5 \mid \sigma(31^b)$, then $17351 \mid \sigma(31^4)$, and if $17351^c \parallel N$, $5^2 \nmid \sigma(17351^c)$ because $101 \cdot 2351 \mid \sigma(17351^{24})$.

Suppose $p^b \parallel N$, $p \equiv -1$ (5), and $p \leq 107$. Then $p = 19, 29, 59, 79$, or 89 , and by Lemma 1 $p \neq \pi$. Hence by Lemma 2 $5 \nmid \sigma(p^b)$.

In summary if $p^b \parallel N$, $p \leq 107$, and if $5 \mid \sigma(p^b)$, then $p = 11$ or 31 , in which case $q^c \parallel N$ where $q = 3221$ or 17351 and $5^2 \nmid \sigma(q^c)$.

Now we will show that $5^{a-8} \mid \pi + 1$. Suppose three $p_i \equiv 1$ (5) for $1 \leq i \leq 7$. Then $p_3 = 11$, $p_6 = 31$ and $p_7 = 41$, and it is easy to show that $41 < p_8 \leq 61$. Hence $5 \nmid \sigma(p_8^a)$. Since $p_{10} \geq 100129$, the above summary shows that $5^2 \nmid \sigma(\prod_{i=1}^8 p_i^{a_i})$. Suppose $5^a \parallel \sigma(p_9^{a_9} p_{10}^{a_{10}})$. By a similar argument used in the proof of Lemma 7 we have for $i = 9$ or 10 $5^{a-4} \mid \sigma(p_i^{a_i})$, $p_i = \pi$, and $5^{a-6} \mid \pi + 1$. Suppose $5^{a-1} \parallel \sigma(p_9^{a_9} p_{10}^{a_{10}})$. Then $p_9 = 3221$ or 17351 and $5^2 \nmid \sigma(p_9^{a_9})$, $5^{a-2} \mid \sigma(p_{10}^{a_{10}})$, $p_{10} = \pi$, and $5^{a-4} \mid \pi + 1$.

Similar arguments show that if two $p_i \equiv 1$ (5) for $1 \leq i \leq 7$, then $5^{a-8} \mid \pi + 1$ and that if $p_i \equiv 1$ (5) for $1 \leq i \leq 7$, then $5^{a-5} \mid \pi + 1$.

Since $a \geq 16$ and $5^{a-8} \mid \pi + 1$, $\pi \nmid \sigma(5^a)$ by Lemma 5, $\pi \geq 2 \cdot 5^{a-8} - 1 > 7 \cdot 10^5 > 579281$, and Lemma 8 follows from Corollary 6. Q.E.D.

COROLLARY 8. *Suppose $5^a \parallel N$, $a \geq 14$, and $579281 \nmid N$ if $41 \mid N$. Then $a \geq 16$, $p_9 \geq 100129$, $p_{10} \geq 2 \cdot 5^{a-8} - 1 > 7 \cdot 10^5$, and $5^{a-8} \mid \pi + 1$.*

The next lemma is due to McDaniel [6].

LEMMA 9. *Suppose $a \geq 2$, $a + 1$ is a prime, and p is a prime.*

- (a) *If $p^2 \mid \sigma(5^a)$, then $p > 2^{29}$.*
- (b) *If $p^2 \mid \sigma(17^a)$, then $p > 2^{27}$ or $p = 48947$.*

LEMMA 10. *Suppose $p^a \parallel N$, $q \mid \sigma(p^b)$, $b + 1 \mid a + 1$, $q \leq 107$, and $q, b + 1$ are primes. Then*

- (a) *If $p = 5$, $q = 11, 31, 59$, or 71 .*
- (b) *If $p = 17$, $q = 47, 59$, or 83 .*

Proof. Suppose $p = 5$ or 17 , and d is the order of p mod q . Then $p^d \equiv 1$ (q), and $d \mid b + 1$. Since $p \not\equiv 1$ (q), $d > 1$, and $d = b + 1$ because $b + 1$ is an odd prime. The order d is not an odd prime except for those q stated above. Q.E.D.

LEMMA 11. Suppose $17^a \parallel N$, $a \geq 8$, and $307 \nmid N$. If $p_8 < 1000$, then $a \geq 10$, $p_9 \geq 25646167$, and $p_{10} > 8 \cdot 10^8$.

Proof. Since $307 \mid \sigma(17^8)$, $a \geq 10$, and by Lemma 7, $17^{a-3} \mid \pi + 1$ and $p_{10} \geq 2 \cdot 17^{a-3} - 1 > 8 \cdot 10^8$. Suppose $p_8 < 1000$ and $100129 \leq p_9 < 25646167 < 2^{27}$. Choose b such that $b + 1$ is a prime and $b + 1 \mid a + 1$. Then $\sigma(17^b) \mid \sigma(17^a)$, and $b \neq 2, 4$, or 6 because $307 \mid \sigma(17^2)$, $88741 \mid \sigma(17^4)$ and $25646167 \mid \sigma(17^6)$. Hence $b \geq 10$. If $1 \leq i \leq 7$ and $p_i \mid \sigma(17^b)$, then by Lemmas 4, $10 \mid i = 7$ and $p_7 = 47, 59$, or 83 . Since $\pi = p_{10} \nmid \sigma(17^a)$ by Lemma 5, we have $\sigma(17^b) \mid p_7 p_8 p_9$ by Lemma 9. Then $\sigma(17^{10}) \leq 83 \cdot 1000 \cdot p_9$, or $p_9 > 25646167$, a contradiction. Hence $p_9 \geq 25646167$. Q.E.D.

COROLLARY 11. If $p_7 \leq 29$ and $p_8 < 6203$, then $a \geq 10$, $p_9 \geq 25646167$, and $p_{10} > 8 \cdot 10^8$.

Proof. As in Lemma 11 $\sigma(17^b) \mid p_8 p_9$. Then $\sigma(17^{10}) \leq 6203 \cdot p_9$, or $p_9 > 25646167$, a contradiction. Q.E.D.

LEMMA 12. Suppose $5^a \parallel N$, $a \geq 14$, $579281 \nmid N$ if $41 \mid N$, and $p \nmid N$ if $p = 31, 71, 191, 409$, or 19531 . If $p_8 \leq 41$, then $a \geq 22$, $p_9 \geq 12207031$, and $p_{10} \geq 2 \cdot 5^{a-8} - 1 > 10^{10}$.

Proof. $a \geq 22$ because $a \neq 14$ as before, $409 \mid \sigma(5^{16})$, $191 \mid \sigma(5^{18})$, and $19531 \mid \sigma(5^{20})$. The rest of the proof is similar to that of Lemma 11 using $\sigma(5^{10}) = 12207031$ and $\sigma(5^{12}) = 307175781$. Q.E.D.

COROLLARY 12. If $p_7 \leq 29$ and $p_8 < 6203$, then $a \geq 22$, $p_9 \geq 12207031$, and $p_{10} \geq 2 \cdot 5^{a-8} - 1 > 10^{10}$.

LEMMA 13. Suppose $5^a \parallel N$, $a = 10$ or 12 , $p_8 \leq 151$, $p_9 > 3011$, at most two $p_i \equiv 1 \pmod{5}$ for $1 \leq i \leq 8$, $p_i = 11, 31, 41$, or 151 if $p_i \equiv 1 \pmod{5}$ and $1 \leq i \leq 8$, $p_i = 19, 29, 59, 79, 89, 109$, or 149 if $p_i \equiv -1 \pmod{5}$ and $1 \leq i \leq 8$, $p \nmid N$ if $p = 131, 3221$, or 17351 , and $579281 \nmid N$ if $41 \mid N$. Then $p_9 \geq 3 \cdot 10^6$, and $p_{10} \geq 12207031$.

Proof. Suppose $3011 < p_9 < 3 \cdot 10^6$. Then $p_{10} = \sigma(5^a) = 12207031$ or 305175781 , and $5 \nmid \sigma(p_{10}^{a_{10}})$ because $131 \mid \sigma(12207031^4)$ and $3011 \mid \sigma(305175781^4)$. Suppose $5 \mid \sigma(p_i^{a_i})$, $1 \leq i \leq 8$. Since $3221 \mid \sigma(11^4)$, $17351 \mid \sigma(31^4)$, $579281 \mid \sigma(41^4)$ and $104670301 \mid \sigma(151^4)$, $p_i \not\equiv 1 \pmod{5}$. Since $p_i \neq \pi$ if $p_i = 19, 29, 59, 79, 89$, and 149 by Lemma 1, we have $p_8 = 109 = \pi$ by Lemmas 2, 4. If $5^4 \mid \sigma(p_8^{a_8})$, then $5^3 \mid a_8 + 1$ by Lemma 2, and N would have at least six prime factors $\equiv 1 \pmod{5}$, a contradiction. Hence $5^4 \nmid \sigma(p_8^{a_8})$, and so $5^{a-3} \mid \sigma(p_9^{a_9})$, $p_9 \neq \pi$, $p_9 \equiv 1 \pmod{5}$ and N would have at least 7 more prime factors $\equiv 1 \pmod{5}$, a contradiction. Hence $5 \nmid \sigma(p_i^{a_i})$ for $1 \leq i \leq 8$.

Then $5^a \mid \sigma(p_9^{a_9})$, $p_9 = \pi$, $5^2 \mid a_9 + 1$, $5^{a-1} \mid p_9 + 1$, and $p_9 \geq 2 \cdot 5^{a-1} - 1 > 3 \cdot 10^6$, a contradiction. Q.E.D.

Definition. Suppose $M = \prod_{i=1}^r p_i^{a_i}$. Then

$$S(M) = \sigma(M)/M,$$

$$a(p_i) = \min\{a_i \mid p_i^{a_i+1} > 10^{10} \text{ where } p_i^{a_i} \text{ satisfies the restrictions}$$

implied by Lemma 1\},

$$b_i = \begin{cases} a_i & \text{if } a_i < a(p_i), \\ a(p_i) & \text{if } a_i \geq a(p_i). \end{cases}$$

LEMMA 14. Suppose $M = \prod_{i=1}^{10} p_i^{b_i}$. Then $0 \leq \log 2 - \log S(M) < 10 \cdot 10^{-10}$.

Proof. Suppose $p^a \parallel N$ and $a \geq a(p)$. Then

$$\begin{aligned} 0 &\leq \log S(p^a) - \log S(p^{a(p)}) < \log \frac{p}{p-1} - \log \frac{p^{a(p)+1} - 1}{p^{a(p)}(p-1)} \\ &= \log \frac{p^{a(p)+1}}{p^{a(p)+1} - 1} = \log \left(1 + \frac{1}{p^{a(p)+1} - 1} \right) \\ &< \frac{1}{p^{a(p)+1} - 1} \leq 10^{-10}. \end{aligned}$$

Hence

$$\begin{aligned} 0 &\leq \log S(N) - \log S(M) \\ &\leq \sum_{i=1}^{10} |\log S(p_i^{a_i}) - \log S(p_i^{b_i})| < 10 \cdot 10^{-10}. \end{aligned}$$

Q.E.D.

The proof of the next two lemmas is easy.

LEMMA 15. If q is a prime, $q \mid \sigma(p_i^{a_i})$ for some $1 \leq i \leq 7$ with $a_i < a(p_i)$, and if $q \leq p_7$, then $q = 2$ or $q = p_i$ for some $1 \leq i \leq 7$.

LEMMA 16. Suppose $M = \prod_{i=1}^7 p_i^{b_i}$ and $L = M \cdot \prod_{j=1}^r q_j^{c_j}$ where q_j is a prime, $q_j > p_7$, $q_j \mid \sigma(p_i^{b_i})$ for some $1 \leq i \leq 7$ with $b_i < a(p_i)$, $q_1 < \dots < q_r$, and c_j is the minimum allowable power of q_j determined by Lemma 1. If there is no such q_j , then $r = 0$ and the product is defined to be 1. Then

(a) $r \leq 3$ and $\log S(L) \leq \log 2$.

(b) If $r = 3$, then $p_8 = q_1, p_9 = q_2, p_{10} = q_3$ and

$$\log 2 < \log S(M) + 7 \cdot 10^{-10} + \sum_{j=1}^3 \log q_j / (q_j - 1).$$

(c) If $r = 2$ and $q_2 < 100129$, then $p_8 = q_1, p_9 = q_2$ and

$$\log 2 < \log S(M) + 7 \cdot 10^{-10} + \sum_{j=1}^2 \log q_j / (q_j - 1) + \log 100129 / 100128.$$

LEMMA 17. $p_8 < 3011$.

Proof. Suppose $p_8 \geq 3011$. We used a computer (PDP 11/70 at the University of Toledo) to find $M = \prod_{i=1}^7 p_i^{b_i}$ satisfying Lemmas 1, 4, 15, 16, $\log S(M) < \log 2$, and

$$\begin{aligned} \log 2 &< \log S(M) + 7 \cdot 10^{-10} + \log 3011 / 3010 \\ &\quad + \log 3019 / 3018 + \log 100129 / 100128. \end{aligned}$$

The results were:

$$\begin{aligned} &5^{14}7^{12}11^{10}13^917^823^{a_6}29^6, \\ &5^{14}7^{12}11^{10}13^917^623^{a_6}29^6, \\ &5^{14}7^{12}11^{10}13^617^823^{a_6}29^6, \\ &5^{12}7^{12}11^{10}13^917^823^{a_6}29^6, \\ &5^{10}7^{12}11^{10}13^917^823^{a_6}29^6, \end{aligned}$$

where $a_6 = 6$ or 8 . Since

$$\frac{5}{4} \frac{7}{6} \frac{11}{10} \frac{13}{12} \frac{17}{16} \frac{23}{22} \frac{29}{28} \frac{6203}{6202} \frac{6211}{6210} \frac{100129}{100128} < 2,$$

$3011 \leq p_8 < 6203$. By Corollaries 11 and 12 $p_9 \geq \min\{25646167, 12207031\}$ and $p_{10} \geq \min\{8 \cdot 10^8, 10^{10}\}$. Then N is not OP because

$$\frac{5}{4} \frac{7}{6} \frac{11}{10} \frac{13}{12} \frac{17}{16} \frac{23}{22} \frac{29}{28} \frac{3011}{3010} \frac{12207031}{12207030} \frac{800000000}{799999999} < 2. \quad \text{Q.E.D.}$$

The proof of the next lemma is also easy.

LEMMA 18. Suppose $M = \prod_{i=1}^9 p_i^{b_i}$, $q = \max\{p \mid p \text{ is a prime and } \log S(M) + \log S(p^a) \geq \log 2 \text{ where } a \text{ is the minimum allowable power of } p\}$ and $r = \min\{p \mid p \text{ is a prime and } \log S(M) + 9 \cdot 10^{-9} + \log p/(p-1) < \log 2\}$. Then $q < p_{10} < r$; in particular, if there are no primes between q and r , N is not OP.

LEMMA 19. $p_9 < 3011$.

Proof. Suppose $p_8 < 3011 \leq p_9$. We used a computer to find $M = \prod_{i=1}^8 p_i^{b_i}$ satisfying Lemmas 1, 4, 7, 8, 15, 16, $\log S(M) < \log 2$, and

$$\log 2 < \log S(M) + 8 \cdot 10^{-10} + \log 3011/3010 + \log 100129/100128.$$

There were seventy-two such M 's. However, none of them satisfied Lemmas 11, 12 and 13 except

$$5^2 7^{12} 11^{10} 13^{10} 19^{10} 23^8 31^6 59^6.$$

It is easy to show that $7753 \leq p_9 \leq 8389$, $a_2 \geq 22$ ($\sigma(7^{12})$ is a prime), $a_3 \geq 16$, $a_4 \geq 16$, $a_5 = 10$ or ≥ 16 (if $\sigma(19^{10})$ is a prime, $a_5 \geq 16$), $a_6 \geq 12$, $a_7 \geq 12$, and $a_8 \geq 12$. Then for each p_9 with $7753 \leq p_9 \leq 8389$ Lemma 18 is not satisfied. Hence N is not OP. Q.E.D.

Proof of Theorem. By Lemma 19 $p_9 < 3011$. We used a computer to find $M = \prod_{i=1}^9 p_i^{b_i}$ satisfying Lemmas 1, 4, 7, 8, 15, 16, $\log S(M) < \log 2$, and

$$\log 2 < \log S(M) + 9 \cdot 10^{-10} + \log 100129/100129.$$

There were thirty-nine such M 's; however, none of them satisfied Lemma 18. Hence N is not OP. Q.E.D.

Acknowledgement. The author would like to thank Professor P. Hagis, Jr. for supplying the table in Lemma 6.

Department of Mathematics
The University of Toledo
Toledo, Ohio 43606

1. J. E. Z. CHEIN, *An Odd Perfect Number Has at Least 8 Prime Factors*, Ph.D. Thesis, Pennsylvania State University, 1979.
2. P. HAGIS, JR., "Outline of a proof that every odd perfect number has at least eight prime factors," *Math. Comp.*, v. 35, 1980, pp. 1027–1032.
3. P. HAGIS, JR. & W. L. MCDANIEL, "On the largest prime divisor of an odd perfect number. II," *Math. Comp.*, v. 29, 1975, pp. 922–924.
4. M. KISHORE, "Odd perfect numbers not divisible by 3 are divisible by at least ten distinct primes," *Math. Comp.*, v. 31, 1977, pp. 274 – 279.
5. M. KISHORE, "Odd integers N with five distinct prime factors for which $2 - 10^{-12} < \sigma(N)/N < 2 + 10^{-12}$," *Math. Comp.*, v. 32, 1978, pp. 303–309.
6. W. L. MCDANIEL, "On multiple prime divisors of cyclotomic polynomials," *Math. Comp.*, v. 28, 1974, pp. 847–850.
7. C. POMERANCE, "Odd perfect numbers are divisible by at least seven distinct primes," *Acta Arith.*, v. 25, 1973/74, pp. 265–300.
8. C. POMERANCE, "The second largest prime factor of an odd perfect number," *Math. Comp.*, v. 29, 1975, pp. 914–921.