A Lower Bound for the Class Number of Certain Cubic Number Fields

By Günter Lettl

Abstract. Let K be a cyclic number field with generating polynomial

$$X^3 - \frac{a-3}{2}X^2 - \frac{a+3}{2}X - 1$$

and conductor m . We will derive a lower bound for the class number of these fields and list all such fields with prime conductor $m = (a^2 + 27)/4$ or $m = (1 + 27b^2)/4$ and small class number.

1. Introduction. Let h_m denote the class number of the cyclotomic field $\mathbf{Q}(\zeta_m)$, and h_m^+ , the class number of its maximal real subfield $\mathbf{Q}(\cos(2\pi/m))$. It is a well-known conjecture of Vandiver that $p \nmid h_p^+$ holds for all primes $p \in \mathbf{P}$. This is a customary assumption for proving the second case of Fermat's Last Theorem (for more details see Washington [16]). Since h_p^+ grows slowly ($h_p^+ = 1$ for $p < 163$ with the use of the Generalized Riemann Hypothesis (GRH), van der Linden [10]), for no p with $h_p^+ > 1$ the exact value of h_p^+ is known without using GRH. Masley suggested that perhaps $h_p^+ < p$ always holds, but a counterexample was found in [3], [12]. The class number of each real subfield of $\mathbf{Q}(\zeta_p)$ divides h_p^+ , and in this way one can find primes with $h_p^+ > 1$. Using the quadratic subfield, Ankeny, Chowla and Hasse [1] showed that $h_p^+ > 1$ if p belongs to certain quadratic sequences in $\mathbf{N}$, and S.-D. Lang [9] and Takeuchi [15] found more such sequences. Similar results were obtained for h_{4p}^+ by Yokoi [17]. Using the cubic subfield of $\mathbf{Q}(\zeta_p)$, which has been thoroughly investigated (e.g., [2], [5], [8]), the theorem of the present paper yields the following results:

If a is an odd integer, $a > 23$, and $p = (a^2 + 27)/4$, a prime, then $h_p^+ > 1$.

If b is an odd integer, $b > 7$, and $p = (1 + 27b^2)/4$, a prime, then $h_p^+ > 1$.

A conjecture about primes in quadratic sequences (Hardy and Wright [7, I.2.8]) implies that there exist infinitely many primes p of each of these two forms, because one can write

$$\frac{a^2 + 27}{4} = \left(\frac{a-3}{2}\right)^2 + 3\left(\frac{a-3}{2}\right) + 9$$

and

$$\frac{1 + 27b^2}{4} = 3\left(\frac{3b-1}{2}\right)^2 + 3\left(\frac{3b-1}{2}\right) + 1.$$

Received November 6, 1984; revised July 15, 1985.

1980 *Mathematics Subject Classification.* Primary 12A30, 12A50.

©1986 American Mathematical Society
0025-5718/86 \$1.00 + \$.25 per page

2. Class Number Bounds and Main Results. Let K be a cyclic cubic number field with conductor m and class number h . It is well known that m is the product of distinct primes, which are congruent to 1 mod (6), and of 9, if 3 ramifies in K . The class number h is congruent to 1 mod (3), if m is a prime or $m = 9$, and h is divisible by 3 otherwise. Set $f(s) = L(s, \chi) \cdot L(s, \bar{\chi})$ for $s \in \mathbb{C}$, where χ and $\bar{\chi}$ are the nontrivial cubic Dirichlet characters modulo (m) belonging to K . Since the discriminant of K equals m^2 , the analytic class number formula yields

$$(1) \quad h = \frac{m \cdot f(1)}{4 \cdot R},$$

where R is the regulator of K . Moser [11] showed that for prime conductors, $h < m/3$ holds, so cubic fields will never lead to a contradiction to Vandiver's conjecture. Our aim is to establish a lower bound for the class number of a special family of cubic fields and to list all fields of some special types with prime conductor and small class number. From a result of Stark [14] one can deduce $f(1) > c/\log m$, where c is effectively computable, but this bound is not suited for our purposes. From the results of the next section we will obtain:

$$(2) \quad \text{If } K \text{ is a cyclic cubic number field with conductor } m > 10^5, \text{ then} \\ f(1) > 0.023 \cdot m^{-0.054}.$$

The harder problem is to find an upper bound for the regulator, which is only achieved for the following family of cyclic cubic fields. The polynomial

$$(3) \quad f_a(X) = X^3 - \frac{a-3}{2}X^2 - \frac{a+3}{2}X - 1, \quad a \in \mathbb{N} \text{ odd},$$

is irreducible over $\mathbb{Q}$, has discriminant $D(f_a) = ((a^2 + 27)/4)^2$, and if ϵ is a zero of f_a , the other zeros are $\epsilon' = -1/(\epsilon + 1)$ and $\epsilon'' = -(\epsilon + 1)/\epsilon$. Therefore, f_a is a generating polynomial of a cyclic cubic field K with conductor m , and we define $k \in \mathbb{N}$ by $\sqrt{D(f_a)} = (a^2 + 27)/4 = km$.

We call the field K of type A, if $k = 1$, and of type B, if $k = 27$ and $a = 27b$ with $b \in \mathbb{N}$ odd, $b \neq 1$ (in this case we have $m = (1 + 27b^2)/4$). It is well known that fields of type A or B have relatively large class numbers (see, for example, the tables of Gras [5]). Shanks [13] states that for cubic fields of type A with prime conductor "a rough mean value for h is given by $h \approx 12m/35(\log m)^2$ ".

LEMMA 1. *Let K be a cyclic cubic field with generating polynomial f_a , conductor m and regulator R . Then,*

$$(4) \quad 4R < \left(\frac{1}{2} \log D(f_a)\right)^2 = (\log(km))^2.$$

Proof of Lemma 1. Since the zeros $\epsilon, \epsilon', \epsilon''$ of f_a are units of K , we can estimate the regulator of K by $R \leq \text{Reg}(\{\epsilon, \epsilon'\}) =: R'$, if $R' \neq 0$ (see Lemma 4.15 in [16]). Choosing

$$\epsilon = \frac{a - 3 + 4\sqrt{km} \cdot \cos(1/3 \cdot \arctan(\sqrt{27}/a))}{6} \sim \sqrt{km}$$

with the principal value of arctan, we obtain

$$R' = \left| \det \begin{pmatrix} \log|\epsilon| & \log|\epsilon'| \\ \log|\epsilon'| & \log|\epsilon''| \end{pmatrix} \right| \\ = (\log|\epsilon + 1|)^2 - \log|\epsilon + 1| \log|\epsilon| + (\log|\epsilon|)^2.$$

Series expansions yield

$$R' = \frac{1}{4}(\log km)^2 - \frac{3 \log(km)}{2km} + \frac{3}{4km} + O\left(\frac{\log km}{(km)^2}\right)$$

and elementary calculus explicitly gives (4).

With (2) and Lemma 1 we immediately obtain from (1):

Let K be a cyclic cubic field with conductor $m > 10^5$ and with generating polynomial f_a . Then,

$$(5) \quad h > 0.023 \frac{m^{0.946}}{(\log km)^2}.$$

THEOREM. (a) Let K be a cyclic cubic field of type A with prime conductor m . Then $h < 16$ holds only for the following values of m :

h	m
1	7, 13, 19, 37, 79, 97, 139
4	163, 349, 607, 709, 937
7	313, 877, 1129, 1567, 1987, 2557
13	1063

(b) Let K be a cyclic cubic field of type B with prime conductor m . Then $h < 43$ holds only for the following values of m :

h	m
1	61, 331
4	547, 1951
7	2437, 3571
13	9241
28	4219, 25117
31	23497
37	8269

Proof of the Theorem. From (5) we obtain $h > 14$ for fields of type A with $m \geq 169339$, and $h > 37.2$ for fields of type B with $m > 10^6$. It is well known (see, e.g., Gras [4]) that primes $q \equiv -1 \pmod{3}$ divide the class number of a cyclic cubic field only with an even exponent. The table of class numbers of Shanks [13], and Table 1 below, complete the proof of the theorem.

TABLE 1
Class numbers of cyclic cubic fields of type B
with prime conductor $m < 10^6$

b	$m = \frac{1 + 27b^2}{4}$	h	b	$m = \frac{1 + 27b^2}{4}$	h
3	61	1	173	202021	$316 = 2^2 \cdot 79$
7	331	1	185	231019	$343 = 7^3$
9	547	$4 = 2^2$	189	241117	$1216 = 2^6 \cdot 19$
17	1951	$4 = 2^2$	191	246247	$175 = 5^2 \cdot 7$
19	2437	7	193	251431	$247 = 13 \cdot 19$
23	3571	7	199	267307	$196 = 2^2 \cdot 7^2$
25	4219	$28 = 2^2 \cdot 7$	205	283669	541
33	7351	$49 = 7^2$	221	329677	$316 = 2^2 \cdot 79$
35	8269	37	227	347821	331
37	9241	13	231	360187	$1732 = 2^2 \cdot 433$
39	10267	$49 = 7^2$	235	372769	$553 = 7 \cdot 79$
45	13669	109	243	398581	$1075 = 5^2 \cdot 43$
59	23497	31	259	452797	769
61	25117	$28 = 2^2 \cdot 7$	261	459817	$2257 = 37 \cdot 61$
91	55897	$133 = 7 \cdot 19$	297	595411	$2299 = 11^2 \cdot 19$
95	60919	193	299	603457	739
105	74419	$688 = 2^4 \cdot 43$	301	611557	$889 = 7 \cdot 127$
115	89269	211	303	619711	$1156 = 2^2 \cdot 17^2$
117	92401	$532 = 2^2 \cdot 7 \cdot 19$	305	627919	$1552 = 2^4 \cdot 97$
123	102121	307	341	784897	$688 = 2^4 \cdot 43$
129	112327	$604 = 2^2 \cdot 151$	347	812761	769
131	115837	$148 = 2^2 \cdot 37$	361	879667	$688 = 2^4 \cdot 43$
137	126691	97	367	909151	787
147	145861	$652 = 2^2 \cdot 163$	371	929077	$1588 = 2^2 \cdot 397$
159	170647	$628 = 2^2 \cdot 157$	373	939121	661
			383	990151	$532 = 2^2 \cdot 7 \cdot 19$

The class numbers of Table 1 were calculated with a "Sirius 1 Personal Computer", using the analytic class number formula (1). We also used that for fields of type B the roots of f_a are already fundamental units, and therefore $R = R'$ can be calculated with the explicit formula for ϵ , given in the proof of Lemma 1. In the following way it can be proved that ϵ is a fundamental unit:

Let K be a field of type B with generating polynomial f_a , $a = 27b$ and $m = (1 + 27b^2)/4$. Hasse [8] investigated the arithmetic of cyclic cubic fields, using the Gauss sums of the corresponding Dirichlet characters. With Hasse's notation, every integer $\alpha \in K$ can be written as $\alpha = [x, y]$ with $x \in \mathbf{Z}$, $y \in \mathbf{Z}[\rho]$, where $\rho^2 + \rho + 1 = 0$, and $x \equiv y \pmod{1 - \rho}$. If α is a unit of K , $N(\alpha) = 1$ implies $x^3 \equiv 27 \pmod{m}$ and $|x| \leq 2\sqrt{my\bar{y}}$ (Satz 8, [8]). For the roots of f_{27b} we have $\epsilon = [(27b - 3)/2, 3i\sqrt{3}]$ and its conjugates. Since Godwin's conjecture about fundamental units holds for cyclic cubic fields with $m > 9$ (see Gras [6]), we have to show:

There exists no unit $\alpha = [x, y] \in K$, $\alpha \neq \pm 1$, with $my\bar{y} = \frac{1}{2} \text{tr}(\alpha - \alpha')^2$
 $< \frac{1}{2} \text{tr}(\epsilon - \epsilon')^2 = 27m$, where tr denotes the trace from K to $\mathbf{Q}$.

Suppose the contrary. Then $x^3 \equiv 27 \pmod{m}$ and $|x| < 2\sqrt{27m}$ imply $x \in \{3, (27b - 3)/2, -(27b + 3)/2\}$ for $b \geq 7$. Considering $0 \equiv x \equiv y \pmod{1 - \rho}$ and $y\bar{y} < 27$ yields only a few possibilities for $y \in \mathbf{Z}[\rho]$, and one can check that for each

of these y , $N(\alpha) = 1$ has no solution $\alpha \neq 1$. For small values of b , one can consult the table in [5].

In the same way, but with much less computation, one can prove that for $k = 1$ (type A) and $k = 3$ the roots of f_a are also fundamental units. In these cases one has $\varepsilon = [(a - 3)/2, \pm 1]$ with $(a - 3)/2 \equiv \pm 1 \pmod{(3)}$, and $\varepsilon = [(9b - 3)/2, i\sqrt{3}]$, respectively.

3. A Lower Bound for $L(1, \chi) \cdot L(1, \bar{\chi})$. Let m be the conductor of a cyclic cubic field K , χ and $\bar{\chi}$ the nontrivial cubic Dirichlet characters modulo m associated with K , and $f(s) = L(s, \chi)L(s, \bar{\chi})$. To find a lower bound for $f(1)$, we first need an upper bound for $|f(s)|$ in a disk in $\mathbb{C}$ containing 1. Consider $C = C(\mu, \rho) = \{s \in \mathbb{C} \mid |s - \mu| < \rho\}$ with $1 < \mu$ and $\mu - 1 < \rho < \mu$, and set $\sigma_0 = \mu - \rho$. Let $s = \sigma + it \in \mathbb{C}$. For $\sigma > 0$ we have the representation

$$L(s, \chi) = \sum_{n=1}^{m-1} \frac{\chi(n)}{n^s} + s \cdot \int_m^\infty \frac{S(x, \chi)}{x^{s+1}} dx \quad \text{with } S(x, \chi) = \sum_{1 \leq n < x} \chi(n)$$

(see [16, p. 211]). The inequality of Pólya-Vinogradov [16, Lemma 11.8] states that $|S(x, \chi)| < \sqrt{m} \cdot \log m$. For $s \in C(\mu, \rho)$, the function $|s|/\sigma = 1/\cos(\arg s)$ attains its maximum $\mu/\sqrt{\mu^2 - \rho^2}$ if s is the point of contact of a tangent of C through 0. Combining these results, we obtain for every $s \in C(\mu, \rho)$:

$$\begin{aligned} |L(s, \chi)| &< 1 + \int_1^m \frac{1}{x^{\sigma_0}} dx + |s| \sqrt{m} \cdot \log m \int_m^\infty \frac{1}{x^{\sigma+1}} dx \\ &< \frac{1}{1 - \sigma_0} m^{1-\sigma_0} + \frac{\mu}{\sqrt{\mu^2 - \rho^2}} \log m \cdot m^{0.5-\sigma_0}. \end{aligned}$$

Since $\log x/\sqrt{x}$ is monotone decreasing for $x \geq e^2$, we conclude that for $m \geq m_0 \geq e^2$,

$$(6) \quad |f(s)| < c_1 \cdot m^{2-2\sigma_0}$$

holds for all $s \in C(\mu, \rho)$, with

$$c_1 = \left(\frac{1}{1 - \sigma_0} + \frac{\mu}{\sqrt{\mu^2 - \rho^2}} \cdot \frac{\log m_0}{\sqrt{m_0}} \right)^2.$$

LEMMA 2. *If K is a cyclic cubic number field with conductor m , then $f(1) > c_6 \cdot m^{-c_7}$, with $c_6, c_7 > 0$ as given in the course of the proof. Furthermore, c_7 can be made arbitrarily small.*

Proof of Lemma 2. The proof follows mainly Washington [16, pp. 212–214]. Let $\zeta(s)$ be the Riemann zeta function and $\zeta_K(s) = \zeta(s)f(s)$ the zeta function of the cyclic cubic field K with conductor m . If $s = \sigma + it \in \mathbb{C}$, we have

$$\zeta_K(s) = 1 + \sum_{n=2}^\infty \frac{a_n}{n^s} \quad \text{for } \sigma > 1,$$

with $a_n \geq 0$, and $a_n \geq 1$ if n is a cube. Developing ζ_K in a power series around $\mu > 1$ gives

$$\zeta_K(s) = \sum_{j=0}^\infty b_j(\mu - s)^j,$$

with

$$(7) \quad b_0 = \zeta_K(\mu) > \zeta(3\mu) > 1 \quad \text{and} \quad b_j = \frac{1}{j!} \sum_{n=2}^{\infty} (\log n)^j \cdot \frac{a_n}{n^\mu} > 0 \quad \text{for } j \geq 1.$$

The integral representation of $\zeta(s)$ for $\sigma > 0$ yields

$$|\zeta(s)| \leq \left| \frac{s}{s-1} \right| + |s| \int_1^\infty \frac{1}{u^{\sigma+1}} du = \left| \frac{s}{s-1} \right| + \frac{|s|}{\sigma}$$

and

$$\begin{aligned} |\zeta(s)| &\leq \left| \frac{s}{s-1} \right| + |s| \cdot \sum_{n=1}^{\infty} \frac{1}{n^{\sigma+1}} \cdot \int_n^{n+1} (u - [u]) du \\ &< \left| \frac{s}{s-1} \right| + \frac{|s|}{2} \left(1 + \frac{1}{\sigma} \right). \end{aligned}$$

Let $C = C(\mu, \rho)$, with $\mu - 1 < \rho < \mu$, be the disk with center μ and radius ρ , and denote its boundary by ∂C . Using (6), we get for all $s \in \partial C$:

$$(8) \quad \left| \zeta_K(s) - \frac{f(1)}{s-1} \right| \leq |\zeta(s)| \cdot |f(s)| + \frac{1}{|s-1|} \cdot |f(1)| < c_2 \cdot m^{2-2\sigma_0},$$

with

$$c_2 = c_1 \cdot \max_{s \in \partial C} \left(\frac{|s|+1}{|s-1|} + |s| \cdot \min \left\{ \frac{1}{\sigma}, \frac{1}{2} \left(1 + \frac{1}{\sigma} \right) \right\} \right).$$

Since $\zeta_K(s) - f(1)/(s-1)$ is holomorphic in the whole complex plane, (8) holds for all $s \in C(\mu, \rho)$. Computing the coefficients of

$$\zeta_K(s) - \frac{f(1)}{s-1} = \sum_{j=0}^{\infty} \left(b_j - \frac{f(1)}{(\mu-1)^{j+1}} \right) \cdot (\mu-s)^j$$

with a Cauchy integral gives

$$\left| b_j - \frac{f(1)}{(\mu-1)^{j+1}} \right| = \left| \frac{1}{2\pi i} \int_{\partial C} \left(\zeta_K(s) - \frac{f(1)}{s-1} \right) \frac{ds}{(s-\mu)^{j+1}} \right| < \frac{c_2}{\rho^j} \cdot m^{2-2\sigma_0}.$$

For $0 < \sigma < 1$, the integral representation of $\zeta(s)$, and $f(\sigma) = |L(\sigma, \chi)|^2$, show that $\zeta_K(\sigma) \leq 0$. So for any α with $\sigma_0 < \alpha < 1$, and any $\nu \in \mathbf{R}^+$ with $1 < \nu$, we have

$$\begin{aligned} -\frac{f(1)}{\alpha-1} &\geq \zeta_K(\alpha) - \frac{f(1)}{\alpha-1} > \sum_{j=0}^{[\nu]-1} \left(b_j - \frac{f(1)}{(\mu-1)^{j+1}} \right) \cdot (\mu-\alpha)^j \\ &\quad - c_2 \cdot m^{2-2\sigma_0} \cdot \sum_{j=[\nu]}^{\infty} \left(\frac{\mu-\alpha}{\rho} \right)^j \\ &\geq c_3 - \frac{f(1)}{\alpha-1} - \frac{f(1)}{1-\alpha} \left(\frac{\mu-\alpha}{\mu-1} \right)^\nu - c_2 \cdot m^{2-2\sigma_0} \left(\frac{\mu-\alpha}{\rho} \right)^{\nu-1} \cdot \frac{\rho}{\alpha-\sigma_0}, \\ &\quad \text{where } \sum_{j=0}^{[\nu]-1} b_j (\mu-\alpha)^j \geq c_3 \geq 1. \end{aligned}$$

From this inequality, we obtain

$$f(1) > c_3(1-\alpha) \left(\frac{\mu-1}{\mu-\alpha} \right)^\nu - c_2 \cdot m^{2-2\sigma_0} \cdot \frac{\rho^2(1-\alpha)}{(\mu-\alpha)(\alpha-\sigma_0)} \left(\frac{\mu-1}{\rho} \right)^\nu.$$

Choosing $\nu = c_4 \cdot \log m + c_5$, with

$$c_4 = \frac{2 - 2\sigma_0}{\log \frac{\rho}{\mu - \alpha}}$$

and

$$c_5 = \frac{\log \frac{c_2 \cdot \rho^2}{(\mu - \alpha)(\alpha - \sigma_0)} + \log \log \frac{\rho}{\mu - 1} - \log \log \frac{\mu - \alpha}{\mu - 1}}{\log \frac{\rho}{\mu - \alpha}},$$

gives $f(1) > c_6 \cdot m^{-c_7}$, with

$$c_6 = c_3(1 - \alpha) \left(\frac{\mu - 1}{\mu - \alpha} \right)^{c_5} - c_2 \cdot \frac{\rho^2(1 - \alpha)}{(\mu - \alpha)(\alpha - \sigma_0)} \cdot \left(\frac{\mu - 1}{\rho} \right)^{c_5}$$

and

$$c_7 = (2 - 2\sigma_0) \cdot \log \frac{\mu - \alpha}{\mu - 1} / \log \frac{\rho}{\mu - \alpha}.$$

Since $c_7 \rightarrow 0$ for $\alpha \rightarrow 1$, the proof of Lemma 2 is completed.

Numerical computations show that for $m_0 = 10^5$ good results can be obtained by choosing $\mu = 10$, $\rho = 9.9$ and $\alpha = 0.975$. With these values we obtain $c_2 = 10.8685$ and $\nu \approx 315$.

Using (7), and $a_n \geq 1$ for n a cube, we obtain the following estimations for c_3 :

$$\begin{aligned} \sum_{j=0}^{[\nu]-1} b_j(\mu - \alpha)^j &\geq \zeta_K(\mu) + \sum_{j=1}^{300} \frac{1}{j!} \sum_{n=2}^{\infty} \frac{a_n}{n^\mu} ((\mu - \alpha) \log n)^j \\ &> \zeta(3\mu) + \sum_{k=2}^{N_0} \frac{1}{k^{3\mu}} \sum_{j=1}^{300} \frac{1}{j!} ((\mu - \alpha) 3 \cdot \log k)^j \\ &> 1 + \sum_{k=2}^{N_0} \frac{1}{k^{3\mu}} \left(k^{3(\mu - \alpha)} - \frac{((\mu - \alpha) 3 \cdot \log k)^{301}}{301!} \cdot \frac{302}{302 - (\mu - \alpha) 3 \cdot \log k} \right), \end{aligned}$$

where $N_0 < e^{302/3(\mu - \alpha)}$. With the special values for μ , ρ and α , and $N_0 = 100$, we obtain

$$\sum_{j=0}^{[\nu]-1} b_j(\mu - \alpha)^j > \sum_{k=1}^{100} k^{-2.925} - 10^{-40} > 1.2175 = c_3.$$

These values yield $c_6 > 0.023$ and $c_7 < 0.054$, and thus (2) is proved.

Institut für Mathematik
Karl-Franzens-Universität
Halbärthgasse 1
A-8010 Graz
Österreich, Austria

1. N. C. ANKENY, S. CHOWLA & H. HASSE, "On the class-number of the maximal real subfield of a cyclotomic field," *J. Reine Angew. Math.*, v. 217, 1965, pp. 217–220.

2. A. CHÂTELET, "Arithmétique des corps abéliens du troisième degré," *Ann. Sci. École Norm. Sup.* (3), v. 63, 1946, pp. 109–160.

3. G. CORNELL & L. C. WASHINGTON, "Class numbers of cyclotomic fields," *J. Number Theory*, v. 21, 1985, pp. 260–274.

4. M.-N. GRAS, *Sur le Nombre de Classes du Sous-Corps Cubique de $\mathbf{Q}^{(p)}$ ($p \equiv 1 \pmod{3}$)*, Thèse, Grenoble, 1971.
5. M.-N. GRAS, "Méthodes et algorithmes pour le calcul numérique du nombre de classes et des unités des extensions cubiques cycliques de $\mathbf{Q}$," *J. Reine Angew. Math.*, v. 277, 1975, pp. 89–116.
6. M.-N. GRAS, "Note a propos d'une conjecture de H. J. Godwin sur les unités des corps cubiques," *Ann. Inst. Fourier (Grenoble)*, v. 30/4, 1980, pp. 1–6.
7. G. H. HARDY & E. M. WRIGHT, *An Introduction to the Theory of Numbers*, 5th ed., Oxford Univ. Press, New York, 1979.
8. H. HASSE, "Arithmetische Bestimmung von Grundeinheit und Klassenzahl in zyklischen kubischen und biquadratischen Zahlkörpern," *Abh. Deutsch. Akad. Wiss. Berlin Math.-Nat. Kl.*, 1948, Nr. 2, 95 pp.
9. S.-D. LANG, "Note on the class-number of the maximal real subfield of a cyclotomic field," *J. Reine Angew. Math.*, v. 290, 1977, pp. 70–72.
10. F. J. VAN DER LINDEN, "Class number computations of real abelian number fields," *Math. Comp.*, v. 39, 1982, pp. 693–707.
11. C. MOSER, *Sur le Nombre de Classes d'un Corps K Réel Cyclique de Conducteur Premier*, $\deg K \geq 4$, Séminaire Théorie des Nombres, Grenoble, 1980, 17 pp.
12. E. SEAH, L. C. WASHINGTON & H. C. WILLIAMS, "The calculation of a large cubic class number with an application to real cyclotomic fields," *Math. Comp.*, v. 41, 1983, pp. 303–305.
13. D. SHANKS, "The simplest cubic fields," *Math. Comp.*, v. 28, 1974, pp. 1137–1152.
14. H. M. STARK, "Some effective cases of the Brauer-Siegel Theorem," *Invent. Math.*, v. 23, 1974, pp. 135–152.
15. H. TAKEUCHI, "On the class-number of the maximal real subfield of a cyclotomic field," *Canad. J. Math.*, v. 33, 1981, pp. 55–58.
16. L. C. WASHINGTON, *Introduction to Cyclotomic Fields*, GTM 83, Springer, New York, 1982.
17. H. YOKOI, "On the diophantine equation $x^2 - py^2 = \pm 4q$ and the class number of real subfields of a cyclotomic field," *Nagoya Math. J.*, v. 91, 1983, pp. 151–161.