

On the Discretization in Time of Semilinear Parabolic Equations with Nonsmooth Initial Data

By Michel Crouzeix and Vidar Thomée

Abstract. Single-step discretization methods are considered for equations of the form $u_t + Au = f(t, u)$, where A is a linear positive definite operator in a Hilbert space H . It is shown that if the method is consistent with the differential equation then the convergence is essentially of first order in the stepsize, even if the initial data v are only in H , but also that, in contrast to the situation in the linear homogeneous case, higher-order convergence is not possible in general without further assumptions on v .

1. Introduction. We shall begin by recalling some results concerning the discretization in time of the linear homogeneous equation

$$(1.1) \quad \begin{aligned} u_t + Au &= 0 \quad \text{for } t > 0, \quad u_t = \partial u / \partial t, \\ u(0) &= v, \end{aligned}$$

where A is a selfadjoint positive definite operator in a Hilbert space H (cf., e.g., Baker, Bramble and Thomée [1]).

Let $r(z)$ be a rational function having no poles for $z \geq 0$, and define an approximate solution U_n at $t = t_n = nk$, where k is the time step, by

$$\begin{aligned} U_{n+1} &= r(kA)U_n \quad \text{for } n = 0, 1, 2, \dots, \\ U_0 &= v. \end{aligned}$$

Assume that the approximation is of order p with $p \geq 1$, or

$$(1.2) \quad r(z) = e^{-z} + O(z^{p+1}) \quad \text{as } z \rightarrow 0,$$

and also that the method is stable in the sense that

$$|r(z)| \leq 1 \quad \text{for } z \geq 0.$$

Then one may show the "smooth data" error estimate

$$\|U_n - u(t_n)\| \leq Ck^p \|A^p v\| \quad \text{for } v \in D(A^p).$$

This follows easily from spectral representations and the fact that under our assumptions

$$|r(z)^n - e^{-nz}| \leq Cz^p \quad \text{for } z \geq 0.$$

In applications, the requirement $v \in D(A^p)$ is quite restrictive. For example, if A is associated with an elliptic partial differential operator in a domain $\Omega \subset \mathbb{R}^d$, it demands not only smoothness of the initial data, but also that they satisfy certain

Received April 1, 1985.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 65M15.

©1987 American Mathematical Society
0025-5718/87 \$1.00 + \$.25 per page

compatibility conditions at the boundary $\partial\Omega$ for $t = 0$. However, under the stronger stability assumption

$$(1.3) \quad |r(z)| < 1 \quad \text{for } z > 0, \text{ and } |r(\infty)| < 1,$$

one can also show the “nonsmooth” data error estimate

$$(1.4) \quad \|U_n - u(t_n)\| \leq Ck^p t_n^{-p} \|v\| \quad \text{for } v \in H, t_n > 0.$$

This follows again by spectral arguments from

$$|r(z)^n - e^{-nz}| \leq Cn^{-p} \quad \text{for } z \geq 0,$$

and shows that even with v only in H , the $O(k^p)$ convergence is retained for $t_n > 0$. It follows also that for $0 \leq q \leq p$ the intermediate estimates

$$(1.5) \quad \|U_n - u(t_n)\| \leq Ck^p t_n^{-q} \|A^{p-q}v\| \quad \text{for } v \in D(A^{p-q})$$

hold.

The question we want to address below is to what extent these error estimates with reduced regularity assumptions carry over to semilinear equations. Thus assume that $f(t, u)$ is a smooth function on $\bar{J} \times H$, where $J = (0, T]$ with $T < \infty$, and consider the semilinear problem

$$(1.6) \quad \begin{aligned} u_t + Au &= f(t, u) \quad \text{for } t \in J, \\ u(0) &= v. \end{aligned}$$

For its approximate solution we will investigate in Section 2 single-step discretization schemes of the form

$$\begin{aligned} U_{n+1} &= r(kA)U_n + kF(k, t_n, U_n) \quad \text{for } t_n \in \bar{J}, \\ U_0 &= v, \end{aligned}$$

where $r(z)$ satisfies (1.2) with $p = 1$ and $F(k, t, v)$ is chosen to be consistent with (1.6) in a sense to be made precise below. As an example of such schemes, consider the standard first-order backward Euler scheme defined by

$$(1.7) \quad U_{n+1} = (I + kA)^{-1}U_n + k(I + kA)^{-1}f(t_{n+1}, U_{n+1}),$$

or the linearized version

$$(1.8) \quad U_{n+1} = (I + kA)^{-1}U_n + k(I + kA)^{-1}f(t_n, U_n),$$

where in the first case $F(k, t_n, U_n)$ is defined implicitly by (1.7). We shall be able to show (Theorem 1) that for such schemes

$$\|U_n - u(t_n)\| \leq Ckt_n^{-1} \log \frac{t_{n+1}}{k} \quad \text{for } t_n \in J,$$

where C depends on an upper bound for $\|v\|$, so that for first-order schemes the estimate (1.4) essentially remains valid in the semilinear case.

In Section 3 we briefly recall the definitions and basic properties of Runge-Kutta methods (cf., e.g., Crouzeix [2]) and show that our result above applies to such methods.

In Section 4 we shall then demonstrate that, more surprisingly, it is not in general possible to generalize the higher-order estimate (1.4) with $p > 1$ to semilinear equations. This will be done by exhibiting a simple system of the form (1.6) such

that, for any choice of a Runge-Kutta method satisfying (1.3), and any $t \in J$, we have

$$\limsup_{n=t/k \rightarrow \infty} \|U_n - u(t_n)\| \geq ck \quad \text{with } c = c(t) > 0.$$

We shall then proceed, in Section 5, within the framework of Runge-Kutta methods satisfying (1.3), to show (Theorem 2) that if the method is accurate of order p , with order $p - 1$ for the intermediate equations (cf. Section 5), then, if $u^{(j)}(t)$ are bounded for $j < p$ together with $tu^{(p)}(t)$, $f(t, u(t))^{(j)}$, $j < p$, and $tf(t, u(t))^{(p)}$, we have

$$\|U_n - u(t_n)\| \leq Ck^p \left(t_n^{-1} \log \frac{t_{n+1}}{k} + \left(\log \frac{t_{n+1}}{k} \right)^2 \right) \quad \text{for } t_n \in J,$$

which is thus an analogue of (1.5) with $q = 1$. Again, in practice, these assumptions will require certain compatibility conditions at $t = 0$.

These investigations are in a sense a continuation of work by Johnson, Larsson, Thomée and Wahlbin [3] concerning finite element type discretization with respect to the space variables of semilinear parabolic equations, and as we shall see below, our present results may be combined with those of [3] to yield error bounds for completely discrete schemes. The fact that the nonsmooth data error estimates for the linear homogeneous equation do not generalize to the semilinear problem for higher-order methods was shown in the case of semidiscretization in space in [3] by a counterexample, which was the starting point of this work.

2. The First-Order Error Estimate for Nonsmooth Data. Consider the initial value problem

$$(2.1) \quad \begin{aligned} u_t + Au &= f(t, u) \quad \text{for } t \in J = (0, T], \\ u(0) &= v, \end{aligned}$$

where A is a selfadjoint positive definite operator in a Hilbert space H and where $f(t, u)$ has values in H and is continuous and bounded together with its first-order derivatives with respect to t and u for $(t, u) \in \bar{J} \times H$. This problem has a unique solution on J for $v \in H$, which satisfies the integral equation

$$u(t) = E(t)v + \int_0^t E(t-s)f(s, u(s))ds,$$

where $E(t)$ is the semigroup generated by $-A$. This semigroup is analytic, since by spectral representation

$$\|tAE(t)v\| \leq \sup_{z \geq 0} (tze^{-tz})\|v\| = C\|v\|,$$

so that, in particular, for the solution of the homogeneous linear problem (1.1),

$$\|u_t(t)\| = \|Au(t)\| = \|AE(t)v\| \leq \frac{C}{t}\|v\| \quad \text{for } t \in J,$$

and for the solution of (1.1) we also have that u_t is bounded in H if $v \in D(A)$, i.e.,

$$\|u_t(t)\| = \|E(t)Av\| \leq \|Av\|.$$

We shall need the corresponding results for the solution of our semilinear problem (2.1).

LEMMA 1. *There are constants $C_i = C_i(\rho)$, $i = 0, 1$, such that the solution of (2.1) satisfies*

$$\|u_t(t)\| \leq C_0 t^{-1} \quad \text{for all } v \in H \text{ with } \|v\| \leq \rho$$

and

$$\|u_t(t)\| \leq C_1 \quad \text{for all } v \in D(A) \text{ with } \|Av\| \leq \rho.$$

The constants C_i , $i = 0, 1$, depend only on ρ and on bounds in H for f , f_t and f_u and are independent of the Hilbert space H and the positive definite operator A .

Proof. We introduce the symmetric, positive definite bilinear form

$$a(v, w) = (Av, w) \quad \text{for } v, w \in D(A),$$

which we may consider extended to the subspace V of H defined by the norm $\|v\|_V = a(v, v)^{1/2}$. We may then write our differential equation in weak form

$$(2.2) \quad (u_t, \varphi) + a(u, \varphi) = (f(t, u), \varphi) \quad \text{for } \varphi \in V.$$

We obtain by differentiation with respect to t , which is legitimate since the equation obtained is linear in u_t , with bounded coefficients,

$$(2.3) \quad (u_{tt}, \varphi) + a(u_t, \varphi) = (f_t(t, u) + f_u(t, u)(u_t), \varphi) \quad \text{for } t \in J,$$

and hence with $\varphi = u_t$,

$$\frac{1}{2} \frac{d}{dt} \|u_t\|^2 + \|u_t\|_V^2 \leq C(\|u_t\| + 1) \|u_t\|.$$

This yields

$$\frac{d}{dt} \|u_t\| \leq C \|u_t\| + C,$$

and, since J is bounded,

$$\|u_t(t)\| \leq C \|u_t(0)\| + C \leq C \|Av\| + C \leq C_1(\rho),$$

which is the second statement of the lemma.

In order to show the first, we choose $\varphi = t^2 u_t$ in (2.3) to obtain

$$\frac{1}{2} \frac{d}{dt} (t^2 \|u_t\|^2) + t^2 \|u_t\|_V^2 = t^2 (f_t + f_u(u_t), u_t) + t \|u_t\|^2 \leq Ct \|u_t\|^2 + C,$$

and thus

$$t^2 \|u_t(t)\|^2 \leq C \int_0^t s \|u_t\|^2 ds + C.$$

Taking $\varphi = tu_t$ in (2.2), we find

$$t \|u_t\|^2 + \frac{1}{2} \frac{d}{dt} (t \|u\|_V^2) = t (f(t, u), u_t) + \frac{1}{2} \|u\|_V^2,$$

and hence

$$\int_0^t s \|u_t(s)\|^2 ds \leq C \int_0^t \|u\|_V^2 ds + C.$$

Finally, $\varphi = u$ in (2.2) gives similarly

$$\int_0^t \|u\|_V^2 ds \leq \|v\|^2 + C \leq C_0(\rho),$$

which completes the proof.

As in the introduction, we consider now a difference scheme for (2.1) of the form

$$(2.4) \quad \begin{aligned} U_{n+1} &= E_k U_n + kF(k, t_n, U_n) \quad \text{for } t_n \in \bar{J}, \\ U_0 &= v, \end{aligned}$$

where $E_k = r(kA)$ for some rational function r satisfying (1.2) with $p = 1$ and (1.3), and where $F(k, t, \varphi)$ is a sufficiently smooth function chosen to make (2.4) consistent with (2.1). For a finite-dimensional problem this would simply mean that

$$(2.5) \quad F(0, t, \varphi) = f(t, \varphi);$$

in the general Hilbert space context we shall need to make this more precise. We shall thus assume that F is uniformly Lipschitz continuous with respect to φ , so that, for some $k_0 > 0$,

$$(2.6) \quad \|F(k, t, \varphi) - F(k, t, \psi)\| \leq C\|\varphi - \psi\| \quad \text{on } [0, k_0] \times \bar{J} \times H,$$

and, in addition, that

$$(2.7) \quad \|A^{-1}(F(k, t, \varphi) - f(t, \varphi))\| \leq Ck(\|A\varphi\| + 1) \quad \text{on } [0, k_0] \times \bar{J} \times D(A).$$

Note that the latter condition follows from (2.5) in the finite-dimensional case if F is Lipschitz continuous with respect to k . Observe also that (2.6) implies

$$(2.8) \quad \|F(k, t, \varphi)\| \leq C(1 + \|\varphi\|) \quad \text{on } [0, k_0] \times \bar{J} \times H.$$

We are now ready to state and prove the main result of this section.

THEOREM 1. *Assume for the difference scheme (2.4) that $E_k = r(kA)$, where r satisfies (1.2) with $p = 1$ and (1.3), and that (2.6) and (2.7) hold. Then there is a constant $C = C(\rho)$ such that for $t_n \in J$*

$$\|U_n - u(t_n)\| \leq C \left(\frac{k}{t_n} \log \frac{t_{n+1}}{k} + k \left(\log \frac{t_{n+1}}{k} \right)^2 \right) \quad \text{for } v \in H \text{ with } \|v\| \leq \rho.$$

The constant C depends, in addition to ρ , only on bounds for f, f_t, f_u and on the constants of (2.6) and (2.7) and is independent of the particular choice of the Hilbert space H and the positive definite operator A .

Proof. We find at once

$$U_n = E_k^n v + k \sum_{j=0}^{n-1} E_k^{n-1-j} F(k, t_j, U_j).$$

For the exact solution we may write similarly, with $I_j = (t_j, t_{j+1})$ and $u_n = u(t_n)$,

$$u_n = E(t_n)v + \sum_{j=0}^{n-1} \int_{I_j} E(t_n - s) f(s, u(s)) ds,$$

so that for the error, $e_n = U_n - u_n$,

$$(2.9) \quad e_n = (E_k^n - E(t_n))v + \sum_{j=0}^{n-1} \int_{I_j} D_{n,j}(s) ds,$$

where

$$D_{n,j}(s) = E_k^{n-1-j} F(k, t_j, U_j) - E(t_n - s) f(s, u(s)).$$

We write this latter expression in the form

$$\begin{aligned} D_{n,j}(s) &= E_k^{n-1-j} (F(k, t_j, U_j) - F(k, t_j, u_j)) \\ &\quad + (E_k^{n-1-j} - E(t_{n-1-j})) F(k, t_j, u_j) \\ &\quad + E(t_{n-1-j}) (F(k, t_j, u_j) - f(t_j, u_j)) \\ &\quad + E(t_{n-1-j}) (f(t_j, u_j) - f(s, u(s))) \\ &\quad + (E(t_{n-1-j}) - E(t_n - s)) f(s, u(s)) \\ &= \sum_{l=1}^5 d_{j,l}(s). \end{aligned}$$

We now proceed to estimate these five terms for $s \in I_j$. We first have, by the stability of E_k and (2.6),

$$\|d_{j,1}\| \leq C \|U_j - u_j\| = C \|e_j\|.$$

For the second term we note that (1.4) may be written

$$(2.10) \quad \|(E_k^n - E(t_n))v\| \leq C \frac{k}{t_n} \|v\| \quad \text{for } t_n > 0,$$

and we conclude, by (2.8),

$$\|d_{j,2}\| \leq C \frac{k}{t_{n-1-j}} \quad \text{for } j \neq n-1.$$

Since $d_{n-1,2} = 0$ we may write

$$\|d_{j,2}\| \leq \frac{Ck}{t_{n-j}} \quad \text{for } 0 \leq j \leq n-1.$$

For the third term we use the consistency condition (2.7), the analyticity of $E(t)$ and Lemma 1 to obtain for $0 < j < n-1$,

$$\begin{aligned} \|d_{j,3}\| &= \|AE(t_{n-1-j})A^{-1}(F(k, t_j, u_j) - f(t_j, u_j))\| \\ &\leq \frac{C}{t_{n-1-j}} k (\|Au_j\| + 1) \leq C \frac{k}{t_{n-1-j}} (\|u_t(t_j)\| + 1) \leq C \frac{k}{t_j t_{n-1-j}}. \end{aligned}$$

For $j = 0$ and $n-1$ we find easily by the boundedness of F and f ,

$$\|d_{j,3}\| \leq C = C \frac{k}{t_1} \leq C \frac{k}{t_1 t_n},$$

so that we may write

$$\|d_{j,3}\| \leq C \frac{k}{t_{j+1} t_{n-j}} = \frac{Ck}{t_n} \left(\frac{1}{t_{j+1}} + \frac{1}{t_{n-j}} \right), \quad 0 \leq j \leq n-1.$$

For the fourth term we have

$$\begin{aligned} \|d_{j,4}\| &\leq \|f(t_j, u_j) - f(s, u(s))\| \leq C(k + \|u_j - u(s)\|) \\ &\leq Ck \left(\sup_{I_j} \|u_t(s)\| + 1 \right) \leq C \frac{k}{t_j} \quad \text{for } 0 < j \leq n-1, \end{aligned}$$

and, since $d_{0,4}$ is bounded,

$$\|d_{j,4}\| \leq C \frac{k}{t_{j+1}} \quad \text{for } 0 \leq j \leq n-1.$$

Finally,

$$\begin{aligned} \|d_{j,5}\| &= \|AE(t_{n-1-j})A^{-1}(E(t_{j+1}-s)-I)f(s, u(s))\| \\ &\leq C \frac{k}{t_{n-1-j}} \quad \text{for } j < n-1, \end{aligned}$$

where we have used the analyticity of $E(t)$ and the fact that

$$\|A^{-1}(E(t_{j+1}-s)-I)v\| \leq \sup_{z>0} \left| \frac{e^{-(t_{j+1}-s)z}}{z} \right| \cdot \|v\| \leq Ck\|v\|.$$

Again,

$$\|d_{n-1,5}\| \leq C = C \frac{k}{t_1},$$

so that

$$\|d_{j,5}\| \leq C \frac{k}{t_{n-j}} \quad \text{for } 0 \leq j \leq n-1.$$

Altogether we have thus from (2.9), using (2.10) to estimate the first term, for $\|v\| \leq \rho$,

$$\|e_n\| \leq C \frac{k}{t_n} + Ck \sum_{j=0}^{n-1} \|e_j\| + C \frac{k}{t_n} \sum_{j=1}^n \frac{k}{t_j} \leq Ck \sum_{j=0}^{n-1} \|e_j\| + C \frac{k}{t_n} \log(n+1);$$

setting $\beta_n = k \sum_{j=0}^n \|e_j\|$, we thus have

$$\beta_n \leq (1 + Ck)\beta_{n-1} + \frac{k}{n} \log(n+1),$$

and hence

$$\beta_n \leq k \sum_{j=1}^n (1 + Ck)^{n-1-j} \frac{1}{j} \log(j+1) \leq Ck \log(n+1) \sum_{j=1}^n \frac{1}{j} \leq Ck (\log(n+1))^2$$

and finally

$$\begin{aligned} \|e_n\| &\leq Ck (\log(n+1))^2 + C \frac{k}{t_n} \log(n+1) \\ &= C \left(\frac{k}{t_n} \log \frac{t_{n+1}}{k} + k \left(\log \frac{t_{n+1}}{k} \right)^2 \right), \end{aligned}$$

which completes the proof of the theorem.

As our first example we consider the linearized modification of the backward Euler scheme defined in (1.8). Here,

$$F(k, t, \varphi) = (I + kA)^{-1} f(t, \varphi),$$

and it is clear from our assumption on f that (2.6) is satisfied. Further, we note that

$$\begin{aligned} A^{-1}(F(k, t, \varphi) - f(t, \varphi)) &= k(kA)^{-1}((I + kA)^{-1} - I)f(t, \varphi) \\ &= k(I + kA)^{-1}f(t, \varphi) \end{aligned}$$

and hence

$$\|A^{-1}(F(k, t, \varphi) - f(t, \varphi))\| \leq Ck,$$

so that (2.7) holds even without the term $\|A\varphi\|$. For schemes with this property the result of Theorem 1 is in fact valid without the term $Ck(\log(t_{n+1}/k))^2$.

Turning now to the standard backward Euler scheme, we have here

$$F(k, t, \varphi) = (I + kA)^{-1}f(t + k, \Psi),$$

where $\Psi = \Psi(\varphi)$ is obtained from the nonlinear equation

$$\Psi = (I + kA)^{-1}(\varphi + kf(t + k, \Psi)).$$

It is clear by the contraction mapping theorem that this equation has a unique solution Ψ for small k and that Ψ depends Lipschitz continuously on φ . Obviously, again (2.6) is satisfied for this scheme. As for (2.7), we have

$$\begin{aligned} \Psi - \varphi &= \Psi - (I + kA)^{-1}\varphi - (I + kA)^{-1}kA\varphi \\ &= k(I + kA)^{-1}f(t + k, \Psi) - (I + kA)^{-1}kA\varphi, \end{aligned}$$

and thus

$$\|\Psi - \varphi\| \leq Ck(1 + \|A\varphi\|).$$

Further,

$$\begin{aligned} F(k, t, \varphi) - f(t, \varphi) &= (I + kA)^{-1}(f(t + k, \Psi) - f(t + k, \varphi)) \\ &\quad + (I + kA)^{-1}(f(t + k, \varphi) - f(t, \varphi)) \\ &\quad - (I + kA)^{-1}kAf(t, \varphi) \end{aligned}$$

and hence, since A^{-1} is bounded,

$$\|A^{-1}(F(k, t, \varphi) - f(t, \varphi))\| \leq C\|\Psi - \varphi\| + Ck \leq Ck(1 + \|A\varphi\|),$$

which is the desired estimate.

The above result may be applied to parabolic problems which have already been discretized in the space variables. For instance, for concreteness, consider the case that $H = L_2(\Omega)$, where Ω is a domain with smooth boundary in R^d , where $A = -\Delta$, with $D(A) = H_0^1(\Omega) \cap H^2(\Omega)$, and where $f(t, u)$ is generated by a smooth function $\tilde{f}(x, t, u)$ on $\bar{\Omega} \times \bar{J} \times R$ which is bounded together with its first-order derivatives with respect to t and u . Now let $S_h \subset H_0^1(\Omega)$ consist of continuous, piecewise linear functions on a partitioning of Ω into simplices and let $u_h: \bar{J} \rightarrow S_h$ be defined by

$$\begin{aligned} (2.11) \quad &(u_{h,t}, \chi) + (\nabla u_h, \nabla \chi) = (\tilde{f}(\cdot, t, u_h), \chi) \quad \text{for } \chi \in S_h, t \in J, \\ &u_h(0) = v_h, \end{aligned}$$

where (\cdot, \cdot) denotes the standard inner products on $L_2(\Omega)$ and $L_2(\Omega)^d$. Defining the discrete solution operator $T_h: L_2(\Omega) \rightarrow S_h$ of the associated elliptic problem by

$$(\nabla T_h w, \nabla \chi) = (w, \chi) \quad \text{for } \chi \in S_h,$$

and setting $\Delta_h = -T_h^{-1}$, (2.11) may be written in the form (2.1) as

$$(2.12) \quad u_{h,t} - \Delta_h u_h = P_0 \tilde{f}(\cdot, t, u_h) \quad \text{for } t \in J,$$

where P_0 denotes the orthogonal L_2 -projection onto S_h . From [3] it is known that, if $v_h = P_0 v$, the error in this space discretization satisfies

$$\|u_h(t) - u(t)\| \leq C(\rho) \frac{h^2}{t} \log \frac{t}{h^2} \quad \text{for all } v \in L_2(\Omega) \text{ with } \|v\| \leq \rho,$$

where h is the maximal diameter of the simplices of the partitioning.

Our above analysis applies to this situation and yields estimates which are uniform in h . In fact, if \tilde{f} is as above, then $P_0 \tilde{f}(x, t, u)$ generates a function $\bar{J} \times S_h \ni (t, u_h) \rightarrow P_0 \tilde{f}(\cdot, t, u_h) \in S_h$ which satisfies our above assumptions with respect to the Hilbert space defined by S_h equipped with the norm of $L_2(\Omega)$. For instance, the derivative of this function with respect to u applied to $w \in S_h$ is $P_0(\tilde{f}_u(\cdot, t, u) \cdot w) \in S_h$, which is clearly bounded in the L_2 -norm, uniformly in h . Also, if $v_h = P_0 v$ and $\|v\| \leq \rho$, we have $\|v_h\| \leq \|v\| \leq \rho$ and hence Theorem 1 implies that for a completely discrete solution obtained by discretization in time of (2.12) by a scheme of the above type, and with $v_h = P_0 v$, we have

$$\|U_n - u(t_n)\| \leq C(\rho) \left\{ \frac{h^2}{t_n} \log \frac{t_n}{h^2} + k \left[t_n^{-1} \log \frac{t_{n+1}}{k} + \left(\log \frac{t_{n+1}}{k} \right)^2 \right] \right\}$$

for $v \in L_2(\Omega)$ with $\|v\| \leq \rho$.

We remark that in interesting applications of the type just described it is generally the case that \tilde{f} and \tilde{f}_u are unbounded for $u \in R$ so that, as u is not necessarily bounded when $u \in L_2(\Omega)$, the above analysis does not apply. However, it is then often the case that by some independent argument, for instance by a maximum principle, the exact solution is known to be uniformly bounded in modulus by some constant M , say, in some interval $\bar{J} = [0, T]$, if the initial data are bounded, and that thus the values of $\tilde{f}(x, t, u)$ for $|u| > M$ do not influence the exact solution of (2.1). One can then modify \tilde{f} for these values of u in such a way that our assumptions become valid, thus changing the equation (2.1) without changing its solution for the initial data under consideration. With $F(k, t, u)$ based on the modified function, our assumptions (2.6) and (2.7) may remain valid. Note that this procedure might lead to a different discrete solution than the one based on the original \tilde{f} . Similarly, such a modification would change the semidiscrete equation (2.11) and thus also the totally discrete solution based on (2.12).

3. Runge-Kutta Methods. We recall (cf., e.g., [2] for details) that a Runge-Kutta method for the initial value problem

$$\begin{aligned} y' &= g(t, y) \quad \text{for } t \geq 0, \\ y(0) &= y_0, \end{aligned}$$

defines an approximate solution Y_n at $t_n = nk$ successively by setting $Y_0 = y_0$ and then determining Y_{n+1} from Y_n for $n \geq 0$ as follows: Let $t_{nj} = t_n + k\tau_j$ be given quadrature points with $\tau_j \geq 0$ for $j = 1, \dots, m$; define intermediate values Y_{nj} , $j = 1, \dots, m$, approximating $y(t_{nj})$ by the nonlinear system

$$Y_{ni} = Y_n + k \sum_{j=1}^m a_{ij} g(t_{nj}, Y_{nj}), \quad i = 1, \dots, m,$$

and set finally

$$Y_{n+1} = Y_n + k \sum_{j=1}^m b_j g(t_{nj}, Y_{nj}).$$

The coefficient matrix $\mathcal{A} = (a_{ij})$ and the vector $b = (b_1, \dots, b_m)^T$ are associated with the quadrature formulae

$$(3.1) \quad \int_0^{\tau_j} \Psi(t) dt \approx \sum_{j=1}^m a_{ij} \Psi(\tau_j)$$

and

$$(3.2) \quad \int_0^1 \Psi(t) dt \approx \sum_{j=1}^m b_j \Psi(\tau_j),$$

respectively, and we shall always assume that the latter is exact for constants, so that

$$(3.3) \quad \sum_{j=1}^m b_j = 1.$$

The method is implicit unless \mathcal{A} is strictly lower triangular.

Applied to the parabolic problem (2.1), the method takes the form

$$(3.4) \quad U_{ni} = U_n + k \sum_{j=1}^m a_{ij} (-AU_{nj} + f(t_{nj}, U_{nj})), \quad j = 1, \dots, m,$$

$$(3.5) \quad U_{n+1} = U_n + k \sum_{j=1}^m b_j (-AU_{nj} + f(t_{nj}, U_{nj})).$$

We shall assume that \mathcal{A} has no eigenvalues α_j with $\alpha_j \leq 0$ so that, in particular, the method is implicit and $I + z\mathcal{A}$ is nonsingular for $z \geq 0$. We set

$$\begin{aligned} \sigma(z) &= (\sigma_{ij}(z)) = (I + z\mathcal{A})^{-1}, \\ s(z) &= (s_1(z), \dots, s_m(z))^T = \sigma(z)e \quad \text{where } e = (1, \dots, 1)^T, \\ S(z) &= (s_{ij}(z)) = \sigma(z)\mathcal{A}, \\ r(z) &= 1 - zb^T s(z) = 1 - zb^T \sigma(z)e, \\ q(z)^T &= (q_1(z), \dots, q_m(z)) = b^T \sigma(z), \end{aligned}$$

and note that all these functions are bounded for $z \geq 0$, and that σ , s , S and q all vanish at $z = \infty$. With this notation, the equations (3.4) and (3.5) may be written

$$(3.6) \quad \begin{aligned} U_{ni} &= s_i(kA)U_n + k \sum_{j=1}^m s_{ij}(kA)f(t_{nj}, U_{nj}), \quad i = 1, \dots, m, \\ U_{n+1} &= r(kA)U_n + k \sum_{j=1}^m q_j(kA)f(t_{nj}, U_{nj}), \end{aligned}$$

where the rational functions of kA are defined by spectral representation and thus, by the above, are all bounded linear operators on H .

Henceforth we shall restrict ourselves, as earlier in Section 2, to schemes such that $r(z)$ satisfies the strong stability property (1.3). Recall also that the method is accurate of order p , and, in particular, that (1.2) holds, if (3.1) and (3.2) are exact for

polynomials of degree $p - 2$ and $p - 1$, respectively (cf. [2, p. 12]). For $p = 1$ this reduces to the condition (3.3).

Since $f(t, \varphi)$ is Lipschitz continuous with respect to φ , it is easy to see that the nonlinear system (3.6) has a unique solution (U_{n1}, \dots, U_{nm}) for U_n given, and that it depends Lipschitz continuously on U_n . We may thus write our method in the form (2.4), where

$$F(k, t, \varphi) = \sum_{j=1}^m q_j(kA) f(t + \tau_j k, \varphi_j),$$

with $\varphi_j = \varphi_j(k, t, \varphi)$ defined by

$$(3.7) \quad \varphi_i = s_i(kA)\varphi + k \sum_{j=1}^m q_j(kA) f(t + \tau_j k, \varphi_j), \quad i = 1, \dots, m.$$

We show that this method satisfies the conditions of Theorem 1. In fact, (2.6) is obvious and it remains only to consider (2.7). We write

$$\begin{aligned} F(k, t, \varphi) - f(t, \varphi) &= \sum_{j=1}^m q_j(kA) (f(t + \tau_j k, \varphi_j) - f(t + \tau_j k, \varphi)) \\ &\quad + \sum_{j=1}^m q_j(kA) (f(t + \tau_j k, \varphi) - f(t, \varphi)) \\ &\quad + \left(\sum_{j=1}^m q_j(kA) - I \right) f(t, \varphi) \\ &= \delta_1 + \delta_2 + \delta_3. \end{aligned}$$

Here

$$\|A^{-1}\delta_1\| \leq C\|\delta_1\| \leq C \sum_{j=1}^m \|\varphi_j - \varphi\|$$

and, noting that $s_i(0) = 1$, we have easily from (3.7)

$$\|\varphi_i - \varphi\| \leq \|(s_i(kA) - I)\varphi\| + Ck \leq Ck(\|A\varphi\| + 1).$$

Further, it is obvious that

$$\|A^{-1}\delta_2\| \leq C\|\delta_2\| \leq Ck.$$

Finally, since by (3.3)

$$\sum_{j=1}^m q_j(0) = \sum_{j=1}^m b_j = 1,$$

we obtain

$$\|A^{-1}\delta_3\| = k \left\| (kA)^{-1} \left(\sum_{j=1}^m q_j(kA) - I \right) f \right\| \leq Ck.$$

Together, these estimates complete the proof that the Runge-Kutta methods under consideration satisfy the assumptions of Theorem 1.

4. The counterexample. We shall now show that, at least for methods of Runge-Kutta type, the estimate of Theorem 1 is essentially best possible. For this purpose we introduce the unidimensional parabolic system

$$(4.1) \quad \begin{aligned} u_t - u_{xx} &= v^2 \\ v_t - v_{xx} &= 0 \end{aligned} \quad \text{in } [0, \pi] \times J,$$

with the boundary and initial conditions

$$(4.2) \quad \begin{aligned} u(0, t) &= u(\pi, t) = v(0, t) = v(\pi, t) = 0, \\ u(x, 0) &= 0, \quad v(x, 0) = w(x), \end{aligned}$$

which we consider in the Hilbert space $H = L_2(0, \pi)^2$, with the obvious corresponding definition of A . We note that by the maximum principle $\|v(t)\|_{L_\infty(\Omega)}$ is nonincreasing and hence $u(t)$ is uniformly bounded on J for all w with a common uniform bound.

We consider now the discrete solution (U_n, V_n) of (4.1), (4.2), defined by a Runge-Kutta method as described above. We shall show that for no $t \in J$ is it possible to find $C = C(t)$ such that the error, measured in the norm in $L_2(0, \pi)^2$, is bounded by $Ck\varepsilon_k$ with $\varepsilon_k \rightarrow 0$ as $k \rightarrow 0$, for all w with $\|w\|_{L_\infty} \leq 1$, say. More precisely, we shall show that there is a positive constant c such that for initial data of the form $w = \sin Nx$ with $N = 1, 2, \dots$ it is possible to find associated sequences $\{n_N\}$ and $\{k_N\}$ with $n_N k_N = t$ such that the error in the u -component satisfies

$$(4.3) \quad \|U_{n_N} - u(t)\|_{L_2(0, \pi)} \geq ck_N \quad \text{for large } N.$$

This will be demonstrated by showing that the corresponding estimate holds for the first Fourier coefficient of the error.

For the v -component of the exact solution of (4.1), (4.2) we have at once

$$v(t) = e^{-N^2 t} \sin Nx,$$

which gives for the determination of u the inhomogeneous linear equation

$$(4.4) \quad \begin{aligned} u_t - u_{xx} &= e^{-2N^2 t} \sin^2 Nx \quad \text{for } t \in J, \\ u(0) &= 0. \end{aligned}$$

Similarly, we have for the corresponding discrete problem, using the Runge-Kutta method, with our above notation,

$$V_n = r(kN^2)^n \sin Nx \quad \text{for } t_n \in \bar{J},$$

and thus also

$$V_{ni} = s_i(kN^2)r(kN^2)^n \sin Nx, \quad i = 1, \dots, m,$$

and hence for the determination of U_n the recursion formula

$$(4.5) \quad \begin{aligned} U_{n+1} &= r(kA)U_n + k \sum_{j=1}^m q_j(kA)s_j(kN^2)^2 r(kN^2)^{2n} \sin^2 Nx, \\ U_0 &= 0. \end{aligned}$$

We now introduce the first Fourier coefficients of $u(t)$ and U_n ,

$$\alpha(t) = \frac{2}{\pi} \int_0^\pi u(x, t) \sin x \, dx \quad \text{and} \quad \bar{\alpha}_n = \frac{2}{\pi} \int_0^\pi U_n(x) \sin x \, dx.$$

We obtain from (4.4)

$$\alpha' + \alpha = \frac{2}{\pi} e^{-2N^2 t} \int_0^\pi \sin x \sin^2 Nx \, dx = \frac{8N^2}{\pi(4N^2 - 1)} e^{-2N^2 t},$$

$$\alpha(0) = 0,$$

which yields

$$\alpha(t) = \frac{8N^2}{\pi(4N^2 - 1)} \frac{1}{2N^2 - 1} (e^{-t} - e^{-2N^2 t}).$$

Correspondingly, from (4.5),

$$\bar{\alpha}_{n+1} = r(k) \bar{\alpha}_n + k \sum_{j=1}^m q_j(k) s_j(kN^2)^2 r(kN^2)^{2n} \int_0^\pi \sin x \sin^2 Nx \, dx,$$

$$\bar{\alpha}_0 = 0,$$

and hence

$$\begin{aligned} \bar{\alpha}_n &= \frac{8N^2}{\pi(4N^2 - 1)} \sum_{i=0}^{n-1} r(k)^{n-1-i} k \sum_{j=1}^m q_j(k) s_j(kN^2)^2 r(kN^2)^{2i} \\ &= \frac{8N^2}{\pi(4N^2 - 1)} k \sum_{j=1}^m q_j(k) s_j(kN^2)^2 \frac{r(k)^n - r(kN^2)^{2n}}{r(k) - r(kN^2)^2}. \end{aligned}$$

We write

$$\begin{aligned} \frac{\bar{\alpha}_n - \alpha(t)}{k} &= \frac{8N^2}{\pi(4N^2 - 1)} \left\{ \sum_{j=1}^m q_j(k) s_j(kN^2)^2 \frac{r(k)^n - r(kN^2)^{2n}}{r(k) - r(kN^2)^2} \right. \\ &\quad \left. - \frac{1}{2N^2 - 1} \frac{1}{k} (e^{-t} - e^{-2N^2 t}) \right\}. \end{aligned}$$

We now fix t positive and set, with M a fixed positive integer, $n_N = MN^2$ and $k_N = t/n_N$ so that $k_N N^2 = t/M = t_0$. We note that since $r(z) = e^{-z} + O(z^2)$ for small z , we have $r(0) = 1$ and $r(k_N)^{n_N} = r(t/n_N)^{n_N} \rightarrow e^{-t}$ as $N \rightarrow \infty$. Further, since $|r(t_0)| < 1$, we have $r(k_N N^2)^{2n_N} = r(t_0)^{2n_N} \rightarrow 0$ as $N \rightarrow \infty$, and since $q_j(0) = b_j$, we conclude

$$(4.6) \quad \lim_{N \rightarrow \infty} \frac{\bar{\alpha}_{n_N} - \alpha(t)}{k_N} = \frac{2}{\pi} e^{-t} \left\{ \sum_{j=1}^m b_j s_j(t_0)^2 \frac{1}{1 - r(t_0)^2} - \frac{1}{2t_0} \right\}.$$

We shall show that for M suitably chosen the last factor is nonzero, which then shows that, for large N ,

$$\|U_{n_N} - u(t)\|_{L_2(0, \pi)} \geq \left(\frac{\pi}{2}\right)^{1/2} |\bar{\alpha}_{n_N} - \alpha(t)| \geq ck_N \quad \text{with } c = c(t) > 0,$$

and thus completes the proof of (4.3). Assume therefore that for any choice of M the expression inside the parentheses in (4.6) vanishes. In such a case, this rational function vanishes identically, and we have

$$2z \sum_{j=1}^m b_j s_j(z)^2 \equiv 1 - r(z)^2.$$

But for large z , $s_j(z) = O(z^{-1})$ so that, by letting z tend to infinity, we obtain $|r(\infty)| = 1$, contrary to our hypothesis. This completes the proof.

We note that the right-hand side of (4.1) is not bounded for $v \in R$ so that, formally, the assumptions of Section 2 are not satisfied for (4.1), (4.2). However, both $|v(t)|$ and $|V_n|$ are bounded by 1 and $|V_{ni}|$ by $K = \max_i \sup_{z \geq 0} |s_i(z)|$. Hence only the values of the right-hand side for $|v| \leq K$ enter the calculations, and we may replace v^2 by a smooth function $f(v)$ which satisfies our previous conditions and agrees with v^2 for $|v| \leq K$, without changing either the exact or the approximate solutions.

5. A Higher-Order Result for a Class of Runge-Kutta Methods. Although for initial data in H it was only possible, above, to show an essentially first-order error estimate, it may still be possible to do better for initial data which are more regular, but not regular enough for optimal order estimates to hold uniformly down to $t = 0$. In this section we shall show a $O(k^p)$ error estimate for a Runge-Kutta type method based on quadrature formulas of orders $p - 1$ for the intermediate points and p for the whole interval, and for the case that $u^{(p)}$ and $f(t, u)^{(p)}$ are of order $O(t^{-1})$ for small t .

THEOREM 2. *Let U_n be the discrete solution of (2.1) by a Runge-Kutta scheme satisfying (1.3) and for which the quadrature values (3.1) and (3.2) are exact for all polynomials of degree $p - 2$ and $p - 1$, respectively. Then there is a constant $C = C(\rho)$ such that*

$$\|U_n - u(t_n)\| \leq Ck^p \left(t_n^{-1} \log \frac{t_{n+1}}{k} + \left(\log \frac{t_{n+1}}{k} \right)^2 \right) \quad \text{for } t_n \in J$$

if, with $\varphi(t) = f(t, u(t))$,

$$(5.1) \quad \max \left(\max_{j \leq p-1} (\|u^{(j)}\|, \|\varphi^{(j)}\|), t\|u^{(p)}\|, t\|\varphi^{(p)}\| \right) \leq \rho.$$

The constant C is independent of the particular choice of the Hilbert space H and the positive definite operator A .

Proof. Let us introduce the error functional for the quadrature formulae (3.1) and (3.2), transformed to the interval I_n , i.e.,

$$\begin{aligned} Q_{n,j}(\Psi) &= \int_{t_n}^{t_{n,j}} \Psi ds - k \sum_{l=1}^m a_{jl} \Psi(t_{nl}), \quad j = 1, \dots, m, \\ Q_n(\Psi) &= \int_{t_n}^{t_{n+1}} \Psi ds - k \sum_{l=1}^m b_l \Psi(t_{nl}). \end{aligned}$$

Recall that our assumptions that (3.1) and (3.2) are exact for polynomials of degree $p - 1$ and $p - 2$, respectively, imply

$$(5.2) \quad \|Q_{n,j}(\Psi)\| \leq Ck^{l+1} \sup_{I_n} \|\Psi^{(l)}\| \quad \text{for } l \leq p - 1,$$

$$(5.3) \quad \|Q_n(\Psi)\| \leq Ck^{l+1} \sup_{I_n} \|\Psi^{(l)}\| \quad \text{for } l \leq p.$$

We note now that $Q_{n_j}(u_t)$ and $Q_n(u_t)$ are the truncation errors in (3.4) and (3.5), or that

$$u(t_{ni}) = u(t_n) + k \sum_{j=1}^m a_{ij}(-Au(t_{nj}) + f(t_{nj}, u(t_{nj}))) + Q_{ni}(u_t),$$

$$i = 1, \dots, m,$$

$$u(t_{n+1}) = u(t_n) + k \sum_{j=1}^m b_j(-Au(t_{nj}) + f(t_{nj}, u(t_{nj}))) + Q_n(u_t).$$

It follows by obvious calculations that

$$\begin{aligned} u(t_{ni}) &= s_i(kA)u(t_n) + k \sum_{j=1}^m s_{ij}(kA)f(t_{nj}, u(t_{nj})) \\ &\quad + \sum_{j=1}^m \sigma_{ij}(kA)Q_{n_j}(u_t), \quad i = 1, \dots, m, \\ u(t_{n+1}) &= r(kA)u(t_n) + k \sum_{j=1}^m q_j(kA)f(t_{nj}, u(t_{nj})) \\ &\quad - \sum_{j=1}^m kAq_j(kA)Q_{n_j}(u_t) + Q_n(u_t). \end{aligned}$$

Hence for the errors $e_n = U_n - u(t_n)$, $e_{n_j} = U_{n_j} - u(t_{n_j})$, we have

$$\begin{aligned} e_{ni} &= s_i(kA)e_n + k \sum_{j=1}^m s_{ij}(kA)(f(t_{nj}, U_{n_j}) - f(t_{nj}, u(t_{n_j}))) \\ &\quad - \sum_{j=1}^m \sigma_{ij}(kA)Q_{n_j}(u_t), \\ (5.4) \quad e_{n+1} &= r(kA)e_n + k \sum_{j=1}^m q_j(kA)(f(t_{n,j}, U_{n,j}) - f(t_{n,j}, u(t_{n_j}))) \\ &\quad + \sum_{j=1}^m kAq_j(kA)Q_{n_j}(u_t) - Q_n(u_t) \\ &= r(kA)e_n + \eta_{n1} + \eta_{n2} + \eta_{n3}, \end{aligned}$$

and, where as in Section 2 we have set $E_k = r(kA)$,

$$(5.5) \quad e_n = \sum_{j=0}^{n-1} E_k^{n-1-j}(\eta_{j1} + \eta_{j2} + \eta_{j3}).$$

We obtain at once from (5.4), for k small,

$$\sum_{i=1}^m \|e_{ni}\| \leq C\|e_n\| + C \sum_{j=1}^m \|Q_{n_j}(u_t)\|,$$

and hence, using also (5.2), for $j = 1, \dots, n-1$,

$$\begin{aligned} \|\eta_{j1}\| &\leq Ck \sum_{l=1}^m \|e_{jl}\| \leq Ck \left(\|e_j\| + \sum_{l=1}^m \|Q_{jl}(u_t)\| \right) \\ &\leq Ck\|e_j\| + Ck^{p+1} \sup_{I_j} \|u^{(p)}\| \leq Ck\|e_j\| + Ck^{p+1}t_j^{-1} \\ &\leq Ck\|e_j\| + Ck^{p+1}t_{j+1}^{-1}, \end{aligned}$$

and, for $j = 0$,

$$\|\eta_{01}\| \leq Ck\|e_0\| + Ck^p \sup_{I_j} \|u^{(p-1)}\| \leq Ck\|e_0\| + Ck^{p+1}t_1^{-1},$$

so that for the sum in (5.5) with η_{j1} ,

$$\begin{aligned} \left\| \sum_{j=0}^{n-1} E_k^{n-1-j} \eta_{j1} \right\| &\leq Ck \sum_{j=0}^{n-1} \|e_j\| + Ck^{p+1} \sum_{j=0}^{n-1} \frac{1}{t_{j+1}} \\ &\leq Ck \sum_{j=0}^{n-1} \|e_j\| + Ck^p \log \frac{t_{n+1}}{k}. \end{aligned}$$

In order to estimate the term involving η_{j2} , we note that since $zq(z)$ is bounded, we have, using the property (1.3),

$$|r(z)^n zq(z)| \leq \frac{C}{n+1} = C \frac{k}{t_{n+1}},$$

and hence, for $j = 1, \dots, n-1$,

$$\begin{aligned} \|E_k^{n-1-j} \eta_{j2}\| &\leq C \frac{k}{t_{n-j}} \sum_{l=1}^m \|Q_{jl}(u_l)\| \leq C \frac{k^{p+1}}{t_{n-j}} \sup_{I_j} \|u^{(p)}\| \\ &\leq Ck^{p+1} \frac{1}{t_{n-j}t_j} = C \frac{k^{p+1}}{t_n} \left(\frac{1}{t_j} + \frac{1}{t_{n-j}} \right), \end{aligned}$$

so that, again with an obvious modification for the term with $j = 0$,

$$\left\| \sum_{j=0}^{n-1} E_k^{n-1-j} \eta_{j2} \right\| \leq C \frac{k^{p+1}}{t_n} \sum_{j=0}^{n-1} \left(\frac{1}{t_{j+1}} + \frac{1}{t_{n-j}} \right) \leq C \frac{k^p}{t_n} \log \frac{t_{n+1}}{k}.$$

It remains to estimate the term in η_{j3} which we write

$$\eta_{j3} = Q_j(u_l) = Q_j(-Au + \varphi(u)) = -AQ_j(u) + Q_j(\varphi).$$

For the second term we have by (5.3) and our assumption (5.1)

$$\|Q_j(\varphi)\| \leq C \frac{k^{p+1}}{t_{j+1}} = C \frac{k^p}{j+1} \quad \text{for } j = 0, \dots, n-1,$$

and hence

$$\left\| \sum_{j=0}^{n-1} E_k^{n-1-j} Q_j(\varphi) \right\| \leq Ck^p \log \frac{t_{n+1}}{k}.$$

To estimate the first, we note that

$$\begin{aligned} \|E_k^n Av\| &\leq \|(r(kA)^n - e^{-nkA})Av\| + \|e^{-nkA}Av\| \\ &\leq C \frac{k}{t_{n+1}} \|Av\| + \frac{C}{t_{n+1}} \|v\| \leq \frac{C}{t_{n+1}} (\|v\| + k\|Av\|) \quad \text{for } n \geq 0 \end{aligned}$$

(if $r(\infty) = 0$ we have more directly $\|E_k^n Av\| = \|r(kA)^n Av\| \leq Ct_n^{-1}\|v\|$), and hence, using (5.3) with $l = p$ and $l = p-1$, and noting also that $tAu^{(p-1)}$ is bounded by our assumptions,

$$\|E_k^{n-1-j} AQ_j(u)\| \leq \frac{C}{t_{n-j}} (\|Q_j(u)\| + k\|Q_j(Au)\|) \leq Ck^{p+1} \frac{1}{t_{n-j}t_{j+1}},$$

so that

$$\sum_{j=0}^{n-1} \|E_k^{n-1-j} A Q_j(u)\| \leq C \frac{k^p}{t_n} \log \frac{t_{n+1}}{k}.$$

Altogether, we thus have

$$\|e_n\| \leq Ck \sum_{j=0}^{n-1} \|e_j\| + Ck^p t_n^{-1} \log \frac{t_{n+1}}{k},$$

from which our result follows in exactly the same way as in the proof of Theorem 1.

As an example where Theorem 2 applies, consider as in Section 2 the case that $H = L_2(\Omega)$, where Ω is a domain with smooth boundary in R^d , now with $d \leq 3$, where $A = -\Delta$ with $D(A) = H_0^1(\Omega) \cap H^2(\Omega)$, and where $f(t, u)$ is generated by a smooth function $\tilde{f}(x, t, u)$ on $\bar{\Omega} \times \bar{J} \times R$ which is bounded together with its derivatives of first and second order in t and u . The equation (2.1) thus reads

$$(5.6) \quad \begin{aligned} u_t - \Delta u &= \tilde{f}(x, t, u) && \text{in } \Omega \times J, \\ u &= 0 && \text{on } \partial\Omega \times J, \\ u(x, 0) &= v(x) && \text{in } \Omega. \end{aligned}$$

Assuming now that $v \in D(A)$ with $\|\Delta v\| = \|\Delta v\|_{L_2(\Omega)} \leq \rho$, it follows from Lemma 1 of Section 2 that u and $u' = u_t$ are bounded in H for $t \in \bar{J}$. To see that also $tu'' = tu_{tt}$ is bounded in H , we differentiate (5.6) twice to obtain

$$(5.7) \quad u_{tt} - \Delta u_t = \tilde{f}_t + \tilde{f}_u \cdot u_t,$$

and

$$(5.8) \quad u_{ttt} - \Delta u_{tt} = \tilde{f}_{tt} + 2\tilde{f}_{tu}u_t + \tilde{f}_{uu} \cdot u_t^2 + \tilde{f}_u u_{tt}.$$

Note that, since $d \leq 3$, we have

$$(5.9) \quad \|\varphi\|_{L_\infty(\Omega)} \leq C\|\varphi\|_{H^2(\Omega)} \leq C\|\Delta\varphi\| \quad \text{for } \varphi \in D(A).$$

Therefore, since u_t is bounded in H , we obtain from (5.7)

$$\|u_t^2\| \leq \|u_t\|_{L_\infty(\Omega)} \|u_t\| \leq C\|\Delta u_t\| \leq C\|u_{tt}\| + C.$$

Multiplication of (5.8) by $t^2 u_{tt}$ and integration over Ω gives

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} (t^2 \|u_{tt}\|^2) + t^2 \|\nabla u_{tt}\|^2 &\leq Ct^2 (\|u_{tt}\|^2 + \|u_{tt}\|) + t \|u_{tt}\|^2 \\ &\leq Ct \|u_{tt}\|^2 + C, \end{aligned}$$

and hence

$$t^2 \|u_{tt}\|^2 \leq C \int_0^t s \|u_{tt}\|^2 ds + C.$$

Multiplication of (5.7) by tu_{tt} and integrating gives similarly

$$\int_0^t s \|u_{tt}\|^2 ds \leq C \int_0^t \|\nabla u_t\|^2 ds + C,$$

and using instead (5.7) multiplied by u_t we have finally

$$\int_0^t \|\nabla u_t\|^2 ds \leq C\|u_t(0)\|^2 + C \leq C\|\Delta v\|^2 + C \leq C.$$

Together, our estimates show the boundedness of tu_{tt} in H . We also have for $\varphi(t) = f(t, u(t))$ that φ and $\varphi' = \tilde{f}_t + \tilde{f}_u u_t$ are bounded. Finally, in order to see that $t\varphi''$ is bounded, we note that by the above,

$$\|\varphi''\| = \|\tilde{f}_{tt} + 2\tilde{f}_{ut}u_t + \tilde{f}_{uu}u_t^2 + \tilde{f}_{uu}u_{tt}\| \leq C(\|u_{tt}\| + 1) \leq Ct^{-1}.$$

The assumptions of Theorem 2 are thus satisfied with $p = 2$, and we conclude that for such methods

$$\|U_n - u(t_n)\| \leq C(\rho)k^2 \left(t_n^{-1} \log \frac{t_{n+1}}{k} + \left(\log \frac{t_{n+1}}{k} \right)^2 \right) \quad \text{if } \|\Delta v\| \leq \rho.$$

The same method can be applied to the discretization in time of the equation obtained from (5.6) by discretization in space. Consider for example, as in Section 2, the semidiscrete equation (2.11) or (2.12) with continuous piecewise linear approximating functions, now on a quasiuniform partitioning of Ω . With $v_h = P_1 v = -T_h \Delta v$ the elliptic projection of v onto S_h , we have for $v \in D(\Delta)$

$$\|u_{h,t}(0)\| = \|\Delta_h v_h + P_0 f(0, v_h)\| \leq \|\Delta v\| + C.$$

It is easy to show that the error in the semidiscrete solution is then bounded as

$$\|u_h(t) - u(t)\| \leq C(\rho)h^2 \quad \text{for } \Delta v \leq \rho,$$

and we also conclude by Lemma 1 that u_h and $u_{h,t}$ are bounded in $L_2(\Omega)$ with Δv . The same arguments as above will then show that $tu_{h,tt}$ is bounded in $L_2(\Omega)$, uniformly in h , provided only that the analogue of (5.9) is valid in the present situation, namely

$$(5.10) \quad \|\chi\|_{L_\infty(\Omega)} \leq C\|\Delta_h \chi\| \quad \text{for } \chi \in S_h,$$

or, equivalently,

$$\|T_h \chi\|_{L_\infty(\Omega_h)} \leq C\|\chi\| \quad \text{for } \chi \in S_h,$$

where $\Omega_h (\subset \Omega)$ is the union of the simplices in the definition of S_h . But with $T = (-\Delta)^{-1}$ we have by (5.9)

$$\|T\chi\|_{L_\infty(\Omega)} \leq C\|\chi\|,$$

and it remains to estimate $(T_h - T)\chi$, the error in the elliptic problem with right-hand side χ . By well-known error and regularity estimates for the elliptic problem (cf. Schatz and Wahlbin [4]) and an inverse estimate to estimate the norm of χ in $H^{1/2+\epsilon}(\Omega)$ by that in $L_2(\Omega)$, we have with $0 < \epsilon < \frac{1}{2}$,

$$\begin{aligned} \|(T_h - T)\chi\|_{L_\infty(\Omega_h)} &\leq C \log \frac{1}{h} \inf_{\tilde{\chi} \in S_h} \|T\chi - \tilde{\chi}\|_{L_\infty(\Omega_h)} \leq Ch \log \frac{1}{h} \|T\chi\|_{W_\infty^1(\Omega)} \\ &\leq Ch \log \frac{1}{h} \|T\chi\|_{H^{5/2+\epsilon}(\Omega)} \leq Ch \log \frac{1}{h} \|\chi\|_{H^{1/2+\epsilon}(\Omega)} \\ &\leq Ch^{1/2-\epsilon} \log \frac{1}{h} \|\chi\| \leq C\|\chi\|. \end{aligned}$$

Together, our estimates show (5.10) and thus complete the proof that $tu_{h,tt}$ is bounded. As above, it follows that $\varphi_h = P_0 \tilde{f}(\cdot, t, u)$ is bounded together with φ'_h and $t\varphi''_h$, uniformly in h , and we conclude by Theorem 2 for the completely discrete

solution U_n obtained by discretization in time of (2.12) that

$$\|U_n - u(t_n)\| \leq C(\rho) \left\{ h^2 + k^2 \left[t_n^{-1} \log \frac{t_{n+1}}{k} + \left(\log \frac{t_{n+1}}{k} \right)^2 \right] \right\}$$

for $v \in D(A)$ with $\|\Delta v\| \leq \rho$.

For methods which are higher-order in space, a higher power of h may be obtained in combination with some negative power of t_n .

We close by exhibiting two examples of methods which satisfy our assumptions with $p = 2$ (cf. [2]). First let

$$\tau_1 = \frac{1}{3}, \quad \tau_2 = 1, \quad \mathcal{A} = \begin{pmatrix} \frac{5}{12} & -\frac{1}{12} \\ \frac{3}{4} & \frac{1}{4} \end{pmatrix}, \quad b^T = \left(\frac{3}{4}, \frac{1}{4} \right).$$

Then $U_{n+1} = U_{n2}$ and $r(z) = (1 - \frac{1}{3}z)/(1 + \frac{2}{3}z + \frac{1}{6}z^2)$ is the Padé approximant of e^{-z} of orders (1, 2) and satisfies $r(\infty) = 0$. Secondly, with

$$\tau_1 = \frac{1}{2} + \frac{1}{2\sqrt{3}}, \quad \tau_2 = \frac{1}{2} - \frac{1}{2\sqrt{3}}, \quad \mathcal{A} = \begin{pmatrix} \tau_1 & 0 \\ -\frac{1}{\sqrt{3}} & \tau_1 \end{pmatrix}, \quad b^T = \left(\frac{1}{2}, \frac{1}{2} \right),$$

we have (cf. Calahan's scheme)

$$r(z) = 1 - \frac{z}{1 + \tau_1 z} - \frac{\sqrt{3}}{6} \left(\frac{z}{1 + \tau_1 z} \right)^2 \quad \text{with } |r(\infty)| < 1.$$

IRISA

UER Mathématiques et Informatique

Université de Rennes I

Campus de Beaulieu

35042 Rennes Cédex, France

Department of Mathematics

Chalmers University of Technology

S-412 96 Göteborg, Sweden

1. G. A. BAKER, J. H. BRAMBLE & V. THOMÉE, "Single step Galerkin approximations for parabolic problems," *Math. Comp.*, v. 31, 1977, pp. 818–847.

2. M. CROUZEIX, *Sur l'Approximation des Équations Différentielles Opérationnelles Linéaires par des Méthodes de Runge-Kutta*, Thèse, Université Paris VI, 1975.

3. C. JOHNSON, S. LARSSON, V. THOMÉE & L. B. WAHLBIN, "Error estimates for spatially discrete approximations of semilinear parabolic equations with nonsmooth initial data," *Math. Comp.*, v. 49, 1987, pp. 331–357.

4. A. H. SCHATZ & L. B. WAHLBIN, "On the quasi-optimality in L_∞ of the \dot{H}^1 -projection into finite element spaces," *Math. Comp.* v. 38, 1982, pp. 1–22.