

COMPUTATIONAL ESTIMATION OF THE ORDER OF $\zeta(\frac{1}{2} + it)$

TADEJ KOTNIK

ABSTRACT. The paper describes a search for increasingly large extrema (ILE) of $|\zeta(\frac{1}{2} + it)|$ in the range $0 \leq t \leq 10^{13}$. For $t \leq 10^6$, the complete set of ILE (57 of them) was determined. In total, 162 ILE were found, and they suggest that $\zeta(\frac{1}{2} + it) = \Omega(t^{2/\sqrt{\log t \log \log t}})$. There are several regular patterns in the location of ILE, and arguments for these regularities are presented. The paper concludes with a discussion of prospects for further computational progress.

1. INTRODUCTION

Riemann's zeta function on the critical line, $\zeta(\frac{1}{2} + it)$, is unbounded. Balasubramanian and Ramachandra have shown in 1977 [1] that

$$\zeta(\tfrac{1}{2} + it) = \Omega(t^{\frac{3}{4\sqrt{\log t \log \log t}}})$$

whereas Huxley proved in 1993 [3] that

$$\zeta(\tfrac{1}{2} + it) = O(t^{\frac{89}{570} + \varepsilon}) \quad \text{for every } \varepsilon > 0.$$

This leaves a considerable gap between the Ω - and O -results. Already in 1908, Lindelöf conjectured a much stronger O -bound [4]

$$\zeta(\tfrac{1}{2} + it) = O(t^\varepsilon) \quad \text{for every } \varepsilon > 0.$$

The truth of this conjecture, known as Lindelöf's hypothesis, would follow from that of Riemann's hypothesis, since the latter can only hold if [8]

$$\zeta(\tfrac{1}{2} + it) = O(t^{\frac{C}{\log \log t}}) \quad \text{for some } C > 0.$$

Since $|\zeta(\frac{1}{2} + it)| = |Z(t)|$, where $Z(t)$ is the Riemann-Siegel Z function, the conjectures and results about the order of $\zeta(\frac{1}{2} + it)$ may, and henceforth, will be stated more compactly in terms of $Z(t)$. As $Z(t)$ is an even function, any discussion about its behavior will be restricted to $t \in \mathbb{R}_+$ without loss of generality, so "at values of t smaller than T " will always mean $0 \leq t < T$. The acronym *ILE* will be used for *increasingly large extrema* of $|Z(t)|$, and an interval bounded by two consecutive zeros of $Z(t)$ will be referred to as an *interzero interval*.

A computational search for large values of $|Z(t)|$ obviously cannot provide rigorous Ω - and O -results. Still, the results presented in this paper show that with a sufficiently comprehensive set of ILE determined in a sufficiently large t -interval, certain regularities in the values of $Z(t)$ at ILE become detectable. The values of ILE in the interval $0 \leq t \leq 10^{13}$ suggest that the Ω -bound of $Z(t)$ could be

Received by the editor April 24, 2002 and, in revised form, October 21, 2002.

2000 *Mathematics Subject Classification*. Primary 11M06, 11Y60; Secondary 11Y35, 65A05.

Key words and phrases. Riemann's zeta function, critical line, Lindelöf's hypothesis.

improved substantially. On the other hand, a much broader t -interval would have to be investigated to suggest potential improvements of the O -bound of $Z(t)$.

2. METHODS OF COMPUTATION

2.1. General. The computations were performed on a PC equipped with a 1700 MHz Intel Pentium 4 processor. The values of $Z(t)$ and $\vartheta(t)$ were computed with Mathematica 4.0 (Wolfram Research, Urbana, IL, USA) using the `RiemannSiegelZ` and `RiemannSiegelTheta` routines, respectively. The search algorithm was run using 16-digit precision, while the values of ILE were determined with 24-digit precision. Least-squares regression was performed with Sigma Plot 6.0 (SPSS Science, Chicago, IL, USA).

2.2. Determination of all ILE for $0 \leq t \leq 10^6$. In $\mathcal{T}_1 := [0, 10^6]$, $Z(t)$ has 1747146 zeros,¹ and Riemann's hypothesis is never violated there [7]. Hence $Z(t)$ has exactly one local extremum in each interzero interval in \mathcal{T}_1 [2]. Together with three extrema below the first zero, there are thus 1747148 local extrema of $Z(t)$ in \mathcal{T}_1 . Of these extrema, 57 are ILE, forming the list \mathcal{Z}_1 (see the Appendix).

Section 4.1 presents two plausible theoretical arguments for the proximity of large extrema of $|Z(t)|$ to the points $t_k := \frac{2k\pi}{\log 2}$, $k \in \mathbb{N}$. This indeed appears to be the case — *each* of the interzero intervals containing an ILE of \mathcal{Z}_1 also contains such a point. Furthermore, in all cases, $|Z(t_k)|$ exceeds 47% of the maximum value of $|Z(t)|$ in the same interzero interval, and on average, it exceeds 91% of that value.

2.3. Search for ILE for $10^6 < t \leq 10^9$. The regularity in the location of ILE in \mathcal{Z}_1 suggests that many large $|Z(t)|$ are located in the interzero intervals containing a point t_k and a relatively large $|Z(t_k)|$. The search for ILE in $\mathcal{T}_2 := (10^6, 10^9]$ was performed as follows:

- (1) $Z(t_k)$ was computed;
- (2) if $|Z(t_k)|$ exceeded 20% of the largest ILE for smaller t , the local extremum was computed;
- (3) if $|Z(t)|$ at the extremum exceeded the largest ILE for smaller t , it was added to the list \mathcal{Z}_2 .

None of the ILE in \mathcal{T}_1 would have been missed by this algorithm. In total, the list \mathcal{Z}_2 consists of 43 extrema, and they are given in the Appendix.

Section 4.2 sketches an argument for another regular pattern in the location of large extrema of $|Z(t)|$. Denoting by d_p the absolute deviation of $\frac{k \log p}{\log 2}$ (p prime) from an integer, a large $|Z(t_k)|$ is likely if d_3, d_5, d_7, \dots , are relatively small. The list \mathcal{Z}_2 provides a sample of d_p for 43 ILE in \mathcal{T}_2 . The increase of d_p with p in \mathcal{Z}_2 is rather rapid; thus $\text{mean}(d_3) = 0.0281\dots$, $\text{max}(d_3) = 0.0861\dots$, and $\text{mean}(d_{47}) = 0.1581\dots$, $\text{max}(d_{47}) = 0.4966\dots$.

2.4. Search for ILE for $10^9 < t \leq 10^{13}$. Since in \mathcal{Z}_2 the d_p for small p are small, ILE near t_k with large d_p are unlikely. The ranges of permitted d_p were chosen on the basis of their respective values in \mathcal{Z}_2 , and the search for ILE in $\mathcal{T}_3 := (10^9, 10^{13}]$ was performed as follows:

- (1) the values of d_p , $3 \leq p \leq 17$, were checked to be within prescribed ranges:
 $d_3 \leq 0.10$, $d_5 \leq 0.15$, $d_7 \leq 0.20$, $d_{11} \leq 0.25$, $d_{13} \leq 0.28$, $d_{17} \leq 0.30$;

¹The list of zeros, accurate to $\pm 10^{-9}$, was kindly provided by Dr. Andrew M. Odlyzko.

- (2) if the value of k qualified, $Z(t_k)$ was computed;
- (3) if $|Z(t_k)|$ exceeded 20% of the largest ILE for smaller t , the local extremum was computed;
- (4) if $|Z(t)|$ at the extremum exceeded the largest ILE for smaller t , it was added to the list \mathcal{Z}_3 .

None of the ILE found in \mathcal{T}_2 would have been missed with this choice of bounds on d_3, \dots, d_{17} . In total, the list \mathcal{Z}_3 consists of 62 extrema, and they are given in the Appendix.

3. RESULTS AND DISCUSSION

Let

$$a(t) := \frac{\log |Z(t)|}{\log t} \quad \text{and} \quad b(t) := \frac{\log |Z(t)|}{\sqrt{\log t}} \frac{\sqrt{\log \log t}}{\sqrt{\log t}}.$$

Denoting $\limsup_{t \rightarrow \infty} a(t) = A$ and $\limsup_{t \rightarrow \infty} b(t) = B$, we have $0 \leq A \leq \frac{89}{570}$ by the theorem of Huxley, and $\frac{3}{4} \leq B \leq \infty$ by the theorem of Balasubramanian and Ramachandra. At sufficiently large t , where large $|Z(t)|$ start to reflect the actual order of $Z(t)$, the values of $a(t)$ and $b(t)$ at ILE should start to approach the true values of A and B , respectively.

Figure 1 shows the values of $a(t)$ and $b(t)$ for ILE in $\mathcal{Z}_1 \cup \mathcal{Z}_2 \cup \mathcal{Z}_3$, excluding $|Z(0)|$, and for $|Z(4.257... \times 10^{15})| = 855.3...$ in the vicinity of a point located by Odlyzko [5]. The values of $a(t)$ at ILE seem to delineate a monotonically decreasing asymptote for $t > 10^3$, but these a -values are too large to suggest a stronger upper bound of A than the value $\frac{89}{570} = 0.1561...$ imposed by the theorem of Huxley. On the other hand, the values of $b(t)$ at ILE seem to delineate a monotonically increasing asymptote for all t , exceeding for $t > 10^2$ the lower bound of B imposed by the theorem of Balasubramanian and Ramachandra. Close to the upper bound of the investigated t -range, we have $b(t) > 2$, and the asymptotic increase of $b(t)$ seems to continue, which suggests that

$$Z(t) = \Omega(t^{2/\sqrt{\log t \log \log t}}).$$

It seems likely that the extension of the range of ILE to larger t would allow to strengthen this tentative estimate.




FIGURE 1.

4. PATTERNS IN THE LOCATION OF LARGE EXTREMA

4.1. Proximity of $\frac{t \log 2}{2\pi}$ to \mathbb{N} . As described in Section 2.2, of the subzero interval and the 55 interzero intervals containing ILE in \mathcal{Z}_1 , *each* also contains a point $t_k := \frac{2k\pi}{\log 2}$. Plausible arguments for this can be derived from at least two starting points.

Argument A. From the well-known formula

$$\zeta(\sigma + it) = \frac{1}{(1 - 2^{1-\sigma-it})} \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^{\sigma+it}} \quad \text{for } \sigma > 0$$

we have

$$|Z(t)| = (3 - 2\sqrt{2} \cos(t \log 2))^{-1/2} \left| \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^{1/2+it}} \right|.$$

Large $|Z(t_k)|$ can then be explained by periodicity of $(3 - 2\sqrt{2} \cos(t \log 2))^{-1/2}$, with maxima of $\sqrt{2} + 1$ at $t = \frac{2k\pi}{\log 2}$ and minima of $\sqrt{2} - 1$ at $t = \frac{(2k-1)\pi}{\log 2}$.

Argument B. We invoke the main sum in the Riemann-Siegel formula

$$Z_0(t) = 2 \sum_{1 \leq n \leq \sqrt{\frac{t}{2\pi}}} \frac{\cos(\vartheta(t) - t \log n)}{\sqrt{n}},$$

where $\vartheta(t)$ is the Riemann-Siegel theta function. At the points $t = t_k$ we have $\cos(\vartheta(t_k) - t_k \log n) = \cos \vartheta(t_k)$ for summands with $n = 2^m$, $m \in \{0\} \cup \mathbb{N}$, which therefore reinforce each other (i.e., have the same sign), and

$$Z_0(t_k) = 2 \cos \vartheta(t_k) \sum_{\substack{1 \leq n \leq \sqrt{\frac{t}{2\pi}} \\ n=2^m}} \frac{1}{\sqrt{n}} + 2 \sum_{\substack{3 \leq n \leq \sqrt{\frac{t}{2\pi}} \\ n \neq 2^m}} \frac{\cos(\vartheta(t_k) - t_k \log n)}{\sqrt{n}}.$$

4.2. Proximity of $k \frac{\log p}{\log 2}$ to \mathbb{N} . For $\{t_{k(3)}\} \subset \{t_k\}$, for which $\frac{k \log 3}{\log 2} \approx l \in \mathbb{N}$, we have $\frac{2k\pi}{\log 2} \approx \frac{2l\pi}{\log 3}$, so $\cos(\vartheta(t_k) - t_k \log n) \approx \cos \vartheta(t_k)$ for summands with $n = 3^m$ and $n = 2^m 3^{m'}$, with $m, m' \in \mathbb{N}$, and these summands are also mutually reinforcing. Denoting $\{t_{k(3,5)}\} \subset \{t_{k(3)}\}$, for which $\frac{k \log 5}{\log 2}$ is also close to an integer, mutual reinforcement also occurs for summands with $n = 5^m$, $n = 2^m 5^{m'}$, $n = 3^m 5^{m'}$, and $n = 2^m 3^{m'} 5^{m''}$. Thus, $\{t_{k(3,5,7)}\}$, $\{t_{k(3,5,7,11)}\}, \dots$ are subsets of points t_k at which large $|Z(t)|$ are increasingly likely.

4.3. Proximity of $\frac{\vartheta(t)}{\pi}$ to \mathbb{N} . The Riemann-Siegel formula provides another hint about the location of large values of $|Z(t)|$. The mutually reinforcing terms (see Section 4.2) are proportional to $|\cos \vartheta(t)|$, which is the largest if t corresponds to a Gram point (a point $t = g_m > 7$ such that $\vartheta(g_m) = m\pi$, $m \in \{-1, 0\} \cup \mathbb{N}$). In fact, for each of the 105 ILE in $\mathcal{Z}_2 \cup \mathcal{Z}_3$, either at the closest Gram point below t_k , or at the closest Gram point above t_k , $|Z(g_m)|$ exceeds 99.2% of the value at the local extremum.² Among the t_k that qualify both by proximity of $\frac{k \log p}{\log 2}$ to integers and by a large $|Z(t_k)|$, further selection of the candidates for ILE can thus be made by computing $|Z(t)|$ at the two Gram points closest to t_k .

²Of the two Gram points closest to t_k , it is *not always* the one closer to t_k at which $|Z(t)|$ is large (e.g., for $k = 954$, the closest Gram point is $t = g_{8571}$, yet $|Z(t)|$ is larger at $t = g_{8570}$).

4.4. Partial Riemann-Siegel sums at large $|Z(t)|$. Let

$${}_rZ_0(t) := 2 \sum_{1 \leq n \leq m} \frac{\cos(\vartheta(t) - t \log n)}{\sqrt{n}}, \quad \text{where } m = \left\lceil \left(\frac{t}{2\pi} \right)^{1/(2+r)} \right\rceil,$$

so that $Z_0(t) \equiv {}_0Z_0(t)$. For 61 of the 62 ILE in \mathcal{Z}_3 , the value of ${}_1Z_0$ at the corresponding point t_k exceeds 9.0% of the value of Z at the extremum. Furthermore, for all 62 ILE in \mathcal{Z}_3 , the value of ${}_1Z_0$ (resp. ${}_2Z_0$) at one of the two Gram points closest to t_k exceeds 39.5% (resp. 19.6%) of the value of Z at the extremum. In all these cases, the sign of ${}_rZ_0$ at the considered point equals the sign of Z at the extremum. Thus, evaluation of ${}_1Z_0$ at points t_k and of either ${}_1Z_0$ or ${}_2Z_0$ at Gram points could be used for elimination of unlikely ILE candidates, significantly reducing the number of complete Z -evaluations.

5. PROSPECTS FOR FURTHER PROGRESS

The analysis of the order of $Z(t)$ by means of the functions $a(t)$ and $b(t)$ is based on the rigorously established results, $Z(t) = O(t^A)$ and $Z(t) = \Omega(t^{B/\sqrt{\log t \log \log t}})$, and as such might be viewed as rather conservative. It would be tempting to evaluate a stronger Ω -conjecture than the one tested through $b(t)$, e.g., by considering the function $g(t) := \log |Z(t)| / \sqrt{\log t}$ to test the conjecture $Z(t) = \Omega(t^{G/\sqrt{\log t}})$ for some $G > 0$. However, the results of such a procedure could be misleading, as we have no knowledge of the multiplicative constant involved in the order of $Z(t)$. For example, the values of $|Z(t)|$ at ILE agree rather well (with the correlation coefficient $R = 0.9994$ for ILE with $t > 10^3$) with the estimate $|Z(t)| = 0.0199t^{3.36/\sqrt{\log t \log \log t}}$. If this were actually the case, then $g(t)$ at ILE would increase up to $t \approx 10^{89}$, and in any computationally accessible t -range one would be led to the wrong conclusion that $G > 0$. In other words, while $g(t)$ at ILE increases for $t \leq 10^{13}$ and exceeds the value of 1, there is no guarantee that $\limsup_{t \rightarrow \infty} g(t) > 0$.

One might also be tempted to extrapolate. That is, if the functional forms of the asymptotes that the values of $a(t)$ and $b(t)$ at ILE seem to outline were identified correctly, say as $a_S(t)$ and $b_S(t)$, the respective limits as $t \rightarrow \infty$ would yield estimates of A and B . Yet, without any theoretical indications with respect to what the functions $a_S(t)$ and $b_S(t)$ should be, such an identification would amount to guessing, and it is unclear how one could assess its correctness. For example, the values of $a(t)$ at ILE agree reasonably well ($R = 0.9985$ for ILE with $t > 10^3$) with the power-decay function $a_S(t) = 0.149 + 0.255t^{-0.0528}$, which would suggest that $Z(t) = \Omega(t^{0.149})$. This estimate would contradict Lindelöf's (and hence Riemann's) hypothesis, and while it also agrees well with the data for $t < 10^{13}$, for sufficiently large t it is destined to run into a complete disagreement with the estimate of $|Z(t)|$ at ILE given in the previous paragraph.

It is sometimes supposed that if any violations of Riemann's hypothesis exist, they could be located close to very large values of $|Z(t)|$. There are no such violations in the vicinity of the 162 ILE determined in this study.

The computations presented in this paper took approximately nine months using a personal computer. At the time of writing, the most powerful supercomputers could have handled this task at least one thousand times faster. It is unlikely that a supercomputer would be dedicated somewhere to the search for further ILE, but this search could also be distributed among a number of personal computers,

with the rate of advancement proportional to the total computing power of the computers involved.³ In addition, the search could be accelerated by selecting ILE candidates through partial Riemann-Siegel sums (Section 4.4) and by computing the extrema using the Odlyzko-Schönhage algorithm [6].

APPENDIX

	t	$Z(t)$
$\overline{\mathcal{Z}}_1$ 1	0.000000	-1.460
2	10.212075	-1.552
3	17.882582	2.341
4	27.735883	2.847
5	35.392730	2.942
6	45.636113	-3.665
7	63.060428	-4.167
8	90.723857	4.477
9	108.986791	5.193
10	171.759106	-4.980
11	199.651794	6.063
12	245.532580	6.069
13	280.810364	-7.003
14	371.545466	7.570
15	480.401432	-8.250
16	652.212123	9.158
17	897.836383	9.406
18	1069.360643	9.851
19	1178.449084	10.355
20	1378.316536	-10.468
21	1550.029928	11.077
22	1967.268238	11.271
23	2030.520469	11.730
24	2447.635780	13.371
25	3099.906368	13.479
26	3825.816853	-13.497
27	3997.707224	-13.575
28	4478.096605	-14.755
29	6726.121510	-15.612
30	6925.621938	-15.955
31	8475.812323	-16.252
32	8647.210888	16.391
33	9173.716528	16.506
34	10025.578053	16.906
35	10677.929307	-17.237
36	11204.207758	17.337

	t	$Z(t)$
37	12645.135236	-18.006
38	13125.470242	18.091
39	14303.975890	19.817
40	22299.074877	21.059
41	24329.633861	21.434
42	30774.966419	23.228
43	50626.478383	23.747
44	55104.583439	-24.830
45	63751.863162	-26.073
46	74956.025038	-27.694
47	77403.722067	28.216
48	105731.032300	28.853
49	130060.556256	31.415
50	152359.757336	32.671
51	260538.282724	34.161
52	314464.228643	34.516
53	328768.228899	-36.689
54	521928.541866	36.739
55	534573.688201	-40.991
56	865898.755362	-42.392
57	929650.688269	-43.107
$\overline{\mathcal{Z}}_2$ 58	1024177.378756	44.063
59	1345367.802772	-47.593
60	1923053.135018	-48.350
61	2186410.518907	-50.879
62	2939652.714358	53.233
63	3268420.883436	-55.204
64	3345824.546021	55.767
65	5419578.489302	-58.425
66	6155416.653707	61.038
67	9850232.528074	-62.448
68	9969615.203761	62.793
69	11026769.624984	-65.674
70	12372137.487612	-67.952
71	15236834.026567	-68.116
72	15457423.712975	74.268

³This strategy is being applied efficiently in an ongoing computation of the zeros of Riemann's zeta function, which has so far shown that Riemann's hypothesis holds for $|t| < 3 \times 10^{10}$ [9].

	t	$Z(t)$		t	$Z(t)$
73	28642802.916415	-75.213	119	21559062801.941668	-192.996
74	28660206.960842	75.625	120	22412382038.812786	-196.059
75	30694257.761606	79.679	121	23165396411.338070	196.477
76	37002034.097306	-80.035	122	25985505104.438565	-197.606
77	42792359.891727	-80.513	123	27279224693.810314	204.462
78	46747714.116054	-82.469	124	27331684151.577735	209.054
79	53325356.508449	-84.321	125	31051083602.364182	213.898
80	60090302.842436	84.715	126	38688523992.011831	224.263
81	81792403.155463	85.761	127	62792807608.657779	-228.392
82	82985411.177787	-86.254	128	79881740253.040389	233.330
83	87568424.951600	91.882	129	102108905446.095547	240.103
84	99273480.761352	-91.989	130	108903432915.370254	242.415
85	102805259.027575	92.643	131	124855728535.680010	-246.885
86	119015924.891142	92.654	132	131443859639.685072	251.267
87	124570459.059572	95.158	133	133159989048.388546	251.576
88	144327207.118141	-95.326	134	165822762086.732367	-254.192
89	151614082.016804	97.031	135	170165889140.424800	-256.095
90	173723252.257957	-101.319	136	192604855973.407448	258.354
91	178900422.227382	103.906	137	197804421842.227818	-262.702
92	244946055.644911	108.011	138	243860776768.360133	-271.338
93	298271412.198149	108.187	139	297280771283.496679	-276.661
94	363991205.176448	-114.451	140	326473979757.428188	-289.781
95	418878041.160027	-118.153	141	461305748544.638105	292.784
96	607838127.431023	118.447	142	472692195365.796730	-293.833
97	631240860.404037	119.782	143	479489261691.339254	293.845
98	673297382.192693	124.043	144	514119669706.650653	295.026
99	868556070.995988	128.017	145	576555893019.852818	295.375
100	900138526.590236	-128.993	146	643049954739.247192	-297.567
\mathcal{Z}_3 101	1189754916.313216	-130.488	147	669980906189.791285	301.088
102	1253191043.688385	133.120	148	722931694992.231828	309.299
103	1387123309.986048	148.728	149	812980259631.147353	-334.401
104	2287261836.552282	149.404	150	1459387308608.408274	349.779
105	3238682014.814266	149.611	151	1765497206246.212277	354.787
106	3443895116.936669	-152.488	152	2515593134489.563683	-361.066
107	4209002696.395103	155.270	153	2589877332690.841810	370.395
108	4266153346.590529	157.986	154	3210707929490.468401	375.250
109	4945603697.701426	-160.578	155	4154422573264.686997	-376.393
110	5230260126.511580	-164.581	156	4778933265685.642359	379.550
111	5272517912.850547	170.199	157	5695465916337.181354	388.067
112	7181324522.908048	-171.458	158	6586779209214.248987	-403.914
113	7965404181.305970	-176.842	159	7709188977559.148583	405.312
114	11166740191.846172	180.227	160	8743721888758.038535	415.783
115	12251628740.237935	181.884	161	9090142088295.475463	416.329
116	13066290725.695175	183.530	162	9918400224732.229613	-441.106
117	18168214001.673350	190.187			
118	19018488753.002784	192.635		4257232978148261.797669	855.364

ACKNOWLEDGMENTS

I would like to thank Dr. Jan van de Lune (Hallum, The Netherlands), Prof. Roger Heath-Brown FRS (Oxford University), and Prof. Andrew M. Odlyzko (University of Minnesota) for many instructive discussions and suggestions.

REFERENCES

- [1] R. Balasubramanian and K. Ramachandra, *On the frequency of Titchmarsh's phenomenon for $\zeta(s)$. III*, Proc. Ind. Acad. Sci. **86A** (1977), 341-351. MR **58**:21968
- [2] H. M. Edwards, *Riemann's Zeta Function*, Academic Press, 1974, pp. 176-177. MR **57**:5922
- [3] M. N. Huxley, *Exponential sums and the Riemann zeta function. IV*, Proc. Lond. Math. Soc. **66** (1993), 1-40. MR **93j**:11056
- [4] E. Lindelöf, *Quelques remarques sur la croissance de la fonction $\zeta(s)$* , Bull. Sci. Math. **32** (1908), 341-356.
- [5] A. M. Odlyzko, *The 10^{20} -th zero of the Riemann zeta function and 175 million of its neighbors*, <http://www.dtc.umn.edu/~odlyzko/unpublished/index.html>
- [6] A. M. Odlyzko and A. Schönhage, *Fast algorithms for multiple evaluations of the Riemann zeta function*, Trans. Am. Math. Soc. **309** (1988), 797-809. MR **89j**:11083
- [7] J. B. Rosser, J. M. Yohe, and L. Schoenfeld, *Rigorous computation and the zeros of the Riemann zeta-function*, Proc. IFIP Congress 1968, North-Holland, 1969, pp. 70-76. MR **41**:2892
- [8] E. C. Titchmarsh and D. R. Heath-Brown, *The Theory of the Riemann Zeta-function*, 2nd ed., Oxford University Press, 1986, p. 354. MR **88c**:11049
- [9] S. Wedeniwski, *ZetaGrid—Verification of the Riemann hypothesis*, <http://www.zetagrid.net/zeta/index.html>

FACULTY OF ELECTRICAL ENGINEERING, UNIVERSITY OF LJUBLJANA, SI-1000 LJUBLJANA, SLOVENIA

E-mail address: `tadej.kotnik@fe.uni-lj.si`