NUMERICAL SIMULATION OF STOCHASTIC EVOLUTION EQUATIONS ASSOCIATED TO QUANTUM MARKOV SEMIGROUPS

CARLOS M. MORA

ABSTRACT. We address the problem of approximating numerically the solutions $(X_t : t \in [0, T])$ of stochastic evolution equations on Hilbert spaces $(\mathfrak{h}, \langle \cdot, \cdot \rangle)$, with respect to Brownian motions, arising in the unraveling of backward quantum master equations. In particular, we study the computation of mean values of $\langle X_t, AX_t \rangle$, where A is a linear operator. First, we introduce estimates on the behavior of X_t . Then we characterize the error induced by the substitution of X_t with the solution $X_{t,n}$ of a convenient stochastic ordinary differential equation. It allows us to establish the rate of convergence of $\mathbf{E} \langle \tilde{X}_{t,n}, A\tilde{X}_{t,n} \rangle$ to $\mathbf{E} \langle X_t, AX_t \rangle$, where $\tilde{X}_{t,n}$ denotes the explicit Euler method. Finally, we consider an extrapolation method based on the Euler scheme. An application to the quantum harmonic oscillator system is included.

1. INTRODUCTION

This paper is concerned with the computation of $\mathbf{E} \langle X_t, AX_t \rangle$, where A is a linear operator and X_t satisfies the following linear stochastic evolution equation on the infinite dimensional separable (complex) Hilbert space $(\mathfrak{h}, \langle \cdot, \cdot \rangle)$

$$(1.1) \quad X_t = x_0 + \int_0^t GX_s ds + \sum_{k=1}^m \int_0^t L_k X_s dW_s^k, \quad t \in [0, T],$$

with $x_0 \in \mathfrak{h}$, $\{W^k\}_{k=1, \dots, m}$ independent standard Brownian motions on a filtered complete probability space $(\Omega, F, (F_t)_{t \geq 0}, \mathbf{P})$, $T \in \mathbb{R}_+$ and $m \in \mathbb{N}$. Here, all integrals are understood in the Itô sense and $L_1, \dots, L_m$, G are general linear operators in $\mathfrak{h}$. Furthermore, we will focus our attention on the case

$$(1.2) \quad G = -iH - \frac{1}{2} \sum_{j=1}^m L_j^* L_j,$$

where H is a self-adjoint operator in $\mathfrak{h}$.

Received by the editor August 25, 2002 and, in revised form, January 7, 2003.

2000 *Mathematics Subject Classification.* Primary 60H35; Secondary 60H10, 60H15, 60H30, 65C30, 65C05.

Key words and phrases. Stochastic evolution equation, numerical solution, rate of convergence, Euler scheme, Galerkin method, quantum dynamical semigroup, quantum master equation.

This research has been partially supported by FONDECYT grant 2000036, a DIPUC Ph.D. grant and the program “Cátedra Presidencial on Qualitative Analysis of Quantum Dynamical Systems”.

Our main motivation came from the numerical simulation of open quantum systems. In particular, we are interested in problems arising in quantum optics. Next, we give a brief introduction to this application. In the framework of quantum mechanics the state of a physical system is defined by specifying an element of an adequate complex Hilbert space [1, 4]. Since $\mathfrak{h}$ plays the role here of this quantum state space, $\mathfrak{h}$ describes, for example, one-particle wave functions [1, 4] or the Fock representation of the states in the second quantization formalism [1]. In this context, x_0 is the initial state. Moreover, $\mathbf{E} \langle X_t, AX_t \rangle$ is interpreted as the mean value of the observable A at the instant t whenever A is self-adjoint. This interpretation is easily obtained in the approach that describes the dynamics of quantum systems by stochastic evolutions of the corresponding state vectors [3, 19]. This technique allows us to characterize, for instance, the dynamics of an individual quantum system which is continuously monitored by some measurement device [14]. Alternatively, in the Heisenberg picture the evolution of the observable A , in systems whose initial state is x_0 , is simulated by $\langle x_0, \Phi_t[A] x_0 \rangle$, where $\Phi_t[A]$ is the minimal solution of the backward quantum master equation

$$\begin{aligned} \Phi_0[A] &= A, \quad \text{and} \quad \forall \psi, \phi \in \mathfrak{D} = \text{Dom}(G) \cap \left(\bigcap_{k=1}^m \text{Dom}(L_k) \right), \\ (1.3) \quad \frac{d}{dt} \langle \psi, \Phi_t[A] \phi \rangle &= \langle G\psi, \Phi_t[A] \phi \rangle + \langle \psi, \Phi_t[A] G\phi \rangle + \sum_{j=1}^m \langle L_j \psi, \Phi_t[A] L_j \phi \rangle. \end{aligned}$$

Here H represents the Hamiltonian and the operators L_j , $j = 1, \dots, m$, describe the effects of the environment. Now, we have that if A is a bounded linear operator, then under weak conditions [10, 2]

$$(1.4) \quad \mathbf{E} \langle X_t, AX_t \rangle = \langle x_0, \Phi_t[A] x_0 \rangle.$$

We conclude this paragraph with some complementary remarks. First, the property (1.4) leads us to solve (1.3) through the weak numerical solution of (1.1). Furthermore, this procedure allows us to overcome the difficulties arising in the direct numerical integration of (1.3) in many cases; see, e.g., [19]. Second, Φ_t commonly define a minimal quantum dynamical semigroup on $B(\mathfrak{h})$ [7]. Third, (1.3) is based on the Born-Markov approximation of quantum phenomena, which has been used in quantum optics with great success. Finally, it is worth pointing out that there is a semigroup Ψ_t in the space of trace-class operators such that Φ_t is the adjoint of Ψ_t [7]; that is, Ψ_t is the predual semigroup of Φ_t . Hence

$$\mathbf{E} \langle X_t, AX_t \rangle = \text{Tr}(\Psi_t[P_{x_0}]A),$$

where Tr means the trace and P_{x_0} is the orthogonal projection of $\mathfrak{h}$ over the linear span of x_0 . Furthermore, Ψ_t solves the forward quantum master equation [10, 7] and represents the evolution of the density operators in the Schrödinger picture.

To compute $\mathbf{E} \langle X_t, AX_t \rangle$, we consider a sequence $(\mathfrak{h}_n)_n$ of finite dimensional Hilbert spaces such that

- P1:** $\mathfrak{h}_n \subset \text{Dom}(H) \cap (\bigcap_{k=1}^m \text{Dom}(L_k)) \cap (\bigcap_{k=1}^m \text{Dom}(L_k^*))$,
- P2:** $\forall x \in \mathfrak{h}, \exists x_n \in \mathfrak{h}_n$, such that $x_n \rightarrow_n x$.

Then we may proceed as follows:

Step 1: $\mathbf{E} \langle X_t, AX_t \rangle$ is approximated by $\mathbf{E} \langle X_{t,n}, AX_{t,n} \rangle$, where $X_{t,n}$ is a continuous adapted stochastic process with values on $\mathfrak{h}_n$ given by

$$(1.5) \quad X_{t,n} = P_n X_0 + \int_0^t G_n X_{s,n} ds + \sum_{k=1}^m \int_0^t L_{k,n} X_{s,n} dW_s^k,$$

with $P_n : \mathfrak{h} \rightarrow \mathfrak{h}_n$ the orthogonal projection of $\mathfrak{h}$ over $\mathfrak{h}_n$, $L_{k,n} = P_n L_k$, whenever $k = 1, \dots, m$, and

$$(1.6) \quad G_n = -iP_n H - \frac{1}{2} \sum_{j=1}^m P_n L_j^* P_n L_j.$$

Step 2: $\mathbf{E} \langle X_{t,n}, AX_{t,n} \rangle$ is simulated numerically using weak schemes for stochastic differential equations (SDE's)

Before going on, we would like to say a few words in relation to the definition of G_n . Fagnola and Chebotarev [5, 7] proved that the minimal quantum semigroup is Markov, i.e., $\Phi_t[I] = I$, under rather general hypotheses. Therefore (1.1) is conservative; that is, $\mathbf{E} \|X_t\|^2 = \|X_0\|^2$, in a wide class of applications. The approximations $X_{t,n}$ preserve this property if G_n is chosen as in (1.6). This is not necessarily the case if $G_n = P_n G$. Notice that the fact that (1.5) is conservative allows us to construct efficient numerical schemes [17].

Our primary objective is to start providing the numerical simulation of quantum phenomena using stochastic evolution equations like (1.1) [9, 3, 19, 14, 22] with a theoretical treatment. Particularly, in this article we are interested in the theoretical understanding of Steps 1 and 2.

This paper is organized in six sections. Section 2 is devoted to introducing notation and preliminary results. In Section 3 we assume analogous hypotheses to those used by Fagnola and Chebotarev to obtain the uniqueness of the solution of (1.3). Let C be a linear operator associated to (1.1) through these conditions. Then we estimate the quantities $\sup_{t \in [0, T]} \mathbf{E} \|C^{1/2} X_t\|^2$ and $\sup_{t \in [0, T]} \mathbf{E} \|C^{1/2} X_{t,n}\|^2$. These bounds, called “a priori” estimates on X_t and $X_{t,n}$, will play an important role in the characterization of the speed of convergence the approximations introduced in Steps 1 and 2. Section 4 provides the rate of convergence of $\mathbf{E} \langle X_{t,n}, AX_{t,n} \rangle$ to $\mathbf{E} \langle X_t, AX_t \rangle$ whenever $t \in [0, T]$. In other words, we address the problem of estimating $|\mathbf{E} \langle X_t, AX_t \rangle - \mathbf{E} \langle X_{t,n}, AX_{t,n} \rangle|$. The primary goal of Section 5 is to study the rate of convergence of $\mathbf{E} \langle Z_{T_j^M, n}^M, AZ_{T_j^M, n}^M \rangle$ to $\mathbf{E} \langle X_{T_j^M}, AX_{T_j^M} \rangle$, where $M \in \mathbb{N}$, $T_j^M = jT/M$, $j = 0, \dots, M$, and $Z_{\cdot, n}^M$ is the Euler approximation of (1.5), i.e., $Z_{0,n}^M = X_{0,n}$ and for any $t \in [T_j^M, T_{j+1}^M]$,

$$Z_{t,n}^M = Z_{T_j^M, n}^M + G_n Z_{T_j^M, n}^M (t - T_j^M) + \sum_{k=1}^m L_{k,n} Z_{T_j^M, n}^M (W_t^k - W_{T_j^M}^k).$$

This is the first step to approaching more complex numerical schemes, for example, those based in the computation of $X_{t,n} / \|X_{t,n}\|$. Finally, in Section 6 we apply our main results to an example of a quantum harmonic oscillator.

2. PRELIMINARIES

2.1. Notation. In the following $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. Let $p \in \mathbb{N}$ and $m = (m^j)_{j=1, \dots, p}$, $n = (n^j)_{j=1, \dots, p} \in \mathbb{Z}^p$. Then $|n| = \sum_{k=1}^p |n^k|$. In addition, we write $n \geq m$ iff

for any $j = 1, \dots, p$, $n^j \geq m^j$. Now we set $[n, m] = \{k \in \mathbb{Z}^p : k \geq n \text{ and } m \geq k\}$. Analogously, we define the partial orders $\leq, <, >$ and the boxes $]\cdot, \cdot]$, $[\cdot, \cdot[$. With each $c \in \mathbb{Z}$ is associated the vector $\vec{c}$ in $\mathbb{Z}^p$ with all coordinates equal to c .

As usual in quantum mechanics, we consider that the scalar product $\langle \cdot, \cdot \rangle$ is linear with respect to the second variable and antilinear with respect to the first one. Let B be a linear operator. We use the notation $\text{Dom}(B)$ for the domain of B , and B^* will be the adjoint of B . If Γ is any logical statement, then we consider the Boolean function

$$\mathbf{1}_\Gamma = \begin{cases} 1, & \text{if } \Gamma \text{ is true,} \\ 0, & \text{if } \Gamma \text{ is false.} \end{cases}$$

For notational clarity we shall often suppress the explicit dependence of T_j^M and $Z_{\cdot, n}^M$ on M . Moreover, the same symbol K will denote various constants in the forthcoming sections. These are always assumed to be independent of $\mathfrak{h}_n$ and M unless it is explicitly stated otherwise.

2.2. Preliminary results. Suppose that

- H1:** G is the infinitesimal generator of a strongly continuous contraction semigroup in $\mathfrak{h}$,
- H2:** $\text{Dom}(G) \subset \text{Dom}(L_k)$, whenever $k = 1, \dots, m$,
- H3:** for each $k = 1, \dots, m$, L_k is closable,
- H4:** there is a core $\mathfrak{D}^*$ for G^* such that $\mathfrak{D}^* \subset \text{Dom}(L_k^*)$, $k = 1, \dots, m$.

Hence from [10] we have that there exists a unique (a.s.) adapted stochastic process X such that

- X is weakly continuous in probability,
- $\mathbf{E} \|X_t\|^2 \leq \|X_0\|^2$,
- for all $\phi \in \mathfrak{D}^*$ and $t \in [0, T]$,

$$(2.1) \quad \langle \phi, X_t \rangle = \langle \phi, X_0 \rangle + \int_0^t \langle G^* \phi, X_s \rangle ds + \sum_{k=1}^m \int_0^t \langle L_k^* \phi, X_s \rangle dW_s^k.$$

In addition, (1.4) defines a quantum dynamical semigroup Φ_t in $B(\mathfrak{h})$, which is the minimal solution of (1.3) when $\mathfrak{D}^*$ is an invariant domain of e^{G^*t} .

2.3. Auxiliary lemmata. For completeness we state and prove the following two lemmata.

Lemma 2.1. *Let $Y \in L^2(\mathbf{P}; \mathfrak{h}) = \{f : \Omega \rightarrow \mathfrak{h}, \mathbf{E} \|f\|^2 < +\infty\}$. Suppose that there is a self-adjoint positive operator C such that $\sup_{\varepsilon > 0} \mathbf{E} \|C_\varepsilon^{1/2} Y\|^2 < +\infty$, where $C_\varepsilon = (I + \varepsilon C)^{-1} C (I + \varepsilon C)^{-1}$. Then $Y \in \text{Dom}(C^{1/2})$ a.s. and*

$$\mathbf{E} \|C^{1/2} Y\|^2 < +\infty.$$

Proof. First, by the Banach-Alaoglu theorem, there are $\varepsilon_n \rightarrow_n 0$ and $g \in L^2(\mathbf{P}; \mathfrak{h})$ such that for any $f \in L^2(\mathbf{P}; \mathfrak{h})$,

$$\mathbf{E} \langle C_{\varepsilon(n)}^{1/2} Y, f \rangle \rightarrow_n \mathbf{E} \langle g, f \rangle.$$

Next, we consider a linear operator $M_{C^{1/2}}$ in $L^2(\mathbf{P}; \mathfrak{h})$ with domain

$$\left\{ f \in L^2(\mathbf{P}; \mathfrak{h}) : \forall w \in \Omega, f(w) \in \text{Dom}(C^{1/2}) \text{ and } \mathbf{E} \|C^{1/2} f\|^2 < +\infty \right\}$$

such that $M_{C^{1/2}}f = C^{\frac{1}{2}}f$ whenever $f \in \text{Dom}(M_{C^{1/2}})$. Since $C^{1/2}$ is self-adjoint [12], from the properties of the resolvent of C it follows that

$$(2.2) \quad \mathbf{E} \langle Y, M_{C^{1/2}}f \rangle = \lim_n \mathbf{E} \left\langle C^{1/2} (I + \varepsilon_n C)^{-1} Y, f \right\rangle = \mathbf{E} \langle g, f \rangle.$$

Therefore $Y \in \text{Dom}(M_{C^{1/2}}^*)$.

Finally, the conclusion follows as in the proof of Lemma 1.1 of [8]. In fact, if $(\phi_k)_{k \geq 1}$ is an orthonormal basis of $L^2(\mathbf{P}; \mathbb{C})$, then there are $y_k, g_k \in \mathfrak{h}$, $k \geq 1$, such that $\bar{Y} = \sum_{k \geq 1} y_k \phi_k$ and $g = \sum_{k \geq 1} g_k \phi_k$. It is worth pointing out that the above equalities are understood in the $L^2(\mathbf{P}; \mathfrak{h})$ sense. From (2.2) we obtain that for any $v \in \text{Dom}(C^{1/2})$, $\langle y_k, C^{1/2}v \rangle = \langle g_k, v \rangle$. Then $g_k = (C^{1/2})^* y_k$. This implies that $C^{1/2}Y = \sum_{k \geq 1} g_k \phi_k$ a.s. and the required result follows. $\square$

Lemma 2.2. *Let α, ψ and $\phi : [a, b] \rightarrow \mathbb{R}_+$ be integrable functions. Assume that ψ is continuous. If α has a nonnegative continuous derivative and for all $t \in [a, b]$,*

$$(2.3) \quad \phi(t) \leq \alpha(t) + \int_a^t \psi(s) \sqrt{\phi(s)} ds,$$

$$\text{then } \phi(t) \leq \left(\frac{1}{2} \int_a^t \psi(s) ds + \sqrt{\alpha(t)} \right)^2.$$

Proof. By (2.3) we have that for any $\varepsilon > 0$ and $t \in [a, b]$

$$\frac{1}{2} \frac{\dot{\alpha}(t) + \psi(t) \sqrt{\phi(t)}}{\left(\varepsilon + \alpha(t) + \int_a^t \psi(s) \sqrt{\phi(s)} ds \right)^{1/2}} \leq \frac{1}{2} \frac{\dot{\alpha}(t)}{(\varepsilon + \alpha(t))^{1/2}} + \frac{\psi(t)}{2}.$$

It follows that $\sqrt{\varepsilon + \phi(t)} \leq \frac{1}{2} \int_a^t \psi(s) ds + \sqrt{\varepsilon + \alpha(t)}$ and the proof is finished taking the limit as $\varepsilon \rightarrow 0$. $\square$

3. A PRIORI ESTIMATES ON X AND $X_{\cdot, n}$

This section has two main objectives: first, to obtain uniform estimates with respect to n on the behavior of $X_{t, n}$; second, to state regularity results for X_t , where X is the stochastic process X given in subsection 2.2.

Theorem 3.1. *Suppose that assumptions H1 to H4 hold and*

H5: *for any $\phi \in \mathfrak{D}^*$, $G^* \phi = \lim_n G_n^* \phi$ and $L_k^* \phi = \lim_n L_{k, n}^* \phi$,*

H6: *$\forall n \in \mathbb{N}$, $\mathfrak{h}_n \subset \mathfrak{D}^*$,*

H7: *there is a self-adjoint positive operator C such that for each $n \geq 0$, $\mathfrak{h}_n \subset \text{Dom}(C^{1/2})$ and $\forall \phi \in \mathfrak{h}_n$,*

$$2 \text{Re} \left\langle C^{1/2} \phi, C^{1/2} G_n \phi \right\rangle + \sum_{k=1}^m \left\| C^{1/2} L_{k, n} \phi \right\|^2 \leq K \left(\left\| C^{1/2} \phi \right\|^2 + \left\| \phi \right\|^2 \right),$$

H8: *$\forall \phi \in \text{Dom}(C^{1/2})$, $\left\| C^{1/2} P_n \phi \right\| \rightarrow_{n \rightarrow +\infty} \left\| C^{1/2} \phi \right\|$.*

If $X_0 \in \text{Dom}(C^{1/2})$ a.s., then $X_t \in \text{Dom}(C^{1/2})$ a.s. and there is a constant K such that

$$(3.1) \quad \sup_{t \in [0, T]} \mathbf{E} \left\| C^{1/2} X_t \right\|^2 \leq K \left(\left\| C^{1/2} X_0 \right\|^2 + \left\| X_0 \right\|^2 \right)$$

and

$$\sup_{t \in [0, T]} \mathbf{E} \|C^{1/2} X_{t,n}\|^2 \leq K \left(\|C^{1/2} P_n X_0\|^2 + \|X_0\|^2 \right), \quad \text{for all } n \in \mathbb{N}.$$

In a wide class of applications assumptions H1 to H8 hold. In fact, these conditions are analogous to those used in [7] to derive that there is a unique σ -weak continuous family $(\Phi_t)_{t \geq 0}$ of positive maps on $B(\mathfrak{h})$ satisfying (1.3).

Notice that the first stage of the Galerkin method [15] is fulfilled in Step 1. Now, we carry out the next three stages of this procedure in the proof of Theorem 3.1. Stage 4 rests on modifications of ideas presented in [8].

Proof of Theorem 3.1. In the second stage of the Galerkin method we have to establish estimates on X_n . Here, since for any $\varphi \in \mathfrak{h}_n$, $2 \operatorname{Re} \langle \varphi, G_n \varphi \rangle + \sum_{k=1}^m \|L_{k,n} \varphi\|^2 = 0$, it follows from Itô's formula that

$$(3.2) \quad \mathbf{E} \|X_{t,n}\|^2 = \|X_{0,n}\|^2.$$

In addition, for any $\phi \in \mathfrak{D}^*$,

$$\langle \phi, X_{t,n} \rangle = \langle \phi, X_{0,n} \rangle + \int_0^t \langle G_n^* \phi, X_{s,n} \rangle ds + \sum_{k=1}^m \int_0^t \langle L_{k,n}^* \phi, X_{s,n} \rangle dW_s^k.$$

Therefore, using (3.2), H5 and the integration by parts formula [20], we obtain that for all $\phi \in \mathfrak{D}^*$ there exists a constant K_ϕ such that

$$(3.3) \quad \mathbf{E} |\langle \phi, X_{t,n} - X_{s,n} \rangle|^2 \leq K_\phi (t - s) \|X_0\|^2.$$

Finally, $C^{1/2} P_n \in B(\mathfrak{h})$ leads to

$$(3.4) \quad C^{1/2} X_{t,n} = C^{1/2} X_{0,n} + \int_0^t C^{1/2} P_n G_n X_{s,n} ds + \sum_{k=1}^m \int_0^t C^{1/2} P_n L_{k,n} X_{s,n} dW_s^k.$$

Therefore for each $t \in [0, T]$,

$$\begin{aligned} \mathbf{E} \|C^{1/2} X_{t,n}\|^2 &= \mathbf{E} \|C^{1/2} X_{0,n}\|^2 + 2\mathbf{E} \int_0^t \operatorname{Re} \langle C^{1/2} X_{s,n}, C^{1/2} G_n X_{s,n} \rangle ds \\ &\quad + \sum_{k=1}^m \mathbf{E} \int_0^t \|C^{1/2} L_{k,n} X_{s,n}\|^2 ds. \end{aligned}$$

Hence, by (3.2) and condition H7

$$\mathbf{E} \|C^{1/2} X_{t,n}\|^2 \leq \mathbf{E} \|C^{1/2} X_{0,n}\|^2 + Kt \|X_0\|^2 + \int_0^t \mathbf{E} \|C^{1/2} X_{s,n}\|^2 ds.$$

This implies

$$(3.5) \quad \mathbf{E} \|C^{1/2} X_{t,n}\|^2 \leq K \left(\|C^{1/2} X_{0,n}\|^2 + \|X_0\|^2 \right).$$

Next, we consider the third stage of the Galerkin method, where we have to study the limit as $k \rightarrow \infty$ of a certain subsequence $(X_{\cdot, n_k})_k$.

Let $\tilde{F}_t$ be the σ -algebra generated by $(W_s)_{s \in [0, t]}$ and the $\mathbf{P}$ -null sets of F . Thus, for each n there is a $(\tilde{F}_t)_{t \geq 0}$ -adapted stochastic process $\tilde{X}_{\cdot, n}$ such that $\tilde{X}_{\cdot, n}$ and

$X_{\cdot, n_k}$ are indistinguishable. Now, by (3.3) we can ensure that for all $\xi \in L^2(\mathbf{P}; \mathbb{C})$ and $\phi \in \mathfrak{D}^*$,

$$\left| \mathbf{E} \xi \left\langle \phi, \tilde{X}_{t,n} - \tilde{X}_{s,n} \right\rangle \right|^2 \leq K_\phi (t-s) \|x_0\|^2 \mathbf{E} |\xi|^2.$$

Thus $\left\{ \mathbf{E} \xi \left\langle \phi, \tilde{X}_{\cdot, n} \right\rangle \right\}_{n \geq 0}$ is a uniformly bounded and equicontinuous family of functions defined on $[0, T]$. Let $(\xi_l \phi_l)_{l \geq 1}$ be the sequence obtained by arranging all products of elements of the dense subset $\{\phi_n\}_{n \geq 1}$ of $\mathfrak{D}^*$ with elements of the dense subset $\{\xi_n\}_{n \geq 1}$ of $L^2(\mathbf{P}; \mathbb{C})$. Hence Arzela-Ascoli's theorem and diagonalization arguments show that there exist a subsequence $(\tilde{X}_{\cdot, n_k})_{k \geq 1}$ and a sequence $(f_l)_{l \geq 1}$ of continuous functions defined on $[0, T]$ such that $\mathbf{E} \xi_l \left\langle \phi_l, \tilde{X}_{\cdot, n_k} \right\rangle \rightrightarrows_k f_l$ and $\left\| \mathbf{E} \xi_l \left\langle \phi_l, \tilde{X}_{\cdot, n_k} \right\rangle - f_l \right\|_\infty < 1/l$, whenever $k \geq l$.

On the other hand, using (3.2), we obtain that for each $t \in [0, T]$ there is $\bar{X}_t \in L^2(\mathbf{P}, \tilde{\mathcal{F}}_t; \mathfrak{h})$ and a subsequence $(\tilde{X}_{t, n_{k(p)}})_{p \geq 0}$, such that

$$\tilde{X}_{t, n_{k(p)}} \xrightarrow[p \rightarrow +\infty]{\text{weak}^*} \bar{X}_t.$$

Then $f_l(t) = \mathbf{E} \xi_l \langle \phi_l, \bar{X}_t \rangle$ and

$$\mathbf{E} \|\bar{X}_t\|^2 \leq \liminf_p \mathbf{E} \|\tilde{X}_{t, n_{k(p)}}\|^2 \leq \|x_0\|^2.$$

Therefore it follows from the previous paragraph that for each $Z \in L^2(\mathbf{P}; \mathfrak{h})$, $\mathbf{E} \langle Z, \tilde{X}_{\cdot, n_k} \rangle \rightrightarrows_{k \rightarrow +\infty} \mathbf{E} \langle Z, \bar{X} \rangle$.

We are ready to proceed with Stage 4 of the Galerkin method, which concerns the properties of $\bar{X}$.

Let $\xi \in L^2(\mathbf{P}, \tilde{\mathcal{F}}_t; \mathbb{C})$. Then there exist predictable processes H^k , $k = 1, \dots, m$, in $L^2([0, t] \times \Omega; \nu \times \mathbf{P}; \mathbb{C})$ such that $\xi = \mathbf{E} \xi + \sum_{k=1}^m \int_0^t H_s^k dW_s^k$. Thus we have that for any $\phi \in \mathfrak{D}^*$ and $j = 1, \dots, m$,

$$\begin{aligned} \mathbf{E} \xi \int_0^t \left\langle L_{j, n_k}^* \phi, \tilde{X}_{s, n_k} \right\rangle dW_s^j - \mathbf{E} \xi \int_0^t \left\langle L_{j, n_k}^* \phi, \bar{X}_s \right\rangle dW_s^j \\ = \mathbf{E} \int_0^t H_s^j \left(\left\langle L_{j, n_k}^* \phi, \tilde{X}_{s, n_k} \right\rangle - \left\langle L_{j, n_k}^* \phi, \bar{X}_s \right\rangle \right) ds. \end{aligned}$$

As a consequence, taking the limit as $k \rightarrow \infty$ in

$$\begin{aligned} \mathbf{E} \xi \left\langle \phi, \tilde{X}_{t, n_k} \right\rangle &= \mathbf{E} \xi \left\langle \phi, \tilde{X}_{0, n_k} \right\rangle + \mathbf{E} \int_0^t \xi \left\langle G_{n_k}^* \phi, \tilde{X}_{s, n_k} \right\rangle ds \\ &\quad + \sum_{j=1}^m \mathbf{E} \xi \int_0^t \left\langle L_{j, n_k}^* \phi, \tilde{X}_{s, n_k} \right\rangle dW_s^j, \end{aligned}$$

we obtain

$$\mathbf{E} \xi \langle \phi, \bar{X}_t \rangle = \mathbf{E} \xi \langle \phi, \bar{X}_0 \rangle + \mathbf{E} \int_0^t \xi \langle G^* \phi, \bar{X}_s \rangle ds + \sum_{j=1}^m \mathbf{E} \xi \int_0^t \langle L_j^* \phi, \bar{X}_s \rangle dW_s^j.$$

Then $\bar{X}$ satisfies (2.1). Since $\bar{X}$ is weakly continuous in the mean-square sense, it follows from subsection 2.2 that $\bar{X} = X$ a.s.

Inspired by the proof of Theorem 2.2 of [8], we now consider the variant of the Yosida approximation $C_\varepsilon = (I + \varepsilon C)^{-1}C(I + \varepsilon C)^{-1}$. Combining (3.5) and $C_\varepsilon^{1/2} \leq C^{1/2}$, we arrive at

$$(3.6) \quad \mathbf{E} \left\| C_\varepsilon^{1/2} X_{t,n} \right\|^2 \leq K \left(\left\| C^{1/2} X_{0,n} \right\|^2 + \|X_0\|^2 \right).$$

On the other hand, for any $Z \in L^2(\mathbf{P}; \mathfrak{h})$ and each $t \in [0, T]$

$$\mathbf{E} \left\langle Z, C_\varepsilon^{1/2} X_{t,n_k} \right\rangle = \mathbf{E} \left\langle C_\varepsilon^{1/2} Z, X_{t,n_k} \right\rangle \rightarrow_k \mathbf{E} \left\langle Z, C_\varepsilon^{1/2} X_t \right\rangle.$$

Then (3.6) leads to

$$(3.7) \quad \mathbf{E} \left\| C_\varepsilon^{1/2} X_t \right\|^2 \leq \liminf_k \mathbf{E} \left\| C_\varepsilon^{1/2} X_{t,n_k} \right\|^2 \leq K \left(\left\| C^{1/2} X_0 \right\|^2 + \|X_0\|^2 \right).$$

By (3.7) and Lemma 2.1, $X_t \in \text{Dom}(C^{1/2})$ a.s. and $\mathbf{E} \|C^{1/2} X_t\|^2 < +\infty$. Finally, using the inequality $\left\| C_\varepsilon^{1/2} X_t \right\| \leq \|C^{1/2} X_t\|$ and the property

$$C_\varepsilon^{1/2} X_t = (I + \varepsilon C)^{-1} C^{1/2} X_t \xrightarrow{\varepsilon \rightarrow 0+} C^{1/2} X_t,$$

one concludes (3.1) from (3.7). $\square$

It follows from Itô's formula in a Hilbert space and Theorem 3.1 that

Corollary 3.2. *Let $\text{Dom}(C^{1/2}) \subset \text{Dom}(G)$. Then there exists a unique strong topology solution of (1.1) which is weakly continuous in probability under the assumptions of Theorem 3.1.*

Remark 3.3. A closer inspection of the proof of Theorem 3.1 reveals that the assertions of this theorem hold when $G_n = P_n G$ and for any $\varphi \in \text{Dom}(G)$,

$$2 \text{Re} \langle \varphi, G\varphi \rangle + \sum_{k=1}^m \|L_k \varphi\|^2 \leq 0.$$

Remark 3.4. Employing similar arguments to those used in the proof of Theorem 3.1, we can generalize the results given in Theorem 3.1 and Remark 3.3 to the case $m = +\infty$ under the hypothesis that for any $\phi \in \mathfrak{D}^*$, $\sum_{j=1}^\infty \|L_j^* \phi\|^2 < +\infty$.

4. RATE OF CONVERGENCE OF FINITE DIMENSIONAL APPROXIMATIONS

This section deals with the last stage of the Galerkin method, i.e., with strong convergence results. More precisely, we derive the rate of convergence of $\mathbf{E} \langle X_{t,n}, AX_{t,n} \rangle$ to $\mathbf{E} \langle X_t, AX_t \rangle$, where A belongs to a certain class of linear operators.

First, let us note that A is an unbounded operator in many physics applications [19, 14]. Therefore we do not restrict ourselves here to the case $A \in B(\mathfrak{h})$. We also point out that the basic idea behind the proof of the main results of this part is the application of Theorem 3.1. Roughly speaking, we will combine the constraint given by the domain of $C^{\frac{1}{2}}$ with estimates introduced in Theorem 3.1.

To be concrete, from now on we will suppose that

S1: There are an orthonormal basis of $\mathfrak{h}$ $\{\varphi_j\}_{j \in \mathbf{Z}_+^p}$, $p \in \mathbb{N}$, and a positive real number l_C such that the operator $C\varphi_j = |j|^{l_C} \varphi_j$ satisfies condition H7.

In addition to the fact that assumption S1 frequently holds in applications in quantum optics (see, e.g., [6, 7] and Section 6), this case allows us to introduce a general procedure which can be useful in other situations. Notice that p is interpreted as the number of modes or degree of freedom of physical systems.

For simplicity, we consider that $\mathfrak{h}_n$ is the linear span of $\{\varphi_j : \vec{0} \leq j \leq \vec{n}\}$. That is, we reduce ourselves to the Galerkin approximation with fixed basis. To state our next results precisely, we introduce the following notation.

Definition 4.1. Let a and b be nonnegative integer numbers. Assume that l is a nonnegative real number. Then $O_{l,a,b}$ denotes the set of all linear operators $B : \text{Dom}(B) \subset \mathfrak{h} \rightarrow \mathfrak{h}$ such that for any $k \in \mathbb{Z}_+^p$, $B\varphi_k$ belongs to the linear span of $\{\varphi_j\}_{j \in [k-a, k+b]}$ and $\|B\varphi_k\| \leq K \max\{|k|^l, 1\}$.

We are now ready to fulfill our objective of characterizing the convergence properties corresponding to Step 1 explained in Section 1.

Theorem 4.2. Let $A \in O_{l_A, n_A, m_A}$, $G \in O_{l_G, n_G, m_G}$ and $L_j \in O_{l_L, n_L, m_L}$, $j = 1, \dots, m$. Suppose that $X_0 \in \text{Dom}(C^{1/2}) \subset \text{Dom}(G)$ and $l_C \geq 2 \max\{l_A, l_G, 2l_L\}$. Then under assumptions S1 and H1 to H6, we have that for any $n > 0$,

$$\sup_{t \in [0, T]} |\mathbf{E} \langle X_t, AX_t \rangle - \mathbf{E} \langle X_{t,n}, AX_{t,n} \rangle| \leq K \left(\|C^{1/2} X_0\|^2 + \|X_0\|^2 \right) \\ * \left(\frac{1}{n^{l_C - l_A}} + \frac{\mathbf{1}_{n_G > 0}}{n^{l_C/2 - l_G}} + \frac{\mathbf{1}_{n_L > 0} + \mathbf{1}_{m_L > 0}}{n^{l_C/2 - 2l_L}} + \frac{\mathbf{1}_{n_G > 0}}{n^{l_C - 2l_G - l_A}} + \frac{\mathbf{1}_{n_L > 0} + \mathbf{1}_{m_L > 0}}{n^{l_C - 4l_L - l_A}} \right).$$

For expository purposes, before proving Theorem 4.2, we give Lemma 4.3.

Lemma 4.3. Let $\Phi \in O_{l,a,b}$. Then $\Phi^* \in O_{l,b,a}$. Furthermore, if $\bar{P}_k$ is the orthogonal projection of $\mathfrak{h}$ over the linear span of $\{\varphi_j\}_{j \in [\vec{0}, k]}$, then $\|\Phi \bar{P}_k\| \leq K |k|^l$ whenever $k \neq \vec{0}$.

Proof. Since $j \in [k-a, k+b]$ iff $k \in [j-b, j+a]$, $\Phi^* \varphi_k$ belongs to the linear span of $\{\varphi_j\}_{j \in [k-b, k+a]}$. Let $k > 0$. By the Cauchy-Schwarz inequality,

$$\|\Phi \bar{P}_k z\|^2 \leq K |k|^{2l} \left(\sum_{i \in [\vec{0}, k]} |z^i|^2 \right)^{\frac{1}{2}} \left(\sum_{i \in [\vec{0}, k]} \sum_{j \in [i-a-b, i+a+b] \cap [\vec{0}, k]} |z^j|^2 \right)^{\frac{1}{2}},$$

where $z \in \mathfrak{h}$. Thus $\|\Phi \bar{P}_k z\|^2 \leq K |k|^{2l} \|z\|^2$. Finally, if S is the linear span of $\{\varphi_j\}_{j \in [0, k+a]}$, then

$$\|\Phi^* \varphi_k\| = \sup_{\phi \in S, \phi \neq 0} \frac{\langle \Phi^* \varphi_k, \phi \rangle}{\|\phi\|} \leq \|\varphi_k\| \|\Phi \bar{P}_{k+a}\|.$$

This implies that $\Phi^* \in O_{l,b,a}$. □

Proof of Theorem 4.2. One easily obtains that we can reduce the study of the truncation error

$$|\mathbf{E} \langle X_t, AX_t \rangle - \mathbf{E} \langle X_{t,n}, AX_{t,n} \rangle|$$

to the analysis of the errors $H_t^1 = |\mathbf{E} \langle X_t, AX_t \rangle - \mathbf{E} \langle P_n X_t, AP_n X_t \rangle|$ and $H_t^2 = |\mathbf{E} \langle P_n X_t, AP_n X_t \rangle - \mathbf{E} \langle X_{t,n}, AX_{t,n} \rangle|$. Then we will concentrate on estimating H_t^1 and H_t^2 .

Next, we will check that

$$(4.1) \quad H_t^1 \leq K n^{l_A - l_C} \left(\|C^{1/2} X_0\|^2 + \|X_0\|^2 \right).$$

To verify (4.1), we decompose H_t^1 as follows:

$$(4.2) \quad H_t^1 \leq |\mathbf{E} \langle P_n X_t, A(X_t - P_n X_t) \rangle| + |\mathbf{E} \langle X_t - P_n X_t, A P_n X_t \rangle| \\ + |\mathbf{E} \langle X_t - P_n X_t, A(X_t - P_n X_t) \rangle|.$$

Now, by $A \in O_{l_A, n_A, m_A}$,

$$|\langle X_t - P_n X_t, A(X_t - P_n X_t) \rangle| \\ \leq K \sum_{j \notin [\vec{0}, \vec{n}]} \left(|X_t^j| |j|^{l_A} \sum_{k \in [j - n_A, j + m_A] \cap [\vec{0}, \vec{n}]^c} |X_t^k| \right).$$

Hence, using the Cauchy-Schwarz inequality and the fact that $k \in [j - n_A, j + m_A]$ iff $j \in [k - m_A, k + n_A]$, one concludes that

$$|\langle X_t - P_n X_t, A(X_t - P_n X_t) \rangle| \\ \leq K \sqrt{\sum_{j \notin [\vec{0}, \vec{n}]} |X_t^j|^2 |j|^{l_A}} \sqrt{\sum_{j \notin [\vec{0}, \vec{n}]} \sum_{k \in [j - n_A, j + m_A] \cap [\vec{0}, \vec{n}]^c} |X_t^k|^2 (|k| + p n_A)^{l_A}} \\ \leq K \sum_{j \notin [\vec{0}, \vec{n}]} |X_t^j|^2 |j|^{l_A}.$$

Since $l_C \geq l_A$,

$$(4.3) \quad |\langle X_t - P_n X_t, A(X_t - P_n X_t) \rangle| \leq K n^{l_A - l_C} \|C^{1/2} X_t\|^2.$$

Repeating the arguments given above, we obtain

$$(4.4) \quad |\langle X_t - P_n X_t, A P_n X_t \rangle| \\ \leq \sum_{k \notin [\vec{0}, \vec{n}]} \sum_{j \in [\vec{0}, \vec{n}]} |X_t^j| |X_t^k| |\langle \varphi_k, A \varphi_j \rangle| \\ \leq K \sqrt{\sum_{k \notin [\vec{0}, \vec{n}]} |X_t^k|^2} \sqrt{\sum_{j \in [0, \vec{n}], |j| > n - m_{Ap}} |X_t^j|^2 \max \{|j|^{2l_A}, 1\}}.$$

It follows from $l_C \geq 2l_A$ that for every $n > m_{Ap}$,

$$(4.5) \quad |\langle X_t - P_n X_t, A P_n X_t \rangle| \leq K n^{l_A - l_C} \|C^{1/2} X_t\|^2.$$

Furthermore, (4.4) implies that for any $n \leq m_{Ap}$,

$$(4.6) \quad |\langle X_t - P_n X_t, A P_n X_t \rangle| \leq K n^{l_A - l_C/2} \|C^{1/2} X_t\| \|X_t\| \\ \leq K n^{l_A - l_C} \|C^{1/2} X_t\| \|X_t\|.$$

Finally, since $\langle P_n X_t, A(X_t - P_n X_t) \rangle = \langle A^* P_n X_t, X_t - P_n X_t \rangle$, combining Lemma 4.3, inequalities (4.5), (4.6), (4.3) and Theorem 3.1, we arrive at (4.1).

Now, we go on estimating H_t^2 . To treat this term, we first note that

$$(4.7) \quad H_t^2 \leq \left(\left(\mathbf{E} \|A^* X_{t,n}\|^2 \right)^{\frac{1}{2}} + \left(\mathbf{E} \|AX_{t,n}\|^2 \right)^{\frac{1}{2}} \right) \left(\mathbf{E} \|P_n X_t - X_{t,n}\|^2 \right)^{\frac{1}{2}} \\ + \|AP_n\| \mathbf{E} \|P_n X_t - X_{t,n}\|^2.$$

Since

$$(4.8) \quad \|A^* X_{t,n}\|^2 \leq K \left(\sum_{j \in [\vec{0}, \vec{n}]} |X_{t,n}^j|^2 \max \{ |j|^{l_A}, 1 \}^2 \right)^{\frac{1}{2}} \\ * \left(\sum_{j \in [\vec{0}, \vec{n}]} \sum_{k \in [j - n_A - m_A, j + n_A + m_A] \cap [\vec{0}, \vec{n}]} |X_{t,n}^k|^2 \max \{ |k|^{l_A}, 1 \}^2 \right)^{\frac{1}{2}}, \\ \|A^* X_{t,n}\|^2 \leq K \sum_{j \in [\vec{0}, \vec{n}]} |X_{t,n}^j|^2 \max \{ |j|^{l_A}, 1 \}^2 \leq K \left(\|C^{\frac{1}{2}} X_{t,n}\|^2 + |X_{t,n}^{\vec{0}}|^2 \right).$$

Therefore, Lemma 4.3 leads to

$$(4.9) \quad \|AX_{t,n}\|^2 \leq K \left(\|C^{\frac{1}{2}} X_{t,n}\|^2 + |X_{t,n}^{\vec{0}}|^2 \right).$$

Using Theorem 3.1, Lemma 4.3, (4.8) and (4.9) in (4.7), it follows that

$$(4.10) \quad H_t^2 \leq K \sqrt{\|C^{\frac{1}{2}} P_n X_0\|^2 + \|X_0\|^2} \sqrt{\mathbf{E} \|P_n X_t - X_{t,n}\|^2} \\ + K n^{l_A} \mathbf{E} \|P_n X_t - X_{t,n}\|^2.$$

Consequently, it remains to bound $h(t) = \mathbf{E} \|P_n X_t - X_{t,n}\|^2$. Applying Corollary 3.2, one concludes that

$$P_n X_t - X_{t,n} = \int_0^t (P_n G X_s - G_n X_{s,n}) ds + \sum_{k=1}^m \int_0^t L_{k,n} (X_s - X_{s,n}) dW_s^k.$$

Therefore, Itô's formula yields

$$h(t) = 2\mathbf{E} \int_0^t \operatorname{Re} \langle P_n X_s - X_{s,n}, P_n G X_s - G_n X_{s,n} \rangle ds \\ + \sum_{k=1}^m \mathbf{E} \int_0^t \|L_{k,n} (X_s - X_{s,n})\|^2 ds.$$

Thus, we deduce from (1.6) that $\mathbf{E} \|P_n X_t - X_{t,n}\|^2 = \sum_{k=1}^4 J_t^k$, where

$$J_t^1 = 2\mathbf{E} \int_0^t \operatorname{Re} \langle P_n X_s - X_{s,n}, (P_n G - G_n) P_n X_s \rangle ds, \\ J_t^2 = 2\mathbf{E} \int_0^t \operatorname{Re} \langle P_n X_s - X_{s,n}, P_n G (X_s - P_n X_s) \rangle ds, \\ J_t^3 = \sum_{k=1}^m \mathbf{E} \int_0^t \|L_{k,n} (X_s - P_n X_s)\|^2 ds,$$

and $J_t^4 = 2 \sum_{k=1}^m 2\mathbf{E} \int_0^t \operatorname{Re} \langle L_{k,n}(X_s - P_n X_s), L_{k,n}(P_n X_s - X_{s,n}) \rangle ds$.

Fix $m_L = 0$. Then $J_t^1 = 0$, because $(P_n G - G_n) P_n X_s = 0$. In the other case, i.e., $m_L \geq 1$, combining Lemma 4.3 and Theorem 3.1, we obtain

$$\begin{aligned} & \mathbf{E} \|(P_n G - G_n) P_n X_s\|^2 \\ & \leq \frac{m}{4} \sum_{k=1}^m \mathbf{E} \|(P_n L_k^* P_n L_k - P_n L_k^* L_k)(P_n X_s - P_{n-m_L} X_s)\|^2 \\ & \leq K n^{4l_L} \mathbf{E} \|P_n X_s - P_{n-m_L} X_s\|^2 \\ & \leq K n^{4l_L - l_C} \left(\|C^{1/2} X_0\|^2 + \|X_0\|^2 \right), \end{aligned}$$

which yields

$$J_t^1 \leq K \mathbf{1}_{m_L > 0} n^{2l_L - \frac{l_C}{2}} \left(\|C^{\frac{1}{2}} X_0\|^2 + \|X_0\|^2 \right)^{\frac{1}{2}} \int_0^t \mathbf{E}^{\frac{1}{2}} (\|P_n X_s - X_{s,n}\|^2) ds.$$

It is worth mentioning that in previous inequalities if $n - m_L < 0$, then we put $P_{n-m_L} = 0$.

Clearly, if $n_G = 0$, resp. $n_L = 0$, then $J_t^2 = 0$, resp. $J_t^3 = 0$. Moreover, from (4.3), Lemma 4.3,

$$\|P_n G(X_s - P_n X_s)\|^2 = \langle X_s - P_n X_s, G^* P_n G(X_s - P_n X_s) \rangle$$

and $\|L_{k,n}(X_s - P_n X_s)\|^2 = \langle X_s - P_n X_s, L_k^* P_n L_k(X_s - P_n X_s) \rangle$, it follows that

$$\begin{aligned} J_t^2 + J_t^3 & \leq \mathbf{1}_{n_L > 0} K n^{2l_L - l_C} \left(\|C^{\frac{1}{2}} X_0\|^2 + \|X_0\|^2 \right) \\ & + \mathbf{1}_{n_G > 0} K n^{l_G - l_C/2} \left(\|C^{\frac{1}{2}} X_0\|^2 + \|X_0\|^2 \right)^{\frac{1}{2}} \int_0^t \mathbf{E}^{\frac{1}{2}} (\|P_n X_s - X_{s,n}\|^2) ds. \end{aligned}$$

Since

$$\begin{aligned} & \langle L_{k,n}(X_s - P_n X_s), L_{k,n}(P_n X_s - X_{s,n}) \rangle \\ & = \langle L_k^* P_n L_k(X_s - P_n X_s), P_n X_s - X_{s,n} \rangle, \end{aligned}$$

we also have

$$J_t^4 \leq \mathbf{1}_{n_L > 0} K n^{2l_L - l_C/2} \left(\|C^{\frac{1}{2}} X_0\|^2 + \|X_0\|^2 \right)^{\frac{1}{2}} \int_0^t \mathbf{E}^{\frac{1}{2}} (\|P_n X_s - X_{s,n}\|^2) ds.$$

To summarize, we have obtained that

$$\begin{aligned} h(t) & \leq K \frac{\mathbf{1}_{n_L > 0}}{n^{l_C - 2l_L}} \left(\|C^{\frac{1}{2}} X_0\|^2 + \|X_0\|^2 \right) \\ & + K \left(\|C^{\frac{1}{2}} X_0\|^2 + \|X_0\|^2 \right)^{\frac{1}{2}} * \left(\frac{\mathbf{1}_{n_L > 0} + \mathbf{1}_{m_L > 0}}{n^{l_C/2 - 2l_L}} + \frac{\mathbf{1}_{n_G > 0}}{n^{l_C/2 - l_G}} \right) \int_0^t \sqrt{h(s)} ds. \end{aligned}$$

Therefore, according to Lemma 2.2

$$(4.11) \quad h(t) \leq K \left(\|C^{\frac{1}{2}} X_0\|^2 + \|X_0\|^2 \right) \left(\frac{\mathbf{1}_{n_L > 0} + \mathbf{1}_{m_L > 0}}{n^{l_C/2 - 2l_L}} + \frac{\mathbf{1}_{n_G > 0}}{n^{l_C/2 - l_G}} \right)^2.$$

Then the required result follows from (4.10). $\square$

To handle the case $A \in B(\mathfrak{h}) \setminus \bigcup_{n_A, m_A=0}^\infty O_{0, n_A, m_A}$, we present the following theorem.

Theorem 4.4. *Let $A \in \mathcal{A} \in B(\mathfrak{h})$, $G \in \mathcal{O}_{l_G, n_G, m_G}$ and $L_j \in \mathcal{O}_{l_L, n_L, m_L}$, $j = 1, \dots, m$. Suppose that $X_0 \in \text{Dom}(C^{1/2}) \subset \text{Dom}(G)$ and $l_C \geq 2 \max\{l_G, 2l_L\}$. Then under assumptions S1 and H1–H6, we have that for any $n > 0$,*

$$\begin{aligned} & \sup_{t \in [0, T]} |\mathbf{E} \langle X_t, AX_t \rangle - \mathbf{E} \langle X_{t,n}, AX_{t,n} \rangle| \\ & \leq K \left(\|C^{1/2} X_0\|^2 + \|X_0\|^2 \right) \left(\frac{1}{n^{l_C/2}} + \frac{\mathbf{1}_{n_G > 0}}{n^{l_C/2 - l_G}} + \frac{\mathbf{1}_{n_L > 0} + \mathbf{1}_{m_L > 0}}{n^{l_C/2 - 2l_L}} \right). \end{aligned}$$

Proof. Let H_t^1 and H_t^2 be defined as in the proof of Theorem 4.2. Hence

$$H_t^1 \leq \|A\| \mathbf{E} \left(2 \|P_n X_t\| \|X_t - P_n X_t\| + \|X_t - P_n X_t\|^2 \right)$$

and

$$\begin{aligned} H_t^2 & \leq \|A\| \left(\left(\mathbf{E} \|X_{t,n}\|^2 \right)^{\frac{1}{2}} + \left(\mathbf{E} \|X_{t,n}\|^2 \right)^{\frac{1}{2}} \right) \left(\mathbf{E} \|P_n X_t - X_{t,n}\|^2 \right)^{\frac{1}{2}} \\ & \quad + \|A\| \mathbf{E} \|P_n X_t - X_{t,n}\|^2. \end{aligned}$$

Therefore, applying (4.3) and (4.11), the theorem follows. $\square$

5. RATE OF CONVERGENCE OF THE EXPLICIT EULER SCHEME

In this section our aim is to progress in the theoretical understanding of Step 2 of the procedure given in Section 1. Indeed, we will address the numerical simulation of $\mathbf{E} \langle X_{t,n}, AX_{t,n} \rangle$ using the explicit Euler scheme, i.e., the scheme $Z_{\cdot, n}$ recalled in Section 1.

Because of its instability, the explicit Euler scheme Z does not present a good performance when (1.5) is a stiff SDE [16, 13]. Though this situation often appears in quantum mechanics, the error analysis presented here can be a useful pattern for studying more complex numerical schemes, such as implicit Euler methods [16, 13, 26], Euler-Exponential methods [17] and those resulting of the application of the above schemes to the solution of the SDE satisfied by $X_{t,n}/\|X_{t,n}\|$.

We start the section with Theorem 5.1 which provides the rate of convergence of the explicit Euler scheme. Next, we focus on a numerical method based on the extrapolation of this scheme.

Theorem 5.1. *Suppose that μ is a positive constant and conditions of Theorem 4.2 hold. Let $(c_n)_n$ be a sequence of real numbers such that*

S2: *for any $n \geq 1$ and $x \in \mathfrak{h}_n$, $\|G_n P_n\|^2 \leq c_n$ and*

$$\|C^{1/2} G_n x\|^2 \leq K c_n \left(\|C^{1/2} x\|^2 + \|x\|^2 \right).$$

Moreover, assume that

S3: *the operator given by $C_1 \varphi_n = |n|^{l_{C_1}} \varphi_n$, where l_{C_1} is a positive real number such that $l_C \geq l_{C_1} + 4 \max\{l_G, l_L\}$ and $l_{C_1} \geq l_A$, fulfills condition H7.*

If $c_n \leq \mu M$, then for any $j = 0, \dots, M$,

$$\begin{aligned} & \left| \mathbf{E} \langle X_{T_j}, AX_{T_j} \rangle - \mathbf{E} \langle Z_{T_j,n}^M, AZ_{T_j,n}^M \rangle \right| \\ & \leq K \left(\left\| C^{1/2} X_0 \right\|^2 + \|X_0\|^2 \right) \\ & \quad * \left(\frac{T}{M} + \frac{\mathbf{1}_{n_G > 0}}{n^{l_C/2 - l_G}} \right. \\ & \quad \left. + \frac{\mathbf{1}_{n_L > 0} + \mathbf{1}_{m_L > 0}}{n^{l_C/2 - 2l_L}} + \frac{\mathbf{1}_{n_G > 0}}{n^{l_C - 2l_G - l_A}} + \frac{\mathbf{1}_{n_L > 0} + \mathbf{1}_{m_L > 0}}{n^{l_C - 4l_L - l_A}} + \frac{1}{n^{l_C - l_A}} \right). \end{aligned}$$

As we will illustrate in Section 6, the hypotheses of Theorem 5.1 are general enough for applications in quantum optics. The underlying idea in the proof of Theorem 5.1 is to refine standard arguments employed to prove the rate of convergence of classical weak numerical schemes for finite dimensional SDEs [23, 24, 16, 13] keeping in mind the characteristics of our case, e.g., Remark 5.2 and Lemma 5.3.

Remark 5.2. One specific feature which is used in the proof is that for each $x \in \mathfrak{h}_n$, $t \in [0, T]$ and $s \in [0, t]$,

$$(5.1) \quad \langle x, \Phi_{t,n}^s x \rangle = \mathbf{E} \langle Y_{t,n}^{s,x}, AY_{t,n}^{s,x} \rangle,$$

where for every $t \in [s, T]$

$$(5.2) \quad Y_{t,n}^{s,x} = x + \int_s^t G_n Y_{u,n}^{s,x} du + \sum_{k=1}^m \int_s^t L_{k,n} Y_{u,n}^{s,x} dW_u^k$$

and

$$(5.3) \quad \Phi_{t,n}^s = P_n A P_n + \int_s^t \mathcal{L}_n(\Phi_{u,n}^s) du$$

with $\mathcal{L}_n : B(\mathfrak{h}_n) \rightarrow B(\mathfrak{h}_n)$ given by

$$\mathcal{L}_n(X) = P_n G_n^* X + X G_n P_n + \sum_{k=1}^m P_n L_k^* P_n X P_n L_k P_n.$$

For the sake of simplicity, we use the shorthand notation $\Phi_{t-s,n}$ to mean $\Phi_{t-s,n}^0$.

Lemma 5.3. *Suppose that the hypotheses of Theorem 5.1 hold. Consider $B_1, B_2 \in O_{l_B, n_B, m_B}$ satisfying $l_C \geq l_{C_1} + 2l_B$. Then*

$$(5.4) \quad \left| \mathbf{E} \langle B_1 Z_{T_j,n}, \Phi_{T_\kappa - s,n} B_2 Z_{T_j,n} \rangle \right| \leq K \left(\left\| C^{\frac{1}{2}} X_0 \right\|^2 + \|X_0\|^2 \right)$$

whenever $T_j \leq s \leq T_\kappa$.

Proof. First, we establish estimates on $Z_{t,n}$. Next, using them, we obtain (5.4).

By Itô's formula, we have that for every $t \in]T_j, T_{j+1}]$

$$\begin{aligned} & \mathbf{E} \|Z_{t,n}\|^2 \\ & = \mathbf{E} \|Z_{T_j,n}\|^2 + \mathbf{E} \int_{T_j}^t 2 \operatorname{Re} \langle Z_{s,n}, G_n Z_{T_j,n} \rangle ds + \sum_{k=1}^m \mathbf{E} \int_{T_j}^t \|L_{k,n} Z_{T_j,n}\|^2 ds \\ & = \mathbf{E} \|Z_{T_j,n}\|^2 + \mathbf{E} \int_{T_j}^t 2 \operatorname{Re} \langle G_n Z_{T_j,n}(s - T_j), G_n Z_{T_j,n} \rangle ds. \end{aligned}$$

Hence $\mathbf{E} \|Z_{t,n}\|^2 \leq \mathbf{E} \|Z_{T_j,n}\|^2 \left(1 + \|G_n P_n\|^2 \frac{(t-T_j)^2}{2}\right)$. Therefore

$$(5.5) \quad \mathbf{E} \|Z_{t,n}\|^2 \leq \|Z_{0,n}\|^2 \left(1 + c_n \frac{T^2}{2M^2}\right)^j \leq \|Z_{0,n}\|^2 e^{\frac{c_n T^2}{2M}} \leq \|Z_{0,n}\|^2 e^{\frac{4}{2} T^2}.$$

On the other hand, for every $t \in]T_j, T_{j+1}]$

$$C^{\frac{1}{2}} Z_{t,n} = C^{\frac{1}{2}} Z_{T_j,n} + \int_{T_j}^t C^{\frac{1}{2}} G_n Z_{T_j,n} ds + \sum_{k=1}^m \int_{T_j}^t C^{\frac{1}{2}} L_{k,n} Z_{T_j,n} dW_s^k.$$

Therefore, from Itô's formula it follows that

$$\begin{aligned} \mathbf{E} \|C^{\frac{1}{2}} Z_{t,n}\|^2 &= \mathbf{E} \|C^{\frac{1}{2}} Z_{T_j,n}\|^2 + \mathbf{E} \int_{T_j}^t 2 \operatorname{Re} \left\langle C^{\frac{1}{2}} Z_{s,n}, C^{\frac{1}{2}} G_n Z_{T_j,n} \right\rangle ds \\ &\quad + \sum_{k=1}^m \mathbf{E} \int_{T_j}^t \|C^{\frac{1}{2}} L_{k,n} Z_{T_j,n}\|^2 ds. \end{aligned}$$

Thus (5.5) and hypothesis S1 yield

$$\begin{aligned} \mathbf{E} \|C^{\frac{1}{2}} Z_{t,n}\|^2 &\leq \left(1 + \frac{KT}{M}\right) \mathbf{E} \|C^{\frac{1}{2}} Z_{T_j,n}\|^2 + \frac{KT}{M} \|Z_{0,n}\|^2 \\ &\quad + \mathbf{E} \int_{T_j}^t 2 \operatorname{Re} \left\langle C^{\frac{1}{2}} (Z_{s,n} - Z_{T_j,n}), C^{\frac{1}{2}} G_n Z_{T_j,n} \right\rangle ds. \end{aligned}$$

Hence combining S2, the linearity of (1.5) and the fact that $c_n \leq \mu M$, we obtain

$$\mathbf{E} \|C^{\frac{1}{2}} Z_{t,n}\|^2 \leq \left(1 + \frac{KT}{M}\right) \mathbf{E} \|C^{\frac{1}{2}} Z_{T_j,n}\|^2 + \frac{KT}{M} \|Z_{0,n}\|^2.$$

Then for any $t \in]T_j, T_{j+1}]$,

$$\begin{aligned} \mathbf{E} \|C^{\frac{1}{2}} Z_{t,n}\|^2 &\leq \left(1 + \frac{KT}{M}\right)^j \mathbf{E} \|C^{\frac{1}{2}} Z_{0,n}\|^2 + \sum_{k=0}^{j-1} \left(1 + \frac{KT}{M}\right)^k \frac{KT}{M} \|Z_{0,n}\|^2 \\ &\leq \mathbf{E} \|C^{\frac{1}{2}} Z_{0,n}\|^2 e^K + K \|Z_{0,n}\|^2 \left(\left(1 + \frac{K}{M}\right)^M - 1 \right) \\ &\leq K \mathbf{E} \|C^{\frac{1}{2}} Z_{0,n}\|^2 + K \|Z_{0,n}\|^2 (e^K - 1). \end{aligned}$$

In the remainder of the proof we verify (5.4). Using the polarization identity, (5.1) and the property $\Phi_{t-s,n} = \Phi_{t,n}^s$, we obtain

$$\begin{aligned} &4 \left| \mathbf{E} \left\langle B_1 Z_{T_j,n}, \Phi_{T_\kappa-s,n} B_2 Z_{T_j,n} \right\rangle \right| \\ &= \left| \sum_{k=0}^3 i^{-k} \mathbf{E} \left\langle (B_1 + i^k B_2) Z_{T_j,n}, \Phi_{T_\kappa-s,n} (B_1 + i^k B_2) Z_{T_j,n} \right\rangle \right| \\ &\leq \sum_{k=0}^3 \left| \mathbf{E} \left\langle \tilde{Y}_{n,k}, A \tilde{Y}_{n,k} \right\rangle \right|, \end{aligned}$$

with $\tilde{Y}_{n,k} = Y_{T_\kappa, n}^{s, (B_1 + i^k B_2) Z_{T_j, n}}$. Proceeding as in the proof of Theorem 4.2, one arrives at

$$\begin{aligned} & \left| \mathbf{E} \left(\langle \tilde{Y}_{n,k}, A\tilde{Y}_{n,k} \rangle / F_s \right) \right| \\ & \leq K \mathbf{E} \left(\sum_{l \in [\vec{0}, \vec{n}]} \left| \tilde{Y}_{n,k}^l \right|^2 \max \{ |l|^{l_A}, 1 \} / F_{T_s} \right) \\ & \leq K \mathbf{E} \left(\left\| C_1^{\frac{1}{2}} \tilde{Y}_{n,k} \right\|^2 + \left\| \tilde{Y}_{n,k}^{\vec{0}} \right\|^2 / F_{T_s} \right). \end{aligned}$$

It follows from Theorem 3.1 and the Markov property of Y that

$$\left| \mathbf{E} \left(\langle \tilde{Y}_{n,k}, A\tilde{Y}_{n,k} \rangle / F_s \right) \right| \leq K \left(\left\| C_1^{\frac{1}{2}} (B_1 + i^k B_2) Z_{T_j, n} \right\|^2 + \left\| (B_1 + i^k B_2) Z_{T_j, n} \right\|^2 \right).$$

Since $l_C \geq l_{C_1} + 2l_B$,

$$\left| \mathbf{E} \left(\langle \tilde{Y}_{n,k}, A\tilde{Y}_{n,k} \rangle / F_s \right) \right| \leq K \left(\left\| C^{\frac{1}{2}} Z_{T_j, n} \right\|^2 + \left\| Z_{T_j, n} \right\|^2 \right).$$

Then we conclude the proof of (5.4) by an application of the estimates obtained in the first part. $\square$

Proof of Theorem 5.1. The error $\left| \mathbf{E} \langle X_{T_j}, AX_{T_j} \rangle - \mathbf{E} \langle Z_{T_j, n}^M, AZ_{T_j, n}^M \rangle \right|$ can be decomposed into the sum of the error $\left| \mathbf{E} \langle X_{T_j}, AX_{T_j} \rangle - \mathbf{E} \langle X_{T_j, n}, AX_{T_j, n} \rangle \right|$, which arises in Step 1, and the discretization error

$$\left| \mathbf{E} \langle X_{T_j, n}, AX_{T_j, n} \rangle - \mathbf{E} \langle Z_{T_j, n}^M, AZ_{T_j, n}^M \rangle \right|.$$

Theorem 4.2 provides estimates for the first term, so that it remains to bound the discretization error.

To this end, we introduce the functions $u_n : [0, T_j] \times \mathfrak{h}_n \times \mathfrak{h}_n \rightarrow \mathbb{C}$, $n \in \mathbb{N}$, defined by

$$u_n(s, x, y) = \langle \bar{x}, \Phi_{T_j, n}^s y \rangle,$$

where $\bar{x} = \sum_{j \in [\vec{0}, \vec{n}]} \bar{x}_j \varphi_j$ whenever $x = \sum_{j \in [\vec{0}, \vec{n}]} x_j \varphi_j$. Recall that $\{\varphi_j : \vec{0} \leq j \leq \vec{n}\}$ is an orthogonal basis of $\mathfrak{h}_n$. It follows from (5.1) that

$$\begin{aligned} & \mathbf{E} \langle Z_{T_j, n}, AZ_{T_j, n} \rangle - \mathbf{E} \langle X_{T_j, n}, AX_{T_j, n} \rangle \\ & = \mathbf{E} (u_n(T_j, \overline{Z_{T_j, n}}, Z_{T_j, n}) - u_n(0, \overline{X_{0, n}}, X_{0, n})) \\ & = \sum_{k=0}^{j-1} \delta_k, \end{aligned}$$

where $\delta_k = \mathbf{E} (u_n(T_{k+1}, \overline{Z_{T_{k+1}, n}}, Z_{T_{k+1}, n}) - u_n(T_k, \overline{Z_{T_k, n}}, Z_{T_k, n}))$.

Now, Itô's formula and (5.3) yield

$$\begin{aligned}\delta_k &= \mathbf{E} \int_{T_k}^{T_{k+1}} \langle G_n(Z_{T_k,n} - Z_{s,n}), \Phi_{T_j-s,n} Z_{s,n} \rangle ds \\ &\quad + \mathbf{E} \int_{T_k}^{T_{k+1}} \langle Z_{s,n}, \Phi_{T_j-s,n} G_n(Z_{T_k,n} - Z_{s,n}) \rangle ds \\ &\quad + \sum_{l=1}^m \mathbf{E} \int_{T_k}^{T_{k+1}} \langle L_{l,n} Z_{T_k,n}, \Phi_{T_j-s,n} L_{l,n} Z_{T_k,n} \rangle ds \\ &\quad - \sum_{l=1}^m \mathbf{E} \int_{T_k}^{T_{k+1}} \langle L_{l,n} Z_{s,n}, \Phi_{T_j-s,n} L_{l,n} Z_{s,n} \rangle ds.\end{aligned}$$

Therefore, a simple calculation leads to

$$\delta_k = - \sum_{(B_1, B_2, r) \in \mathcal{S}} \int_{T_k}^{T_{k+1}} \mathbf{E} \langle B_1 Z_{T_k,n}, \Phi_{T_j-s,n} B_2 Z_{T_k,n} \rangle (s - T_k)^r ds,$$

where $\mathcal{S}$ is the set formed by the following triplets: $(G_n^2, I, 1)$, $(I, G_n^2, 1)$, $(G_n^2, G_n, 2)$, $(G_n, G_n^2, 2)$, $(G_n L_{l,n}, L_{l,n}, 1)$, $(L_{l,n}, G_n L_{l,n}, 1)$, $(L_{l,n}, L_{l,n} G_n, 1)$, $(L_{l,n} G_n, L_{l,n}, 1)$, $(L_{l,n} G_n, L_{l,n} G_n, 2)$ and $(L_{l,n} L_{l_1,n}, L_{l,n} L_{l_1,n}, 1)$, with $l, l_1 = 1, \dots, m$. Thus Lemma 5.3 completes the proof. $\square$

To improve the weak convergence order of the Euler scheme, Talay and Tubaro generalized in [25] the Romberg extrapolation methods to the context of stochastic differential equations. Inspired by this work, we next develop a second-order scheme.

Theorem 5.4. *Suppose that in addition to the hypotheses of Theorem 5.1 we have*

S4: *the operator given by $C_2 \varphi_n = |n|^{l_{C_2}} \varphi_n$, where l_{C_2} is a positive real number such that $l_{C_1} \geq l_{C_2} + 4 \max\{l_G, l_L\}$ and $l_{C_2} \geq l_A$, fulfills condition H7.*

Then for any $j = 0, \dots, M$,

$$\begin{aligned}& \left| \mathbf{E} \langle X_{T_j^M}, A X_{T_j^M} \rangle - 2 \mathbf{E} \langle Z_{T_{2j}^{2M},n}^{2M}, A Z_{T_{2j}^{2M},n}^{2M} \rangle + \mathbf{E} \langle Z_{T_j^M,n}^M, A Z_{T_j^M,n}^M \rangle \right| \\ & \leq K \left(\|C^{1/2} X_0\|^2 + \|X_0\|^2 \right) \left(\frac{T^2}{M^2} + \frac{\mathbf{1}_{n_G > 0}}{n^{l_C/2 - l_G}} + \frac{\mathbf{1}_{n_L > 0} + \mathbf{1}_{m_L > 0}}{n^{l_C/2 - 2l_L}} \right. \\ & \quad \left. + \frac{\mathbf{1}_{n_G > 0}}{n^{l_C - 2l_G - l_A}} + \frac{\mathbf{1}_{n_L > 0} + \mathbf{1}_{m_L > 0}}{n^{l_C - 4l_L - l_A}} + \frac{1}{n^{l_C - l_A}} \right)\end{aligned}$$

provided that $c_n \leq \mu M$.

For expository purposes, before proving Theorem 5.4, we give the following lemma.

Lemma 5.5. *Suppose that the hypotheses of Theorem 5.4 hold. Assume that $B \in O_{l_B, n_B, m_B}$ and that $B_k \in O_{l_{B_1}, n_{B_1}, m_{B_1}}$ for $k = 1, \dots, 4$. If $l_C \geq l_{C_1} + 2l_{B_1}$ and $l_{C_1} \geq l_{C_2} + 2l_B$, then*

$$\left| \mathbf{E} \langle B Y_{T_j,n}^{s, B_1 Z_{s,n} + B_2 Z_{T_j,n}}, \Phi_{T_\kappa - \bar{s}, n} B Y_{T_j,n}^{s, B_3 Z_{s,n} + B_4 Z_{T_j,n}} \rangle \right| \leq K \left(\|C^{\frac{1}{2}} X_0\|^2 + \|X_0\|^2 \right)$$

whenever $0 \leq s \leq T_j \leq \bar{s} \leq T_\kappa$.

Proof. The linearity of (5.2) and the polarization identity allow us to assume without loss of generality that $B_1 = B_3$ and $B_2 = B_4$. It follows from (5.1) that

$$\mathbf{E} \left\langle BY_{T_j,n}^{s,B_1 Z_{s,n}+B_2 Z_{T_j,n}}, \Phi_{T_\kappa-\bar{s},n} BY_{T_j,n}^{s,B_3 Z_{s,n}+B_4 Z_{T_j,n}} \right\rangle = \mathbf{E} \left\langle \tilde{Y}, A\tilde{Y} \right\rangle,$$

with $\tilde{Y} = Y_{T_\kappa,n}^{\bar{s},BY_{T_j,n}^{s,B_1 Z_{s,n}+B_2 Z_{T_j,n}}}$. Hence proceeding as in the proof of Lemma 5.3, we obtain

$$\begin{aligned} \left| \mathbf{E} \left(\left\langle \tilde{Y}, A\tilde{Y} \right\rangle / F_{\bar{s}} \right) \right| &\leq K \left(\left\| C_2^{\frac{1}{2}} BY_{T_j,n}^{s,B_1 Z_{s,n}+B_2 Z_{T_j,n}} \right\|^2 + \left\| BY_{T_j,n}^{s,B_1 Z_{s,n}+B_2 Z_{T_j,n}} \right\|^2 \right) \\ &\leq K \left(\left\| C_1^{\frac{1}{2}} Y_{T_j,n}^{s,B_1 Z_{s,n}+B_2 Z_{T_j,n}} \right\|^2 + \left\| Y_{T_j,n}^{s,B_1 Z_{s,n}+B_2 Z_{T_j,n}} \right\|^2 \right). \end{aligned}$$

Therefore

$$\begin{aligned} \left| \mathbf{E} \left(\left\langle \tilde{Y}, A\tilde{Y} \right\rangle / F_{\bar{s}} \right) \right| &\leq K \left(\left\| C_1^{\frac{1}{2}} (B_1 Z_{s,n} + B_2 Z_{T_j,n}) \right\|^2 + \|B_1 Z_{s,n} + B_2 Z_{T_j,n}\|^2 \right) \\ &\leq K \left(\left\| C^{\frac{1}{2}} Z_{s,n} \right\|^2 + \left\| C^{\frac{1}{2}} Z_{T_j,n} \right\|^2 + \|Z_{s,n}\|^2 + \|Z_{T_j,n}\|^2 \right). \end{aligned}$$

Thus the desired result follows from the estimates given in the first part of the proof of Lemma 5.3. $\square$

Proof of Theorem 5.4. Returning to the proof of Theorem 5.1, one may notice that the polarization identity leads to

$$\delta_k = -\frac{T}{2M} \sum_{(B_1, B_2, 1) \in \mathcal{S}} \int_{T_k}^{T_{k+1}} \mathbf{E} \langle B_1 X_{s,n}, \Phi_{T_j-s,n} B_2 X_{s,n} \rangle ds - \tilde{\delta}_k^1 - \tilde{\delta}_k^2,$$

with

$$\begin{aligned} (5.6) \quad \tilde{\delta}_k^1 &= \sum_{(B, \bar{k}, r) \in \tilde{\mathcal{S}}} i^{-\bar{k}} \int_{T_k}^{T_{k+1}} \mathbf{E} \langle B Z_{T_k,n}, \Phi_{T_j-s,n} B Z_{T_k,n} \rangle (s - T_k)^r ds \\ &\quad - \sum_{(B, \bar{k}, r) \in \tilde{\mathcal{S}}} i^{-\bar{k}} \int_{T_k}^{T_{k+1}} \mathbf{E} \langle B X_{T_k,n}, \Phi_{T_j-s,n} B X_{T_k,n} \rangle (s - T_k)^r ds, \\ \tilde{\delta}_k^2 &= \sum_{(B, \bar{k}, r) \in \tilde{\mathcal{S}}} i^{-\bar{k}} \int_{T_k}^{T_{k+1}} \mathbf{E} \langle B X_{T_k,n}, \Phi_{T_j-s,n} B X_{T_k,n} \rangle (s - T_k)^r ds \\ &\quad - \sum_{(B, \bar{k}, 1) \in \tilde{\mathcal{S}}} i^{-\bar{k}} \frac{T}{2M} \int_{T_k}^{T_{k+1}} \mathbf{E} \langle B X_{s,n}, \Phi_{T_j-s,n} B X_{s,n} \rangle ds, \end{aligned}$$

where $\tilde{\mathcal{S}} = \left\{ (B, \bar{k}, r) : \bar{k} = 0, \dots, 3, B = B_1 + i^{\bar{k}} B_2 \text{ with } (B_1, B_2, r) \in \mathcal{S} \right\}$.

We now focus on the estimation of $\tilde{\delta}_k^1$. For any B present in (5.6), let us define the function v_n by

$$v_n(s, x, y) = \mathbf{E} \left\langle BY_{T_k,n}^{s,\bar{x}}, \Phi_{T_j-\bar{s},n} BY_{T_k,n}^{s,y} \right\rangle,$$

provided that $0 \leq s \leq T_k \leq \bar{s} \leq T_j$ and $x, y \in \mathfrak{h}_n$. Thus for any $\bar{s} \in [T_k, T_{k+1}]$

$$\begin{aligned} & \mathbf{E} \langle BZ_{T_k, n}, \Phi_{T_j - \bar{s}, n} BZ_{T_k, n} \rangle - \mathbf{E} \langle BX_{T_k, n}, \Phi_{T_j - \bar{s}, n} BX_{T_k, n} \rangle \\ &= \sum_{l=0}^{k-1} \mathbf{E} v_n (T_{l+1}, \overline{Z_{T_{l+1}, n}}, Z_{T_{l+1}, n}) - \mathbf{E} v_n (T_l, \overline{Z_{T_l, n}}, Z_{T_l, n}). \end{aligned}$$

It follows from Itô's formula that

$$\begin{aligned} & \mathbf{E} (v_n (T_{l+1}, \overline{Z_{T_{l+1}, n}}, Z_{T_{l+1}, n}) - v_n (T_l, \overline{Z_{T_l, n}}, Z_{T_l, n})) \\ &= \mathbf{E} \int_{T_l}^{T_{l+1}} \left(\frac{\partial}{\partial r} v_n (r, \overline{Z_{r, n}}, Z_{r, n}) + \mathcal{L}_{\overline{Z_{T_l, n}}, Z_{T_l, n}} (v_n) (r, \overline{Z_{r, n}}, Z_{r, n}) \right) dr, \end{aligned}$$

with

$$\begin{aligned} \mathcal{L}_{\alpha, \beta} (v_n) (s, x, y) &= \sum_{l \in [\vec{0}, \vec{n}]} \frac{\partial v_n}{\partial x^l} (s, x, y) (\overline{G_n \alpha})^l + \sum_{l \in [\vec{0}, \vec{n}]} \frac{\partial v_n}{\partial y^l} (s, x, y) (G_n \beta)^l \\ &+ \frac{1}{2} \sum_{l=1}^m \sum_{\bar{k}, \bar{j} \in [\vec{0}, \vec{n}]} \frac{\partial^2 v_n}{\partial x^{\bar{k}} \partial x^{\bar{j}}} (s, x, y) (\overline{L_{l, n} \alpha})^{\bar{k}} (\overline{L_{l, n} \alpha})^{\bar{j}} \\ &+ \frac{1}{2} \sum_{l=1}^m \sum_{\bar{k}, \bar{j} \in [\vec{0}, \vec{n}]} \frac{\partial^2 v_n}{\partial x^{\bar{k}} \partial y^{\bar{j}}} (s, x, y) (\overline{L_{l, n} \alpha})^{\bar{k}} (L_{l, n} \beta)^{\bar{j}} \\ &+ \frac{1}{2} \sum_{l=1}^m \sum_{\bar{k}, \bar{j} \in [\vec{0}, \vec{n}]} \frac{\partial^2 v_n}{\partial y^{\bar{k}} \partial x^{\bar{j}}} (s, x, y) (L_{l, n} \beta)^{\bar{k}} (\overline{L_{l, n} \alpha})^{\bar{j}} \\ &+ \frac{1}{2} \sum_{l=1}^m \sum_{\bar{k}, \bar{j} \in [\vec{0}, \vec{n}]} \frac{\partial^2 v_n}{\partial y^{\bar{k}} \partial y^{\bar{j}}} (s, x, y) (L_{l, n} \beta)^{\bar{k}} (L_{l, n} \beta)^{\bar{j}}, \end{aligned}$$

where for any $O \in B(\mathfrak{h}_n)$, the operator $\bar{O}$ is defined by $\bar{O}\varphi_k = \overline{O\varphi_k}$, $k \in [\vec{0}, \vec{n}]$.

Furthermore, again using Itô's formula, we arrive at

$$\begin{aligned} & \mathbf{E} (v_n (T_{l+1}, \overline{Z_{T_{l+1}, n}}, Z_{T_{l+1}, n}) - v_n (T_l, \overline{Z_{T_l, n}}, Z_{T_l, n})) \\ &= \mathbf{E} \int_{T_l}^{T_{l+1}} \int_{T_l}^r \left(\mathcal{L}_{\overline{Z_{T_l, n}}, Z_{T_l, n}} \left(\mathcal{L}_{\overline{Z_{T_l, n}}, Z_{T_l, n}} (v_n) \right) (s, \overline{Z_{s, n}}, Z_{s, n}) \right) ds dr \\ &- 2\mathbf{E} \int_{T_l}^{T_{l+1}} \int_{T_l}^r \left(\mathcal{L}_{\overline{Z_{T_l, n}}, Z_{T_l, n}} (\mathcal{L} (v_n)) (s, \overline{Z_{s, n}}, Z_{s, n}) \right) ds dr \\ &+ \mathbf{E} \int_{T_l}^{T_{l+1}} \int_{T_l}^r (\mathcal{L} (\mathcal{L} (v_n)) (s, \overline{Z_{s, n}}, Z_{s, n})) ds dr, \end{aligned}$$

with $\mathcal{L} (v_n) (s, x, y) = \mathcal{L}_{x, y} (v_n) (s, x, y)$.

Now, a long calculation leads to

$$\mathcal{L}_{\bar{\alpha}, \alpha} (\mathcal{L}_{\bar{\alpha}, \alpha} (v_n)) (s, \bar{x}, x) = 2\mathbf{E} \langle BY_{T_k, n}^{s, G\alpha}, \Phi_{T_k - \bar{s}, n} BY_{T_k, n}^{s, G\alpha} \rangle$$

and $\mathcal{L}_{\bar{\alpha}, \alpha} (\mathcal{L} (v_n)) (s, \bar{x}, x) = \sum_{(y, z) \in \mathcal{I}} \mathbf{E} \langle BY_{T_k, n}^{s, y}, \Phi_{T_k - \bar{s}, n} BY_{T_k, n}^{s, z} \rangle$, where $\mathcal{I}$ is the set formed by the following pairs: $(G_n^2 \alpha, x)$, $(G_n \alpha, G_n x)$, $(G_n x, G_n \alpha)$, $(x, G_n^2 \alpha)$,

$(L_{l,n}G_n\alpha, L_{l,n}x), (L_{l,n}x, L_{l,n}G_n\alpha), (L_{l,n}\alpha, G_nL_{l,n}\alpha), (L_{l_1,n}L_{l,n}\alpha, L_{l_1,n}L_{l,n}\alpha)$ and $(G_nL_{l,n}\alpha, L_{l,n}\alpha)$ with $l, l_1 = 1, \dots, m$. Therefore, Lemma 5.5 yields

$$|\tilde{\delta}_k^1| \leq K \left(\frac{T}{M} \right)^3 \left(\|C^{\frac{1}{2}}X_0\|^2 + \|X_0\|^2 \right).$$

It remains to estimate $\tilde{\delta}_k^2$. Combining the integration by parts formula and (5.3), we have that

$$\begin{aligned} & \int_{T_k}^{T_{k+1}} \mathbf{E} \langle BX_{T_k,n}, \Phi_{T_j-s,n} BX_{T_k,n} \rangle (s - T_k) ds \\ &= \frac{1}{2} \left(\frac{T}{M} \right)^2 \mathbf{E} \langle BX_{T_k,n}, \Phi_{T_j-T_{k+1},n} BX_{T_k,n} \rangle \\ &+ \frac{1}{2} \int_{T_k}^{T_{k+1}} \mathbf{E} \langle G_n BX_{T_k,n}, \Phi_{T_j-s,n} BX_{T_k,n} \rangle (s - T_k)^2 ds \\ &+ \frac{1}{2} \int_{T_k}^{T_{k+1}} \mathbf{E} \langle BX_{T_k,n}, \Phi_{T_j-s,n} G_n BX_{T_k,n} \rangle (s - T_k)^2 ds \\ &+ \frac{1}{2} \sum_{l=1}^m \int_{T_k}^{T_{k+1}} \mathbf{E} \langle L_{l,n} BX_{T_k,n}, \Phi_{T_j-s,n} L_{l,n} BX_{T_k,n} \rangle (s - T_k)^2 ds, \end{aligned}$$

where $(B, \bar{k}, 1) \in \tilde{\mathcal{S}}$ for some $\bar{k}$. Since

$$\begin{aligned} \mathbf{E} \langle BX_{s,n}, \Phi_{T_j-s,n} BX_{s,n} \rangle &= \mathbf{E} \langle BX_{T_k,n}, \Phi_{T_j-s,n} BX_{T_k,n} \rangle \\ &+ \int_{T_k}^s \mathbf{E} \langle BG_n X_{r,n}, \Phi_{T_j-s,n} BX_{r,n} \rangle dr \\ &+ \int_{T_k}^s \mathbf{E} \langle BX_{r,n}, \Phi_{T_j-s,n} BG_n X_{r,n} \rangle dr \\ &+ \sum_{l=1}^m \int_{T_k}^s \mathbf{E} \langle BL_{l,n} X_{r,n}, \Phi_{T_j-s,n} BL_{l,n} X_{r,n} \rangle dr, \end{aligned}$$

we conclude the proof by applying the integration by parts formula and (5.3) to $\int_{T_k}^{T_{k+1}} \mathbf{E} \langle BX_{T_k,n}, \Phi_{T_j-s,n} BX_{T_k,n} \rangle ds$ and arguing similarly as in the last part of the proof of Lemma 5.3. $\square$

6. APPLICATION

This section is devoted to illustrating the main results of this paper. More precisely, in view of the harmonic oscillators' considerable importance in quantum mechanics [4, 21], we develop here an example of a forced and damped quantum harmonic oscillator in the interaction representation. Then this concrete situation allows us to give the flavor of the hypotheses that support our presentation. We begin by establishing the corresponding notation.

In what follows, we assume $\mathfrak{h} = l^2(\mathbb{Z}_+)$ and that $(\varphi_k)_{k \geq 0}$ is the canonical orthonormal basis on the space $l^2(\mathbb{Z}_+)$. Moreover, we denote by $a^\dagger, a$ the creation and annihilation operators; that is, the domain of $a^\dagger, a$ is

$$\left\{ x \in l^2(\mathbb{Z}_+) : \sum_{k \geq 0} k |x_k|^2 < +\infty \right\}$$

and $a^\dagger \varphi_n = \sqrt{n+1} \varphi_{n+1}$,

$$a \varphi_n = \begin{cases} 0, & \text{if } n = 0, \\ \sqrt{n} \varphi_{n-1}, & \text{if } n > 0. \end{cases}$$

The number operator is given by $N = a^\dagger a$.

From now on we consider the system defined by the effective Hamiltonian

$$H = ik_1 (a^\dagger - a) + k_2 N$$

and the Lindblad operators $L_1 = c_1 a$, $L_2 = c_2 a^2$, $L_3 = c_3 N$, $L_4 = c_4 a^\dagger$, where $k_1, k_2 \in \mathbb{R}$ and $c_1, \dots, c_4 \in \mathbb{C}$.

The model presented here describes many physical phenomena, for instance a single mode of a quantized electromagnetic field. In this framework, the vectors φ_k , $k = 0, 1, \dots$, characterize the levels of energy. Moreover, the operator $a^\dagger$ creates a photon, whereas the operator a destroys a photon. Then the term $ik_1 (a^\dagger - a)$ describes a driving force or lineal pumping and L_1 describes the damping due to photon emission. $\mathbf{E} \langle X_t, N X_t \rangle$ is the mean photon number or energy at time t .

Let $\mathfrak{h}_n$ be equal to the linear manifold spanned by $\{\varphi_j : 0 \leq j \leq n\}$. Then $\mathfrak{D}^* = \bigcup_{n=0}^\infty \mathfrak{h}_n$ satisfies hypothesis H4. In addition, for any $\phi = (\phi_0, \dots, \phi_n, 0, \dots) \in \mathfrak{h}_n$,

$$\begin{aligned} & 2 \operatorname{Re} \langle N^p \phi, N^p G_n \phi \rangle + \sum_{k=1}^m \|N^p L_{k,n} \phi\|^2 \\ &= \sum_{j=1}^n |c_1|^2 |\phi_j|^2 j \left(-j^{2p} + (j-1)^{2p} \right) + |c_2|^2 |\phi_j|^2 j (j-1) \left(-j^{2p} + (j-2)^{2p} \right) \\ &+ \sum_{j=0}^{n-1} |c_4|^2 |\phi_j|^2 (j+1) (-j^{2p} + (j+1)^{2p}) \\ &+ 2k_1 \sqrt{j+1} \operatorname{Re} (\phi_{j+1} \overline{\phi_j}) ((j+1)^{2p} - j^{2p}). \end{aligned}$$

Hence H7 is fulfilled by any $C = N^{2p}$, with $p \in \mathbb{Z}_+$. Since for any $\phi \in \operatorname{Dom}(G)$, $\operatorname{Re} \langle \phi, G \phi \rangle = -\frac{1}{2} \sum_{k=1}^m \|L_k \phi\|^2 \leq 0$, G is a dissipative linear operator on $\mathfrak{h}$. Furthermore, $\|GP_n - P_n GP_n\| \leq |k_1| \sqrt{n+1} + \frac{|c_4|^2}{2} (n+1)$. Thus H1 follows from Theorem 2 of [11]. Notice that $G \in O_{2,1,1}$ and $L_j \in O_{1,2,1}$, $j = 1, \dots, 4$. Therefore, Theorem 4.2 implies

Proposition 6.1. *Let $A \in O_{l_A, n_A, m_A}$ and $p \in \mathbb{N}$ such that $p \geq \max\{l_A, 2\}$. If $X_0 \in \operatorname{Dom}(N^p)$, then for any $n > 0$,*

$$\begin{aligned} & \sup_{t \in [0, T]} |\mathbf{E} \langle X_t, A X_t \rangle - \mathbf{E} \langle X_{t,n}, A X_{t,n} \rangle| \\ & \leq K \left(\|N^p X_0\|^2 + \|X_0\|^2 \right) \left(\frac{1}{n^{p-2}} + \frac{1}{n^{2p-4-l_A}} \right). \end{aligned}$$

Furthermore, for any $\phi \in \mathfrak{h}_n$ a straightforward computation yields that $\|N^p G_n \phi\|^2$ is equal to

$$\begin{aligned} & (k_1 |\phi_{n-1}|)^2 n^{2p+1} + n^{2p} |\phi_n|^2 \left| ik_2 n + \frac{|c_1|^2 n}{2} + \frac{|c_2|^2 n(n-1)}{2} + \frac{|c_3|^2 n^2}{2} \right|^2 \\ & + \sum_{j=1}^{n-1} j^{2p} \left| k_1 \phi_{j-1} \sqrt{j} - k_1 \phi_{j+1} \sqrt{j+1} \right. \\ & \quad \left. - \phi_j \left(ik_2 j + \frac{|c_1|^2 j}{2} + \frac{|c_2|^2 j(j-1)}{2} + \frac{|c_3|^2 j^2}{2} + \frac{|c_4|^2 (j+1)}{2} \right) \right|^2. \end{aligned}$$

Thus choosing $c_n = n^4$, we can obtain condition S2. Then Proposition 6.2 follows from Theorem 5.1

Proposition 6.2. *Let $A \in O_{l_A, n_A, m_A}$ and $p \in \mathbb{N}$ such that $p \geq \max\{l_A/2 + 4, l_A\}$. If there exists a constant μ such that $n^4/M \leq \mu$, then*

$$\begin{aligned} & \left| \mathbf{E} \langle X_{T_j}, AX_{T_j} \rangle - \mathbf{E} \langle Z_{T_j, n}^M, AZ_{T_j, n}^M \rangle \right| \\ & \leq K_p \left(\|N^p X_0\|^2 + \|X_0\|^2 \right) \left(\frac{T}{M} + \frac{1}{n^{p-2}} + \frac{1}{n^{2p-4-l_A}} \right), \end{aligned}$$

where the constant K_p depends on p . Furthermore, for any $p \geq \{l_A/2 + 8, l_A\}$,

$$\begin{aligned} & \left| \mathbf{E} \langle X_{T_j^M}, AX_{T_j^M} \rangle - 2\mathbf{E} \langle Z_{T_j^M, n}^{2M}, AZ_{T_j^M, n}^{2M} \rangle + \mathbf{E} \langle Z_{T_j^M, n}^M, AZ_{T_j^M, n}^M \rangle \right| \\ & \leq K_p \left(\|N^p X_0\|^2 + \|X_0\|^2 \right) \left(\left(\frac{T}{M} \right)^2 + \frac{1}{n^{p-2}} + \frac{1}{n^{2p-4-l_A}} \right). \end{aligned}$$

ACKNOWLEDGMENTS

I am indebted to Prof. R. Rebolledo for his valuable advice. I thank Prof. F. Fagnola, Prof. C. Saavedra and Prof. D. Talay for many helpful comments and suggestions. I also thank the anonymous referee and editor for their helpful comments. I acknowledge the warm hospitality of the “Facultad de Matemáticas, Pontificia Universidad Católica de Chile”, where a big part of this research was performed.

REFERENCES

- [1] R. Balescu, *Equilibrium and nonequilibrium statistical mechanics*. Krieger Publishing Company (1991) (original edition 1975). MR **53**:12405
- [2] A. Barchielli, A.M. Paganoni, F. Zucca, *On stochastic differential equations and semigroups of probability operators in quantum probability*. Stochastic Proc. and their Appl. **73** (1998), 69–86. MR **99k**:81138
- [3] H. Carmichael, *An open systems approach to quantum optics*. Lecture Notes in Physics. Springer-Verlag (1993).
- [4] C. Cohen-Tannoudji, B. Diu, F. Laloë, *Quantum mechanics. Volume 1*. Hermann (1977).
- [5] A.M. Chebotarev, F. Fagnola, *Sufficient conditions for conservativity of minimal quantum dynamical semigroups*. J. Funct. Anal. **153** (1998), 382–404. MR **99d**:81064
- [6] F. Fagnola, R. Rebolledo, C. Saavedra, *Quantum flows associated to master equations in quantum optics*. J. Math. Phys. **35** (1994), 1–12. MR **94k**:81353
- [7] F. Fagnola, *Quantum markov semigroups and quantum flows*. Proyecciones **18** (1999), 1–144. MR **2002f**:81054

- [8] F. Fagnola, S. J. Wills, *Solving quantum stochastic differential equations with unbounded coefficients*. J. Funct. Anal. **198** (2003), 279–310.
- [9] N. Gisin, I.C. Percival, *The quantum-state diffusion model applied to open systems*. J. Phys. A **25** (1992), 5677–5691. MR **94b**:82028
- [10] A.S. Holevo, *On dissipative stochastic equations in a Hilbert space*. Probab. Theory Related Fields **104** (1996), 483–500. MR **98c**:60068
- [11] P.T. Jorgensen, *Approximately reducing subspaces for unbounded linear operators*. J. Funct. Anal. **23** (1976), 392–414. MR **98c**:60068
- [12] T. Kato, *Perturbation theory for linear operators. Corrected printing of the second edition*. Springer-Verlag (1980). MR **96a**:47025
- [13] P.E. Kloeden, E. Platen, *Numerical solution of stochastic differential equations*. Springer, Berlin (1992). MR **94b**:60069
- [14] Tarso B.L. Kist, M. Orszag, T.A. Brun, L. Davidovich, *Stochastic Schrödinger equations in cavity QDE: physical interpretation and localization*. J. Opt. B: Quantum Semiclass. Opt. **1** (1999), 251–263.
- [15] R. Dautray, J.-L. Lions, *Mathematical analysis and numerical methods for science and technology. Vol. 5, Evolution Problems I*, Springer-Verlag (1992). MR **92k**:00006
- [16] G.N. Milstein, *The numerical integration of stochastic differentials equations*. Ural University Press (1988).
- [17] C. Mora, *Solución numérica de ecuaciones diferenciales estocásticas mediante métodos exponenciales*. Ph.D. Thesis, Pontificia Universidad Católica de Chile (2002).
- [18] A. Pazy, *Semigroups of linear operators and applications to partial differential equations*. Springer-Verlag (1975). MR **85g**:47061
- [19] I.C. Percival, *Quantum state diffusion*. Cambridge University Press (1998). MR **2000f**:81023
- [20] P. Protter, *Stochastic integration and differential equations*. Springer (1990). MR **91i**:60148
- [21] M.O. Scully, M.S. Zubairy, *Quantum optics*. Cambridge University Press (1997).
- [22] W. T. Strunz, L. Diósi, N. Gisin, T. Yu, *Quantum trajectories for Brownian motion*. Phys. Rev. Lett. **83** (1999), 4909–4913. MR **2000h**:82042
- [23] D. Talay, *Efficient numerical schemes for the approximation of expectations of functionals of the solution of an SDE and applications*. Springer Lecture Notes in Control and Inform. Sc. **61** (1984), 294–313. MR **88b**:60140
- [24] D. Talay, *Discrétisation d'une E.D.S. et calcul approché d'espérances de fonctionnelles de la solution*. Math. Mod. Numer. Anal. **20** (1986), 141–179. MR **87k**:60153
- [25] D. Talay, L. Tubaro, *Expansion of the global error for numerical schemes solving stochastic differential equations*. Stochastic Anal. Appl. **8** (1990), 483–509. MR **92e**:60124
- [26] D. Talay, *Stochastic Hamiltonian systems: exponential convergence to the invariant measure, and discretization by the implicit Euler scheme*. Markov Processes. Related Fields **8** (2002), 163–198. MR **2003e**:60129

DEPARTAMENTO DE INGENIERÍA MATEMÁTICA, UNIVERSIDAD DE CONCEPCIÓN, CONCEPCIÓN, CHILE

Current address: Departamento de Ingeniería Matemática, Universidad de Concepción, Casilla 160 C, Concepción, Chile

E-mail address: cmora@ing-mat.udec.cl