ON THE MINIMAL POLYNOMIAL OF GAUSS PERIODS FOR PRIME POWERS

S. GURAK

ABSTRACT. For a positive integer m , set $\zeta_m = \exp(2\pi i/m)$ and let $\mathbf{Z}_m^*$ denote the group of reduced residues modulo m . Fix a congruence group H of conductor m and of order f . Choose integers $t_1, \dots, t_e$ to represent the $e = \phi(m)/f$ cosets of H in $\mathbf{Z}_m^*$. The Gauss periods

$$\theta_j = \sum_{x \in H} \zeta_m^{t_j x} \quad (1 \leq j \leq e)$$

corresponding to H are conjugate and distinct over $\mathbf{Q}$ with minimal polynomial

$$g(x) = x^e + c_1 x^{e-1} + \cdots + c_{e-1} x + c_e.$$

To determine the coefficients of the period polynomial $g(x)$ (or equivalently, its reciprocal polynomial $G(X) = X^e g(X^{-1})$) is a classical problem dating back to Gauss. Previous work of the author, and Gupta and Zagier, primarily treated the case $m = p$, an odd prime, with $f > 1$ fixed. In this setting, it is known for certain integral power series $A(X)$ and $B(X)$, that for any positive integer N

$$G(X) \equiv A(X) \cdot B(X)^{\frac{p-1}{f}} \pmod{X^N}$$

holds in $\mathbf{Z}[X]$ for all primes $p \equiv 1 \pmod{f}$ except those in an effectively determinable finite set. Here we describe an analogous result for the case $m = p^\alpha$, a prime power ($\alpha > 1$). The methods extend for odd prime powers p^α to give a similar result for certain twisted Gauss periods of the form

$$\psi_j = i^* \sqrt{p} \sum_{x \in H} \left(\frac{t_j x}{p}\right) \zeta_{p^\alpha}^{t_j x} \quad (1 \leq j \leq e),$$

where $\left(\frac{\cdot}{p}\right)$ denotes the usual Legendre symbol and $i^* = i^{\frac{(p-1)^2}{4}}$.

1. INTRODUCTION

For any positive integer m , set $\zeta_m = \exp(2\pi i/m)$ and let $\mathbf{Z}_m^*$ denote the group of reduced residues modulo m . Fix a congruence group H defined modulo m of order f and conductor m . Choose integers $t_1 = 1, t_2, \dots, t_e$ to represent the $e = \phi(m)/f$ cosets of H in $\mathbf{Z}_m^*$. The Gauss periods

$$(1) \quad \theta_j = \sum_{x \in H} \zeta_m^{t_j x} \quad (1 \leq j \leq e)$$

corresponding to H lie in the subfield K of $\mathbf{Q}(\zeta_m)$ fixed by the Galois actions induced by sending $\zeta_m \rightarrow \zeta_m^x$ for $x \in H$. They are conjugate and distinct over $\mathbf{Q}$

Received by the editor June 2, 2005.

2000 *Mathematics Subject Classification*. Primary 11L05, 11T22, 11T23.

©2006 American Mathematical Society
Reverts to public domain 28 years from publication

[8, 11], and have minimal polynomial

$$(2) \quad g(x) = g_H(x) = x^e + c_1 x^{e-1} + \cdots + c_{e-1} x + c_e$$

of degree e . For any numerical character χ defined modulo m , the Gauss sum $G(\chi)$ is given by

$$(3) \quad G(\chi) = \sum_{x \in \mathbf{Z}_m^*} \chi(x) \zeta_m^x.$$

The Gauss periods (1) are intimately related with the sums (3); namely,

$$(4) \quad G(\chi) = \sum_{j=1}^e \chi(t_j) \theta_j$$

for any character χ annihilating H , and

$$(5) \quad \theta_j = \frac{1}{e} \sum_{\chi} \bar{\chi}(t_j) G(\chi) \quad (1 \leq j \leq e),$$

where χ runs through the characters defined modulo m which annihilate H . (Here $\bar{\chi}$ denotes the multiplicative inverse of the χ .)

It is well known from the theory of equations [4] that the coefficients c_r of $g(x)$ in (2) can be computed in terms of the symmetric power sums $S_n = \sum (\theta_j)^n$ using Newton's identities

$$(6) \quad S_r + c_1 S_{r-1} + \cdots + c_{r-1} S_1 + r c_r = 0 \quad (1 \leq r \leq e),$$

$$S_n + c_1 S_{n-1} + \cdots + c_{e-1} S_{n-e+1} + c_e S_{n-e} = 0 \quad (n > e).$$

Alternatively, if

$$(7) \quad G(X) = G_H(X) = X^e g(X^{-1}) = 1 + c_1 X + \cdots + c_e X^e$$

is the reciprocal polynomial of $g(x)$, its logarithm is formally expressed as

$$(8) \quad \log G(X) = \log \prod_{j=1}^e (1 - \theta_j X) = - \sum_{n=1}^{\infty} \frac{S_n X^n}{n}$$

in terms of the power sums S_n .

To determine the coefficients of the period polynomial $g(x)$ in (2) (or equivalently its reciprocal polynomial in (7)) is a classical problem dating back to Gauss [5]. When $f = 1$ one has $\theta_1 = \zeta_m$ in (1) whose minimal polynomial is the well-known cyclotomic polynomial

$$(9) \quad \psi_m(x) = \prod_{d|m} (x^d - 1)^{\mu(m/d)},$$

$$\text{with } \psi_{p^\alpha}(x) = x^{p^{\alpha-1}(p-1)} + \cdots + x^{p^\alpha} + 1$$

when $m = p^\alpha$, a prime power. Henceforth, we shall assume $f > 1$ throughout this paper.

Previous work of the author [7] and Gupta and Zagier [6] primarily treated the case $m = p$, an odd prime, with $f > 1$ fixed. In this case, the beginning coefficients of $g(x)$ exhibit a polynomial dependence on p for all sufficiently large primes $p \equiv 1 \pmod{f}$. More precisely, Gupta and Zagier [6] showed for certain integral

power series $A(X)$ and $B(X)$ depending only on simple arithmetic properties of the f -roots of unity ζ_f^v ($1 \leq v \leq f$), that for any positive integer N

$$(10) \quad G(X) \equiv A(X) \cdot B(X)^{(p-1)/f} \pmod{X^N}$$

holds in $\mathbf{Z}[X]$ for all primes $p \equiv 1 \pmod{f}$ except for those in an effectively determinable finite set. Moreover, they give an elegant algorithm to compute the exceptional sets. For $f = 2$, the congruence (10) holds for $N = p$, so $G(X)$ is uniquely determined. Indeed,

$$(11) \quad G(X) = \sum_{r=0}^e (-1)^{[r/2]} \binom{[e-r/2]}{[r/2]} X^r$$

exactly in closed form, a formula known to Gauss [5]. (Here $[]$ denotes the greatest integer function.) Some extensions of these results for certain congruence groups H of square-free conductor $m = p_1 \cdots p_r$ with a fixed number of prime factors of specified types have been suggested (see [8] and also [6]). However, the prime power case $m = p^\alpha$, $\alpha > 1$, seems to have been overlooked. It is this situation we describe here. The case $p = 2$ is treated first, separately in Section 2, where closed form formulas are known for $G_H(X)$ for any congruence group H of conductor 2^α . In Section 3 we give an analog of (10) for Gauss periods corresponding to congruence groups H of odd prime power conductor p^α , $\alpha > 1$, together with an adaptation of Gupta and Zagier's algorithm to determine exceptional prime powers. When $f = 2$ this congruence is shown to hold modulo $X^{p^{\alpha-1}}$, not enough though to completely determine $G(X)$. However, we have recently found [9] closed form formulas in this case for $G(X)$ which generalize (11). In the final section, we extend the results for odd prime powers p^α to certain twisted Gauss periods of the form

$$(12) \quad \psi_j = i * \sqrt{p} \sum_{x \in H} \left(\frac{t_j x}{p} \right) \zeta_{p^\alpha}^{t_j x} \quad (1 \leq j \leq e),$$

where $(\frac{\cdot}{p})$ denotes the usual Legendre symbol and $i * = i^{(p-1)^2/4}$. Such quadratic twists or integer multiples of them arise classically as values of Kloosterman sums [10], [13] for odd prime powers p^α , $\alpha > 1$.

2. GAUSS PERIODS FOR 2^α

Throughout this section H is a congruence group of conductor 2^α ($\alpha > 1$) and order $f > 1$. It is known that there are only two possible choices for H . For the sake of completeness we include a brief rationale for this fact.

Proposition 1. *A congruence group H of conductor 2^α , $\alpha > 1$, and order $f > 1$ must be either $\{1, 2^\alpha - 1\}$ or $\{1, 2^{\alpha-1} - 1\}$ modulo 2^α .*

Proof. Since H has conductor 2^α with $f > 1$, it is known from class field theory that the field K corresponding to H is a proper subfield of $\mathbf{Q}(\zeta_{2^\alpha})$ not contained in $\mathbf{Q}(\zeta_{2^{\alpha-1}})$, and so $\alpha > 2$. Since $2|f$, H contains an element of order two modulo 2^α —either $2^\alpha - 1$, $2^{\alpha-1} - 1$ or $2^{\alpha-1} + 1$. The last choice can be eliminated since H has conductor 2^α . Thus K is contained in one of the cyclic fields $\mathbf{Q}(\zeta_{2^\alpha} + \zeta_{2^\alpha}^{-1})$ or $\mathbf{Q}(\zeta_{2^\alpha} - \zeta_{2^\alpha}^{-1})$ which correspond to the congruence groups $\{1, 2^\alpha - 1\}$ or $\{1, 2^{\alpha-1} - 1\}$ modulo 2^α , respectively. But any proper subfield of $\mathbf{Q}(\zeta_{2^\alpha} + \zeta_{2^\alpha}^{-1})$ or $\mathbf{Q}(\zeta_{2^\alpha} - \zeta_{2^\alpha}^{-1})$ is a subfield of $\mathbf{Q}(\zeta_{2^{\alpha-1}} + \zeta_{2^{\alpha-1}}^{-1})$ contained in $\mathbf{Q}(\zeta_{2^{\alpha-1}})$. Thus either $K = \mathbf{Q}(\zeta_{2^\alpha} + \zeta_{2^\alpha}^{-1})$ or $\mathbf{Q}(\zeta_{2^\alpha} - \zeta_{2^\alpha}^{-1})$, so H equals $\{1, 2^\alpha - 1\}$ or $\{1, 2^{\alpha-1} - 1\}$. $\square$

From the proposition above the only possibilities for θ_1 in (1) here is $\zeta_{2^\alpha} + \zeta_{2^\alpha}^{-1}$ or $\zeta_{2^\alpha} - \zeta_{2^\alpha}^{-1}$, each with $f = 2$ and $\alpha > 2$. The corresponding minimal polynomials are classically known from the properties of Chebyshev polynomials. We quote results from [9] relevant to the discussion here.

Theorem 1. *The minimal polynomial for $\zeta_{2^\alpha} + \zeta_{2^\alpha}^{-1}$ for $\alpha > 2$ is equivalently characterized by*

$$g(x) = \sum_{n=0}^{2^{\alpha-3}} (-1)^n \frac{2^{\alpha-2}}{2^{\alpha-2} - n} \binom{2^{\alpha-2} - n}{n} x^{2^{\alpha-2} - 2n}$$

or the power sums

$$S_n = \begin{cases} 2^{\alpha-2} \binom{n}{n/2} & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd,} \end{cases}$$

for $1 \leq n \leq 2^{\alpha-2}$, or the congruence

$$G(X) \equiv B_+(X)^{2^{\alpha-2}} \pmod{X^{2^{\alpha-1}}},$$

where

$$B_+(X) = \exp\left(-\sum_{n=1}^{\infty} \binom{2n}{n} \frac{X^{2n}}{2n}\right) = \frac{1}{2}(1 + \sqrt{1 - 4X^2}).$$

The minimal polynomial for $\zeta_{2^\alpha} - \zeta_{2^\alpha}^{-1}$ for $\alpha > 2$ is characterized equivalently by

$$g(x) = \sum_{n=0}^{2^{\alpha-3}} \frac{2^{\alpha-2}}{2^{\alpha-2} - n} \binom{2^{\alpha-2} - n}{n} x^{2^{\alpha-2} - 2n}$$

or the power sums

$$S_n = \begin{cases} -2^{\alpha-2} \binom{n}{n/2} & \text{if } 2 \parallel n \\ 2^{\alpha-2} \binom{n}{n/2} & \text{if } 4 \mid n \\ 0 & \text{if } n \text{ is odd,} \end{cases}$$

for $1 \leq n \leq 2^{\alpha-2}$, or the congruence

$$G(X) \equiv B_-(X)^{2^{\alpha-2}} \pmod{X^{2^{\alpha-1}}},$$

where

$$B_-(X) = \exp\left(-\sum_{n=1}^{\infty} (-1)^n \binom{2n}{n} \frac{X^{2n}}{2n}\right) = \frac{1}{2}(1 + \sqrt{1 + 4X^2}).$$

We treat the case when H has an odd prime power conductor next.

3. GAUSS PERIODS FOR ODD PRIME POWERS

Throughout this section we assume the congruence group H in (1) has conductor $m = p^\alpha$ with p odd and $\alpha > 1$. Since $\mathbf{Z}_{p^\alpha}^*$ is cyclic, $H = (\mathbf{Z}_{p^\alpha}^*)^{\phi(p^\alpha)/f}$, or equivalently H equals the group of f -roots of unity in $\mathbf{Z}_{p^\alpha}^*$. In particular, since H has conductor p^α , $p \nmid f$ so $p \equiv 1 \pmod{f}$. To compute the symmetric power sums S_n , we introduce certain counting functions $T_n(p^\gamma)$ for $0 < \gamma \leq \alpha$ as in [8]. Specifically, let $T_n(p^\gamma)$ count the number of times

$$(13) \quad x_1 + \cdots + x_n \equiv 0 \pmod{p^\gamma}$$

for choice of tuples $(x_1, \dots, x_n)$ with x_i in $(\mathbf{Z}_{p^\gamma}^*)^{\phi(p^\gamma)/f}$ ($1 \leq i \leq n$). These counting functions possess the following useful property.

Lemma 1. *For $\gamma > 1$, $T_n(p^{\gamma-1}) - T_n(p^\gamma)$ equals the number of tuples $(x_1, \dots, x_n)$ with x_i in $(\mathbf{Z}_{p^\gamma}^*)^{\phi(p^\gamma)/f}$ for which $p^{\gamma-1} \mid (x_1 + \cdots + x_n)$.*

Proof. First observe that any f -root of unity $x \pmod{p^{\gamma-1}}$ lifts to a unique f -root of unity $x' \pmod{p^\gamma}$, since

$$(x + tp^{\gamma-1})^f \equiv x^f + ftp^{\gamma-1} \equiv 1 \pmod{p^\gamma}$$

has a unique solution t satisfying $-ft \equiv (x^f - 1)/p^{\gamma-1} \pmod{p}$. Thus each solution $x_1 + \cdots + x_n \equiv 0 \pmod{p^{\gamma-1}}$ with x_i in $(\mathbf{Z}_{p^{\gamma-1}}^*)^{\phi(p^{\gamma-1})/f}$ lifts to a unique solution $x'_1 + \cdots + x'_n \equiv 0 \pmod{p^{\gamma-1}}$ with x'_i in $(\mathbf{Z}_{p^\gamma}^*)^{\phi(p^\gamma)/f}$. The statement of the lemma now readily follows.

Now observe in the expansion (8)

$$\log G(X) = - \sum_{n=1}^{\infty} S_n X^n / n = - \sum_{n=1}^{\infty} \sum_{j=1}^e \left(\sum_{x \in H} \zeta_{p^\alpha}^{t_j x} \right)^n \frac{X^n}{n},$$

that S_n equals the inner sum

$$\sum_{j=1}^e \left(\sum_{x \in H} \zeta_{p^\alpha}^{t_j x} \right)^n = \frac{1}{f} \sum_{r \in \mathbf{Z}_{p^\alpha}^*} \sum_{x_i \in H} \zeta_{p^\alpha}^{r(x_1 + \cdots + x_n)}.$$

But in view of the lemma above, and since

$$\begin{aligned} \sum_{r \in \mathbf{Z}_{p^\alpha}^*} \zeta_{p^\alpha}^{rx} &= \begin{cases} \phi(p^\alpha) & \text{if } p^\alpha \mid x \\ -p^{\alpha-1} & \text{if } p^{\alpha-1} \parallel x \\ 0 & \text{otherwise,} \end{cases} \\ \sum_{r \in \mathbf{Z}_{p^\alpha}^*} \sum_{x_i \in H} \zeta_{p^\alpha}^{r(x_1 + \cdots + x_n)} &= \phi(p^\alpha) T_n(p^\alpha) - p^{\alpha-1} (T_n(p^{\alpha-1}) - T_n(p^\alpha)) \\ &= p^\alpha T_n(p^\alpha) - p^{\alpha-1} T_n(p^{\alpha-1}). \end{aligned}$$

Thus,

$$(14) \quad S_n = \frac{p^{\alpha-1}}{f} (p T_n(p^\alpha) - T_n(p^{\alpha-1})).$$

Now let $\beta_f(n)$ count the number of times $\epsilon_1 + \cdots + \epsilon_n = 0$ for f -roots of unity ϵ_i in $\mathbf{Q}(\zeta_f)$, so $T_n(p^{\alpha-1}) \geq T_n(p^\alpha) \geq \beta_f(n)$. Thus as long as $T_n(p^{\alpha-1}) = \beta_f(n)$,

$$(15) \quad S_n = \frac{\phi(p^\alpha)}{f} \beta_f(n)$$

above. Letting $\xi_f(n)$ be the set of odd prime powers p^α ($\alpha > 1$), $p \equiv 1 \pmod{f}$, for which $T_n(p^{\alpha-1}) > \beta_f(n)$ and putting

$$(16) \quad B(X) = \exp\left(-\sum_{n=1}^{\infty} \beta_f(n) \frac{X^n}{n}\right),$$

one obtains from (8) and (15) the following analog of Gupta and Zagier's result [6, Theorem 1]. $\square$

Theorem 2. *For each natural number N , the congruence*

$$G(X) \equiv B(X)^{\phi(p^\alpha)/f} \pmod{X^N}$$

holds in $\mathbf{Z}[X]$ for all odd prime powers p^α ($\alpha > 1$), $p \equiv 1 \pmod{f}$, except those lying in $\xi_f(n)$ for some $n < N$.

Before giving some examples, some comments concerning the result above and the computation of the exceptional sets $\xi_f(n)$ are in order. The power series $B(X)$ in (16) is the same integral power series appearing in [6]. The counting function $\beta_f(n)$ has a nice expression when f is a prime power or twice a prime (chiefly, Theorem 2 in [6]). In particular, one has

$$(17) \quad \beta_l(n) = \begin{cases} 0 & \text{if } l \nmid n \\ \frac{n!}{((n/l)!)^l} & \text{if } l \mid n \end{cases}$$

when l is a prime and

$$(18) \quad \beta_4(n) = \begin{cases} 0 & \text{if } n \text{ is odd} \\ \left(\binom{n}{n/2}\right)^2 & \text{if } n \text{ is even} \end{cases}$$

(see also Corollary 2 in [8]).

Gupta and Zagier [6] give an elegant alternative expression for $\beta_f(n)$ which is useful in computing the exceptional sets $\xi_f(n)$. For any tuple $\bar{i} = (i_1, \dots, i_n)$ in $(\mathbf{Z}/f\mathbf{Z})^n$, let $P_{\bar{i}}(x)$ denote the polynomial $x^{i_1} + \dots + x^{i_n}$ with i_v ($1 \leq v \leq n$) chosen to be the least nonnegative representative of the class $i_v \pmod{f}$. Then

$$(19) \quad \beta_f(n) = \text{card}(\{\bar{i} = (i_1, \dots, i_n) \in (\mathbf{Z}/f\mathbf{Z})^n \mid \psi_f(x) \text{ divides } P_{\bar{i}}(x)\}),$$

where $\psi_f(x)$ denotes the f th cyclotomic polynomial (9). If one fixes a prime $\tilde{p}$ lying above p in $\mathbf{Q}(\zeta_f)$, then

$$(20) \quad T_n(p^\gamma) = \text{card}(\{\bar{i} = (i_1, \dots, i_n) \in (\mathbf{Z}/f\mathbf{Z})^n \mid \tilde{p}^\gamma \text{ divides } \zeta_f^{i_1} + \dots + \zeta_f^{i_n}\})$$

in (13), independent of the choice of $\tilde{p}$. Clearly $T_n(p^{\alpha-1}) > \beta_f(n)$ if and only if for some tuple $\bar{i} = (i_1, \dots, i_n)$ in $(\mathbf{Z}/f\mathbf{Z})^n$

$$(21) \quad \tilde{p}^{\alpha-1} \mid (\zeta_f^{i_1} + \dots + \zeta_f^{i_n}) \text{ but } \zeta_f^{i_1} + \dots + \zeta_f^{i_n} \neq 0.$$

Thus $\xi_f(n)$ consists of all odd prime powers p^α , $p \equiv 1 \pmod{f}$ for which (21) holds for some tuple $\bar{i}$ in $(\mathbf{Z}/f\mathbf{Z})^n$. Now a necessary condition for $\bar{i}$ to satisfy (21) is $R_{\bar{i}} \equiv 0 \pmod{p^{\alpha-1}}$, but $R_{\bar{i}} \neq 0$, where $R_{\bar{i}}$ denotes the resultant of $P_{\bar{i}}(x)$ and $\psi_f(x)$. The resultant $R_{\bar{i}} = 0$ if $\psi_f(x)$ divides $P_{\bar{i}}(x)$. When $\psi_f(x) \nmid P_{\bar{i}}(x)$ (so $(\psi_f, P_{\bar{i}}) = 1$), $R_{\bar{i}}$ is essentially the smallest positive integer R for which the equation $g(x)\psi_f(x) + h(x)P_{\bar{i}}(x) = R$ is solvable with polynomials $g, h \in \mathbf{Z}[x]$, or equivalently, the norm $N_{\mathbf{Q}(\zeta_f)/\mathbf{Q}} P_{\bar{i}}(\zeta_f)$. Thus to determine which prime powers lie in $\xi_f(n)$, one

TABLE 1.

$f = 4, n = 7 : \tilde{p}_5 = 2 - \zeta_5 \text{ where } \zeta_4 \equiv 7(\text{mod } \tilde{p}_5^2)$		
$\bar{i}$	$R_{\bar{i}}$	$\zeta_f^{i_1} + \cdots + \zeta_f^{i_n}$ factors
0000002	25	$5 = (2 + \zeta_4)(2 - \zeta_4)$
0000111	25	$4 + 3\zeta_4 = \zeta_4(2 - \zeta_4)^2$
0000333	25	$4 - 3\zeta_4 = -\zeta_4(2 + \zeta_4)^2$
0111112	25	$5\zeta_4 = \zeta_4(2 + \zeta_4)(2 - \zeta_4)$
$f = 5, n = 7 : \tilde{p}_{11} = 2 + \zeta_5^2 \text{ where } \zeta_5 \equiv 3(\text{mod } \tilde{p}_{11}^2)$		
$\bar{i}$	$R_{\bar{i}}$	$\zeta_f^{i_1} + \cdots + \zeta_f^{i_n}$ factors
0000123	121	$3 - \zeta_5^4 = -\zeta_5^4(1 + \zeta_5^4)(2 + \zeta_5^2)^2$
0000124	121	$3 - \zeta_5^3 = -\zeta_5^3(1 + \zeta_5^3)(2 + \zeta_5)^2$
0000134	121	$3 - \zeta_5^2 = -\zeta_5^2(1 + \zeta_5^2)(2 + \zeta_5^4)^2$
0000234	121	$3 - \zeta_5 = -\zeta_5(1 + \zeta_5)(2 + \zeta_5^2)^2$
0001113	121	$3 + 3\zeta_5 + \zeta_5^3 = -\zeta_5^2(1 + \zeta_5^2)(2 + \zeta_5)(2 + \zeta_5^4)$
0001222	121	$3 + 3\zeta_5^2 + \zeta_5 = -\zeta_5^4(1 + \zeta_5^4)(2 + \zeta_5^2)(2 + \zeta_5^3)$
$f = 3, n = 11 : \tilde{p}_7 = \zeta_3 - 2 \text{ where } \zeta_3 \equiv -19(\text{mod } \tilde{p}_7^2)$		
$\bar{i}$	$R_{\bar{i}}$	$\zeta_f^{i_1} + \cdots + \zeta_f^{i_n}$ factors
00000000111	49	$8 + 3\zeta_3 = -\zeta_3^2(\zeta_3 - 2)^2$
00000000222	49	$8 + 3\zeta_3^2 = -\zeta_3(\zeta_3^2 - 2)^2$

first restricts to those prime powers p^α , $p \equiv 1(\text{mod } f)$ for which $R_{\bar{i}} \equiv 0(\text{mod } p^{\alpha-1})$ but $R_{\bar{i}} \not\equiv 0$ for some tuple $\bar{i}$ in $(\mathbf{Z}/f\mathbf{Z})^n$. For such a prime power p^α , select a primitive f -root of unity g modulo $p^{\alpha-1}$ for which $\tilde{p}^{\alpha-1} | (\zeta_f - g)$. Then in view of (21), $p^\alpha \in \xi_f(n)$ if and only if

$$g^{i_1} + \cdots + g^{i_n} \equiv 0 \pmod{p^{\alpha-1}} \text{ with } \zeta_f^{i_1} + \cdots + \zeta_f^{i_n} \neq 0$$

for some tuple $\bar{i} = (i_1, \dots, i_n)$ in $(\mathbf{Z}/f\mathbf{Z})^n$. It is enough to consider tuples $\bar{i}$, inequivalent under translation and permutation.

Table 1 shows the results of this computation for finding exceptional prime powers p^3 among the inequivalent n -tuples $\bar{i}$ for which $R_{\bar{i}} \equiv 0(\text{mod } p^2)$, $R_{\bar{i}} \neq 0$ in cases $(f, n) = (4, 7), (5, 7)$ and $(3, 11)$.

To illustrate when $f = 5$ and $n = 7$, one finds six inequivalent classes of 7-tuples $\bar{i}$ with $p^2 | R_{\bar{i}}$ and $R_{\bar{i}} \neq 0$. Representatives for each such class are given in Table 1; all have resultant $R_{\bar{i}} = 121$. To check if $11^3 \in \xi_5(7)$ one may choose $\tilde{p}_{11} = 2 + \zeta_5^2$, where $\zeta_5 \equiv 3(\text{mod } \tilde{p}_{11}^2)$. One of these tuples satisfies (21) with $\tilde{p}_{11} = 2 + \zeta_5^2$ and $\alpha = 3$; namely, 0000234, so $11^3 \in \xi_5(7)$. There are $\frac{7!}{4!} = 210$ permutations of 0000234 with five translations each, so

$$T_7(121) = \beta_5(7) + 210 \cdot 5 = 1250$$

as $\beta_5(7) = 0$.

The “new” exceptional prime powers for given n are the elements of $\xi_f(n)$ which do not appear in $\xi_f(n')$ for some $n' < n$. Table 2 lists the “new” exceptional prime powers for $3 \leq f \leq 8$ and small values of n .

TABLE 2.

$\underline{n}$	$f = 3$	$f = 4$	$f = 6$
3	— — —	5^2	7^2
4	7^2	— — —	13^2
5	13^2	$13^2, 17^2$	19^2
6	— — —	— — —	31^2
7	$19^2, 31^2$	$5^3, 29^2, 37^2$	$37^2, 43^2$
8	43^2	— — —	7^3
9	— — —	$41^2, 53^2$	$61^2, 67^2, 73^2$
10	$37^2, 73^2$	— — —	79^2
11	$7^3, 67^2$	$61^2, 73^2, 101^2$	$97^2, 103^2$
$\underline{n}$	$f = 5$	$f = 7$	$f = 8$
3	11^2	43^2	17^2
4	61^2	$29^2, 71^2, 547^2$	41^2
5	41^2	$113^2, 197^2,$ $421^2, 463^2$	$73^2, 89^2, 97^2$ $113^2, 257^2$
6	$31^2, 71^2$ $181^2, 521^2$	$211^2, 379^2, 449^2$ $751^2, 2689^2$	$137^2, 313^2$
7	$11^3, 101^2, 151^2$ $191^2, 461^2$	$239^2, 281^2, 337^2$ $1583^2, 1597^2$	$17^3, 193^2, 233^2$ $241^2, 281^2, 337^2$ $353^2, 409^2, 433^2,$ $641^2, 1297^2$

Here are a couple of examples to illustrate Theorem 2.

Example 1. Consider the case $p = 7$ with $f = 3$, so

$$\beta_3(n) = \begin{cases} n!/((n/3)!)^3 & \text{if } 3|n, \\ 0 & \text{otherwise} \end{cases}$$

from (17) with

$$B(X) = 1 - 2X^3 - 13X^6 - 158X^9 - 2431X^{12} - \dots$$

in (16), The power 49 first appears in the exceptional set $\xi_3(4)$ with $T_4(7) = 12 > \beta_3(4) = 0$, whereas 343 first appears in the exceptional set $\xi_3(11)$ with $T_{11}(49) = 495 > \beta_3(11) = 0$. The minimal polynomials $G(X)$ for $\theta_{49}^{-1} = (\zeta_{49} + \zeta_{49}^{18} + \zeta_{49}^{-19})^{-1}$ begins

$$1 - 28X^3 + \underline{7}X^4 + \underline{14}X^5 + \underline{189}X^6 + \dots,$$

whereas that for $\theta_1^{-1} = (\zeta_{343} + \zeta_{343}^{18} + \zeta_{343}^{-19})^{-1}$ begins

$$1 - 196X^3 + 7^2 \cdot 362X^6 - 7^3 \cdot 2872X^9 + \underline{7}^2 \cdot \underline{15}X^{11} + 7^3 \cdot 109733X^{12} + \dots$$

The underscored coefficients deviate as expected from the pattern of the beginning coefficients given by Theorem 2.

Example 2. Consider next $p = 5$ with $f = 4$, so $\beta_4(n)$ is given by (18) with

$$B(X) = 1 - 2X^2 - 7X^4 - 50X^6 - 456X^8 - \dots$$

in (16). The power 25 first appears in $\xi_4(3)$ with $T_3(5) = 12 > \beta_4(3) = 0$, whereas 125 first appears in the exceptional set $\xi_4(7)$ with $T_7(25) = 140 > \beta_4(7) = 0$. Here

one may take $H = \{\pm 1, \pm 7\}$ for $m = 25$ and $H = \{\pm 1, \pm 57\}$ for $m = 125$. The minimal polynomial $G(X)$ for $\theta_1^{-1} = (\zeta_{25} + \zeta_{25}^{-1} + \zeta_{25}^{57} + \zeta_{25}^{-57})^{-1}$ is

$$1 - 10X^2 + \underline{5}X^3 + \underline{10}X^4 + \underline{1}X^5.$$

That for $\theta_1^{-1} = (\zeta_{125} + \zeta_{125}^{-1} + \zeta_{125}^{57} + \zeta_{125}^{-57})^{-1}$ begins

$$1 - 50X^2 + 1025X^4 - 11250X^6 + \underline{125}X^7 + \dots$$

The underscored coefficients deviate as expected from the pattern of the beginning coefficients given by Theorem 2.

We conclude this section with a discussion of the case $f = 2$ where a closed formula for $G(X)$ has been obtained [9]. The power series $B(X)$ in (16) is just

$$B(X) = \frac{1}{2}(1 + \sqrt{1 - 4X^2}) = 1 - \sum_{n=0}^{\infty} \binom{2n}{n} \frac{X^{2n+2}}{n+1}$$

with $\beta_2(n)$ given by (17). The minimal polynomial $G(X)$ for $(2\cos 2\pi/p^\alpha)^{-1}$ is seen to have the form

$$(22) \quad G(X) = \left(\frac{1 + \sqrt{1 - 4X^2}}{2} \right)^{\phi(p^\alpha)/2} \cdot \frac{1 - \left(\frac{1 - \sqrt{1 - 4X^2}}{2X} \right)^{p^\alpha}}{1 - \left(\frac{1 - \sqrt{1 - 4X^2}}{2X} \right)^{p^{\alpha-1}}}$$

with power sums S_n satisfying

$$(23) \quad S_n = p^\alpha \sum_{t=1, t \text{ odd}}^{[np^{-\alpha}]} \binom{n}{(n - p^\alpha t)/2} - p^{\alpha-1} \sum_{t=1, t \text{ odd}}^{[np^{1-\alpha}]} \binom{n}{(n - p^{\alpha-1} t)/2}$$

if n is odd, or

$$\frac{\phi(p^\alpha)}{2} \binom{n}{n/2} + p^\alpha \sum_{t=1}^{[np^{-\alpha}/2]} \binom{n}{n/2 - p^\alpha t} - p^{\alpha-1} \sum_{t=1}^{[np^{1-\alpha}/2]} \binom{n}{n/2 - p^{\alpha-1} t}$$

if n is even. A closed form formula for $G(X)$ is

$$(24) \quad G(X) = X^{\phi(p^\alpha)/2} + \sum_{j=0}^{(p-3)/2} X^{p^{\alpha-1}j} C_{p^{\alpha-1}(\frac{p-1}{2}-j)}(X),$$

where

$$C_d(X) = \sum_{n=0}^{[d/2]} (-1)^n \frac{d}{d-n} \binom{d-n}{n} X^{2n},$$

or equivalently with coefficients in (7) satisfying for $1 \leq r < \phi(p^\alpha)/2$,

$$(25) \quad c_r = \sum_{j=0, j \equiv r \pmod{2}}^{[rp^{1-\alpha}]} (-1)^{t_j} \frac{p^{\alpha-1}(\frac{p-1}{2}-j)}{p^{\alpha-1}(\frac{p-1}{2}-j)-t_j} \binom{p^{\alpha-1}(\frac{p-1}{2}-j)-t_j}{t_j}$$

with $c_{\phi(p^\alpha)/2} = \left(\frac{-2}{p} \right)$, where $t_j = (r - p^{\alpha-1}j)/2$.

From (22) or (24), it follows that the congruence

$$G(X) \equiv B(X)^{\phi(p^\alpha)/2} \pmod{X^N}$$

holds in $\mathbf{Z}[X]$ for $N = p^{\alpha-1}$ but not for $N > p^{\alpha-1}$, and thus fails to determine $G(X)$ which is of degree $\phi(p^\alpha)/2$. Indeed, one can show p^α first appears in the exceptional set $\xi_2(n)$ for $n = p^{\alpha-1}$.

4. TWISTED GAUSS PERIODS FOR p^α

Here we consider quadratic twists of Gauss periods for prime powers p^α , $\alpha > 1$. When $p = 2$ with $\alpha > 3$, the twisted Gauss periods have the form $\sqrt{2}(\zeta_{2^\alpha}^v + \zeta_{2^\alpha}^{-v})$ or $\sqrt{2}(\zeta_{2^\alpha}^v - \zeta_{2^\alpha}^{-v})$ for some odd integer v , and their corresponding minimal polynomials are easily obtained from the Theorem 1 in Section 2. With no loss of generality then we restrict attention to the case in which p is odd, and consider the twisted Gauss periods

$$(26) \quad \psi_j = i^* \sqrt{p} \sum_{x \in H} \left(\frac{t_j x}{p}\right) \zeta_{p^\alpha}^{t_j x} \quad (1 \leq j \leq e),$$

with H a congruence group of conductor p^α ($\alpha > 1$) and order $f > 1$ as in (1). The twisted Gauss periods (26) also lie in the subfield K of $\mathbf{Q}(\zeta_{p^\alpha})$ corresponding to H with $[K : \mathbf{Q}] = \phi(p^\alpha)/f$. In fact $\psi_j = \text{Tr}_{\mathbf{Q}(\zeta_{p^\alpha})/K}(i^* \sqrt{p} \zeta_{p^\alpha}^{t_j})$ ($1 \leq j \leq e$), and each is seen to generate K .

Proposition 2. *Each twisted Gauss period ψ_j in (26) generates the field K over $\mathbf{Q}$.*

Proof. It suffices to show the conjugates ψ_j ($1 \leq j \leq e$) are all distinct. For this purpose set

$$T(\chi) = \sum_{j=1}^e \chi(t_j) \psi_j$$

for any numerical character χ annihilating H . Then

$$\psi_j = \frac{1}{e} \sum_{\chi} \bar{\chi}(t_j) T(\chi) \quad (1 \leq j \leq e),$$

the sum taken over the characters χ annihilating H . Generalizing the argument in the Appendix of [8] one finds in view of the lemma there that the ψ_j ($1 \leq j \leq e$) will be distinct provided $T(\chi) \neq 0$ for all χ annihilating H with conductor $f(\chi)$ satisfying $(p^\alpha/f(\chi), f(\chi)) = 1$, where $p^\alpha/f(\chi)$ is square-free. We assert that this hypothesis above holds here so that the ψ_j are distinct. Indeed for any character χ annihilating H , $T(\chi)$ equals

$$\sum_{j=1}^e \chi(t_j) i^* \sqrt{p} \sum_{x \in H} \left(\frac{t_j x}{p}\right) \zeta_{p^\alpha}^{t_j x} = i^* \sqrt{p} \sum_{x \in Z_{p^\alpha}^*} \chi(x) \left(\frac{x}{p}\right) \zeta_{p^\alpha}^x,$$

or $i^* \sqrt{p} G(\chi\psi)$ in terms of a Gauss sum (3), where ψ is the quadratic character $(\frac{x}{p})$ considered modulo p^α . But any such character χ with $(p^\alpha/f(\chi), f(\chi)) = 1$ and $p^\alpha/f(\chi)$ square-free must have conductor $f(\chi) = p^\alpha$. Hence $\chi\psi$ must also have conductor p^α so $G(\chi\psi) \neq 0$, and thus $T(\chi) \neq 0$.

The proof of the proposition is now complete. $\square$

Our goal here is to give an analog of Theorem 2 in Section 3 for the minimal polynomial $G(X)$ for the reciprocals of the ψ_j in (26). Two situations arise naturally depending on whether $(p-1)/f$ is even or odd. To deal with the case $(p-1)/f$ is even, we set

$$(27) \quad \hat{\beta}_f(n) = \begin{cases} \beta_f(n) & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd,} \end{cases}$$

where $\beta_f(n)$ is the counting function in (15). In terms of this adjusted counting function $\hat{\beta}_f(n)$ and the exceptional sets $\xi_f(n)$ as before, we find

Theorem 3. Suppose $(p-1)/f$ is even where $\hat{B}(X) = \exp(-\sum_{n=1}^{\infty} \hat{\beta}_f(n) X^n/n)$ with $\hat{\beta}_f(n)$ defined by (27). Then for each natural number N , the congruence

$$G(X) \equiv \hat{B}(i^* \sqrt{p} X)^{\phi(p^\alpha)/f} \pmod{X^N}$$

holds in $\mathbf{Z}[X]$ for all odd prime powers p^α ($\alpha > 1$) except those lying in $\xi_f(n)$ for some $n < N$.

Proof. Here $\log G(X) = -\sum_{n=1}^{\infty} S_n X^n/n$ takes the form

$$-\sum_{n=1}^{\infty} \frac{(i^* \sqrt{p} X)^n}{n} \left(\sum_{j=1}^e \left(\sum_H \left(\frac{t_j x}{p} \right) \zeta_{p^\alpha}^{t_j x} \right)^n \right)$$

with inner sum

$$\sum_{j=1}^e \left(\sum_H \left(\frac{t_j x}{p} \right) \zeta_{p^\alpha}^{t_j x} \right)^n = \frac{1}{f} \sum_{r \in Z_{p^\alpha}^*} \left(\sum_{x \in H} \left(\frac{rx}{p} \right) \zeta_{p^\alpha}^{rx} \right)^n = \frac{1}{f} \sum_{r \in Z_{p^\alpha}^*} \left(\frac{r}{p} \right)^n \left(\sum_H \zeta_{p^\alpha}^{rx} \right)^n,$$

since H is contained in $Z_{p^\alpha}^{*2}$ as $(p-1)/f$ is even. When n is even, this last sum is just

$$\frac{1}{f} \sum_{r \in Z_{p^\alpha}^*} \left(\sum_H \zeta_{p^\alpha}^{rv} \right)^n = \frac{1}{f} (p^\alpha T_n(p^\alpha) - p^{\alpha-1} T_n(p^{\alpha-1})) = \frac{\phi(p^\alpha)}{f} \beta_f(n)$$

as in the proof of Theorem 2 when p^α does not lie in $\xi_f(n)$ for $n < N$. When n is odd this sum becomes

$$\frac{1}{f} \sum_{r \in Z_{p^\alpha}^*} \sum_{x_i \in H} \left(\frac{r}{p} \right) \zeta_{p^\alpha}^{r(x_1 + \dots + x_n)} = \frac{1}{f} \sum_{x_i \in H} \sum_{r \in Z_{p^\alpha}^*} \left(\frac{r}{p} \right) \zeta_{p^\alpha}^{r(x_1 + \dots + x_n)} = 0$$

for any prime power p^α not in $\xi_f(n)$ for some $n < N$, since $\sum_{r \in Z_{p^\alpha}^*} \left(\frac{r}{p} \right) \zeta_{p^\alpha}^{rv} = 0$ unless $p^{\alpha-1} \mid v$. This establishes the theorem with $\hat{B}(X)$ defined in terms of the adjusted counting function $\hat{\beta}_f(n)$ in (27). $\square$

Example 3. Consider $p = 7$ with $f = 3$, so

$$\tilde{\beta}_3(n) = \begin{cases} n!/((n/3)!)^3 & \text{if } 6 \mid n, \\ 0 & \text{otherwise,} \end{cases}$$

in (27) with

$$\tilde{B}(X) = 1 - 15X^6 - 2775X^{12} - 910202X^{18} - \dots$$

in Theorem 3. Then

$$\tilde{B}(i\sqrt{7}X) = 1 + 7^3 \cdot 15X^6 - 7^6 \cdot 2775X^{12} + 7^9 \cdot 910202X^{18} - \dots$$

One may again take $H = \{1, 18, -19\}$ for $m = 49$ or 343 as in Example 1, where 49 first appears in $\xi_3(4)$ and 343 first appears in $\xi_3(11)$. The minimal polynomial for the reciprocal of $\psi_1 = i * \sqrt{7}(\zeta_{49} + \zeta_{49}^{18} + \zeta_{49}^{-19})$ begins

$$1 + \underline{7}^3 X^4 + \underline{7}^4 \cdot \underline{2} X^5 + \underline{7}^4 \cdot \underline{29} X^6 + \dots,$$

whereas that for the reciprocal of $\psi_1 = i * \sqrt{7}(\zeta_{343} + \zeta_{343}^{18} + \zeta_{343}^{-19})$ begins

$$1 + 7^5 \cdot 36X^6 - \underline{7}^8 \cdot \underline{165} X^{11} + 7^9 \cdot 2325X^{12} + \dots$$

The underscored coefficients deviate as expected from the pattern given in Theorem 3.

Now consider the case $(p-1)/f$ is odd, so f must be even. For any positive integer n , let $\beta_f^+(n)$ count the number of times a sum satisfies $\epsilon_1 + \cdots + \epsilon_n = 0$ for f -roots of unity ϵ_i in $\mathbf{Q}(\zeta_f)$ with an **even** number of the ϵ_i actually being $f/2$ -roots of unity. Similarly, let $\beta_f^-(n)$ count the number of times a sum satisfies $\epsilon_1 + \cdots + \epsilon_n = 0$ for f -roots of unity ϵ_i in $\mathbf{Q}(\zeta_f)$ with an **odd** number of the ϵ_i actually being $f/2$ -roots of unity. Clearly $\beta_f^+(n) + \beta_f^-(n) = \beta_f(n)$ for any n , and also $\beta_f^+(n) = \beta_f^-(n)$ if n is odd. Define an adjusted counting function $\hat{\beta}_f(n)$ by

$$(28) \quad \hat{\beta}_f(n) = \beta_f^+(n) - \beta_f^-(n)$$

for any positive integer n . Note that $\hat{\beta}_f(n) = 0$ if n is odd.

In terms of this adjusted counting function and exceptional sets $\xi_f(n)$ as before, we find

Theorem 4. *Suppose $(p-1)/f$ is odd where $\hat{B}(X) = \exp(-\sum_{n=1}^{\infty} \hat{\beta}_f(n)X^n/n)$ with $\hat{\beta}_f(n)$ defined by (28). Then for each natural number N , the congruence*

$$G(X) \equiv \hat{B}(i^* \sqrt{p}X)^{\phi(p^\alpha)/f} \pmod{X^N}$$

holds in $\mathbf{Z}[X]$ for all odd primes powers p^α ($\alpha > 1$) except those lying in $\xi_f(n)$ for some $n < N$.

Proof. In this case the inner sum in the expansion for $\log G(X)$ in the proof of Theorem 3 can be expressed

$$(29) \quad \begin{aligned} \sum_{j=1}^e \left(\sum_H \left(\frac{t_j x}{p} \right) \zeta_{p^\alpha}^{t_j x} \right)^n &= \frac{1}{f} \sum_{r \in Z_{p^\alpha}^*} \left(\sum_{x \in H} \left(\frac{rx}{p} \right) \zeta_{p^\alpha}^{rx} \right)^n \\ &= \frac{1}{f} \sum_{x_i \in H} \sum_{r \in Z_{p^\alpha}^*} \left(\frac{r}{p} \right)^n \zeta_{p^\alpha}^{r(x_1 + \cdots + x_n)} - \frac{1}{f} \sum_{x_i \in H} \sum_{r \in Z_{p^\alpha}^*} \left(\frac{r}{p} \right)^n \zeta_{p^\alpha}^{r(x_1 + \cdots + x_n)} \end{aligned}$$

since $(\frac{x}{p}) = -1$ for any $x \in H - H^2$. Here H^2 is just the set of $f/2$ -roots of unity in $\mathbf{Z}_{p^\alpha}^*$, $\sum^+$ is the sum over tuples $(x_1, \dots, x_n)$ with an **even** number of components lying in $H - H^2$ and $\sum^-$ is the analogous sum over tuples $(x_1, \dots, x_n)$ with an **odd** number of components lying in $H - H^2$. If p^α is not in any $\xi_f(n')$ for $n' < N$, then each term $\sum_{r \in Z_{p^\alpha}^*} (\frac{r}{p}) \zeta_{p^\alpha}^{rx} = 0$ in (29) and hence $S_n = 0$ when n is odd, since no sum $x_1 + \cdots + x_n$ is exactly divisible by $p^{\alpha-1}$. When n is even, there may be sums $x_1 + \cdots + x_n$ divisible by p^α , each such sum $\sum_{r \in Z_{p^\alpha}^*} \zeta_{p^\alpha}^{r(x_1 + \cdots + x_n)}$ contributing $\phi(p^\alpha)$ in (29). Overall, there are $\beta_f^+(n)$ of these with a $+$ sign and $\beta_f^-(n)$ with a $-$ sign, so (29) matches $\frac{\phi(p^\alpha)}{f}(\beta_f^+(n) - \beta_f^-(n))$. This establishes the assertion of the theorem with $\hat{B}(X)$ defined in terms of the adjusted sums $\hat{\beta}_f(n)$ in (28). $\square$

Example 4. Consider $p = 7$ with $f = 6$. Here one finds $\beta_6^+(1) = \beta_6^-(1) = 0$, $\beta_6^+(2) = 0$, $\beta_6^-(2) = 6$, $\beta_6^+(3) = \beta_6^-(3) = 6$, $\beta_6^+(4) = 90$, $\beta_6^-(4) = 0$, $\beta_6^+(5) = \beta_6^-(5) = 180$, $\beta_6^+(6) = 180$ and $\beta_6^-(6) = -1860$ to obtain values $\hat{\beta}_6(2) = -6$, $\hat{\beta}_6(4) = 90$ and $\hat{\beta}_6(6) = -1680$ in (28). Then

$$\hat{B}(X) = 1 + 3X^2 - 18X^4 + 217X^6 + \dots$$

in Theorem 4 so

$$\hat{B}(i\sqrt{7}X) = 1 - 21X^2 - 18 \cdot 7^2X^4 - 217 \cdot 7^3X^6 - \dots.$$

One may take $H = \{\pm 1, \pm 18, \pm 19\}$ for $m = 49$ or 343 . From Table 2 one finds that 49 first appears in $\xi_6(3)$, whereas 343 first appears in $\xi_6(8)$. The minimal polynomial for the reciprocal of $\psi_1 = i\sqrt{7}(\zeta_{49} + \zeta_{49}^{18} + \zeta_{49}^{-19} - \zeta_{49}^{-1} - \zeta_{49}^{-18} - \zeta_{49}^{19})$ in (26) is

$$1 - 7^2 \cdot 3X^2 - \underline{7}^3X^3 + \underline{7}^4X^4 + \underline{2} \cdot \underline{7}^4X^5 - \underline{7}^4X^7,$$

whereas that for the reciprocal of $\psi_1 = i\sqrt{7}(\zeta_{343} + \zeta_{343}^{18} + \zeta_{343}^{-19} - \zeta_{343}^{-1} - \zeta_{343}^{-18} - \zeta_{343}^{19})$ begins

$$1 - 7^3 \cdot 3X^2 - 7^4 \cdot 198X^4 - 7^6 \cdot 1111X^6 + \underline{7}^7 \cdot \underline{29027}X^8 + \underline{7}^8 \cdot \underline{6}X^9 + \dots.$$

The underscored coefficients deviate as expected from the pattern of the beginning coefficients given in Theorem 4.

Example 5. Next consider $p = 5$ with $f = 4$. One readily finds that

$$\hat{\beta}_4(n) = \begin{cases} \binom{n}{n/2}^2 & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd} \end{cases}$$

in (28) so

$$\hat{B}(X) = 1 - 2X^2 - 7X^4 - 50X^6 - 456X^8 - \dots$$

in Theorem 4 and hence

$$\hat{B}(\sqrt{5}X) = 1 - 10X^2 - 5^2 \cdot 7X^4 - 5^5 \cdot 2X^6 - 5^4 \cdot 456X^8 - \dots.$$

Here we take $H = \{\pm 1, \pm 7\}$ for $m = 25$ and $H = \{\pm 1, \pm 57\}$ for $m = 125$ again as in Example 2, where 25 first appears in $\xi_4(3)$ and 125 first appears in $\xi_4(7)$. The minimal polynomial for the reciprocal of $\psi_1 = \sqrt{5}(\zeta_{25} + \zeta_{25}^{-1} - \zeta_{25}^7 - \zeta_{25}^{-7})$ is

$$1 - 50X^2 + \underline{125}X^3 + \underline{125}X^4 - \underline{500}X^5,$$

whereas that for the reciprocal of $\psi_1 = \sqrt{5}(\zeta_{125} + \zeta_{125}^{-1} - \zeta_{125}^{57} - \zeta_{125}^{-57})$ begins

$$1 - 250X^2 + 5^4 \cdot 41X^4 - 5^7 \cdot 18X^7 - \underline{5}^7X^7 + \dots.$$

The underscored coefficients deviate as expected from the pattern given in Theorem 4.

One may have noticed that $\hat{\beta}_4(n) = \beta_4(n)$ in the last example. More generally,

Proposition 3. For $\nu > 1$, $\hat{\beta}_{2^\nu}(n)$ in (28) satisfies $\hat{\beta}_{2^\nu}(n) = \beta_{2^\nu}(n)$.

Proof. We first note that no sum of an **odd** number of 2^ν -roots of unity for $\nu \geq 1$ can vanish since each such root of unity is congruent to 1 mod π_2 , where π_2 is the unique prime above 2 in $\mathbf{Q}(\zeta_{2^\nu})$. Now we assert no sum of an **odd** number of primitive 2^ν -roots of unity can lie in $\mathbf{Q}(\zeta_{2^{\nu-1}})$ for $\nu > 1$. Suppose to the contrary a sum $\zeta_{2^\nu}^{i_1} + \cdots + \zeta_{2^\nu}^{i_n}$ lies in $\mathbf{Q}(\zeta_{2^{\nu-1}})$, where $\nu > 1$ and the i_j and n are odd. Then $\zeta_{2^\nu}(\zeta_{2^{\nu-1}}^{(i_1-1)/2} + \cdots + \zeta_{2^{\nu-1}}^{(i_n-1)/2})$ lies in $\mathbf{Q}(\zeta_{2^{\nu-1}})$ so $\zeta_{2^{\nu-1}}^{(i_1-1)/2} + \cdots + \zeta_{2^{\nu-1}}^{(i_n-1)/2} = 0$ contradicting our first remark. It follows easily now that for $\nu > 1$ any sum of an even number of 2^ν -roots of unity which vanishes must always contain an even number of primitive 2^ν -roots of unity. Hence $\beta_{2^\nu}^-(n) = 0$ for n even. The statement of the proposition readily follows. $\square$

We conclude this section with a remark concerning the case $f = 2$ which is not governed by the last proposition. One finds in this case that the counting function $\hat{\beta}_2(n) = \beta_2(n)$ in (27) whenever $p \equiv 1 \pmod{4}$, whereas $\hat{\beta}_2(n) = (-1)^{[n/2]} \beta_2(n)$ in (28) for $p \equiv 3 \pmod{4}$. In each case, the respective power series $\hat{B}(X) = \exp(-\sum_{n=1}^{\infty} \hat{\beta}_2(n) X^n/n)$ in Theorems 3 or 4 satisfies

$$\hat{B}(i^*X) = \exp\left(-\sum_{n=1}^{\infty} \binom{2n}{n} \frac{x^{2n}}{2n}\right) = \frac{1}{2}(1 + \sqrt{1 - 4X^2}).$$

In particular for the case $f = 2$, one finds for any odd prime power p^α , $\alpha > 1$ that the congruence

$$(30) \quad G(X) \equiv B(\sqrt{p}X)^{\phi(p^\alpha)/2} \pmod{X^N}$$

holds in $\mathbf{Z}[X]$ for $N = p^{\alpha-1}$ but not for $N > p^{\alpha-1}$, where

$$B(X) = \exp\left(-\sum_{n=1}^{\infty} \binom{2n}{n} \frac{X^{2n}}{2n}\right).$$

Indeed, we had previously remarked in Section 3 that p^α first appears in an exceptional set $\xi_2(n)$ for $n = p^{\alpha-1}$. One consequence is that the first $p^{\alpha-1}$ coefficients of $G(X)$ satisfy

$$(31) \quad c_r = \begin{cases} (-1)^{r/2} p^{r/2} \frac{\phi(p^\alpha)}{\phi(p^\alpha)-r} \binom{\phi(p^\alpha)/2 - r/2}{r/2} & \text{if } r \text{ is even} \\ 0 & \text{if } r \text{ is odd} \end{cases}$$

for $1 \leq r < p^{\alpha-1}$ (chiefly, Corollary 3 in [9]). While (31) is not enough to determine $G(X)$, we have recently found a closed form formula for the coefficients of $G(X)$ analogous to (25) expressed in terms of an Aurifeuillian factor [3] of the cyclotomic polynomial ψ_p . The reader is referred to [9] for details.

REFERENCES

- [1] B.C. Berndt, R.J. Evans and K.S. Williams, *Gauss and Jacobi sums*, Wiley-Interscience, New York, (1998). MR1625181 (99d:11092)
- [2] Z. Borevich and I. Shafarevich. *Number Theory*, Academic Press, New York, (1966). MR0195803 (33:4001)
- [3] R.P. Brent, "On computing factors of cyclotomic polynomials," Math. Comp. 61 (1993), 131-149. MR1205459 (93m:11131)
- [4] L.E. Dickson, *Elementary Theory of Equations*, Wiley, New York.
- [5] C.F. Gauss, *Disquisitiones Arithmeticae*, Yale University Press, New Haven, (1966). MR0197380 (33:5545)

- [6] S. Gupta and D. Zagier, "On the coefficients of the minimal polynomial of Gaussian periods," *Math. Comp.* 60 (1993), 385-398. MR1155574 (93d:11086)
- [7] S. Gurak, "Minimal polynomials for Gauss circulants and cyclotomic units," *Pac. J. Math.* 102 (1982), 347-353. MR0686555 (84c:10032)
- [8] S. Gurak, "Minimal polynomials for circular numbers," *Pac. J. Math.* 112 (1984), 313-331. MR0743988 (85i:11107)
- [9] S. Gurak, "Minimal polynomials for Gauss periods with $f=2$," *Acta Arith.* 121 (2006), 233-257.
- [10] S. Gurak, "Explicit evaluation of multi-dimensional Kloosterman sums for prime powers" (to appear).
- [11] H. Hasse, *Vorlesungen über Zahlentheorie*, Springer-Verlag, Berlin, (1950). MR0051844 (14:534c)
- [12] J. Neukirch, *Class Field Theory*, Springer-Verlag, New York, (1986). MR0819231 (87i:11005)
- [13] H. Salie, "Über die Kloostermanschen Summen $S(u,v;q)$," *Math. Z.* 34 (1932), 91-109. MR1545243

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SAN DIEGO, SAN DIEGO, CALIFORNIA 92110
E-mail address: `gurak@sandiego.edu`