THE EULER IMPLICIT/EXPLICIT SCHEME FOR THE 2D TIME-DEPENDENT NAVIER-STOKES EQUATIONS WITH SMOOTH OR NON-SMOOTH INITIAL DATA

YINNAN HE

ABSTRACT. This paper considers the stability and convergence results for the Euler implicit/explicit scheme applied to the spatially discretized two-dimensional (2D) time-dependent Navier-Stokes equations. A Galerkin finite element spatial discretization is assumed, and the temporal treatment is implicit/explicit scheme, which is implicit for the linear terms and explicit for the nonlinear term. Here the stability condition depends on the smoothness of the initial data $u_0 \in H^\alpha$, i.e., the time step condition is $\tau \leq C_0$ in the case of $\alpha = 2$, $\tau |\log h| \leq C_0$ in the case of $\alpha = 1$ and $\tau h^{-2} \leq C_0$ in the case of $\alpha = 0$ for mesh size h and some positive constant C_0 . We provide the H^2 -stability of the scheme under the stability condition with $\alpha = 0, 1, 2$ and obtain the optimal $H^1 - L^2$ error estimate of the numerical velocity and the optimal L^2 error estimate of the numerical pressure under the stability condition with $\alpha = 1, 2$.

1. INTRODUCTION

Let Ω be a bounded domain in R^2 assumed to have a Lipschitz continuous boundary $\partial\Omega$ and to satisfy a further condition stated in **(A1)** below. We consider the time-dependent Navier-Stokes problem

$$(1.1) \quad \begin{cases} u_t - \nu \Delta u + (u \cdot \nabla)u + \nabla p = f, \operatorname{div} u = 0, & (x, t) \in \Omega \times (0, T]; \\ u(x, 0) = u_0(x), \quad x \in \Omega; \quad u(x, t)|_{\partial\Omega} = 0, & t \in [0, T], \end{cases}$$

where $u = u(x, t) = (u_1(x, t), u_2(x, t))$ represents the velocity vector, $p = p(x, t)$ the pressure, $f = f(x, t)$ the prescribed body force, $u_0(x)$ the initial velocity, $\nu > 0$ the viscosity, and $T > 0$ a finite time.

There are numerous works devoted to the development of efficient schemes for the Navier-Stokes equations [3, 4, 9, 10, 11, 13, 14, 15, 16, 19, 20, 21, 23, 27, 6, 30, 32, 31, 37], fully implicit, semi-implicit and implicit/explicit scheme. A key issue is the stability conditions of schemes. Usually the fully implicit schemes are unconditionally stable. However, at each time step, one has to solve a system of nonlinear equations. An explicit scheme is much easier in computation. But it suffers the severely restricted time step size from stability requirement. A popular approach is based on an implicit scheme for the linear terms and a semi-implicit

Received by the editor February 26, 2007 and, in revised form, September 17, 2007.

2000 *Mathematics Subject Classification.* Primary 35L70, 65N30, 76D06.

Key words and phrases. Navier-Stokes equations, mixed finite element, Euler implicit/explicit scheme, Smooth or non-smooth initial data.

This research was subsidized by the NSF of China 10671154 and the National Basic Research Program under the grant 2005CB321703.

©2008 American Mathematical Society
Reverts to public domain 28 years from publication

scheme or an explicit scheme for the nonlinear term. A semi-implicit scheme for the nonlinear term results in a linear system with a variable coefficient matrix of time, and an explicit treatment for the nonlinear term gives a constant matrix. Stability and convergence conditions of schemes have been studied by many authors. The main results are summarized below, where we set $\Omega \subset R^d$ with $d = 2, 3$, and $0 < h < 1$ denotes the mesh size in the spatial direction and $0 < \tau = \frac{T}{N} < 1$ denotes the step size in the time direction, which may change. However, $T > 0$ is fixed throughout this paper.

- For the Crank-Nicolson scheme (fully implicit), Heywood and Rannacher [23] proved that it is almost unconditionally stable and convergent, i.e. stable and convergent when

$$(1.2) \quad \tau \leq C_0,$$

for some positive constant C_0 depending on the data (ν, Ω, T, u_0, f) in the case of $d = 2, 3$.

- For a two-step scheme (semi-implicit), He and Li [14] gave the following convergence condition:

$$(1.3) \quad \tau h^{-1/2} \leq C_0.$$

- For the Crank-Nicolson extrapolation scheme (semi-implicit), He [15] has proved that (1.2) is the stability and convergence condition of the scheme in the case of $d = 2$.
- For the Crank-Nicolson/Adams-Bashforth scheme (implicit/explicit), Marion and Temam provided in [32] the following stability condition:

$$(1.4) \quad \tau h^{-d} \leq C_0, \quad d = 2, 3,$$

and recently, Tone [37] proved the convergence under the condition

$$(1.5) \quad \tau h^{-2-d/2} \leq C_0, \quad d = 2, 3.$$

- A modified Crank-Nicolson/Adams-Bashforth scheme (implicit/explicit) was proposed by Johnston and Liu [26], in which the nonlinear term and pressure term are discretized explicitly. They claimed in their numerical simulations that the scheme is stable under the standard stability condition

$$(1.6) \quad \|u\|_{L^\infty} \tau h^{-1} \leq 1, \quad d = 2, 3.$$

No theoretical analysis has been given.

- For a three-step backward extrapolating scheme (implicit/explicit), Baker et al. [4] gave the convergence condition

$$(1.7) \quad \tau h^{-4/7} \leq C_0,$$

in the case of $d = 2, 3$.

- Clearly, the time-step condition

$$(1.8) \quad \tau h^{-r} \leq C_0,$$

for some $r > 0$ was imposed in these previous works when an implicit/explicit scheme is applied.

Recently, He and Sun [19] have improved the result of (1.8) and proved that the stability and convergence condition of the Crank-Nicolson/Adams-Bashforth scheme is (1.2).

This paper focuses on the Euler implicit/explicit scheme with a finite element approximation in spatial direction for solving the time-dependent Navier-Stokes equations in the case of $d = 2$, which were studied by Marion and Temam [32], Tone [37], Kim and Moin [27] and Issacson and Keller [25]. The scheme consists of using a finite element pair (X_h, M_h) for the spatial discretization, the implicit scheme for the linear term and the explicit scheme for the nonlinear term for the time discretization. Under the assumptions (A1), (A2) in §2 with $u^0 \in D(A^{\alpha/2})$, $\alpha = 0, 1, 2$ and (A3) in §3, we prove that the scheme is stable, i.e.,

$$(1.9) \quad \sigma^{2-\alpha}(t_m) \left(\left\| \frac{u_h^m - u_h^{m-1}}{\tau} \right\|_0^2 + \nu^2 \|A_h u_h^m\|_0^2 + \|p_h^m\|_0^2 \right) \leq \kappa, \quad 1 \leq m \leq N,$$

when the stability condition

$$(1.10) \quad \begin{cases} \tau \leq C_0, & \alpha = 2, \\ \tau |\log h| \leq C_0, & \alpha = 1, \\ \tau h^{-2} \leq C_0, & \alpha = 0. \end{cases}$$

is satisfied. Under the stability condition (1.10) with $\alpha = 1, 2$, we also provide the $H^1 - L^2$ optimal error estimate for the numerical velocity and the L^2 -optimal error estimate for the numerical pressure:

$$(1.11) \quad \|u(t_m) - u_h^m\|_{L^2}^2 \leq \kappa(\sigma^{-(2-\alpha)}(t_m)\tau^2 + \sigma^{-(2-\alpha)}(t_m)h^4),$$

$$(1.12) \quad \|u(t_m) - u_h^m\|_{H^1}^2 \leq \kappa(\sigma^{-(3-\alpha)}(t_m)\tau^2 + \sigma^{-(2-\alpha)}(t_m)h^2),$$

$$(1.13) \quad \|p(t_m) - p_h^m\|_{L^2}^2 \leq \kappa(\sigma^{-(4-\alpha)}(t_m)\tau^2 + \sigma^{-(3-\alpha)}(t_m)h^2),$$

for all $1 \leq m \leq N$. Here $\sigma(t) = \min\{1, t\}$, κ is some positive constant depending on the data (ν, Ω, T, u_0, f) , and A_h is a discrete Stokes operator.

Moreover, similar results were proved for the Euler implicit/explicit scheme which is applied to the spatial discretization based on the spectral Galerkin method by He [11, 12].

Remark 1.1. In the case of $\alpha = 2$, for the first order scheme (the Euler implicit/explicit scheme) we obtain the same H^1 -error bound of the numerical velocity and a better L^2 -error bound of the numerical pressure than the second order scheme (Crank-Nicolson scheme), excepting the L^2 -error estimate for the numerical velocity. Previously, Heywood and Rannacher in [23] provided the following error estimates for the numerical velocity and pressure:

$$(1.14) \quad \|u(t_m) - u_h^m\|_{H^1}^2 \leq \kappa(\sigma^{-1}(t_m)\tau^2 + h^2), \quad 1 \leq m \leq N,$$

$$(1.15) \quad \|p(t_m) - p_h^m\|_{L^2}^2 \leq \kappa(\sigma^{-3}(t_m)\tau^2 + \sigma^{-1}(t_m)h^2), \quad 1 \leq m \leq N,$$

and the L^2 -error estimate for the numerical velocity:

$$(1.16) \quad \|u(t_m) - u_h^m\|_{L^2}^2 \leq \kappa(\sigma^{-2}(t_m)\tau^4 + h^4), \quad t_m \in (0, T], \quad 1 \leq m \leq N.$$

This paper is organized as follows. In §2 an abstract functional setting of the Navier-Stokes problem is given together with some basic assumptions (A1) and (A2) with $\alpha = 0, 1, 2$. In §3 we set out our assumption (A3) concerning the finite element spaces X_h and M_h , finite element Galerkin approximation in space and some properties on the trilinear form $b(\cdot, \cdot, \cdot)$. Section 3 contains the optimal error estimate and a priori estimate results of the finite element solution $(u_h(t), p_h(t))$. In §4 we describe the Euler implicit/explicit scheme and prove the stability result of the scheme. In §5 we describe the dual scheme and prove its stability result. In §6 we obtain the optimal $H^1 - L^2$ -error estimate of the numerical velocity and the

optimal L^2 -error estimate of the numerical pressure under the stability condition (1.10) with $\alpha = 1, 2$.

2. FUNCTIONAL SETTING OF THE NAVIER-STOKES EQUATIONS

For the mathematical setting of problem (1.1), we introduce the following Hilbert spaces:

$$X = H_0^1(\Omega)^2, \quad Y = L^2(\Omega)^2, \quad M = L_0^2(\Omega) = \left\{ q \in L^2(\Omega); \int_{\Omega} q dx = 0 \right\}.$$

The space $L^2(\Omega)^d$, $d = 1, 2, 4$, is associated with the usual L^2 -scalar product $(\cdot, \cdot)$ and L^2 -norm $\|\cdot\|_0$. The space X is associated with its usual scalar product and equivalent norm

$$((u, v)) = (\nabla u, \nabla v), \quad \|u\|_X = \|\nabla u\|_0.$$

Next, let the closed subset V of X be given by

$$V = \{v \in X; \operatorname{div} v = 0\}$$

and denote by H the closed subset of Y , i.e.,

$$H = \{v \in Y; \operatorname{div} v = 0, v \cdot n|_{\partial\Omega} = 0\}.$$

We refer readers to [1, 10, 22, 36] for details on these spaces. We denote the Stokes operator by $A = -P\Delta$, where P is the L^2 -orthogonal projection of Y onto H and the domain of A by $D(A) = H^2(\Omega)^2 \cap V$. As mentioned above, we need a further assumption on Ω provided in [23].

(A1) Assume that Ω is smooth so that the unique solution $(v, q) \in (X, M)$ of the steady Stokes problem

$$-\nu\Delta v + \nabla q = g, \quad \operatorname{div} v = 0 \quad \text{in } \Omega, \quad v|_{\partial\Omega} = 0,$$

for any prescribed $g \in Y$, exists and satisfies

$$\|v\|_{H^2} + \|q\|_{H^1} \leq c\|g\|_0,$$

where $c > 0$ is a generic constant depending on Ω and ν , and may take different values at its different occurrences.

We remark that the validity of assumption (A1) is known (see [10, 22, 28, 36]) if $\partial\Omega$ is of C^2 or if Ω is a two-dimensional convex polygon. From the assumption (A1), it is well known [1, 22, 29] that

$$(2.1) \quad \|v\|_{H^2} \leq c\|Av\|_0, \quad v \in D(A),$$

$$(2.2) \quad \|v\|_0 \leq \gamma_0\|\nabla v\|_0, \quad v \in X, \quad \|\nabla v\|_0 \leq \gamma_0\|Av\|_0, \quad v \in D(A),$$

where γ_0 is a positive constant depending only on Ω . We usually make the following assumption about the prescribed data for problem (1.1).

(A2) The initial velocity $u_0(x)$ and the force $f(x, t)$ are such that $u_0 \in D(A^{\alpha/2})$, $f, f_t, f_{tt} \in L^\infty(0, T; Y)$ with

$$\|A^{\alpha/2}u_0\|_0 + \sup_{0 \leq t \leq T} \{\|f(t)\|_0 + \|f_t(t)\|_0 + \|f_{tt}(t)\|_0\} \leq C$$

for some positive constant C , and $\alpha = 0, 1, 2$, where $D(A^{\frac{1}{2}}) = V$ and $D(A^0) = H$.

Moreover, we define the continuous bilinear forms $a(\cdot, \cdot)$ and $d(\cdot, \cdot)$ on $X \times X$ and $X \times M$, respectively, by

$$a(u, v) = \nu((u, v)), \quad u, v \in X, \quad d(v, q) = (q, \operatorname{div} v), \quad v \in X, \quad q \in M,$$

and a trilinear form on $X \times X \times X$ by

$$\begin{aligned} b(u, v, w) &= ((u \cdot \nabla)v + \frac{1}{2}(\operatorname{div} u)v, w) \\ &= \frac{1}{2}((u \cdot \nabla)v, w) - \frac{1}{2}((u \cdot \nabla)w, v), \quad u, v, w \in X. \end{aligned}$$

With the above notation, the variational formulation of problem (1.1) reads as follows: Find $(u, p) \in (X, M)$ for all $t \in [0, T]$ such that for all $(v, q) \in (X, M)$,

$$(2.3) \quad (u_t, v) + a(u, v) - d(v, p) + d(u, q) + b(u, u, v) = (f, v),$$

$$(2.4) \quad u(0) = u_0.$$

In order to proceed the theoretical and numerical analysis for the variational formulation (2.3)-(2.4), we need to introduce the following existence, uniqueness and modified regularity results.

Theorem 2.1. *Under the assumptions (A1) and (A2), the problem (2.3)-(2.4) admits a unique solution $u \in L^\infty(0, T; H) \cap L^2(0, T; V)$ satisfying the following regularities:*

$$(2.5) \quad \begin{aligned} &\|u(t)\|_0^2 + \sigma^{\frac{1}{2}(1-\alpha)(2-\alpha)}(t) \|\nabla u(t)\|_0^2 + \sigma^{2-\alpha}(t) (\|Au(t)\|_0^2 + \|\nabla p(t)\|_0^2 + \|u_t(t)\|_0^2) \\ &+ \sigma^{3-\alpha}(t) \|\nabla u_t(t)\|_0^2 \leq \kappa, \end{aligned}$$

$$(2.6) \quad \begin{aligned} &\int_0^t \{ \|\nabla u\|_0^2 + \sigma^{\frac{1}{2}(1-\alpha)(2-\alpha)}(s) (\|u_t\|_0^2 + \|Au\|_0^2 + \|\nabla p\|_0^2) \} ds \\ &+ \int_0^t \{ \sigma^{2-\alpha}(s) \|\nabla u_t\|_0^2 + \sigma^{3-\alpha}(s) (\|u_{tt}\|_0^2 + \|Au_t\|_0^2 + \|\nabla p_t\|_0^2) \} ds \leq k, \end{aligned}$$

for all $0 \leq t \leq T$.

Proof. For the existence and uniqueness of the solution in the case of $\alpha = 0$, the reader may refer to Temam [36]. For the regularity results related to $\alpha = 2$, the reader may refer to Heywood and Rannacher [22], and for the regularity results related to $\alpha = 1$, the reader may refer to Hill and Süli [24] and He [11] and He et al. [17].

The case $\alpha = 0$ has been proved in [12], except for the estimates of $\|\nabla p(t)\|_0^2$ and $\|\nabla p_t\|_0^2$. However, these can be done by using (1.1) and some nonlinear term estimates. $\square$

3. FINITE ELEMENT GALERKIN APPROXIMATION

Let $h > 0$ be a real positive parameter. The finite element subspace (X_h, M_h) of (X, M) is characterized by $J_h = J_h(\Omega)$, a partitioning of Ω into triangles K or quadrilaterals K , assumed to be uniformly regular as $h \rightarrow 0$. For further details, the reader may refer to Ciarlet [7] and Girault and Raviart [10].

We define the subspace V_h of X_h given by

$$(3.1) \quad V_h = \left\{ v_h \in X_h; d(v_h, q_h) = 0, \forall q_h \in M_h \right\}.$$

Let $P_h : Y \rightarrow V_h$ denote the L^2 -orthogonal projection defined by

$$(P_h v, v_h) = (v, v_h), \quad v \in Y, \quad v_h \in V_h.$$

We assume that the couple (X_h, M_h) satisfies the following approximation properties:

(A3) For each $v \in H^2(\Omega)^2 \cap X$ and $q \in H^1(\Omega) \cap M$, there exist approximations $\pi_h v \in X_h$ and $\rho_h q \in M_h$ such that

$$(3.2) \quad \|\nabla(v - \pi_h v)\|_0 \leq ch\|Av\|_0, \quad \|q - \rho_h q\|_0 \leq ch\|\nabla q\|_0.$$

For each $v_h \in X_h$, one has the inverse inequality

$$(3.3) \quad \|\nabla v_h\|_0 \leq c_1 h^{-1} \|v_h\|_0, \quad v_h \in X_h;$$

and the so-called inf-sup inequality: For each $q_h \in M_h$, there exists $v_h \in X_h, v_h \neq 0$, such that

$$(3.4) \quad d(v_h, q_h) \geq c_2 \|q_h\|_0 \|\nabla v_h\|_0,$$

where c_1 and c_2 are positive constants depending on Ω .

We give an example of the spaces X_h and M_h such that the assumption (A3) is satisfied. Let Ω be a convex, polygonal domain in plane and $J_h = J_h(\Omega)$, a partitioning of $\bar{\Omega}$ into triangles K , assumed to be uniformly regular as $h \rightarrow 0$. For any nonnegative integer l , we denote by $P_l(K)$ the space of polynomials of degrees less than or equal to l on K .

Example 1 (Girault-Raviart [10]).

$$\begin{aligned} X_h &= \{v_h \in C^0(\bar{\Omega})^2 \cap X; v_h|_K \in P_2(K)^2, \forall K \in J_h\}, \\ M_h &= \{q_h \in M; q_h|_K \in P_0(K), \forall K \in J_h\}. \end{aligned}$$

Example 2 (Bercovier-Pironneau [5]). We consider the triangulation $J_{h/2}$ obtained by dividing each triangle of J_h into four triangles (by joining the mid-sides). We set

$$\begin{aligned} X_h &= \{v_h \in C^0(\bar{\Omega})^2 \cap X; v_h|_K \in P_1(K)^2, \forall K \in J_{h/2}\}, \\ M_h &= \{q_h \in C^0(\bar{\Omega}) \cap M; q_h|_K \in P_1(K), \forall K \in J_h\}. \end{aligned}$$

The following properties are classical (see [2, 10, 22, 24]):

$$(3.5) \quad \|\nabla P_h v\|_0 \leq c \|\nabla v\|_0, \quad v \in X,$$

$$(3.6) \quad \|v - P_h v\|_0 + h \|\nabla(v - P_h v)\|_0 \leq ch^2 \|Av\|_0, \quad v \in D(A),$$

$$(3.7) \quad \|v - P_h v\|_0 \leq ch \|\nabla(v - P_h v)\|_0, \quad v \in X.$$

The standard finite element Galerkin approximation of (2.3)–(2.4) based on (X_h, M_h) reads as follows: Find $(u_h, p_h) \in (X_h, M_h)$ such that for all $0 < t \leq T$ and $(v_h, q_h) \in (X_h, M_h)$,

$$(3.8) \quad (u_{ht}, v_h) + a(u_h, v_h) - d(v_h, p_h) + d(u_h, q_h) + b(u_h, u_h, v_h) = (f, v_h),$$

$$(3.9) \quad u_h(0) = u_0h = P_h u_0.$$

With the above statements, a discrete analogue $A_h = -P_h \Delta_h$ of the Stokes operator A is defined through the condition that $(-\Delta_h u_h, v_h) = ((u_h, v_h))$ for all $u_h, v_h \in X_h$. The restriction of A_h to V_h is invertible, with the inverse A_h^{-1} . Since A_h^{-1} is self-adjoint and positive definite, we may define “discrete” Sobolev norms on V_h , of any order $r \in R$, by setting

$$\|v_h\|_r = \|A_h^{r/2} v_h\|_0, \quad v_h \in V_h.$$

These norms will be assumed to have various properties similar to their continuous counterparts, an assumption that implicitly imposes conditions on the structure of the spaces X_h and M_h . In particular, it holds that

$$\|v_h\|_1 = \|\nabla v_h\|_0, \quad \|v_h\|_2 = \|A_h v_h\|_0, \quad v_h \in V_h.$$

By the way, we derive from (2.2) that

$$(3.10) \quad \|v_h\|_0 \leq \gamma_0 \|\nabla v_h\|_0, \quad \|\nabla v_h\|_0 \leq \gamma_0 \|A_h v_h\|_0, \quad v_h \in V_h,$$

where $\gamma_0 > 0$ is a constant depending only on Ω .

This section considers preliminary estimates which are useful in the error estimates of finite element solution. Some estimates of the trilinear form b are given in the following lemma and the proof can be found in [15, 16, 24].

Lemma 3.1. *The trilinear form b satisfies the following estimates:*

$$(3.11) \quad b(u, v_h, w_h) = ((u \cdot \nabla) v_h, w_h) = -((u \cdot \nabla) w_h, v_h),$$

$$(3.12) \quad b(u_h, v_h, w_h) = -b(u_h, w_h, v_h),$$

$$(3.13) \quad \begin{aligned} |b(u_h, v_h, w_h)| &+ |b(v_h, u_h, w_h)| + |b(w_h, u_h, v_h)| \\ &\leq c_0 |\log h|^{1/2} \|u_h\|_1 \|v_h\|_1 \|w_h\|_0, \\ |b(u_h, v_h, w_h)| &+ |b(v_h, u_h, w_h)| + |b(w_h, u_h, v_h)| \\ &\leq \frac{c_0}{2} \|u_h\|_0^{1/2} \|u_h\|_1^{1/2} \|v_h\|_1 \|w_h\|_0^{1/2} \|w_h\|_1^{1/2} \\ (3.14) \quad &+ \frac{c_0}{2} \|u_h\|_1 \|v_h\|_0^{1/2} \|v_h\|_1^{1/2} \|w_h\|_0^{1/2} \|w_h\|_1^{1/2}, \end{aligned}$$

for all $u \in V$, $u_h, v_h, w_h \in X_h$ and

$$(3.15) \quad \begin{aligned} |b(u_h, v_h, w_h)| &+ |b(v_h, u_h, w_h)| + |b(w_h, u_h, v_h)| \\ &\leq \frac{1}{2} c_0 \|A_h v_h\|_0^{1/2} \|v_h\|_1^{1/2} \|u_h\|_0^{1/2} \|u_h\|_1^{1/2} \|w_h\|_0 \\ &+ \frac{1}{2} c_0 \|A_h v_h\|_0^{1/2} \|v_h\|_0^{1/2} \|u_h\|_1 \|w_h\|_0, \end{aligned}$$

for all $u_h, v_h \in V_h$, $w_h \in X_h$, where $c_0 > 0$ is a constant depending only on Ω .

Before we proceed further, we need some continuous and discrete Gagliardo-Nirenberg estimates (see Temam [36] and Hill and Süli [24]).

Lemma 3.2. *It holds that*

$$(3.16) \quad \begin{aligned} \|v\|_{L^4} &\leq c \|v\|_0^{1/2} \|\nabla v\|_0^{1/2}, \quad \forall v \in X, \quad \|\nabla v\|_{L^4} \leq c \|\nabla v\|_0^{1/2} \|Av\|_0^{1/2}, \quad \forall v \in D(A), \\ \|v_h\|_{L^\infty} &\leq c \|v_h\|_0^{1/2} \|A_h v_h\|_0^{1/2}, \quad \|v_h\|_{L^\infty} \leq c |\log h|^{1/2} \|\nabla v_h\|_0, \quad \forall v_h \in V_h, \\ \|\nabla v_h\|_{L^4} &\leq c \|\nabla v_h\|_0^{1/2} \|A_h v_h\|_0^{1/2}, \quad \forall v_h \in V_h. \end{aligned}$$

In order to perform our error analysis for time discretization, we recall the following smooth properties of (u_h, p_h) .

Theorem 3.3. Assume that assumptions (A1)–(A3) are valid. Then the finite element solution (u_h, p_h) satisfies the following estimates:

$$(3.17) \quad \begin{aligned} & \|u_h(t)\|_0^2 + \sigma^{\frac{1}{2}(1-\alpha)(2-\alpha)}(t) \|\nabla u_h(t)\|_0^2 + \sigma^{2-\alpha}(t) \|A_h u_h(t)\|_0^2 \\ & + \int_0^t \{ \|\nabla u_h\|_0^2 + \sigma^{\frac{1}{2}(1-\alpha)(2-\alpha)}(s) \|A_h u_h\|_0^2 \} ds \leq \kappa, \end{aligned}$$

$$(3.18) \quad \begin{aligned} & \sigma^{2+r-\alpha}(t) \|u_{ht}(t)\|_r^2 \leq \kappa, \quad r = 0, 1, 2, \\ & \int_0^t \{ \sigma^{\frac{1}{2}(1-\alpha)(2-\alpha)}(s) \|u_{ht}\|_0^2 + \sigma^{1+r-\alpha}(s) \|u_{ht}\|_r^2 \} ds \leq k, \quad r = 1, 2, \end{aligned}$$

for all $0 \leq t \leq T$.

For the proof of Theorem 3.3 in the case of $\alpha = 2$, the reader is referred to Heywood and Rannacher [23] and He and Sun [19]. Theorem 3.3 with $\alpha = 1, 0$ can be proved in a manner similar to the one used in [23, 19].

Next, we can provide some bounds of the error $(u - u_h, p - p_h)$.

Theorem 3.4. Under the assumptions (A1), (A2) with $\alpha = 1, 2$ and (A3), it holds that

$$(3.19) \quad \begin{aligned} & \sigma^{2-\alpha}(t) \|u(t) - u_h(t)\|_0^2 + h^2 \sigma^{2-\alpha}(t) \|\nabla(u(t) - u_h(t))\|_0^2 \\ & + \sigma^{3-\alpha}(t) h^2 \|p(t) - p_h(t)\|_0^2 \leq \kappa h^4, \end{aligned}$$

for all $t \in (0, T]$.

Proof. For the case $\alpha = 2$, Heywood and Rannacher [22] have proved (3.19). For the case $\alpha = 1$, Hill and Süli [24] have proved

$$(3.20) \quad \begin{aligned} & (\sigma(t) + h^2) \|u(t) - u_h(t)\|_0^2 + h^2 \sigma(t) \|\nabla(u(t) - u_h(t))\|_0^2 \\ & + h^2 \int_0^t \|\nabla(u - u_h)\|_0^2 ds \leq \kappa h^4, \end{aligned}$$

for all $t \in (0, T]$.

Hence, it is sufficient to prove

$$(3.21) \quad \sigma(t) \|p(t) - p_h(t)\|_0 \leq \kappa h, \quad \forall t \in (0, T],$$

for $\alpha = 1$.

We set $e_h = P_h u - u_h$ and subtract (3.8) from (2.3) with $v = v_h$ to obtain

$$(3.22) \quad \begin{aligned} & (u_t - u_{ht}, v_h) + a(u - u_h, v_h) - d(v_h, p - p_h) + b(u, u - u_h, v_h) \\ & + b(u - u_h, u, v_h) - b(u - u_h, u - u_h, v_h) = 0, \quad \forall v_h \in X_h. \end{aligned}$$

Taking $v_h = 2e_{ht} \in V_h$ in (3.22) yields

$$(3.23) \quad \begin{aligned} & 2\|e_{ht}\|_0^2 + 2\nu \frac{d}{dt} \|\nabla(u - u_h)\|_0^2 + 2b(u, u - u_h, e_{ht}) \\ & + 2b(u - u_h, u, e_{ht}) - 2b(u - u_h, u - u_h, e_{ht}) \\ & = 2a(u - u_h, u_t - P_h u_t) + 2 \frac{d}{dt} d(e_h, p - p_h p) - 2d(e_h, p_t - \rho_h p_t). \end{aligned}$$

Due to (2.2), (3.2)-(3.3), (3.6) and Lemmas 3.1 and 3.2, we have

$$\begin{aligned}
2a(u - u_h, u_t - P_h u_t) &\leq 2\nu \|\nabla(u - u_h)\|_0 \|\nabla(u_t - P_h u_t)\|_0 \\
&\leq ch \|\nabla(u - u_h)\|_0 \|Au_t\|_0, \\
2|d(e_h, p_t - \rho_h p_t)| &\leq 2\sqrt{2} \|\nabla e_h\|_0 \|p_t - \rho_h p_t\|_0 \\
&\leq ch (\|\nabla(u - u_h)\|_0 + h \|Au\|_0) \|\nabla p_t\|_0, \\
2|b(u, u - u_h, e_{ht})| + 2|b(u - u_h, u, e_{ht})| \\
&\leq 4(\|u\|_{L^\infty} \|\nabla(u - u_h)\|_0 + \|\nabla u\|_{L^4} \|u - u_h\|_{L^4}) \|e_{ht}\|_0 \\
&\leq \frac{1}{2} \|e_{ht}\|_0^2 + c \|Au\|_0^2 \|\nabla(u - u_h)\|_0^2, \\
2|b(u - u_h, u - u_h, e_{ht})| &\leq c \|\nabla(u - u_h)\|_0^2 \|\nabla e_{ht}\|_0 \\
&\leq \frac{1}{2} \|e_{ht}\|_0^2 + ch^{-2} \|\nabla(u - u_h)\|_0^4.
\end{aligned}$$

Combining this inequality with (3.23) gives

$$\begin{aligned}
\|e_{ht}\|_0^2 &+ 2\nu \frac{d}{dt} \|\nabla(u - u_h)\|_0^2 \leq 2 \frac{d}{dt} d(e_h, p - \rho_h p) \\
&+ ch \|\nabla(u - u_h)\|_0 \|Au_t\|_0 + ch (\|\nabla(u - u_h)\|_0 + h \|Au\|_0) \|\nabla p_t\|_0 \\
(3.24) \quad &+ c(\|Au\|_0^2 + h^{-2} \|\nabla(u - u_h)\|_0^2) \|\nabla(u - u_h)\|_0^2.
\end{aligned}$$

Multiplying (3.24) by $\sigma(t)$, and integrating with respect to time and then using Theorem 2.1 and (3.20), we obtain

$$\begin{aligned}
\int_0^t \sigma(s) \|e_{ht}\|_0^2 ds &\leq 2\sigma(t) d(e_h(t), p(t) - \rho_h p(t)) \\
&+ 2 \int_0^t |d(e_h, p - \rho_h p)| ds + 2\nu \int_0^t \|\nabla(u - u_h)\|_0^2 ds + \kappa h^2 \\
&\leq c\sigma(t) h (\|\nabla(u - u_h)\|_0 + h \|Au\|_0) \|\nabla p\|_0 \\
(3.25) \quad &+ h \int_0^t (\|\nabla(u - u_h)\|_0 + h \|Au\|_0) \|\nabla p\|_0 ds + \kappa h^2 \leq \kappa h^2,
\end{aligned}$$

for all $t \in (0, T]$.

Differentiating (3.22) with respect to time gives

$$\begin{aligned}
(u_{tt} - u_{htt}, v_h) &+ a(u_t - u_{ht}, v_h) - d(v_h, p_t - \rho_h p_t) + b(u_t, u - u_h, v_h) \\
&+ b(u - u_h, u_t, v_h) + b(u, u_t - u_{ht}, v_h) + b(u_t - u_{ht}, u, v_h) \\
(3.26) \quad &- b(u_t - u_{ht}, u - u_h, v_h) - b(u - u_h, u_t - u_{ht}, v_h) = 0, \quad \forall v_h \in V_h.
\end{aligned}$$

Taking $v_h = 2e_{ht} \in V_h$ in (3.26) and using Lemma 3.1, one finds

$$\begin{aligned}
\frac{d}{dt} \|e_{ht}\|_0^2 &+ \nu \|\nabla(u_t - u_{ht})\|_0^2 + \nu \|\nabla e_{ht}\|_0^2 + 2b(u_t, u - u_h, e_{ht}) + 2b(u - u_h, u_t, e_{ht}) \\
&+ 2b(u, u_t - P_h u_t, e_{ht}) + 2b(u_t - P_h u_t, u, e_{ht}) + 2b(e_{ht}, u_h, e_{ht}) \\
&- 2b(u_t - P_h u_t, u - u_h, e_{ht}) - 2b(u - u_h, u_t - P_h u_t, e_{ht}) \\
(3.27) \quad &= \nu \|\nabla(u_t - P_h u_t)\|_0^2 + 2d(e_{ht}, p_t - \rho_h p_t).
\end{aligned}$$

Due to (2.2), (3.2), (3.6) and Lemma 3.1, we have

$$\begin{aligned}
& 2|b(u_t, u - u_h, e_{ht})| + 2|b(u - u_h, u_t, e_{ht})| \\
& \leq 8\gamma_0 \|\nabla u_t\|_0 \|\nabla(u - u_h)\|_0 \|\nabla e_{ht}\|_0 \\
& \leq \frac{\nu}{8} \|\nabla e_{ht}\|_0^2 + c \|\nabla u_t\|_0^2 \|\nabla(u - u_h)\|_0^2, \\
& 2|b(e_{ht}, u_h, e_{ht})| \leq 4 \|e_{ht}\|_0^{1/2} \|\nabla e_{ht}\|_0^{3/2} \|u_h\|_0^{1/2} \|\nabla u_h\|_0^{1/2} \\
& \leq \frac{\nu}{8} \|\nabla e_{ht}\|_0^2 + c \|u_h\|_0^2 \|\nabla u_h\|_0^2 \|e_{ht}\|_0^2, \\
& 2|b(u, u_t - P_h u_t, e_{ht})| + 2|b(u_t - P_h u_t, u, e_{ht})| \\
& \leq 8\gamma_0 \|\nabla(u_t - P_h u_t)\|_0 \|\nabla u\|_0 \|\nabla e_{ht}\|_0 \\
& \leq \frac{\nu}{8} \|\nabla e_{ht}\|_0^2 + ch^2 \|\nabla u\|_0^2 \|Au_t\|_0^2, \\
& 2|b(u_t - P_h u_t, u - u_h, e_{ht})| + 2|b(u - u_h, u_t - P_h u_t, e_{ht})| \\
& \leq 8\gamma_0 \|\nabla(u - u_h)\|_0 \|\nabla(u_t - P_h u_t)\|_0 \|\nabla e_{ht}\|_0 \\
& \leq \frac{\nu}{8} \|\nabla e_{ht}\|_0^2 + ch^2 \|\nabla(u - u_h)\|_0^2 \|Au_t\|_0^2, \\
& \nu \|\nabla(u_t - P_h u_t)\|_0^2 + 2d(e_{ht}, p_t - \rho_h p_t) \leq ch^2 \|Au_t\|_0^2 + ch \|\nabla e_{ht}\|_0 \|\nabla p_t\|_0 \\
& \leq \frac{\nu}{8} \|\nabla e_{ht}\|_0^2 + ch^2 (\|Au_t\|_0^2 + \|\nabla p_t\|_0^2).
\end{aligned}$$

Combining (3.27) with the above estimates yields

$$\begin{aligned}
(3.28) \quad \frac{d}{dt} \|e_{ht}\|_0^2 & \leq c \|u_h\|_0^2 \|\nabla u_h\|_0^2 \|e_{ht}\|_0^2 + c \|\nabla u_t\|_0^2 \|\nabla(u - u_h)\|_0^2 \\
& + ch^2 (1 + \|\nabla u\|_0^2 + \|\nabla u_h\|_0^2) (\|Au_t\|_0^2 + \|\nabla p_t\|_0^2),
\end{aligned}$$

Multiplying (3.28) by $\sigma^2(t)$, and integrating with respect to time, we obtain

$$\begin{aligned}
(3.29) \quad \sigma^2(t) \|e_{ht}(t)\|_0^2 & \leq c \int_0^t \sigma(s) (1 + \|u_h\|_0^2 \|\nabla u_h\|_0^2) \|e_{ht}\|_0^2 ds \\
& + c \int_0^t \sigma^2(s) \|\nabla u_t\|_0^2 \|\nabla(u - u_h)\|_0^2 ds \\
& + ch^2 \int_0^t \sigma^2(s) (1 + \|\nabla u_h\|_0^2 + \|\nabla u\|_0^2) (\|Au_t\|_0^2 + \|\nabla p_t\|_0^2) ds.
\end{aligned}$$

Using (3.20), (3.25), Theorem 2.1 and Theorem 3.3 in (3.29), we obtain

$$\sigma^2(t) \|e_{ht}(t)\|_0^2 \leq \kappa h^2,$$

which yields

$$\begin{aligned}
(3.30) \quad \sigma^2(t) \|u_t - u_{ht}\|_0^2 & \leq 2\sigma^2(t) \|e_{ht}(t)\|_0^2 + 2\sigma^2(t) \|u_t(t) - P_h u_t(t)\|_0^2 \\
& \leq 2\sigma^2(t) \|e_{ht}(t)\|_0^2 + ch^2 \sigma^2(t) \|\nabla u_t(t)\|_0^2 \leq \kappa h^2.
\end{aligned}$$

Finally, by using (2.2), (3.2), (3.4), (3.22) and Lemma 3.2, one finds

$$\begin{aligned}
(3.31) \quad \sigma(t) \|p(t) - p_h(t)\|_0 & \leq \sigma(t) (\|\rho_h p(t) - p(t)\|_0 + \|\rho_h p(t) - p_h(t)\|_0) \\
& \leq c\sigma(t) \|u_t(t) - u_{ht}(t)\|_0 + ch\sigma(t) \|\nabla p(t)\|_0 \\
& + c\sigma(t) (1 + \|\nabla u\|_0 + \|\nabla(u - u_h)\|_0) \|\nabla(u(t) - u_h(t))\|_0.
\end{aligned}$$

Using (3.20), (3.30) and Theorem 2.1 in (3.31), we obtain (3.21). $\square$

We will frequently use a discrete version of the Gronwall lemmas used in [13] and [34].

Lemma 3.5. *Let C , τ , and a_n , b_n , d_n , for integers $n \geq 0$, be nonnegative numbers such that*

$$(3.32) \quad a_m + \tau \sum_{n=1}^m b_n \leq \tau \sum_{n=0}^{m-1} a_n d_n + C, \quad m \geq 1.$$

Then

$$(3.33) \quad a_m + \tau \sum_{n=1}^m b_n \leq C \exp \left(\tau \sum_{n=0}^{m-1} d_n \right), \quad m \geq 1.$$

Theorem 3.6. *Under the assumptions (A1), (A2) with $\alpha = 1, 2$ and (A3), u_{htt} and u_{httt} satisfy the following bounds:*

$$(3.34) \quad \int_0^t \sigma^{3-r-\alpha}(s) \|A_h^{-r/2} u_{htt}\|_0^2 ds \leq \kappa, \quad r = 0, 1, 2, \quad \alpha = 1 \text{ or } r = 0, 1, \quad \alpha = 2,$$

$$(3.35) \quad \sigma^{4-\alpha}(t) \|u_{htt}(t)\|_0^2 + \int_0^t \sigma^{4-\alpha}(s) (\|u_{htt}\|_1^2 + \|A_h^{-1/2} u_{httt}\|_0^2) ds \leq \kappa,$$

for all $0 \leq t \leq T$.

Proof. Differentiating (3.8) with respect to t gives

$$(3.36) \quad (u_{htt}, v_h) + a(u_{ht}, v_h) + b(u_{ht}, u_h, v_h) + b(u_h, u_{ht}, v_h) = (f_t, v_h),$$

for all $v_h \in V_h$.

In view of (3.10) and Lemma 3.1, we deduce from (3.36) that

$$\|A_h^{-r/2} u_{htt}\|_0 \leq (\nu + c_0 \gamma_0 \|\nabla u_h\|_0) \|A_h^{1-r/2} u_{ht}\|_0 + \gamma_0^r \|f_t\|_0,$$

which yields

$$(3.37) \quad \begin{aligned} \int_0^t \sigma^{3-r-\alpha}(s) \|A_h^{-r/2} u_{htt}\|_0^2 ds &\leq c \int_0^t (1 + \|\nabla u_h\|_0^2) \sigma^{3-r-\alpha}(s) \|A_h^{1-r/2} u_{ht}\|_0^2 ds \\ &+ c \int_0^t \|f_t\|_0^2 ds, \end{aligned}$$

for $r = 0, 1, 2$, $\alpha = 1$ or $r = 0, 1$, $\alpha = 2$. Using Theorem 3.3 in (3.37) gives (3.34).

Furthermore, by differentiating (3.36) with respect to t gives

$$(3.38) \quad \begin{aligned} (u_{httt}, v_h) &+ a(u_{htt}, v_h) + 2b(u_{ht}, u_{ht}, v_h) + b(u_{htt}, u_h, v_h) \\ &+ b(u_h, u_{htt}, v_h) = (f_{tt}, v_h), \end{aligned}$$

for all $v_h \in V_h$.

Taking $v_h = 2u_{htt}$ in (3.38) and using (3.10) and Lemma 3.1, we deduce

$$(3.39) \quad \begin{aligned} \frac{d}{dt} \|u_{htt}\|_0^2 &+ 2\nu \|u_{htt}\|_1^2 + 4b(u_{ht}, u_{ht}, u_{htt}) + 2b(u_{htt}, u_h, u_{htt}) \\ &\leq \frac{\nu}{4} \|u_{htt}\|_1^2 + 4\nu^{-1} \gamma_0^2 \|f_{tt}\|_0^2. \end{aligned}$$

In view of (3.10) and Lemma 3.1, we deduce from (3.36) that

$$\begin{aligned} 4|b(u_{ht}, u_{ht}, u_{htt})| &\leq 2c_0\gamma_0\|u_{ht}\|_1^2\|u_{htt}\|_1 \\ &\leq \frac{\nu}{4}\|u_{htt}\|_1^2 + 4\nu^{-1}c_0^2\gamma_0^2\|u_{ht}\|_1^4, \\ 2|b(u_{htt}, u_h, u_{htt})| &\leq c_0\gamma_0\|u_{htt}\|_0\|u_{htt}\|_1\|A_h u_h\|_0 \\ &\leq \frac{\nu}{4}\|u_{htt}\|_1^2 + \nu^{-1}c_0^2\gamma_0^2\|A_h u_h\|_0^2\|u_{htt}\|_0^2. \end{aligned}$$

Combining these inequalities with (3.39) gives

$$\begin{aligned} \frac{d}{dt}\|u_{htt}\|_0^2 + \nu\|u_{htt}\|_1^2 &\leq 4\nu^{-1}\gamma_0^2\|f_{tt}\|_0^2 \\ (3.40) \quad &+ 4\nu^{-1}c_0^2\gamma_0^2\|u_{ht}\|_1^4 + \nu^{-1}c_0^2\gamma_0^2\|A_h u_h\|_0^2\|u_{htt}\|_0^2. \end{aligned}$$

Multiplying (3.40) by $\sigma^{4-\alpha}(t)$ yields

$$\begin{aligned} \frac{d}{dt}(\sigma^{4-\alpha}(t)\|u_{htt}\|_0^2) + \nu\sigma^{4-\alpha}(t)\|u_{htt}\|_1^2 &\leq 4\nu^{-1}\gamma_0^2\|f_{tt}\|_0^2 \\ (3.41) \quad &+ c\sigma^{4-\alpha}(t)\|u_{ht}\|_1^4 + (4-\alpha + c\sigma(t)\|A_h u_h\|_0^2)\sigma^{3-\alpha}(t)\|u_{htt}\|_0^2. \end{aligned}$$

Integrating (3.41) from 0 to t and using (3.34) and Theorem 3.3, we deduce

$$(3.42) \quad \sigma^{4-\alpha}(t)\|u_{htt}\|_0^2 + \nu \int_0^t \sigma^{4-\alpha}(s)\|u_{htt}\|_1^2 ds \leq \kappa, \quad \forall t \in (0, T].$$

Finally, it follows from (3.38), (3.10) and Lemma 3.1 that

$$\begin{aligned} \int_0^t \sigma^{4-\alpha}(s)\|A_h^{-1/2}u_{htt}\|_0^2 ds &\leq c \int_0^t (1 + \|u_h\|_1^2)\sigma^{4-\alpha}(s)\|u_{htt}\|_1^2 ds \\ (3.43) \quad &+ c \int_0^t \{\sigma^{4-\alpha}(s)\|u_{ht}\|_1^4 + \|f_{tt}\|_0^2\} ds. \end{aligned}$$

Using Theorem 3.3 in (3.43), together with (3.42), gives (3.35) for $\alpha = 1, 2$. $\square$

4. THE EULER IMPLICIT/EXPLICIT SCHEME

In this section we consider the time discretization of the finite element Galerkin approximation (3.8)-(3.9). Usually for the fully implicit scheme, at each time step, one has to solve a system of nonlinear equations. An explicit scheme is much easier in computation. But it suffers the severely restricted time step size from stability requirement. A popular approach is based on an implicit scheme for the linear terms and an explicit scheme for the nonlinear term. An explicit scheme for the nonlinear term results in a linear system with a constant coefficient matrix such that the computation is easy and the time step restriction is $\tau \leq C_0$ which will be proved in this section and Section 6.

Let $t_n = n\tau$ ($n = 0, 1, \dots, N$), $\tau = \frac{T}{N}$ the time step size, and N an integer. We define $u_h^0 = u_{0h} = P_h u_0$ and $(u_h^n, p_h^n) \in (X_h, M_h)$ by the Euler implicit/explicit scheme:

$$(4.1) \quad (d_t u_h^n, v_h) + a(u_h^n, v_h) - d(v_h, p_h^n) + d(u_h^n, q_h) + b(u_h^{n-1}, u_h^{n-1}, v_h) = (f(t_n), v_h),$$

here $d_t u_h^n = \frac{1}{\tau}(u_h^n - u_h^{n-1})$.

We see from (3.3) and (3.5)-(3.6) that

$$(4.2) \quad \|u_h^0\|_\alpha = \|u_{0h}\|_\alpha = \|P_h u_0\|_\alpha \leq c_\alpha \|A^{\alpha/2} u_0\|_0,$$

if $u_0 \in D(A^{\alpha/2})$ for some constants c_α with $\alpha = 0, 1, 2$.

The following theorem provides the stability of the scheme (4.1).

Theorem 4.1. *Suppose that the assumptions (A1)-(A3) are valid and $0 < \tau < 1$ satisfies the following stability condition:*

$$(4.3) \quad G_h \tau \leq \nu, \quad G_h = \begin{cases} 4^2 \nu^{-3/2} c_0^2 \gamma_0 \kappa_1^{1/2} \kappa_2^{1/2}, & \alpha = 2, \\ 4^2 c_0^2 \nu^{-1} \kappa_1 |\log h|, & \alpha = 1, \\ 4^2 c_0^2 c_1^2 \kappa_0 h^{-2}, & \alpha = 0. \end{cases}$$

Then the following hold:

$$(4.4) \quad \|u_h^m\|_0^2 + \nu \tau \sum_{n=1}^m \|u_h^n\|_1^2 \leq \kappa_0,$$

$$(4.5) \quad \begin{aligned} & \tau \sum_{n=1}^m \sigma^{\frac{1}{2}(1-\alpha)(2-\alpha)}(t_n) (\nu^2 \|A_h u_h^n\|_0^2 + \nu \|d_t u_h^n\|_1^2 \tau + \|d_t u_h^n\|_0^2) \\ & + \sigma^{\frac{1}{2}(1-\alpha)(2-\alpha)}(t_m) \nu \|u_h^m\|_1^2 \leq \kappa_1, \end{aligned}$$

$$(4.6) \quad \begin{aligned} & \sigma^{2-\alpha}(t_m) (\|d_t u_h^m\|_0^2 + \|p_h^m\|_0^2) + \nu \tau \sum_{n=1}^m \sigma^{2-\alpha}(t_n) \|d_t u_h^n\|_1^2 \leq \kappa_2, \\ & \sigma^{2-\alpha}(t_m) \nu^2 \|A_h u_h^m\|_0^2 \leq \kappa_2, \end{aligned}$$

for all $0 \leq m \leq N$, where $\kappa_\alpha \geq c_\alpha^2 \nu^\alpha \|A^{\alpha/2} u_0\|_0^2$ are some positive constants depending on the data (ν, Ω, T, u_0, f) .

Proof. First, taking $v_h = 2u_h^n \tau \in V_h$ and $v_h = A_h u_h^n \tau + \nu^{-1} d_t u_h^n \tau \in V_h$, respectively, and $q_h = 0$ in (4.1) and using (3.12) and the relation

$$(4.7) \quad 2(x - y)x = |x|^2 - |y|^2 + |x - y|^2, \quad \forall x, y \in R^2,$$

we obtain

$$(4.8) \quad \begin{aligned} \|u_h^n\|_0^2 & - \|u_h^{n-1}\|_0^2 + \|d_t u_h^n\|_0^2 \tau^2 + 2\nu \|u_h^n\|_1^2 \tau + 2b(u_h^{n-1}, u_h^n, d_t u_h^n) \tau^2 \\ & = 2(f(t_n), u_h^n) \tau, \end{aligned}$$

$$(4.9) \quad \begin{aligned} \|u_h^n\|_1^2 & - \|u_h^{n-1}\|_1^2 + \|d_t u_h^n\|_1^2 \tau^2 + \nu^{-1} \|d_t u_h^n\|_0^2 \tau + \nu \|A_h u_h^n\|_0^2 \tau \\ & + b(u_h^{n-1}, u_h^{n-1}, A_h u_h^n + \nu^{-1} d_t u_h^n) \tau \\ & = (f(t_n), A_h u_h^n + \nu^{-1} d_t u_h^n) \tau. \end{aligned}$$

In view of Lemma 3.1 and (3.10), it holds that

$$\begin{aligned} 2|b(u_h^{n-1}, u_h^{n-1}, u_h^n)| \tau & = 2|b(u_h^{n-1}, u_h^n, d_t u_h^n)| \tau^2 \leq \frac{1}{2} G^{1/2}(u_h^{n-1}) \|u_h^n\|_1 \|d_t u_h^n\|_0 \tau^2 \\ & \leq \frac{1}{2} \|d_t u_h^n\|_0^2 \tau^2 + \frac{1}{4} G(u_h^{n-1}) \|u_h^n\|_1^2 \tau^2, \\ 2|(f(t_n), u_h^n)| \tau & \leq \frac{\nu}{4} \|u_h^n\|_1^2 \tau + 4\nu^{-1} \gamma_0^2 \|f(t_n)\|_0^2 \tau, \end{aligned}$$

and

$$\begin{aligned}
& |b(u_h^{n-1}, u^n - u_h^{n-1}, A_h u_h^n + \nu^{-1} d_t u_h^n)|\tau \\
& \leq \frac{1}{2} G^{1/2}(u_h^{n-1}) \|d_t u_h^n\|_1 \|A_h u_h^n + \nu^{-1} d_t u_h^n\|_0 \tau^2 \\
& \leq \frac{1}{2} \|d_t u_h^n\|_1^2 \tau^2 + \frac{1}{4} G(u_h^{n-1}) (\|A_h u_h^n\|_0^2 + \nu^{-2} \|d_t u_h^n\|_0^2) \tau^2, \\
& |b(u_h^{n-1}, u_h^n, A_h u_h^n + \nu^{-1} d_t u_h^n)|\tau \\
& \leq c_0 (\|u_h^{n-1}\|_0^{1/2} \|u_h^{n-1}\|_1^{1/2} \|u_h^n\|_1^{1/2} + \|u_h^{n-1}\|_1 \|u_h^n\|_0^{1/2}) \\
& \quad \times \|A_h u_h^n\|_0^{1/2} (\|A_h u_h^n\|_0 + \nu^{-1} \|d_t u_h^n\|_0) \tau \\
& \leq \frac{\nu}{8} \|A_h u_h^n\|_0^2 \tau + \frac{1}{8\nu} \|d_t u_h^n\|_0^2 \tau \\
& \quad + 2\left(\frac{4}{\nu}\right)^3 c_0^4 (\|u_h^{n-1}\|_0^2 \|u_h^n\|_1^2 + \|u_h^{n-1}\|_1^2 \|u_h^n\|_0^2) \|u_h^{n-1}\|_1^2 \tau, \\
& |(f(t_n), A_h u_h^n + \nu^{-1} d_t u_h^n)|\tau \leq \frac{\nu}{8} \|A_h u_h^n\|_0^2 \tau + \frac{1}{8\nu} \|d_t u_h^n\|_0^2 \tau + 4\nu^{-1} \|f(t_n)\|_0^2 \tau,
\end{aligned}$$

where

$$(4.10) \quad G(u_h^n) = \begin{cases} 4^2 c_0^2 \gamma_0 \|u_h^n\|_1 \|A_h u_h^n\|_0, & \alpha = 2, \\ 4^2 c_0^2 |\log h| \|u_h^n\|_1^2, & \alpha = 1, \\ 4^2 c_0^2 c_1^2 h^{-2} \|u_h^n\|_0^2, & \alpha = 0. \end{cases}$$

Combining these inequalities with (4.8) and (4.9) yields

$$(4.11) \quad \begin{aligned} & \|u_h^n\|_0^2 - \|u_h^{n-1}\|_0^2 + \frac{1}{2} \|d_t u_h^n\|_0^2 \tau^2 + \nu \|u_h^n\|_1^2 \tau + \frac{1}{2} (\nu - G(u_h^{n-1})\tau) \|u_h^n\|_1^2 \tau \\ & \leq 4\nu^{-1} \gamma_0^2 \|f(t_n)\|_0^2 \tau, \end{aligned}$$

$$(4.12) \quad \begin{aligned} & 2\nu \|u_h^n\|_1^2 - 2\nu \|u_h^{n-1}\|_1^2 + \|d_t u_h^n\|_1^2 \tau^2 + \|d_t u_h^n\|_0^2 \tau + \nu^2 \|A_h u_h^n\|_0^2 \tau \\ & \quad + \frac{\nu}{2} (\nu - G(u_h^{n-1})\tau) (\|A_h u_h^n\|_0^2 + \nu^{-2} \|d_t u_h^n\|_0^2) \tau \\ & \leq d_{n-1} \nu \|u_h^{n-1}\|_1^2 \tau + 8\nu^{-1} \|f(t_n)\|_0^2 \tau, \end{aligned}$$

where

$$d_{n-1} = 4\left(\frac{4}{\nu}\right)^3 c_0^4 (\|u_h^{n-1}\|_0^2 \|u_h^n\|_1^2 + \|u_h^{n-1}\|_1^2 \|u_h^n\|_0^2).$$

Now, we define $d_t u_h^0 = \lim_{t \rightarrow 0} u_{ht}(t)$ through (3.8), i.e.,

$$(4.13) \quad (d_t u_h^0, v_h) + a(u_h^0, v_h) + b(u_h^0, u_h^0, v_h) = (f(t_0), v_h),$$

for all $v_h \in V_h$. Then, we deduce from (4.1) and (4.13) that

$$(4.14) \quad (d_{tt} u_h^1, v_h) + a(d_t u_h^1, v_h) = \frac{1}{\tau} \int_{t_0}^{t_1} (f_t(t), v_h) dt,$$

and

$$(4.15) \quad \begin{aligned} & (d_{tt} u_h^n, v_h) + a(d_t u_h^n, v_h) + b(d_t u_h^{n-1}, u_h^{n-1}, v_h) + b(u_h^{n-2}, d_t u_h^{n-1}, v_h) \\ & = \frac{1}{\tau} \int_{t_{n-1}}^{t_n} (f_t(t), v_h) dt, \end{aligned}$$

for all $2 \leq n \leq N$. Hence, it follows from (4.14) that

$$(4.16) \quad \|d_t u_h^1\|_0^2 + \|d_{tt} u_h^1\|_0^2 \tau^2 + \nu \|d_t u_h^1\|_1^2 \tau \leq \|d_t u_h^0\|_0^2 + \frac{\gamma_0^2}{\nu} \int_{t_0}^{t_1} \|f_t\|_0^2 dt.$$

Next, by taking $v_h = 2d_t u_h^n \tau$ in (4.15) and using (3.12), we deduce

$$(4.17) \quad \begin{aligned} \|d_t u_h^n\|_0^2 & - \|d_t u_h^{n-1}\|_0^2 + \|d_{tt} u_h^n\|_0^2 \tau^2 + 2\nu \|d_t u_h^n\|_1^2 \tau \\ & + 2b(d_t u_h^{n-1}, u_h^{n-1}, d_t u_h^n) \tau + 2b(u_h^{n-2}, d_t u_h^n, d_{tt} u_h^n) \tau^2 \\ & = 2 \left(\int_{t_{n-1}}^{t_n} f_t(t), d_t u_h^n \right) dt. \end{aligned}$$

In view of Lemma 3.1 and (3.10), it holds that

$$\begin{aligned} 2|b(d_t u_h^n, u_h^{n-1}, d_t u_h^n)| \tau & \leq c_0 \|u_h^{n-1}\|_0^{1/2} \|u_h^{n-1}\|_1^{1/2} \|d_t u_h^n\|_0^{1/2} \|d_t u_h^n\|_1^{3/2} \tau \\ & \leq \frac{\nu}{4} \|d_t u_h^n\|_1^2 \tau + \left(\frac{2}{\nu}\right)^3 c_0^4 \|u_h^{n-1}\|_0^2 \|u_h^{n-1}\|_1^2 \|d_t u_h^n\|_0^2 \tau, \\ 2|b(d_t u_h^n - d_t u_h^{n-1}, u_h^{n-1}, d_t u_h^n)| \tau & \leq \frac{1}{2} G^{1/2}(u_h^{n-1}) \|d_t u_h^n\|_1 \|d_{tt} u_h^n\|_0 \tau^2 \\ & \leq \frac{1}{4} \|d_{tt} u_h^n\|_0^2 \tau^2 + \frac{1}{4} G(u_h^{n-1}) \|d_t u_h^n\|_1^2 \tau^2, \\ 2|b(u_h^{n-2}, d_t u_h^n, d_{tt} u_h^n)| \tau^2 & \leq \frac{1}{2} G^{1/2}(u_h^{n-2}) \|d_t u_h^n\|_1 \|d_{tt} u_h^n\|_0 \tau^2 \\ & \leq \frac{1}{4} \|d_{tt} u_h^n\|_0^2 \tau^2 + \frac{1}{4} G(u_h^{n-2}) \|d_t u_h^n\|_1^2 \tau^2, \\ 2 \left| \int_{t_{n-1}}^{t_n} (f_t(t), d_t u_h^n) dt \right| & \leq \frac{\nu}{4} \|d_t u_h^n\|_1^2 \tau + \frac{4\gamma_0^2}{\nu} \int_{t_{n-1}}^{t_n} \|f_t(t)\|_0^2 dt. \end{aligned}$$

Combining these inequalities with (4.17) yields

$$(4.18) \quad \begin{aligned} \|d_t u_h^n\|_0^2 & - \|d_t u_h^{n-1}\|_0^2 + \frac{1}{2} \|d_{tt} u_h^n\|_0^2 \tau^2 + \nu \|d_t u_h^n\|_1^2 \tau \\ & + \frac{1}{4} (2\nu - G(u_h^{n-1}) \tau - G(u_h^{n-2}) \tau) \|d_t u_h^n\|_1^2 \tau \\ & \leq 2 \left(\frac{4}{\nu}\right)^3 c_0^4 \|u_h^{n-1}\|_0^2 \|u_h^{n-1}\|_1^2 \|d_t u_h^n\|_0^2 \tau + \frac{4\gamma_0^2}{\nu} \int_{t_{n-1}}^{t_n} \|f_t(t)\|_0^2 dt, \end{aligned}$$

for all $2 \leq n \leq N$.

Next, we deduce from (4.1) and Lemma 3.1 that

$$(4.19) \quad \begin{aligned} 2\nu \|A_h u_h^n\|_0 & \leq 2\|d_t u_h^n\|_0 + 2\|f(t_n)\|_0 + c_0 \|u_h^{n-1}\|_0^{1/2} \|u_h^{n-1}\|_1 \|A_h u_h^{n-1}\|_0^{1/2} \\ & \leq 2\|d_t u_h^n\|_0 + 2\|f(t_n)\|_0 + \nu \|A_h u_h^{n-1}\|_0 + \nu^{-1} c_0^2 \|u_h^{n-1}\|_0 \|u_h^{n-1}\|_1^2. \end{aligned}$$

Moreover, we deduce from (2.2), (3.4), (4.1) and Lemma 3.1 that

$$(4.20) \quad \|p_h^n\|_0 \leq c\nu \|u_h^n\|_1 + c\|d_t u_h^n\|_0 + c\|f(t_n)\|_0 + c\|u_h^{n-1}\|_1^2.$$

Now, we will prove (4.4)-(4.6) by induction. For $\alpha = 0, 1, 2$, we deduce from (4.3) that

$$(4.21) \quad G(u_h^0) \tau \leq G_h \tau \leq \nu.$$

Due to (4.2), (4.4)-(4.6) hold for $m = 0$. For $\alpha = 0, 1$, we can obtain (4.4)-(4.6) with $m = 1$ by using (4.11)-(4.12), (4.20)-(4.21). For $\alpha = 2$, (4.13) and Lemma 3.1

can yield

$$(4.22) \quad \|d_t u_h^0\|_0 \leq 2\nu \|A_h u_h^0\|_0 + \|f(t_0)\|_0 + G^{1/2}(u_h^0) \|u_h^0\|_1.$$

Hence, we imply (4.4)-(4.6) with $m = 1$ by using (4.2), (4.11)-(4.12), (4.16) and (4.19)-(4.22). Assuming that (4.4)-(4.6) hold for $m = 0, 1, \dots, J$, we want to prove that they hold for $m = J + 1$.

Proof of (4.4). In view of the induction assumption and (4.3), it holds that

$$(4.23) \quad G(u_h^{n-1})\tau \leq G_h\tau \leq \nu, \quad 1 \leq n \leq J+1, \quad G(u_h^{n-2})\tau \leq G_h\tau \leq \nu, \quad 2 \leq n \leq J,$$

for $\alpha = 0, 1, 2$. Summing (4.11) from $n = 1$ to $J+1$ and using (4.23), we obtain (4.4) for $m = J+1$ in the case of $\alpha = 0, 1, 2$.

Proof of (4.5). For $\alpha = 1, 2$, by summing (4.12) from $n = 1$ to $n = J+1$ and using (4.23), we obtain

$$(4.24) \quad \begin{aligned} & \nu \|u_h^{J+1}\|_1^2 + \tau \sum_{n=1}^{J+1} (\|d_t u_h^n\|_0^2 + \nu \|d_t u_h^n\|_1^2 \tau + \nu^2 \|A_h u_h^n\|_0^2) \\ & \leq \tau \sum_{n=0}^J d_n \nu \|u_h^n\|_1^2 + 8\nu^{-1} T \sup_{0 \leq t \leq T} \|f(t)\|_0^2 + 2\nu \|u_h^0\|_1^2. \end{aligned}$$

We set

$$\begin{aligned} a_n &= \nu \|u_h^n\|_1^2, \quad C = 8\nu^{-1} T \sup_{0 \leq t \leq T} \|f(t)\|_0^2 + 2\nu \|u_h^0\|_1^2, \\ b_n &= \|d_t u_h^n\|_0^2 + \nu \|d_t u_h^n\|_1^2 \tau + \nu^2 \|A_h u_h^n\|_0^2. \end{aligned}$$

Applying Lemma 3.5 to (4.24) and using (4.4), we obtain (4.5) with $m = J+1$.

For $\alpha = 0$, multiplying (4.12) by $\sigma(t_n)$, using (4.23) and noting $\sigma(t_n) \leq \sigma(t_{n-1}) + \tau$, which will often be used later, we obtain

$$(4.25) \quad \begin{aligned} & 2\sigma(t_n)\nu \|u_h^n\|_1^2 - 2\nu\sigma(t_{n-1})\|u_h^{n-1}\|_1^2 + \sigma(t_n)(\|d_t u_h^n\|_0^2 + \nu \|d_t u_h^n\|_1^2 \tau + \nu^2 \|A_h u_h^n\|_0^2)\tau \\ & \leq 2\nu \|u_h^{n-1}\|_1^2 \tau + d_{n-1}(\tau + \sigma(t_{n-1}))\nu \|u_h^{n-1}\|_1^2 \tau + 8\nu^{-1} \|f(t_n)\|_0^2 \tau, \end{aligned}$$

for all $1 \leq n \leq J+1$. Summing (4.25) from $n = 1$ to $n = J+1$, we deduce

$$(4.26) \quad \begin{aligned} & \sigma(t_{J+1})\nu \|u_h^{J+1}\|_1^2 + \tau \sum_{n=1}^{J+1} \sigma(t_n)(\|d_t u_h^n\|_0^2 + \nu \|d_t u_h^n\|_1^2 \tau + \nu^2 \|A_h u_h^n\|_0^2) \\ & \leq 4\tau \sum_{n=0}^J d_n \sigma(t_n)\nu \|u_h^n\|_1^2 + 2\nu\tau \sum_{n=0}^J \|u_h^n\|_1^2 \\ & + 8\nu^{-1} T \sup_{0 \leq t \leq T} \|f(t)\|_0^2 + 2\nu\tau^2 d_0 \|u_h^1\|_1^2. \end{aligned}$$

Setting

$$\begin{aligned} a_n &= \sigma(t_n)\nu \|u_h^n\|_1^2, \quad C = 8\nu^{-1} T \sup_{0 \leq t \leq T} \|f(t)\|_0^2 \tau, \\ b_n &= \sigma(t_n)(\|d_t u_h^n\|_0^2 + \nu \|d_t u_h^n\|_1^2 \tau + \nu^2 \|A_h u_h^n\|_0^2). \end{aligned}$$

Applying Lemma 3.5 to (4.26) and using (4.3)-(4.4), we arrive at (4.5) for $m = J+1$.

Proof of (4.6). If $\|A_h u_h^{J+1}\|_0 \leq \|A_h u_h^J\|_0$, then the induction assumption yields

$$\sigma^{2-\alpha}(t) \nu^2 \|A_h u_h^{J+1}(t)\|_0^2 \leq \kappa_2, \quad 1 \leq J \leq N-1,$$

for $\alpha = 0, 1, 2$. Hence, we always assume that

$$(4.27) \quad \|A_h u_h^{J+1}\|_0 \geq \|A_h u_h^J\|_0, \quad 1 \leq J \leq N-1.$$

For $\alpha = 2$, summing (4.18) from $n = 2$ to $n = J+1$, adding (4.16) and using (4.4)-(4.5) and (4.23), we deduce

$$(4.28) \quad \begin{aligned} & \|d_t u_h^{J+1}\|_0^2 + \tau \sum_{n=1}^{J+1} (\nu \|d_t u_h^n\|_1^2 + \|d_{tt} u_h^n\|_0^2) \\ & \leq 2\left(\frac{4}{\nu}\right)^4 c_0^4 \kappa_0 \kappa_1^2 + \frac{8\gamma_0^2}{\nu} \int_0^T \|f_t(t)\|_0^2 dt + 2\|d_t u_h^0\|_0^2. \end{aligned}$$

Thus, by combining (4.27)-(4.28) with (4.19)-(4.20) with $n = J+1$, we deduce (4.6) for $m = J+1$.

For $\alpha = 1$, by multiplying (4.18) by $\sigma(t_n)$ and summing from $n = 2$ to $n = J+1$ and using (4.12) with $n = 1$, we find

$$(4.29) \quad \begin{aligned} & \sigma(t_{J+1}) \|d_t u_h^{J+1}\|_0^2 + \nu \tau \sum_{n=1}^{J+1} \sigma(t_n) \|d_t u_h^n\|_1^2 \\ & \leq \tau \sum_{n=1}^{J+1} \left(1 + 2\left(\frac{4}{\nu}\right)^3 c_0^4 \|u_h^{n-1}\|_0^2 \|u_h^{n-1}\|_1^2\right) \|d_t u_h^n\|_0^2 \\ & + \frac{4\gamma_0^2}{\nu} \int_0^T \|f_t(t)\|_0^2 dt + 2(1 + d_0 \tau) \nu \|u_h^0\|_1^2 + 16\nu^{-1} \|f(t_1)\|_0^2 \tau. \end{aligned}$$

Now, by using (4.27), (4.24) and (4.19)-(4.20) in (4.29), we obtain (4.6) for $m = J+1$.

Finally, for $\alpha = 0$, by multiplying (4.18) by $\sigma^2(t_n)$, noting $\sigma^2(t_n) \leq \sigma^2(t_{n-1}) + 3\sigma(t_{n-1})\tau$, which will often be used later, summing from $n = 2$ to $n = J+1$ and using (4.12) with $n = 1$, we find

$$(4.30) \quad \begin{aligned} & \sigma^2(t_{J+1}) \|d_t u_h^{J+1}\|_0^2 + \nu \tau \sum_{n=1}^{J+1} \sigma^2(t_n) \|d_t u_h^n\|_1^2 \\ & \leq \tau \sum_{n=1}^{J+1} \sigma(t_n) \left(2 + 2\left(\frac{4}{\nu}\right)^3 c_0^4 \|u_h^{n-1}\|_0^2 \|u_h^{n-1}\|_1^2\right) \|d_t u_h^n\|_0^2 \\ & + \frac{4\gamma_0^2}{\nu} T \sup_{0 \leq t \leq T} \|f_t(t)\|_0^2 + 2(1 + d_0 \tau) \nu \|u_h^0\|_1^2 + 16\nu^{-1} \|f(t_1)\|_0^2. \end{aligned}$$

Hence, by using (4.30), (4.26)-(4.27) and (4.19)-(4.20), we obtain (4.6) for $m = J+1$. $\square$

Theorem 4.2. *Under the assumptions of Theorem 4.1, it holds that*

$$(4.31) \quad \sigma^{3-\alpha}(t_m) \|d_t u_h^m\|_1^2 + \nu \tau \sum_{n=2}^m \sigma^{3-\alpha}(t_n) \|A_h d_t u_h^n\|_0^2 \leq \kappa_3,$$

$$(4.32) \quad \tau \sum_{n=2}^m \sigma^{3-\alpha}(t_n) \|d_{tt} u_h^n\|_0^2 \leq \kappa_4,$$

for all $2 \leq m \leq N$ and $\alpha = 1, 2$, where κ_3 and κ_4 are some positive constants depending on the data (ν, Ω, T, u_0, f) .

Proof. First, taking $v_h = 2A_h d_t u_h^n \tau \in V_h$ in (4.15), we deduce

$$\begin{aligned}
 \|d_t u_h^n\|_1^2 &= \|d_t u_h^{n-1}\|_1^2 + \|d_{tt} u_h^n\|_1^2 \tau^2 + 2\nu \|A_h d_t u_h^n\|_0^2 \tau \\
 &\quad + 2b(d_t u_h^{n-1}, u_h^{n-1}, A_h d_t u_h^n) \tau + 2b(u_h^{n-2}, d_t u_h^{n-1}, A_h d_t u_h^n) \tau \\
 (4.33) \quad &= 2 \int_{t_{n-1}}^{t_n} (f_t(t), A_h d_t u_h^n) dt.
 \end{aligned}$$

In view of Lemma 3.1 and (3.10), it holds that

$$\begin{aligned}
 2|b(d_t u_h^{n-1}, u_h^{n-1}, A_h d_t u_h^n)| \tau &\leq 2c_0 \gamma_0 \|A_h u_h^{n-1}\|_0 \|d_t u_h^{n-1}\|_1 \|A_h d_t u_h^n\|_0 \tau \\
 &\leq \frac{\nu}{8} \|A_h d_t u_h^n\|_0^2 \tau + 8\nu^{-1} c_0^2 \gamma_0^2 \|A_h u_h^{n-1}\|_0^2 \|d_t u_h^{n-1}\|_1^2 \tau, \\
 2|b(u_h^{n-2}, d_t u_h^n, A_h d_t u_h^n)| \tau &\leq 2c_0 \gamma_0^{1/2} \|u_h^{n-2}\|_1 \|d_t u_h^n\|_1^{1/2} \|A_h d_t u_h^n\|_0^{3/2} \tau \\
 &\leq \frac{\nu}{2} \|A_h d_t u_h^n\|_0^2 \tau + 2\left(\frac{2}{\nu}\right)^3 c_0^4 \gamma_0^2 \|u_h^{n-2}\|_1^4 \|d_t u_h^n\|_1^2 \tau, \\
 2|b(u_h^{n-2}, d_t u_h^n - d_t u_h^{n-1}, A_h d_t u_h^n)| \tau &\leq \frac{1}{2} G^{1/2}(u_h^{n-2}) \|d_{tt} u_h^n\|_1 \|A_h d_t u_h^n\|_0 \tau^2 \\
 &\leq \frac{1}{2} \|d_{tt} u_h^n\|_1^2 \tau^2 + \frac{1}{4} G(u_h^{n-2}) \|A_h d_t u_h^n\|_0^2 \tau^2, \\
 2 \left| \int_{t_{n-1}}^{t_n} (f_t(t), A_h d_t u_h^n) dt \right| &\leq \frac{\nu}{8} \|A_h d_t u_h^n\|_0^2 \tau + \frac{8}{\nu} \int_{t_{n-1}}^{t_n} \|f_t(t)\|_0^2 dt.
 \end{aligned}$$

Combining these inequalities with (4.33) yields

$$\begin{aligned}
 \|d_t u_h^n\|_1^2 &= \|d_t u_h^{n-1}\|_1^2 + \nu \|A_h d_t u_h^n\|_0^2 \tau + \frac{1}{4} (\nu - G(u_h^{n-2}) \tau) \|A_h d_t u_h^n\|_0^2 \tau \\
 &\leq 8\nu^{-1} c_0^2 \gamma_0^2 \|A_h u_h^{n-1}\|_0^2 \|d_t u_h^{n-1}\|_1^2 \tau + 2\left(\frac{2}{\nu}\right)^3 c_0^4 \gamma_0^2 \|u_h^{n-2}\|_1^4 \|d_t u_h^n\|_1^2 \tau \\
 (4.34) \quad &+ \frac{8}{\nu} \int_{t_{n-1}}^{t_n} \|f_t(t)\|_0^2 dt,
 \end{aligned}$$

for all $2 \leq n \leq N$. Multiplying (4.34) by $\sigma^{3-\alpha}(t_n)$ and using (4.23), we deduce

$$\begin{aligned}
 \sigma^{3-\alpha}(t_n) \|d_t u_h^n\|_1^2 - \sigma^{3-\alpha}(t_{n-1}) \|d_t u_h^{n-1}\|_1^2 + \nu \sigma^{3-\alpha}(t_n) \|A_h d_t u_h^n\|_0^2 \tau \\
 \leq c \sigma^{3-\alpha}(t_{n-1}) \|A_h u_h^{n-1}\|_0^2 \|d_t u_h^{n-1}\|_1^2 \tau \\
 (4.35) \quad + c \sigma^{3-\alpha}(t_n) \|u_h^{n-2}\|_1^4 \|d_t u_h^n\|_0^2 \tau \\
 + c \sigma^{2-\alpha}(t_{n-1}) \|d_t u_h^{n-1}\|_1^2 \tau + c \int_{t_{n-1}}^{t_n} \|f_t(t)\|_0^2 dt,
 \end{aligned}$$

for all $2 \leq n \leq N$. Summing (4.35) from $n = 2$ to $n = m$ and using (4.6), we obtain (4.31).

Then, we deduce from (4.15), (3.10) and Lemma 3.1 that

$$\begin{aligned}
 \|d_{tt} u_h^n\|_0 &\leq \nu \|A_h d_t u_h^n\|_0 + c \|d_t u_h^{n-1}\|_1 (\|A_h u_h^{n-1}\|_0 + H(n-3) \|A_h u_h^{n-2}\|_0) \\
 (4.36) \quad &+ \frac{1}{2} H(2-n) G^{1/2}(u_h^0) \|d_t u_h^1\|_1 + \tau^{-1/2} \left(\int_{t_{n-1}}^{t_n} \|f_t\|_0^2 dt \right)^{1/2}
 \end{aligned}$$

for all $2 \leq n \leq N$, where $H(t) = 1$, as $t \geq 0$ and $H(t) = 0$, as $t < 0$. Thus, we deduce from (4.36) that

$$\begin{aligned} & \sigma^{3-\alpha}(t_n) \|d_{tt}u_h^n\|_0^2 \tau \\ & \leq c\sigma^{3-\alpha}(t_n) \|A_h d_t u_h^n\|_0^2 \tau + H(2-n)G(u_h^0)(2\tau)^{3-\alpha} \|d_t u_h^1\|_1^2 \tau + c \int_{t_{n-1}}^{t_n} \|f_t\|_0^2 dt \\ & \quad + c\sigma^{2-\alpha}(t_{n-1}) \|d_t u_h^{n-1}\|_1^2 (\sigma(t_{n-1}) \|A_h u_h^{n-1}\|_0^2 + H(n-3)\sigma(t_{n-2}) \|A_h u_h^{n-2}\|_0^2) \tau. \end{aligned}$$

Summing the above inequality from $n = 2$ to $n = m$ and using Theorem 4.1, (4.21) and (4.31), we get (4.32). $\square$

5. DUAL EULER SCHEME: STABILITY ANALYSIS

In order to derive the L^2 -bound on the error $u_h(t_n) - u_h^n$ in the case of $\alpha = 1$, we employ a parabolic argument that has already been used in [23] for the Crank-Nicolson scheme of the time-dependent Navier-Stokes equation. Let $1 \leq m \leq N$ be given. We consider the linearized “backward” counterpart of the discrete Navier-Stokes (4.1): For $\xi^n \in V_h$, $1 \leq n \leq m$, find $\Phi_h^{n-1} \in V_h$ such that

$$(5.1) \quad (v_h, d_t \Phi_h^n) - a(v_h, \Phi_h^{n-1}) - b(u_h^n, v_h, \Phi_h^{n-1}) - b(v_h, u_h^n, \Phi_h^{n-1}) = (v_h, \xi^n),$$

for $v_h \in V_h$ with an initial value $\Phi_h^m = 0$.

Here, we need to introduce the following discrete dual Gronwall lemma provided in [11].

Lemma 5.1. *Let $C > 0$ and let a_n, b_n, d_n , for integers $0 \leq n \leq m$, be nonnegative numbers such that*

$$(5.2) \quad a_k + \tau \sum_{n=k}^m b_n \leq \tau \sum_{n=k+1}^m d_n a_n + C, \quad 0 \leq k \leq m.$$

Then

$$(5.3) \quad a_k + \tau \sum_{n=k}^m b_n \leq C \exp(\tau \sum_{n=k+1}^m d_n), \quad 0 \leq k \leq m,$$

where we assume that $\tau \sum_{n=m+1}^m d_n = 0$.

The following lemma provides the stability of the scheme (5.1).

Lemma 5.2. *Under the assumptions of Theorem 4.1, the following a priori estimate holds:*

$$(5.4) \quad \|\Phi_h^k\|_1^2 + \nu \tau \sum_{n=k}^m \|A_h \Phi_h^n\|_0^2 \leq \kappa \tau \sum_{n=1}^m \|\xi^n\|_0^2,$$

for all $0 \leq k \leq m$.

Proof. The proof follows the line of argument used in the proofs of Theorem 4.1. In view of Lemma 3.1 and (4.3), we can prove that (5.1) admits a unique solution sequence $\{\Phi_h^k\}_0^m$.

Moreover, by taking $v_h = -2A_h\Phi_h^{n-1}\tau$ in (5.1), we obtain

$$\begin{aligned}
 \|\Phi_h^{n-1}\|_1^2 &= \|\Phi_h^n\|_1^2 + \|d_t\Phi_h^n\|_1^2\tau^2 + 2\nu\|A_h\Phi_h^{n-1}\|_0^2\tau \\
 &\quad + 2b(A_h\Phi_h^{n-1}, u_h^n, \Phi_h^{n-1})\tau + 2b(u_h^n, A_h\Phi_h^{n-1}, \Phi_h^{n-1})\tau \\
 (5.5) \quad &\leq \frac{\nu}{4}\|A_h\Phi_h^{n-1}\|_0^2\tau + \frac{4}{\nu}\|\xi^n\|_0^2\tau.
 \end{aligned}$$

From Lemma 3.1 and (3.10), we have

$$\begin{aligned}
 2|b(A_h\Phi_h^{n-1}, u_h^n, \Phi_h^n)|\tau &+ 2|b(u_h^n, A_h\Phi_h^{n-1}, \Phi_h^n)|\tau \\
 &\leq 2c_0\gamma_0\|A_hu_h^n\|_0\|\Phi_h^n\|_1\|A_h\Phi_h^{n-1}\|_0\tau \\
 &\leq \frac{\nu}{4}\|A_h\Phi_h^{n-1}\|_0^2\tau + 4\nu^{-1}c_0^2\gamma_0^2\|A_hu_h^n\|_0^2\|\Phi_h^n\|_1^2\tau, \\
 2|b(A_h\Phi_h^{n-1}, u_h^n, \Phi_h^n - \Phi_h^{n-1})|\tau &+ 2|b(u_h^n, A_h\Phi_h^{n-1}, \Phi_h^n - \Phi_h^{n-1})|\tau \\
 &\leq \frac{1}{2}G^{1/2}(u_h^n)\|d_t\Phi_h^n\|_1\|A_h\Phi_h^{n-1}\|_0\tau^2 \\
 &\leq \frac{1}{2}\|d_t\Phi_h^n\|_1^2\tau^2 + \frac{1}{4}G(u_h^n)\|A_h\Phi_h^{n-1}\|_0^2\tau^2.
 \end{aligned}$$

Combining (5.5) with the above estimate gives

$$\begin{aligned}
 \|\Phi_h^{n-1}\|_1^2 &= \|\Phi_h^n\|_1^2 + \nu\|A_h\Phi_h^{n-1}\|_0^2\tau + \frac{1}{4}(\nu - G(u_h^n)\tau)\|A_h\Phi_h^{n-1}\|_0^2\tau \\
 (5.6) \quad &\leq 4\nu^{-1}c_0^2\gamma_0^2\|A_hu_h^n\|_0^2\|\Phi_h^n\|_1^2\tau + 4\nu^{-1}\|\xi^n\|_0^2\tau,
 \end{aligned}$$

for all $1 \leq n \leq m$. Using (4.3) and Theorem 4.1 with $\alpha = 1, 2$, we have

$$(5.7) \quad \nu - G(u_h^n)\tau \geq \nu - G_h\tau \geq 0, \quad \forall 0 \leq n \leq N.$$

Summing (5.6) from $k+1$ to m and using (5.7) and Theorem 4.1, we obtain

$$\begin{aligned}
 \|\Phi_h^k\|_1^2 &+ \nu\tau \sum_{n=k}^{m-1} \|A_h\Phi_h^n\|_0^2 \\
 (5.8) \quad &\leq 4\tau \sum_{n=k+1}^m \nu^{-1}c_0^2\gamma_0^2\|A_hu_h^n\|_0^2\|\Phi_h^n\|_1^2 + 4\nu^{-1}\tau \sum_{n=1}^m \|\xi^n\|_0^2,
 \end{aligned}$$

for all $0 \leq k \leq m-1$. Applying Lemma 5.1 to (5.8) and using Theorem 4.1 yields (5.4). $\square$

6. ERROR ANALYSIS

In this section, we establish the H^1 - and L^2 -bounds of the error $e^n = u_h(t_n) - u_h^n$ and the L^2 -bound of the error $\eta^n = p_h(t_n) - p_h^n$ for all $1 \leq n \leq N$. To do this, we take $t = t_n$ in (3.8) and note

$$u_{ht}(t_n) - d_t u_h(t_n) = \tau^{-1} \int_{t_{n-1}}^{t_n} (u_{ht}(t_n) - u_{ht}(t))dt = \tau^{-1} \int_{t_{n-1}}^{t_n} (t - t_{n-1})u_{htt}dt,$$

to obtain

$$\begin{aligned}
 (d_t u_h(t_n), v_h) &+ a(u_h(t_n), v_h) - d(v_h, p_h(t_n)) + d(u_h(t_n), q_h) + b(u_h(t_n), u_h(t_n), v_h) \\
 (6.1) \quad &= (f(t_n), v_h) - \frac{1}{\tau} \int_{t_{n-1}}^{t_n} (t - t_{n-1})(u_{htt}(t), v_h)dt.
 \end{aligned}$$

Subtracting (4.1) from (6.1), we obtain

$$(6.2) \quad \begin{aligned} & (d_t e^n, v_h) + a(e^n, v_h) - d(v_h, \eta^n) + d(e^n, q_h) + b(e^n, u_h(t_n), v_h) \\ & + b(u_h^n, e^n, v_h) = (E_n, v_h), \end{aligned}$$

for all $(v_h, q_h) \in (X_h, M_h)$ with

$$(6.3) \quad \begin{aligned} (E_n, v_h) &= -\frac{1}{\tau} \int_{t_{n-1}}^{t_n} (t - t_{n-1})(u_{htt}(t), v_h) dt \\ &+ b(u_h^{n-1} - u_h^n, u_h^{n-1}, v_h) + b(u_h^n, u_h^{n-1} - u_h^n, v_h). \end{aligned}$$

Lemma 6.1. *Under the assumptions of Theorem 4.1 with $\alpha = 1, 2$, the error E_n satisfies the following bounds:*

$$(6.4) \quad \tau \sum_{n=1}^m \|A_h^{-1} P_h E_n\|_0^2 \leq \kappa \tau^2,$$

$$(6.5) \quad \tau \sum_{n=1}^m \|A_h^{-1/2} P_h E_h\|_0^2 \leq \kappa \tau^\alpha,$$

$$(6.6) \quad \tau \sum_{n=1}^m \sigma^{2-\alpha}(t_n) \|A_h^{-1/2} P_h E_h\|_0^2 \leq \kappa \tau^2,$$

for all $1 \leq m \leq N$, and

$$(6.7) \quad \sigma^{4-\alpha}(t_m) \|E_m\|_0^2 + \tau \sum_{n=2}^m \sigma^{3-\alpha}(t_n) \|E_n\|_0^2 \leq \kappa \tau^2, \quad 2 \leq m \leq N,$$

$$(6.8) \quad \tau \sum_{n=3}^m \sigma^{4-\alpha}(t_n) \|A_h^{-1/2} P_h d_t E_n\|_0^2 \leq \kappa \tau^2, \quad 3 \leq m \leq N.$$

Proof. First, it follows from (6.3), (3.10) and Lemma 3.1 that

$$(6.9) \quad \begin{aligned} \|A_h^{-1} P_h E_n\|_0^2 \tau &\leq c \tau^2 \int_{t_{n-1}}^{t_n} \|A_h^{-1} u_{htt}\|_0^2 dt \\ &+ c \|d_t u_h^n\|_0^2 (\|u_h^n\|_1^2 + \|u_h^{n-1}\|_1^2) \tau^3, \end{aligned}$$

$$(6.10) \quad \begin{aligned} \|A_h^{-1/2} P_h E_n\|_0^2 \tau &\leq c \tau^\alpha \int_{t_{n-1}}^{t_n} \sigma^{2-\alpha}(t) \|A_h^{-1/2} u_{htt}\|_0^2 dt \\ &+ c \|d_t u_h^n\|_1^2 (\|u_h^{n-1}\|_1^2 + \|u_h^n\|_1^2) \tau^3, \end{aligned}$$

$$(6.11) \quad \begin{aligned} \sigma^{2-\alpha}(t_n) \|A_h^{-1/2} P_h E_n\|_0^2 \tau &\leq c \tau^2 \int_{t_{n-1}}^{t_n} \sigma^{2-\alpha}(t) \|A_h^{-1/2} u_{htt}\|_0^2 dt \\ &+ c \sigma^{2-\alpha}(t_n) \|d_t u_h^n\|_1^2 (\|u_h^{n-1}\|_1^2 + \|u_h^n\|_1^2) \tau^3. \end{aligned}$$

Summing (6.9), (6.10) and (6.11) from 1 to m , respectively, noting $\tau^2 \leq \sigma^{2-\alpha}(t_n) \tau^\alpha$ and using (2.2), Theorem 3.6 and Theorem 4.1, we deduce (6.4)-(6.6) for $\alpha = 1, 2$.

Next, by using (3.10) and Lemma 3.1, we deduce from (6.3) that

$$(6.12) \quad \begin{aligned} \|E_n\|_0 &\leq c \tau^{-1/2} \left(\int_{t_{n-1}}^{t_n} (t - t_{n-1})^2 \|u_{htt}\|_0^2 dt \right)^{1/2} \\ &+ c (\|A_h u_h^n\|_0 + \|A_h u_h^{n-1}\|_0) \|d_t u_h^n\|_0 \tau, \end{aligned}$$

for all $2 \leq n \leq N$. Hence, we deduce from (6.12) that

$$\begin{aligned} \sigma^{3-\alpha}(t_n)\|E_n\|_0^2\tau &\leq c\tau^2 \int_{t_{n-1}}^{t_n} \sigma^{3-\alpha}(t)\|u_{htt}\|_0^2 dt \\ (6.13) \quad &+ c(\sigma(t_n)\|A_h u_h^n\|_0^2 + \sigma(t_{n-1})\|A_h u_h^{n-1}\|_0^2)\sigma^{2-\alpha}(t_n)\|d_t u_h^n\|_1^2 \tau^3, \end{aligned}$$

$$\begin{aligned} \sigma^{4-\alpha}(t_n)\|E_n\|_0^2 &\leq c\tau^2 \int_{t_{n-1}}^{t_n} \sigma^{3-\alpha}(t)\|u_{htt}\|_0^2 dt \\ (6.14) \quad &+ c(\sigma(t_n)\|A_h u_h^n\|_0^2 + \sigma(t_{n-1})\|A_h u_h^{n-1}\|_0^2)\sigma^{3-\alpha}(t_n)\|d_t u_h^n\|_1^2 \tau^2, \end{aligned}$$

for all $2 \leq n \leq N$. Summing (6.13) from $n = 2$ to $n = m$ and using (6.14) and Theorems 3.6, 4.1 and 4.2, we deduce (6.7).

Moreover, we deduce from (6.3) that

$$\begin{aligned} (d_t E_n, v_h) &= -\tau^{-2} \int_{t_{n-1}}^{t_n} (t - t_{n-1}) \int_{t-\tau}^t (u_{httt}(s), v_h) ds dt \\ &\quad - b(d_{tt} u_h^n, u_h^{n-1}, v_h)\tau - b(u_h^{n-1}, d_{tt} u_h^n, v_h)\tau \\ (6.15) \quad &\quad - b(d_t u_h^n, d_t u_h^n, v_h)\tau - b(d_t u_h^{n-1}, d_t u_h^{n-1}, v_h)\tau, \end{aligned}$$

for all $3 \leq n \leq N$. Using (3.10) and Lemma 3.1, we deduce from (6.15) that

$$\begin{aligned} \|A_h^{-1/2} P_h d_t E_n\|_0 &\leq c\tau^{-3/2} \left(\int_{t_{n-1}}^{t_n} (t - t_{n-1})^2 \left\| \int_{t-\tau}^t u_{httt}(s) ds \right\|_{-1}^2 dt \right)^{1/2} \\ &\quad + c\|d_{tt} u_h^n\|_0 \|A_h u_h^{n-1}\|_0 \tau + c\|d_t u_h^n\|_0 \|A_h d_t u_h^n\|_0 \tau \\ &\quad + c\|d_t u_h^{n-1}\|_0 \|A_h d_t u_h^{n-1}\|_0 \tau, \end{aligned}$$

which yields

$$\begin{aligned} \sigma^{4-\alpha}(t_n)\|A_h^{-1/2} P_h d_t E_n\|_0^2\tau &\leq c\tau^2 \int_{t_{n-2}}^{t_n} \sigma^{4-\alpha}(t)\|A_h^{-1/2} u_{httt}\|_0^2 dt \\ &\quad + c\tau^3 \sigma^{3-\alpha}(t_n)\sigma(t_{n-1})\|d_{tt} u_h^n\|_0^2 \|A_h u_h^{n-1}\|_0^2 \\ &\quad + c\tau^3 \sigma^{4-\alpha}(t_n)\|d_t u_h^n\|_0^2 \|A_h d_t u_h^n\|_0^2 \\ (6.16) \quad &\quad + c\tau^3 \sigma^{4-\alpha}(t_{n-1})\|d_t u_h^{n-1}\|_0^2 \|A_h d_t u_h^{n-1}\|_0^2, \end{aligned}$$

for all $3 \leq n \leq N$. Summing (6.16) from 3 to m and using Theorems 3.6, 4.1 and 4.2, we deduce (6.8). $\square$

Lemma 6.2. *Under the assumptions of Theorem 4.1 with $\alpha = 1, 2$, we have*

$$(6.17) \quad \|e^m\|_0^2 + \tau \sum_{n=1}^m (\|d_t e^n\|_0^2 \tau + \nu \|e^n\|_1^2) \leq \kappa \tau^\alpha,$$

for all $1 \leq m \leq N$.

Proof. Taking $v_h = 2e^n \tau \in V_h$ and $q_h = 0$ in (6.2), we obtain

$$\begin{aligned} \|e^n\|_0^2 &- \|e^{n-1}\|_0^2 + \|d_t e^n\|_0^2 \tau^2 + 2\nu \|e^n\|_1^2 \tau + 2b(e^n, u_h^n, e^n)\tau \\ (6.18) \quad &\leq \frac{\nu}{4} \|e^n\|_1^2 \tau + 4\nu^{-1} \|A_h^{-1/2} P_h E_n\|_0^2 \tau. \end{aligned}$$

Using Lemma 3.1 and (3.10), one finds

$$\begin{aligned}
2|b(e^{n-1}, u_h^n, e^n)|\tau &\leq 2c_0\gamma_0^{1/2}\|e^{n-1}\|_0^{1/2}\|e_h^{n-1}\|_1^{1/2}\|u_h^n\|_1\|e^n\|_1\tau \\
&\leq \frac{\nu}{4}(\|e^n\|_1^2 + \|e^{n-1}\|_1^2)\tau + 2\left(\frac{2}{\nu}\right)^3 c_0^4\gamma_0^2\|u_h^n\|_1^4\|e^{n-1}\|_0^2\tau, \\
2|b(e^n - e^{n-1}, u_h^n, e^n)|\tau &\leq \frac{1}{2}G^{1/2}(u_h^n)\|e^n\|_1\|d_te^n\|_0\tau^2 \\
&\leq \frac{1}{2}\|d_te^n\|_0^2\tau^2 + \frac{1}{4}G(u_h^n)\|e^n\|_1^2\tau^2.
\end{aligned}$$

Hence, by combining the above inequalities with (6.18), we obtain

$$\begin{aligned}
\|e^n\|_0^2 &- \|e^{n-1}\|_0^2 + \frac{1}{2}\|d_te^n\|_0^2\tau^2 + \nu\|e^n\|_1^2\tau \\
&+ \frac{\nu}{4}(\|e^n\|_1^2 - \|e^{n-1}\|_1^2)\tau + \frac{1}{4}(\nu - G(u_h^n)\tau)\|e^n\|_1^2\tau \\
(6.19) \quad &\leq 2\left(\frac{2}{\nu}\right)^3 c_0^4\gamma_0^2\|u_h^n\|_1^4\|e^{n-1}\|_0^2\tau + 4\nu^{-1}\|A_h^{-1/2}P_hE_n\|_0^2\tau,
\end{aligned}$$

for all $1 \leq n \leq N$. Moreover, summing (6.19) from 1 to m and using (5.7), we have

$$\begin{aligned}
\|e^m\|_0^2 &+ \tau \sum_{n=1}^m \left(\frac{1}{2}\|d_te^n\|_0^2\tau + \nu\|e^n\|_1^2 \right) \\
(6.20) \quad &\leq \tau \sum_{n=0}^{m-1} d_n\|e^n\|_0^2 + 4\nu^{-1}\tau \sum_{n=1}^N \|A_h^{-1/2}P_hE_n\|_0^2,
\end{aligned}$$

where $d_n = 2\left(\frac{2}{\nu}\right)^3 c_0^4\gamma_0^2\|u_h^{n+1}\|_1^4$. We set

$$a_n = \|e^n\|_0^2, \quad b_n = \frac{1}{2}\|d_te^n\|_0^2\tau + \nu\|e^n\|_1^2, \quad C = 4\nu^{-1}\tau \sum_{n=1}^N \|A_h^{-1/2}P_hE_n\|_0^2,$$

and apply Lemma 3.5 to (6.20) and use Theorem 4.1 and Lemma 6.1 to deduce (6.17). $\square$

With the aid of Lemma 6.2, we obtain the following error estimate.

Lemma 6.3. *Under the assumptions of Theorem 4.1 with $\alpha = 1, 2$, we have*

$$(6.21) \quad \sigma^{2-\alpha}(t_m)\|e^m\|_0^2 + \tau \sum_{n=1}^m \sigma^{2-\alpha}(t_n)\|e^n\|_1^2 \leq \kappa\tau^2,$$

for all $1 \leq m \leq N$.

Proof. For $\alpha = 2$, Lemma 6.2 yields (6.21). For $\alpha = 1$, we let $\{\Phi_h^n\}_0^m$ be the solution of (5.1), corresponding to the initial value $\Phi_h^m = 0$ and the right-hand side of $\{\xi^n\}_1^m = \{e^n\}_1^m$. Then, by construction, it holds that

$$\begin{aligned}
\|e^n\|_0^2\tau &= (e^n, d_t\Phi_h^n)\tau - a(e^n, \Phi_h^{n-1})\tau \\
(6.22) \quad &- b(u_h^n, e^n, \Phi_h^{n-1})\tau - b(e^n, u_h^n, \Phi_h^{n-1})\tau.
\end{aligned}$$

Taking $v_h = \Phi_h^{n-1}\tau$ in (6.2) and adding (6.22), we obtain

$$(6.23) \quad \|e^n\|_0^2\tau = (e^n, \Phi_h^n) - (e^{n-1}, \Phi_h^{n-1}) - (E_n, \Phi_h^{n-1})\tau + b(e^n, e^n, \Phi_h^{n-1})\tau.$$

Summing (6.23) for $1 \leq n \leq m$ and using Lemmas 5.2, 6.1 and 6.2, we have

$$\begin{aligned}
 \tau \sum_{n=1}^m \|e^n\|_0^2 &\leq (\tau \sum_{n=1}^m \|A_h^{-1} P_h E_n\|_0^2)^{1/2} (\tau \sum_{n=1}^m \|A_h \Phi_h^{n-1}\|_0^2)^{1/2} \\
 &\quad + c(\tau \sum_{n=1}^m \|e^n\|_0^2 \|e^n\|_1^2)^{1/2} (\tau \sum_{n=1}^m \|A_h \Phi_h^{n-1}\|_0^2)^{1/2} \\
 (6.24) \quad &\leq \kappa \tau (\tau \sum_{n=1}^m \|e^n\|_0^2)^{1/2} \leq \frac{1}{2} \tau \sum_{n=1}^m \|e^n\|_0^2 + \kappa \tau^2.
 \end{aligned}$$

Next, multiplying (6.19) by $\sigma(t_n)$, we deduce

$$\begin{aligned}
 \sigma(t_n) \|e^n\|_0^2 - \sigma(t_{n-1}) \|e^{n-1}\|_0^2 + \sigma(t_n) \nu \|e^n\|_1^2 \tau + \frac{\nu}{4} (\sigma(t_n) \|e^n\|_1^2 - \sigma(t_{n-1}) \|e^{n-1}\|_1^2) \tau \\
 \leq \|e^{n-1}\|_0^2 \tau + \frac{\nu}{4} \|e_h^{n-1}\|_1^2 \tau^2 + 2(\frac{2}{\nu})^3 c_0^4 \gamma_0^2 \|u_h^n\|_1^4 \|e^{n-1}\|_0^2 \tau \\
 (6.25) \quad + 4\nu^{-1} \sigma(t_n) \|A_h^{-1/2} P_h E_n\|_0^2 \tau,
 \end{aligned}$$

for all $1 \leq n \leq N$. Summing (6.25) from $n = 1$ to $n = m$, we have

$$\begin{aligned}
 \sigma(t_m) \|e^m\|_0^2 + \nu \tau \sum_{n=1}^m \sigma(t_n) \|e^n\|_1^2 + \frac{\nu}{4} \sigma(t_m) \|e^m\|_1^2 \tau \\
 \leq \tau \sum_{n=1}^m (\|e^n\|_0^2 + \tau \nu \|e^n\|_1^2) + 2\tau \sum_{n=1}^m (\frac{2}{\nu})^3 c_0^4 \gamma_0^2 \|u_h^n\|_1^4 \|e^{n-1}\|_0^2 \\
 + 4\tau \sum_{n=1}^m \nu^{-1} \sigma(t_n) \|A_h^{-1/2} P_h E_n\|_0^2.
 \end{aligned}$$

Using (6.24), Theorem 4.1 and Lemmas 6.1 and 6.2 in the above inequality gives (6.21) for $\alpha = 1$. $\square$

Lemma 6.4. *Under the assumptions of Theorem 4.1 with $\alpha = 1, 2$, we have*

$$(6.26) \quad \sigma^{3-\alpha}(t_m) \|e^m\|_1^2 + \tau \sum_{n=2}^m \sigma^{3-\alpha}(t_n) (\|d_t e^n\|_0^2 + \nu^2 \|A_h e^n\|_0^2) \leq \kappa \tau^2,$$

for all $1 \leq m \leq N$.

Proof. Taking $v_h = 2A_h e^n \tau \in V_h$ and $q_h = 0$ in (6.2), we obtain

$$\begin{aligned}
 \|e^n\|_1^2 - \|e^{n-1}\|_1^2 + \|d_t e^n\|_1^2 \tau^2 + 2\nu \|A_h e^n\|_0^2 \tau + 2b(e^n, u_h(t_n), A_h e^n) \tau \\
 (6.27) \quad + 2b(u_h^n, e_h^n, A_h e^n) \tau \leq \frac{\nu}{4} \|A_h e^n\|_0^2 \tau + 4\nu^{-1} \|E_n\|_0^2 \tau.
 \end{aligned}$$

In view of Lemma 3.1 and (3.10), we have

$$\begin{aligned}
 2|b(e^n, u_h(t_n), e^n)| \tau + 2|b(u_h^n, e^n, A_h e^n)| \tau \\
 \leq 2c_0 \gamma_0^{1/2} \|e^n\|_1 (\|A_h u_h^n\|_0 + \|A_h u_h(t_n)\|_0) \|A_h e^n\|_0 \tau \\
 \leq \frac{\nu}{4} \|A_h e^n\|_0^2 \tau + c(\|A_h u_h(t_n)\|_0^2 + \|A_h u_h^n\|_0^2) \|e_h^n\|_1^2 \tau.
 \end{aligned}$$

Hence, by combining the above inequality with (6.27), we obtain

$$\begin{aligned}
 \|e^n\|_1^2 - \|e^{n-1}\|_1^2 + \nu \|A_h e^n\|_0^2 \tau \\
 (6.28) \quad \leq c(\|A_h u_h^n\|_0^2 + \|A_h u_h(t_n)\|_0^2) \|e^n\|_1^2 \tau + 4\nu^{-1} \|E_n\|_0^2 \tau,
 \end{aligned}$$

for all $1 \leq n \leq N$. Multiplying (6.28) by $\sigma^{3-\alpha}(t_n)$, we find

$$\begin{aligned} \sigma^{3-\alpha}(t_n)\|e^n\|_1^2 &= \sigma^{3-\alpha}(t_{n-1})\|e^{n-1}\|_1^2 + \nu\sigma^{3-\alpha}(t_n)\|A_h e^n\|_0^2\tau \\ &\leq c\sigma^{3-\alpha}(t_n)(\|A_h u_h^n\|_0^2 + \|A_h u_h(t_n)\|_0^2)\|e^n\|_1^2\tau \\ &\quad + c\sigma^{2-\alpha}(t_{n-1})\|e^{n-1}\|_1^2\tau + 4\nu^{-1}\sigma^{3-\alpha}(t_n)\|E_n\|_0^2\tau, \end{aligned} \quad (6.29)$$

for all $1 \leq n \leq N$. Summing (6.29) from 2 to m , and using Theorems 3.3 and 4.1 and Lemmas 6.1, 6.2 and 6.3, we deduce

$$\sigma^{3-\alpha}(t_m)\|e^m\|_1^2 + \nu\tau \sum_{n=2}^m \sigma^{3-\alpha}(t_n)\|A_h e^n\|_0^2 \leq \kappa\tau^2, \quad (6.30)$$

for all $1 \leq m \leq N$.

Finally, we deduce from (6.2), (3.10) and Lemma 3.1 that

$$\begin{aligned} \sigma^{3-\alpha}(t_n)\|d_t e^n\|_0^2 &\leq c\sigma^{3-\alpha}(t_n)(1 + \|u_h^n\|_1^2 + \|e_h^n\|_1^2)\|A_h e^n\|_0^2\tau \\ &\quad + c\sigma^{3-\alpha}(t_n)\|E_n\|_0^2\tau, \end{aligned} \quad (6.31)$$

for all $2 \leq n \leq N$. Summing (6.31) from $n = 2$ to $n = m$ and using (6.30), Theorem 4.1 and Lemmas 6.1 and 6.2, we deduce (6.26). $\square$

It remains to prove the error estimate for the discrete pressure p_h^m . To do this, we need to estimate $d_t e^n$. It follows from (6.2) that

$$\begin{aligned} (d_{tt}e^n, v_h) &+ a(d_t e^n, v_h) + b(d_t e^n, u_h(t_n), v_h) + b(e^{n-1}, d_t u_h(t_n), v_h) \\ &+ b(d_t u_h^n, e^n, v_h) + b(u_h^{n-1}, d_t e^n, v_h) = (d_t E_n, v_h), \end{aligned} \quad (6.32)$$

for all $v_h \in V_h$ and $1 \leq n \leq N$. Taking $v_h = 2d_t e^n \tau$ in (6.32) and using (3.12), we get

$$\begin{aligned} \|d_t e^n\|_0^2 - \|d_t e^{n-1}\|_0^2 + 2\nu\|d_t e^n\|_1^2\tau + 2b(d_t e^n, u_h(t_n), d_t e^n)\tau \\ + 2b(e^{n-1}, d_t u_h^n, d_t e^n)\tau \\ + 2b(d_t u_h^n, e^n, d_t e^n)\tau \leq \frac{\nu}{4}\|d_t e^n\|_1^2\tau + 4\nu^{-1}\|A_h^{-1/2}P_h d_t E_n\|_0^2\tau. \end{aligned} \quad (6.33)$$

In view of (3.10) and Lemma 3.1, we deduce

$$\begin{aligned} 2|b(d_t e^n, u_h(t_n), d_t e^n)|\tau &\leq c_0\gamma_0^{1/2}\|d_t e^n\|_0^{1/2}\|d_t e^n\|_1^{3/2}\|u_h(t_n)\|_1\tau \\ &\leq \frac{\nu}{4}\|d_t e^n\|_1^2\tau + \left(\frac{2}{\nu}\right)^3 c_0^4\gamma_0^2\|u_h(t_n)\|_1^4\|d_t e^n\|_0^2\tau, \\ 2|b(e^{n-1}, d_t u_h^n, d_t e^n)|\tau &+ 2b(d_t u_h^n, e^n, d_t e^n)\tau \\ &\leq 2c_0\gamma_0(\|A_h e^{n-1}\|_0 + \|A_h e^n\|_0)\|d_t u_h^n\|_0\|d_t e^n\|_1\tau \\ &\leq \frac{\nu}{4}\|d_t e^n\|_1^2\tau + 8\nu^{-1}c_0^2\gamma_0^2(\|A_h e^{n-1}\|_0^2 + \|A_h e^n\|_0^2)\|d_t u_h^n\|_0^2\tau. \end{aligned}$$

Combining these inequalities with (6.33) gives

$$\begin{aligned} \sigma^{4-\alpha}(t_n)\|d_t e^n\|_0^2 - \sigma^{4-\alpha}(t_{n-1})\|d_t e^{n-1}\|_0^2 &\leq \sigma^{4-\alpha}(t_n)\left(\frac{2}{\nu}\right)^3 c_0^4\gamma_0^2\|u_h(t_n)\|_1^4\|d_t e^n\|_0^2\tau \\ &\quad + (\sigma^{3-\alpha}(t_{n-1})\|A_h e^{n-1}\|_0^2 + \sigma^{3-\alpha}(t_n)\|A_h e^n\|_0^2)\sigma(t_n)\|d_t u_h^n\|_0^2\tau \\ &\quad + c\sigma^{3-\alpha}(t_{n-1})\|d_t e^{n-1}\|_0^2\tau + c\sigma^{4-\alpha}(t_n)\|A_h^{-1/2}P_h d_t E_n\|_0^2\tau. \end{aligned} \quad (6.34)$$

Summing (6.34) from 3 to m and using Theorem 3.3, Theorem 4.1, Lemma 6.1, Lemma 6.3 with $m = 1, 2$ and Lemma 6.4, we obtain

$$\sigma^{4-\alpha}(t_m)\|d_t e^m\|_0^2 \leq \kappa\tau^2, \quad 1 \leq m \leq N. \quad (6.35)$$

Moreover, we deduce from (6.2), (3.10) and Lemma 3.1 that

$$\begin{aligned} \|E_1\|_0 &\leq \|u_{ht}(t_1)\|_0 + \tau^{-1/2} \left(\int_{t_0}^{t_1} \|u_{ht}\|_0^2 dt \right)^{1/2} \\ &\quad + \frac{1}{2} (G^{1/2}(u_h^0) + G^{1/2}(u_h^1)) \|d_t u_h^1\|_1 \tau, \end{aligned}$$

which yields

$$\begin{aligned} \sigma^{4-\alpha}(t_1) \|E_1\|_0^2 &\leq c \sigma^{4-\alpha}(t_1) \|u_{ht}(t_1)\|_0^2 + c \sigma^{3-\alpha}(t_1) \int_{t_0}^{t_1} \|u_{ht}\|_0^2 dt \\ (6.36) \quad &\quad + c \tau^2 (G(u_h^0) \tau + G(u_h^1) \tau) \sigma^{3-\alpha}(t_1) \|d_t u_h^1\|_1^2. \end{aligned}$$

Using (5.7), Theorem 3.3 and Theorem 4.2 in (6.36), we obtain

$$(6.37) \quad \sigma^{4-\alpha}(t_1) \|E_1\|_0^2 \leq \kappa \tau^2.$$

By (3.4), (3.10), (6.2) and Lemma 3.1, we deduce

$$\begin{aligned} \|\eta^m\|_0 &\leq c(\|d_t e^m\|_0 + \|e^m\|_1) + c\|e^m\|_1(\|u_h(t_m)\|_1 + \|u_h^m\|_1) \\ &\quad + c\|E_m\|_0, \end{aligned}$$

which together with Theorems 3.3 and 4.1 yield

$$\begin{aligned} \sigma^{4-\alpha}(t_m) \|\eta^m\|_0^2 &\leq \kappa \sigma^{4-\alpha}(t_m) \|d_t e^m\|_0^2 \\ (6.38) \quad &\quad + \kappa \sigma^{3-\alpha}(t_m) \|e^m\|_1^2 + \sigma^{4-\alpha}(t_m) \|E_m\|_0^2. \end{aligned}$$

Using (6.35), (6.37), Lemma 6.1 and Lemma 6.4 in (6.38) yields

$$(6.39) \quad \sigma^{4-\alpha}(t_m) \|\eta^m\|_0^2 \leq \kappa \tau^2, \quad 1 \leq m \leq N.$$

Combining (6.39) with Lemma 6.3 and Lemma 6.4 yields the following error estimates results.

Theorem 6.5. *Under the assumptions of Theorem 4.1, the following error estimates hold:*

$$(6.40) \quad \sigma^{2-\alpha}(t_m) \|u_h(t_m) - u_h^m\|_0^2 + \sigma^{3-\alpha}(t_m) \|u_h(t_m) - u_h^m\|_1^2 \leq \kappa \tau^2, \quad t_m \in (0, T],$$

$$(6.41) \quad \sigma^{4-\alpha}(t_m) \|p_h(t_m) - p_h^m\|_0^2 \leq \kappa \tau^2, \quad t_m \in (0, T].$$

Remark. Combining Theorem 6.5 with (3.19) yields (1.11)-(1.13).

ACKNOWLEDGEMENTS

The author would like to thank the referee for valuable comments which helped us to improve the result of this paper.

REFERENCES

- [1] R. A. ADAMS, *Sobolev Spaces*, Academic Press, New York, 1975. MR0450957 (56:9247)
- [2] A. AIT OU AMMI AND M. MARION, *Nonlinear Galerkin methods and mixed finite elements: Two-grid algorithms for the Navier-Stokes equations*, Numer. Math., 68 (1994), pp. 189–213. MR1283337 (95c:65174)
- [3] G. A. BAKER, *Galerkin approximations for the Navier-Stokes equations*, manuscript, Harvard University, Cambridge, MA, 1976.
- [4] G. A. BAKER, V. A. DOUGALIS, AND O. A. KARAKASHIAN, *On a high order accurate fully discrete Galerkin approximation to the Navier-Stokes equations*, Math. Comp., 39 (1982), pp. 339–375. MR669634 (84h:65096)

- [5] J. BERCOVIER AND O. PIRONNEAU, *Error estimates for finite element solution of the Stokes problem in the primitive variables*, Numer. Math., 33 (1979), pp. 211–224. MR549450 (81g:65145)
- [6] J. R. CANNON AND YANPING LIN, *A priori L^2 error estimates for finite-element methods for nonlinear diffusion equations with memory*, SIAM J. Numer. Anal., 27 (1990), pp. 595–607. MR1041253 (91b:65118)
- [7] P. G. CIARLET, *The Finite Element Method for Elliptic Problems*, North-Holland, Amsterdam, 1978. MR0520174 (58:25001)
- [8] W. E AND J.-G. LIU, *Projection methods I: Convergence and numerical boundary layers*, SIAM J. Numer. Anal., 32 (1995), 1017–1057. MR1342281 (96e:65061)
- [9] G. FAIRWEATHER, H. MA AND W. SUN, *Orthogonal Spline Collocation Methods for the Navier-Stokes Equations in Stream Function and Vorticity Formulation*, submitted.
- [10] V. GIRAULT AND P. A. RAVIART, *Finite Element Method for Navier-Stokes Equations: Theory and algorithms*, Springer-Verlag, Berlin, Heidelberg, 1987. MR851383 (88b:65129)
- [11] YINNAN HE, *Stability and error analysis for a spectral Galerkin method for the Navier-Stokes equations with H^2 or H^1 initial data*, Numer. Methods for PDEs, 21 (2005), pp. 875–904. MR2154224 (2006d:65108)
- [12] YINNAN HE, *Stability and error analysis for a spectral Galerkin method for the Navier-Stokes equations with L^2 initial data*, Numer. Methods for PDEs, 24 (2008), pp. 79–103. MR2371349
- [13] YINNAN HE AND KAITAI LI, *Convergence and stability of finite element nonlinear Galerkin method for the Navier-Stokes equations*, Numer. Math., 79 (1998), pp. 77–106. MR1608417 (99c:65165)
- [14] YINNAN HE AND KAITAI LI, *Nonlinear Galerkin method and two-step method for the Navier-Stokes equations*, Numer. Methods for PDEs, 12 (1996):283–305. MR1388441 (97g:65192)
- [15] YINNAN HE, *Two-level method based on finite element and Crank-Nicolson extrapolation for the time-dependent Navier-Stokes equations*, SIAM J. Numer. Anal., 2003, 41(4):1263–1285. MR2034880 (2004k:65173)
- [16] YINNAN HE AND K. M. LIU, *A multi-level finite element method for the time-dependent Navier-Stokes equations*, Numer. Methods for PDEs, 21 (2005), pp. 1052–1068. MR2169167 (2006h:76079)
- [17] YINNAN HE, K. M. LIU AND WEIWEI SUN, *Multi-level spectral Galerkin method for the Navier-Stokes equations I: spatial discretization*, Numer. Math., 101 (2005), pp. 501–522. MR2194826 (2006j:76089)
- [18] YINNAN HE, YANPING LIN AND WEIWEI SUN, *Stabilized finite element methods for the nonstationary Navier-Stokes problem*, Discrete and Continuous Dynamical Systems-Series B, 6 (2006), pp. 41–68. MR2172195 (2006g:65152)
- [19] YINNAN HE AND WEIWEI SUN, *Stability and convergence of the Crank-Nicolson/Adams-Bashforth scheme for the time-dependent Navier-Stokes equations*, SIAM J. Numer. Anal., 2007, 45(2):837–869. MR2300299
- [20] YINNAN HE, *Optimal error estimate of the penalty finite element method for the time-dependent Navier-Stokes problem*, Mathematics of Computation, 74 (2005), pp. 1201–1216. MR2136999 (2006i:65150)
- [21] YINNAN HE, HUANLING MIAO, R. M. M. MATTHEIJ AND ZHANGXIN CHEN, *Numerical analysis of a modified finite element nonlinear Galerkin method*, Numer. Math., 97 (2004), pp. 725–756. MR2127930 (2005m:65188)
- [22] J. G. HEYWOOD AND R. RANNACHER, *Finite-element approximations of the nonstationary Navier-Stokes problem. Part I: Regularity of solutions and second-order spatial discretization*, SIAM J. Numer. Anal., 19 (1982), pp. 275–311. MR650052 (83d:65260)
- [23] J. G. HEYWOOD AND R. RANNACHER, *Finite-element approximations of the nonstationary Navier-Stokes problem. Part IV: Error estimates for second-order time discretization*, SIAM J. Numer. Anal., 27 (1990), pp. 353–384. MR1043610 (92c:65133)
- [24] A. T. HILL AND E. SÜLI, *Approximation of the global attractor for the incompressible Navier-Stokes equations*, IMA J. Numer. Anal., 20 (2000), pp. 633–667. MR1795301 (2001j:37138)
- [25] E. ISSACSON AND H. B. KELLER, *Analysis of Numerical Methods*, Wiley, New York, 1966. MR0201039 (34:924)

- [26] H. JOHNSTON AND J.-G. LIU, *Accurate, stable and efficient Navier-Stokes solvers based on explicit treatment of the pressure term*, J. Computational Physics, 199 (2004), pp. 221–259. MR2081004 (2005b:76093)
- [27] J. KIM AND P. MOIN, *Application of a fractional-step method to incompressible Navier-Stokes equations*, J. Comput. Phys., 59 (1985), pp. 308–323. MR796611 (87a:76046)
- [28] R. B. KELLOGG AND J. E. OSBORN, *A regularity result for the Stokes problem in a convex polygon*, J. Functional Anal., 21 (1976), pp. 397–431. MR0404849 (53:8649)
- [29] S. LARSSON, *The long-time behavior of finite-element approximations of solutions to semi-linear parabolic problems*, SIAM J. Numer. Anal., 26 (1989), pp. 348–365. MR987394 (90g:65124)
- [30] YANPING LIN, *Galerkin methods for nonlinear parabolic integrodifferential equations with nonlinear boundary conditions*, SIAM J. Numer. Anal., 27 (1990), pp. 608–621. MR1041254 (91e:65128)
- [31] HEPING MA AND WEIWEI SUN, *Optimal error estimates of the Legendre Petro-Galerkin and pseudospectral methods for the generalized Korteweg-de Vries Equation*, SIAM J. Numer. Anal., 39 (2001), pp. 1380–1394. MR1870846 (2002k:65162)
- [32] M. MARION AND R. TEMAM, *Navier-Stokes equations: Theory and approximation*, in: Handbook of Numerical Analysis, Vol. VI, pp. 503–688, North-Holland, Amsterdam, 1998. MR1665429 (2000a:76002)
- [33] R. H. NOCHETTO AND J.-H. PYO, *A finite element Gauge-Uzawa method Part I: Navier-Stokes equations*, SIAM J. Numer. Anal., 43 (2005), pp. 1043–1068. MR2177795 (2006m:65206)
- [34] J. SHEN, *Long time stability and convergence for fully discrete nonlinear Galerkin methods*, Appl. Anal., 38 (1990), pp. 201–229. MR1116181 (93a:65130)
- [35] J. C. SIMO AND F. ARMERO, *Unconditional stability and long-term behavior of transient algorithms for the incompressible Navier-Stokes and Euler equations*, Comput. Methods Appl. Mech. Engrg., 111 (1994), pp. 111–154. MR1259618 (94k:76078)
- [36] R. TEMAM, *Navier-Stokes Equations, Theory and Numerical Analysis*, 3rd ed., North-Holland, Amsterdam, 1983. MR769654 (86m:76003)
- [37] F. TONE, *Error analysis for a second scheme for the Navier-Stokes equations*, Applied Numerical Mathematics, 50 (2004), pp. 93–119. MR2060828 (2005b:65101)

FACULTY OF SCIENCE, XI'AN JIAOTONG UNIVERSITY, XI'AN 710049, PEOPLE'S REPUBLIC OF CHINA

E-mail address: heyn@mail.xjtu.edu.cn