THE GENERATING FUNCTION FOR THE DIRICHLET SERIES $L_m(s)$

WILLIAM Y. C. CHEN, NEIL J. Y. FAN, AND JEFFREY Y. T. JIA

ABSTRACT. The Dirichlet series $L_m(s)$ are of fundamental importance in number theory. Shanks defined the generalized Euler and class numbers in connection with the Dirichlet series, denoted by $\{s_{m,n}\}_{n \geq 0}$. We obtain a formula for the exponential generating function $s_m(x)$ of $s_{m,n}$, where m is an arbitrary positive integer. In particular, for $m > 1$, say, $m = bu^2$, where b is square-free and $u > 1$, we show that $s_m(x)$ can be expressed as a linear combination of the four functions $w(b, t) \sec(btx)(\pm \cos((b-p)tx) \pm \sin(ptx))$, where p is a nonnegative integer not exceeding b , $t|u^2$ and $w(b, t) = K_b t/u$ with K_b being a constant depending on b . Moreover, the Dirichlet series $L_m(s)$ can be easily computed from the generating function formula for $s_m(x)$. Finally, we show that the main ingredient in the formula for $s_{m,n}$ has a combinatorial interpretation in terms of the Λ -alternating augmented m -signed permutations defined by Ehrenborg and Readdy. More precisely, when m is square-free, this answers a question posed by Shanks concerning a combinatorial interpretation of the numbers $s_{m,n}$. When m is not square-free, say $m = bu^2$, the numbers $K_b^{-1} s_{m,n}$ can be written as a linear combination of the numbers of Λ -alternating augmented bt -signed permutations with integer coefficients, where $t|u^2$.

1. INTRODUCTION

The Dirichlet series

$$(1.1) \quad L_m(s) = \sum_{\substack{l > 0 \\ \text{odd } l}} \left(\frac{-m}{l} \right) \frac{1}{l^s},$$

where $(-m/l)$ is the Jacobi symbol, originate in the distribution of primes into arithmetic progressions, the class number of binary quadratic forms, as well as the distribution of the Legendre and Jacobi symbols. They play a crucial role in the computation of certain number-theoretic constants; see [3, 6, 12, 14]. Several approaches have been developed for the computation of $L_m(s)$; see, for example, Shanks [15, 16, 18].

The generalized Euler and class numbers were introduced by Shanks for the computation of the Dirichlet series $L_m(s)$ [15, 17]. These numbers are also related to derivative polynomials and Euler polynomials; see Hoffman [7] and Shanks [17].

Received by the editor April 13, 2010 and, in revised form, December 26, 2010.

2010 *Mathematics Subject Classification.* Primary 11B68, 05A05.

Key words and phrases. Dirichlet series, generalized Euler and class number, Λ -alternating augmented m -signed permutation, r -cubical lattice, Springer number.

©2011 American Mathematical Society
 Reverts to public domain 28 years from publication

In this paper, we obtain the generating functions for the generalized Euler and class numbers. Let us recall the definition of the generalized Euler and class numbers $s_{m,n}$ ($m \geq 1, n \geq 0$), introduced by Shanks,

$$s_{m,n} = \begin{cases} c_{m, \frac{n}{2}} & \text{if } n \text{ is even,} \\ d_{m, \frac{n+1}{2}} & \text{if } n \text{ is odd,} \end{cases}$$

where $c_{m,n}$ and $d_{m,n}$ are given by

$$(1.2) \quad c_{m,n} = (2n)! L_m(2n+1) (K_m \sqrt{m})^{-1} \left(\frac{\pi}{2m} \right)^{-2n-1},$$

$$(1.3) \quad d_{m,n} = (2n-1)! L_{-m}(2n) (K_m \sqrt{m})^{-1} \left(\frac{\pi}{2m} \right)^{-2n},$$

in which

$$(1.4) \quad K_m = \begin{cases} \frac{1}{2} & \text{if } m = 1, \\ 1 & \text{otherwise.} \end{cases}$$

Set

$$c_m(x) = \sum_{n \geq 0} c_{m,n} \frac{x^{2n}}{(2n)!},$$

$$d_m(x) = \sum_{n \geq 1} d_{m,n} \frac{x^{2n-1}}{(2n-1)!},$$

$$s_m(x) = \sum_{n \geq 0} s_{m,n} \frac{x^n}{n!}.$$

Clearly,

$$s_m(x) = c_m(x) + d_m(x).$$

By the definitions (1.2) and (1.3), we have

$$\begin{aligned} \sum_{n \geq 0} L_m(2n+1) x^{2n} &= \frac{K_m \pi \sqrt{m}}{2m} c_m \left(\frac{\pi}{2m} x \right), \\ \sum_{n \geq 1} L_{-m}(2n) x^{2n-1} &= \frac{K_m \pi \sqrt{m}}{2m} d_m \left(\frac{\pi}{2m} x \right). \end{aligned}$$

Therefore, if we set

$$\hat{L}_m(s) = \begin{cases} L_{-m}(s+1) & \text{if } s \text{ is odd,} \\ L_m(s+1) & \text{if } s \text{ is even,} \end{cases}$$

then

$$(1.5) \quad \sum_{s \geq 0} \hat{L}_m(s) x^s = \frac{K_m \pi \sqrt{m}}{2m} s_m \left(\frac{\pi}{2m} x \right).$$

It follows from (1.5) that $\hat{L}_m(s)$ is determined by $s_m(x)$. In other words, the generating function $s_m(x)$ leads to a quick way to compute $\hat{L}_m(s)$.

Consider $c_{m,n}$ and $d_{m,n}$ as entries of the infinite matrices C and D , respectively. Then the first column of C forms the sequence of class numbers in connection with primitive binary quadratic forms, and the first row of C forms the sequence of secant numbers, corresponding to up-down permutations, also called alternating

permutations, of even length. Meanwhile, the first row of D forms the sequence of tangent numbers corresponding to up-down permutations of odd length. Recall that both secant numbers and tangent numbers are called Euler numbers. This is why the numbers $s_{m,n}$ are called generalized Euler and class numbers.

Shanks [17] found recurrence relations for $c_{m,n}$ and $d_{m,n}$ with respect to the index n , from which it follows that $c_{m,n}$ and $d_{m,n}$ are integers. For example, we have

$$\sum_{i=0}^n (-4)^i \binom{2n}{2i} c_{2,n-i} = (-1)^n$$

and

$$\sum_{i=0}^{n-1} (-4)^i \binom{2n-1}{2i} d_{2,n-i} = (-1)^{n-1}.$$

In fact, due to the well-known Euler product of the Dirichlet series $L_m(s)$ (see [8, 11]), it can be easily shown that $c_{m,n}$ and $d_{m,n}$ are positive.

Shanks [17] raised the question of whether the generalized Euler and class numbers may have some combinatorial interpretations. The combinatorial interpretations of $s_{m,n}$ for $m = 1, 2, 3, 4$, have been found. Let $(s_{m,n})_{n \geq 0}$ denote the conjoined sequence

$$(c_{m,0}, d_{m,1}, c_{m,1}, d_{m,2}, c_{m,2}, d_{m,3}, \dots).$$

For $m = 1$, the sequence $(1, 1, 1, 2, 5, 16, \dots)$ is listed as A000111 in Sloane [19], which is called the sequence of Euler numbers, enumerating alternating permutations on $[n] = \{1, 2, \dots, n\}$.

For $m = 2$, the sequence $(1, 1, 3, 11, 57, 361, \dots)$ is numbered A001586 in [19], which is also called the sequence of Springer numbers which arise in the work of Springer on the theory of Weyl groups.

For $m = 3$, the sequence $(1, 2, 8, 46, 352, 3362, \dots)$ is listed as A007289 in [19], and we call it the sequence of Ehrenborg and Readdy numbers. The combinatorial interpretation of this sequence in terms of alternating 3-signed permutations has been given by Ehrenborg and Readdy [4].

For $m = 4$, a combinatorial interpretation of the sequence $(1, 4, 16, 128, 1280, 16384, \dots)$ has been given implicitly by Ehrenborg and Readdy [5] in terms of nonaugmented André R -signed permutations on $[n]$ with $R = (4, 4, \dots, 4)$.

For $m \geq 5$, we shall give formulas for the generating function for $s_m(x)$. For the case when m is square-free, we shall give a combinatorial interpretation for $s_{m,n}$. In the general case, we see that the number $s_{m,n}$ or $2s_{m,n}$ is a linear combination with integer coefficients of some numbers with a combinatorial interpretation.

For $m = 1, 2, 3, 4$, it is known that

$$\begin{aligned} s_1(x) &= \sec x + \tan x, \\ s_2(x) &= \frac{\cos x + \sin x}{\cos 2x}, \\ s_3(x) &= \frac{\sin 2x + \cos x}{\cos 3x}, \\ s_4(x) &= \sec 4x + \tan 4x. \end{aligned}$$

From our formulas, we get the following expressions for $m = 5, 6, 7$:

$$\begin{aligned} s_5(x) &= \frac{\cos 4x + \sin x}{\cos 5x} + \frac{\cos 2x + \sin 3x}{\cos 5x}, \\ s_6(x) &= \frac{\cos 5x + \sin x}{\cos 6x} + \frac{\cos x + \sin 5x}{\cos 6x}, \\ s_7(x) &= \frac{\cos 3x + \sin 4x}{\cos 7x} + \frac{\cos x + \sin 6x}{\cos 7x} - \frac{\cos 5x + \sin 2x}{\cos 7x}. \end{aligned}$$

This paper is organized as follows. In Section 2, we compute the generating function $s_m(x)$ when m is square-free, while in Section 3 we consider the case when m is not square-free. Section 4 is devoted to the combinatorial interpretation of the numbers $s_{m,n}$ in terms of m -signed permutations as introduced by Ehrenborg and Readdy when m is square-free.

2. COMPUTATION FOR $s_m(x)$ WHEN m IS SQUARE-FREE

In this section, we compute the generating function $s_m(x)$ when $m > 1$ and m is square-free. For $0 \leq p \leq m$, we adopt the following notation introduced in [4],

$$(2.6) \quad \Lambda_{m,p}(x) := \frac{\cos((m-p)x) + \sin(px)}{\cos(mx)}.$$

When m is square-free, we shall not encounter the case that m is a multiple of 4. We shall have three formulas for $s_m(x)$ depending on the residue of m modulo 4.

Theorem 2.1. *Assume that m is square-free and $m = 4t + 3$. Then*

$$(2.7) \quad s_m(x) = \sum_{k=1}^t \left(\frac{k}{m}\right) \Lambda_{m,4k}(x) + \sum_{k=t+1}^{2t+1} \left(\frac{k}{m}\right) \Lambda_{m,2m-4k}(x).$$

Theorem 2.2. *Assume that m is square-free and $m = 4t + 1$. Then*

$$(2.8) \quad s_m(x) = \sum_{k=1}^t \left(\frac{k}{m}\right) \Lambda_{m,m-4k}(x) - \sum_{k=t+1}^{2t} \left(\frac{k}{m}\right) \Lambda_{m,4k-m}(x).$$

Theorem 2.3. *Assume that m is square-free and $m = 4t + 2$. Then*

$$(2.9) \quad s_m(x) = \sum_{\substack{k=1 \\ \text{odd}}}^{4t+1} \left(\frac{-m}{k}\right) \Lambda_{m,k}(x).$$

To prove the above theorems, let us first recall the following formula of $L_m(2n+1)$ obtained by Shanks [15, 17].

Lemma 2.4. *Suppose that $m > 1$ and m is square-free. Then $L_m(2n+1)$ can be expressed as a linear combination of the Fourier series $S_{2n+1}(x)$. More precisely,*

$$L_m(2n+1) = \frac{2}{\sqrt{m}} \sum_k \epsilon_k S_{2n+1}(y_k),$$

where the Jacobi symbols ϵ_k and rational numbers y_k are uniquely determined by m , and $S_{2n+1}(x)$ is defined by

$$S_{2n+1}(x) = \sum_{k=0}^{\infty} \frac{\sin 2\pi(2k+1)x}{(2k+1)^{2n+1}}.$$

Furthermore, we have

$$c_m(x) = \frac{1}{\cos(mx)} \sum_k \epsilon_k \cos(mx(1 - 4y_k)).$$

In fact, Shanks has given an explicit procedure to determine the constants ϵ_k and y_k . To compute ϵ_k and y_k , we use the definition (1.1) of the series $L_m(s)$ and express the Jacobi symbol $\left(\frac{-m}{l}\right)$ as a linear combination of sines according to the following expansion; see [9].

Proposition 2.5. *Assume that l is odd and m satisfies the following two conditions: $m \equiv 1 \pmod{4}$ or $m \equiv 8$ or $12 \pmod{16}$ and $p^2 \nmid m$ for any odd prime p . Then we have*

$$(2.10) \quad \left(\frac{m}{l}\right) = \frac{1}{\sqrt{m}} \sum_{r=1}^{|m|} \left(\frac{m}{r}\right) e^{2\pi i l r / |m|}.$$

In particular, when $m \equiv 3 \pmod{4}$, we have $-m \equiv 1 \pmod{4}$ and we can use the above expansion for $\left(\frac{-m}{l}\right)$. Similarly, when $m \equiv 1 \pmod{4}$, we see that $-4m \equiv 12 \pmod{16}$ so that we can compute $\left(\frac{-4m}{l}\right)$ by using the above formula. Finally, when $m \equiv 2 \pmod{4}$, we find $-4m \equiv 8 \pmod{16}$ so that we can compute $\left(\frac{-4m}{l}\right)$. Note that when l is odd, we have

$$\left(\frac{-4m}{l}\right) = \left(\frac{-m}{l}\right).$$

Thus, the Jacobi symbol $\left(\frac{-m}{l}\right)$ can be determined by the above procedure for $m > 1$.

On the other hand, Shanks [15, 17] provided the following formula for $L_{-m}(2n)$.

Lemma 2.6. *Suppose that $m > 1$ and m is square-free. Then $L_{-m}(2n)$ can be expressed as a linear combination of the Fourier series $C_{2n}(x)$. To be more specific,*

$$L_{-m}(2n) = \frac{2}{\sqrt{m}} \sum_k \epsilon'_k C_{2n}(y'_k),$$

where the Jacobi symbols ϵ'_k and rational numbers y'_k are uniquely determined by m , and $C_{2n}(x)$ is defined by

$$C_{2n}(x) = \sum_{k=0}^{\infty} \frac{\cos 2\pi(2k+1)x}{(2k+1)^{2n}}.$$

Moreover,

$$d_m(x) = \frac{1}{\cos(mx)} \sum_k \epsilon'_k \sin(mx(1 - 4y'_k)).$$

Similarly, Shanks has shown how to compute the constants ϵ'_k and y'_k . In order to compute ϵ'_k and y'_k , recall that

$$L_{-m}(s) = \sum_{\substack{l>0 \\ \text{odd } l}} \left(\frac{m}{l}\right) \frac{1}{l^s},$$

where the Jacobi symbol $\left(\frac{m}{l}\right)$ can be expressed as a linear combination of cosines, resorting to Proposition 2.5.

For the case $m \equiv 3 \pmod{4}$, we find $4m \equiv 12 \pmod{16}$ so that we can use the above expansion for $\left(\frac{4m}{l}\right)$. When $m \equiv 1 \pmod{4}$, we can also compute $\left(\frac{m}{l}\right)$ by using the above expansion. Finally, when $m \equiv 2 \pmod{4}$, we see that $4m \equiv 8 \pmod{16}$ and $\left(\frac{4m}{l}\right)$ can be determined in the same manner. Note that when l is odd, the following relation holds:

$$\left(\frac{4m}{l}\right) = \left(\frac{m}{l}\right).$$

Thus, the Jacobi symbol $\left(\frac{m}{l}\right)$ can be determined for $m > 1$.

Keep in mind that m is assumed to be square-free. Set

$$\hat{c}_m(x) = \cos(mx)c_m(x), \quad \hat{d}_m(x) = \cos(mx)d_m(x), \quad \hat{s}_m(x) = \cos(mx)s_m(x).$$

Proof of Theorem 2.1. Since $m \equiv 3 \pmod{4}$, by using the expansion (2.10) for $\left(\frac{-m}{l}\right)$ and $\left(\frac{4m}{l}\right)$, we have

$$\epsilon_k = \left(\frac{k}{m}\right), \quad y_k = \frac{k}{m}, \quad \epsilon'_k = \left(\frac{m}{k}\right), \quad y'_k = \frac{k}{4m}.$$

Thus,

$$\begin{aligned} L_m(2n+1) &= \frac{2}{\sqrt{m}} \sum_{k=1}^{(m-1)/2} \left(\frac{k}{m}\right) S_{2n+1}\left(\frac{k}{m}\right), \\ L_{-m}(2n) &= \frac{2}{\sqrt{m}} \sum_{\substack{\text{odd } k < m}} \left(\frac{m}{k}\right) C_{2n}\left(\frac{k}{4m}\right). \end{aligned}$$

Therefore, by Lemmas 2.4 and 2.6, we obtain

$$\begin{aligned} \hat{s}_m(x) = \hat{c}_m(x) + \hat{d}_m(x) &= \sum_{k=1}^{(m-1)/2} \left(\frac{k}{m}\right) \cos(m-4k)x \\ &\quad + \sum_{\substack{\text{odd } k < m}} \left(\frac{m}{k}\right) \sin(m-k)x. \end{aligned}$$

Suppose that $m = 4t + 3$. It follows that

$$\begin{aligned} \hat{c}_m(x) &= \sum_{k=1}^t \left(\frac{k}{m}\right) \cos(m-4k)x + \sum_{k=t+1}^{2t+1} \left(\frac{k}{m}\right) \cos(m-4k)x \\ &= \sum_{k=1}^t \left(\frac{k}{m}\right) \cos(m-4k)x + \sum_{k=t+1}^{2t+1} \left(\frac{k}{m}\right) \cos(4k-m)x, \\ \hat{d}_m(x) &= \sum_{\substack{k=1 \\ \text{odd } k}}^{4t+1} \left(\frac{m}{k}\right) \sin(m-k)x. \end{aligned}$$

Thus, we obtain

$$\begin{aligned} \hat{s}_m(x) &= \sum_{k=1}^t \left(\left(\frac{k}{m}\right) \cos(m-4k)x + \left(\frac{m}{m-4k}\right) \sin(4k)x \right) \\ &\quad + \sum_{k=t+1}^{2t+1} \left(\left(\frac{k}{m}\right) \cos(4k-m)x + \left(\frac{m}{4k-m}\right) \sin(2m-4k)x \right). \end{aligned}$$

It remains to verify that

$$(2.11) \quad \left(\frac{k}{m}\right) = \left(\frac{m}{m-4k}\right)$$

for $1 \leq k \leq t$ and

$$(2.12) \quad \left(\frac{k}{m}\right) = \left(\frac{m}{4k-m}\right)$$

for $t+1 \leq k \leq 2t+1$.

As for (2.11), since both m and $m-4k$ are odd positive numbers, if they are coprime, then by the law of quadratic reciprocity, we find

$$\begin{aligned} \left(\frac{m}{m-4k}\right) &= -\left(\frac{m-4k}{m}\right) = -\left(\frac{-4k}{m}\right) = -\left(\frac{-4}{m}\right) \left(\frac{k}{m}\right) \\ &= -\left(\frac{-1}{m}\right) \left(\frac{4}{m}\right) \left(\frac{k}{m}\right) = \left(\frac{2}{m}\right)^2 \left(\frac{k}{m}\right) = \left(\frac{k}{m}\right). \end{aligned}$$

Otherwise, suppose that the greatest common divisor of m and $m-4k$ is greater than 1, i.e., $(m, m-4k) > 1$. From the definition of the Jacobi symbol, it is clear that

$$\left(\frac{m}{m-4k}\right) = 0.$$

We wish to show that

$$\left(\frac{k}{m}\right) = 0.$$

Since $(m, m-4k)|m$, $(m, m-4k)|(m-4k)$ and $m - (m-4k) = 4k$, we deduce that

$$(2.13) \quad (m, m-4k)|4k.$$

But the fact that m is odd implies that $((m, m-4k), 4) = 1$. Thus from (2.13) it follows that $(m, m-4k)|k$. So we conclude that $(m, m-4k)|(m, k)$ and $(m, k) > 1$, which yields

$$\left(\frac{k}{m}\right) = \left(\frac{m}{m-4k}\right) = 0.$$

Similarly, (2.12) can be checked via the following steps if $4k-m$ and m are coprime,

$$\begin{aligned} \left(\frac{m}{4k-m}\right) &= \left(\frac{4k-m}{m}\right) = \left(\frac{-(m-4k)}{m}\right) = \left(\frac{-1}{m}\right) \left(\frac{m-4k}{m}\right) \\ &= -\left(\frac{m-4k}{m}\right) = -\left(\frac{-4k}{m}\right) = \left(\frac{k}{m}\right). \end{aligned}$$

For the case $(m, 4k-m) > 1$, by the definition of the Jacobi symbol, it is evident that

$$\left(\frac{m}{4k-m}\right) = 0.$$

In this case, we claim that

$$\left(\frac{k}{m}\right) = 0.$$

Since $(m, 4k-m)|m$, $(m, 4k-m)|(4k-m)$ and $m + (4k-m) = 4k$, we deduce that

$$(2.14) \quad (m, 4k-m)|4k.$$

Because m is odd, we see that $((m, 4k - m), 4) = 1$. Thus from (2.14) it follows that $(m, 4k - m) | k$. Therefore, we obtain that $(m, 4k - m) | (m, k)$ and $(m, k) > 1$, which implies

$$\left(\frac{k}{m}\right) = \left(\frac{m}{4k - m}\right) = 0.$$

So we find that

$$\begin{aligned} s_m(x) &= \sum_{k=1}^t \left(\frac{k}{m}\right) \frac{\cos(m - 4k)x + \sin(4k)x}{\cos mx} \\ &\quad + \sum_{k=t+1}^{2t+1} \left(\frac{k}{m}\right) \frac{\cos(4k - m)x + \sin(2m - 4k)x}{\cos mx}. \end{aligned}$$

This completes the proof. $\square$

Proof of Theorem 2.2. Under the condition $m \equiv 1 \pmod{4}$, by applying the expansion (2.10) to $\left(\frac{-m}{l}\right)$ and $\left(\frac{m}{l}\right)$, we get

$$\epsilon_k = \left(\frac{-m}{k}\right), \quad y_k = \frac{k}{4m}, \quad \epsilon'_k = \left(\frac{k}{m}\right), \quad y'_k = \frac{k}{m}.$$

It follows that

$$\begin{aligned} L_m(2n + 1) &= \frac{2}{\sqrt{m}} \sum_{\text{odd } k < m} \left(\frac{-m}{k}\right) S_{2n+1}\left(\frac{k}{4m}\right), \\ L_{-m}(2n) &= \frac{2}{\sqrt{m}} \sum_{k=1}^{(m-1)/2} \left(\frac{k}{m}\right) C_{2n}\left(\frac{k}{m}\right). \end{aligned}$$

In view of Lemmas 2.4 and 2.6, we find

$$\hat{s}_m(x) = \hat{c}_m(x) + \hat{d}_m(x) = \sum_{\text{odd } k < m} \left(\frac{-m}{k}\right) \cos(m - k)x + \sum_{k=1}^{(m-1)/2} \left(\frac{k}{m}\right) \sin(m - 4k)x.$$

Writing $m = 4t + 1$, we obtain

$$\begin{aligned} \hat{d}_m(x) &= \sum_{k=1}^t \left(\frac{k}{m}\right) \sin(m - 4k)x + \sum_{k=t+1}^{2t} \left(\frac{k}{m}\right) \sin(m - 4k)x \\ &= \sum_{k=1}^t \left(\frac{k}{m}\right) \sin(m - 4k)x - \sum_{k=t+1}^{2t} \left(\frac{k}{m}\right) \sin(4k - m)x, \\ \hat{c}_m(x) &= \sum_{\substack{k=1 \\ \text{odd } k}}^{4t-1} \left(\frac{-m}{k}\right) \cos(m - k)x. \end{aligned}$$

Consequently,

$$\begin{aligned} \hat{s}_m(x) &= \sum_{k=1}^t \left(\left(\frac{k}{m}\right) \sin(m - 4k)x + \left(-\frac{m}{m - 4k}\right) \cos(4k)x \right) \\ &\quad + \sum_{k=t+1}^{2t} \left(-\left(\frac{k}{m}\right) \sin(4k - m)x + \left(-\frac{m}{4k - m}\right) \cos(2m - 4k)x \right). \end{aligned}$$

Finally, we need to show that

$$(2.15) \quad \left(\frac{k}{m}\right) = \left(-\frac{m}{m-4k}\right)$$

for $1 \leq k \leq t$ and

$$(2.16) \quad -\left(\frac{k}{m}\right) = \left(-\frac{m}{4k-m}\right)$$

for $t+1 \leq k \leq 2t$.

To verify (2.15), we consider two cases. If $(m, m-4k) = 1$, since both m and $m-4k$ are odd, we may employ the law of quadratic reciprocity to deduce that

$$\begin{aligned} \left(-\frac{m}{m-4k}\right) &= \left(\frac{-1}{m-4k}\right) \left(\frac{m}{m-4k}\right) = \left(\frac{m}{m-4k}\right) = \left(\frac{m-4k}{m}\right) = \left(\frac{-4k}{m}\right) \\ &= \left(\frac{-4}{m}\right) \left(\frac{k}{m}\right) = \left(\frac{-1}{m}\right) \left(\frac{2}{m}\right)^2 \left(\frac{k}{m}\right) = \left(\frac{k}{m}\right). \end{aligned}$$

For the case $(m, m-4k) > 1$, by definition,

$$\left(-\frac{m}{m-4k}\right) = 0.$$

We wish to show that

$$\left(\frac{k}{m}\right) = 0.$$

Since $(m, m-4k)|m$, $(m, m-4k)|(m-4k)$ and $m-(m-4k) = 4k$, we get

$$(2.17) \quad (m, m-4k)|4k.$$

However, the fact that m is odd implies that $((m, m-4k), 4) = 1$. Thus, from (2.17) we see that $(m, m-4k)|k$. It follows that $(m, m-4k)|(m, k)$ and $(m, k) > 1$. Hence

$$\left(\frac{k}{m}\right) = \left(-\frac{m}{m-4k}\right) = 0.$$

To prove (2.16), we also consider two cases. If m and $4k-m$ are coprime, then we have

$$\begin{aligned} -\left(-\frac{m}{4k-m}\right) &= -\left(\frac{-1}{4k-m}\right) \left(\frac{m}{4k-m}\right) = \left(\frac{m}{4k-m}\right) = \left(\frac{4k-m}{m}\right) \\ &= \left(\frac{-(m-4k)}{m}\right) = \left(\frac{-1}{m}\right) \left(\frac{m-4k}{m}\right) = \left(\frac{-4k}{m}\right) = \left(\frac{k}{m}\right). \end{aligned}$$

If $(m, 4k-m) > 1$, then we get

$$-\left(-\frac{m}{4k-m}\right) = 0.$$

We need to show that

$$\left(\frac{k}{m}\right) = 0.$$

Since $(m, 4k-m)|m$, $(m, 4k-m)|(4k-m)$ and $m+(4k-m) = 4k$, we find

$$(2.18) \quad (m, 4k-m)|4k.$$

But m is odd, we see that $((m, 4k - m), 4) = 1$. From (2.18) it follows that $(m, 4k - m) \nmid k$. This implies that $(m, 4k - m) \mid (m, k)$ and $(m, k) > 1$. Consequently,

$$\left(\frac{k}{m}\right) = -\left(-\frac{m}{4k - m}\right) = 0.$$

Thus we obtain

$$\begin{aligned} s_m(x) &= \sum_{k=1}^t \left(\frac{k}{m}\right) \frac{\cos(4k)x + \sin(m - 4k)x}{\cos mx} \\ &\quad - \sum_{k=t+1}^{2t} \left(\frac{k}{m}\right) \frac{\cos(2m - 4k)x + \sin(4k - m)x}{\cos mx}. \end{aligned}$$

This completes the proof. $\square$

Proof of Theorem 2.3. Since $m \equiv 2 \pmod{4}$, by using the expansion (2.10) for $\left(\frac{-4m}{l}\right)$ and $\left(\frac{4m}{l}\right)$, we obtain

$$\epsilon_k = \left(\frac{-m}{k}\right), \quad y_k = \frac{k}{4m}, \quad \epsilon'_k = \left(\frac{m}{k}\right), \quad y'_k = \frac{k}{4m}.$$

Hence

$$\begin{aligned} L_m(2n + 1) &= \frac{2}{\sqrt{m}} \sum_{\text{odd } k < m} \left(\frac{-m}{k}\right) S_{2n+1}\left(\frac{k}{4m}\right), \\ L_{-m}(2n) &= \frac{2}{\sqrt{m}} \sum_{\text{odd } k < m} \left(\frac{m}{k}\right) C_{2n}\left(\frac{k}{4m}\right). \end{aligned}$$

By Lemmas 2.4 and 2.6, we see that

$$\hat{s}_m(x) = \hat{c}_m(x) + \hat{d}_m(x) = \sum_{\text{odd } k < m} \left(\frac{-m}{k}\right) \cos(m - k)x + \sum_{\text{odd } k < m} \left(\frac{m}{k}\right) \sin(m - k)x.$$

Writing $m = 4t + 2$, since

$$\left(\frac{4t + 2}{2t + 1}\right) = 0,$$

we obtain

$$\hat{s}_m(x) = \sum_{\substack{k=1 \\ \text{odd } k}}^{4t+1} \left(\left(\frac{-m}{k}\right) \cos(m - k)x + \left(\frac{m}{m - k}\right) \sin(k)x \right).$$

Finally, it follows from $-k \equiv m - k \pmod{m}$ that

$$(2.19) \quad \left(\frac{-m}{k}\right) = \left(\frac{m}{m - k}\right)$$

for $1 \leq k \leq 4t + 1$. Hence we conclude that

$$s_m(x) = \sum_{\substack{k=1 \\ \text{odd } k}}^{4t+1} \left(\frac{-m}{k}\right) \frac{\cos(m - k)x + \sin(k)x}{\cos mx}.$$

This completes the proof. $\square$

For $m = 5, 6, 7$, the generating functions $s_m(x)$ have been given in the Introduction.

3. COMPUTATION FOR $s_m(x)$ WHEN m IS NOT SQUARE-FREE

In this section, we obtain an expression for $s_m(x)$ for the case when m is not square-free. Assume that m can be divided by a square u^2 and $u > 1$. Recall that the constant K_b is given by (1.4).

Theorem 3.1. *Suppose that $m = bu^2$, where b is square-free and $u > 1$. Then we can express $s_m(x)$ as a linear combination of the four functions*

$$w(b, t) \sec(btx)(\pm \cos((b-p)tx) \pm \sin(ptx)),$$

where p is a nonnegative integer not exceeding b , $t|u^2$ and $w(b, t) = K_b t/u$.

The idea of the proof is to establish two recursive relations (3.21) and (3.22) between $s_{m,n}$ and $s_{b,n}$. Then we express $s_m(x)$ as a linear combination of the functions $c_b(tx)$ and $d_b(tx)$ by considering the two cases according to whether there exist odd prime factors u_i of u with residues 3 modulo 4. Since b is square-free, $c_b(tx)$ and $d_b(tx)$ can be evaluated by using the formulas in the previous section.

Proof. Let us start with the following relation given by Shanks [15]

$$(3.20) \quad L_m(s) = L_b(s) \prod_{u_i|u} \left(1 - \left(\frac{-b}{u_i} \right) \frac{1}{u_i^s} \right),$$

where the product ranges over odd primes u_i (if any) that divide u . To be precise, in case there are no odd prime factors, the empty product is defined to be 1. From the definitions (1.2) and (1.3) it follows that

$$(3.21) \quad c_{m,n} = K_b u(u^2)^{2n} \prod_i \left(u_i^{2n+1} - \left(\frac{-b}{u_i} \right) \right) \left(\prod_i \frac{1}{u_i} \right)^{2n+1} c_{b,n},$$

$$(3.22) \quad d_{m,n} = K_b u(u^2)^{2n-1} \prod_i \left(u_i^{2n} - \left(\frac{b}{u_i} \right) \right) \left(\prod_i \frac{1}{u_i} \right)^{2n} d_{b,n}.$$

For the purpose of computing $s_m(x)$ for the case when m is not square-free, we need to consider two cases according to whether there exist $u_i \equiv 3 \pmod{4}$ among the k odd factors $u_1, u_2, \dots, u_k$ of u .

Case 1. $u_i \equiv 1 \pmod{4}$ for $1 \leq i \leq k$. In this case, we see that

$$\left(\frac{-b}{u_i} \right) = \left(\frac{b}{u_i} \right).$$

Suppose that among the factors $u_1, u_2, \dots, u_k$ there are k_1 primes $u_1, u_2, \dots, u_{k_1}$ satisfying $\left(\frac{b}{u_i} \right) = 1$ for $1 \leq i \leq k_1$, k_2 primes $u_{k_1+1}, u_{k_1+2}, \dots, u_{k_1+k_2}$ satisfying $\left(\frac{b}{u_{k_1+j}} \right) = -1$ for $1 \leq j \leq k_2$, and k_3 primes $u_{k_1+k_2+1}, u_{k_1+k_2+2}, \dots, u_{k_1+k_2+k_3}$ satisfying $\left(\frac{b}{u_{k_1+k_2+l}} \right) = 0$ for $1 \leq l \leq k_3$, where $k_1 + k_2 + k_3 = k$. From (3.21) it is clear that

$$(3.23) \quad c_{m,n} = K_b u(u^2)^{2n} \frac{\prod_{i=1}^{k_1} (u_i^{2n+1} - 1) \times \prod_{j=1}^{k_2} (u_{k_1+j}^{2n+1} + 1)}{\prod_{i=1}^{k_1+k_2} u_i^{2n+1}} c_{b,n}.$$

Let

$$f_c = \frac{\prod_{i=1}^{k_1} (u_i^{2n+1} - 1) \times \prod_{j=1}^{k_2} (u_{k_1+j}^{2n+1} + 1)}{\prod_{i=1}^{k_1+k_2} u_i^{2n+1}}.$$

In this notation, (3.23) can be rewritten as

$$c_{m,n} = K_b u (u^2)^{2n} f_c c_{b,n},$$

which implies that

$$(3.24) \quad c_m(x) = K_b u \sum_{n \geq 0} f_c c_{b,n} \frac{(u^2 x)^{2n}}{(2n)!}.$$

Since

$$\prod_{i=1}^{k_1} (u_i^{2n+1} - 1) = \prod_{i=1}^{k_1} u_i^{2n+1} - \sum_{i=1}^{k_1} \left(\prod_{j=1}^{k_1} u_j \cdots u_{i-1} u_{i+1} \cdots u_{k_1} \right)^{2n+1} + \cdots + (-1)^{k_1}$$

and

$$\begin{aligned} \prod_{j=1}^{k_2} (u_{k_1+j}^{2n+1} + 1) &= \prod_{j=1}^{k_2} u_{k_1+j}^{2n+1} \\ &+ \sum_{j=1}^{k_2} \left(\prod_{i=1}^{k_2} u_{k_1+i} \cdots u_{k_1+j-1} u_{k_1+j+1} \cdots u_{k_1+k_2} \right)^{2n+1} + \cdots + 1, \end{aligned}$$

we can expand f_c as follows

$$(3.25) \quad \begin{aligned} f_c &= 1 + \sum_{j=1}^{k_2} \frac{1}{u_{k_1+j}^{2n+1}} + \cdots + \frac{1}{\prod_{j=1}^{k_2} u_{k_1+j}^{2n+1}} - \sum_{i=1}^{k_1} \frac{1}{u_i^{2n+1}} - \sum_{i=1}^{k_1} \sum_{j=1}^{k_2} \frac{1}{u_i^{2n+1} u_{k_1+j}^{2n+1}} - \cdots \\ &- \sum_{i=1}^{k_1} \frac{1}{u_i^{2n+1} \prod_{j=1}^{k_2} u_{k_1+j}^{2n+1}} + \cdots + \frac{(-1)^{k_1}}{\prod_{i=1}^{k_1+k_2} u_i^{2n+1}}. \end{aligned}$$

Plugging (3.25) into (3.24), we find that $c_m(x)$ is a linear combination of the functions $c_b(tx)$, where $t|u^2$ and the coefficient of $c_b(tx)$ equals $K_b t/u$.

Similarly, we have

$$(3.26) \quad d_{m,n} = K_b u (u^2)^{2n-1} \frac{\prod_{i=1}^{k_1} (u_i^{2n} - 1) \times \prod_{j=1}^{k_2} (u_{k_1+j}^{2n} + 1)}{\prod_{i=1}^{k_1+k_2} u_i^{2n}} d_{b,n}.$$

Let

$$f_d = \frac{\prod_{i=1}^{k_1} (u_i^{2n} - 1) \times \prod_{j=1}^{k_2} (u_{k_1+j}^{2n} + 1)}{\prod_{i=1}^{k_1+k_2} u_i^{2n}},$$

then (3.26) can be rewritten as

$$d_{m,n} = K_b u (u^2)^{2n-1} f_d d_{b,n},$$

which leads to the relation

$$(3.27) \quad d_m(x) = K_b u \sum_{n \geq 1} f_d d_{b,n} \frac{(u^2 x)^{2n-1}}{(2n-1)!}.$$

Again, we may expand f_d as follows:

$$(3.28) \quad \begin{aligned} f_d = & 1 + \sum_{j=1}^{k_2} \frac{1}{u_{k_1+j}^{2n}} + \cdots + \frac{1}{\prod_{j=1}^{k_2} u_{k_1+j}^{2n}} - \sum_{i=1}^{k_1} \frac{1}{u_i^{2n}} - \sum_{i=1}^{k_1} \sum_{j=1}^{k_2} \frac{1}{u_i^{2n} u_{k_1+j}^{2n}} - \cdots \\ & - \sum_{i=1}^{k_1} \frac{1}{u_i^{2n} \prod_{j=1}^{k_2} u_{k_1+j}^{2n}} + \cdots + \frac{(-1)^{k_1}}{\prod_{i=1}^{k_1+k_2} u_i^{2n}}. \end{aligned}$$

Substituting (3.28) into (3.27), we find that $d_m(x)$ is a linear combination of the functions $d_b(tx)$, where $t|u^2$ and the coefficient of $d_b(tx)$ equals $K_b t/u$. Furthermore, we see that $c_m(x)$ and $d_m(x)$ have the same coefficients for the linear combinations. In other words, the relation for $c_m(x)$ and $c_b(tx)$ is still valid after changing $c_m(x)$ and $c_b(tx)$ to $d_m(x)$ and $d_b(tx)$, respectively. Therefore, in this case, $s_m(x)$ can be expressed as a sum of the functions $w(b, t)(c_b(tx) + d_b(tx))$, where $t|u^2$ and $w(b, t) = K_b t/u$.

Case 2. Among the k primes $u_1, u_2, \dots, u_k$, there exist q primes $u_{i_1}, u_{i_2}, \dots, u_{i_q}$ with residue 3 modulo 4. To compute $s_m(x)$, we first consider the case when $q = 1$, and then argue that the case $q > 1$ can be dealt with in the same way.

Since $q = 1$, we assume that the first $k-1$ odd primes $u_1, u_2, \dots, u_{k-1}$ satisfy $u_i \equiv 1 \pmod{4}$ for $1 \leq i \leq k-1$, and suppose that the last prime u_k satisfies $u_k \equiv 3 \pmod{4}$, or

$$\left(\frac{-b}{u_k}\right) = -\left(\frac{b}{u_k}\right).$$

We now define the indices k_1, k_2 and k_3 as in Case 1 except that $k_1 + k_2 + k_3 = k-1$ since there are $k-1$ primes with residue 1 modulo 4. This leads us to consider two subcases according to whether $\left(\frac{-b}{u_k}\right)$ equals 0. Keep in mind that we have $q = 1$ in these two subcases.

On one hand, in order to use the two recursive relations (3.21) and (3.22) between $s_{m,n}$ and $s_{b,n}$, we may assume that $\left(\frac{-b}{u_k}\right) = 0$. Therefore, the term u_k on the denominator and the same term on the numerator cancel each other in (3.21). This argument also applies to the relation (3.22). In other words, there is no need to consider the occurrence of the term u_k^{2n+1} in (3.21) and the term u_k^{2n} in (3.22). In this sense, it remains to consider the other $k-1$ primes $u_1, u_2, \dots, u_{k-1}$ such that $u_i \equiv 1 \pmod{4}$ for $1 \leq i \leq k-1$. By the argument in Case 1, we see again that $s_m(x)$ can be expressed as a sum of the terms $w(b, t)(c_b(tx) + d_b(tx))$, where $t|u^2$ and $w(b, t) = K_b t/u$.

On the other hand, we should consider the case when $\left(\frac{-b}{u_k}\right) = 1$ or $\left(\frac{-b}{u_k}\right) = -1$. Since the proofs for these two cases are similar, we only give the proof for the case

$\left(\frac{-b}{u_k}\right) = -1$. From (3.21) it follows that

$$\begin{aligned} c_{m,n} &= K_b u(u^2)^{2n} \frac{\prod_{i=1}^{k_1} (u_i^{2n+1} - 1) \prod_{j=1}^{k_2} (u_{k_1+j}^{2n+1} + 1)}{\prod_{i=1}^{k_1+k_2} u_i^{2n+1}} \times \frac{u_k^{2n+1} + 1}{u_k^{2n+1}} c_{b,n} \\ &= K_b u(u^2)^{2n} \left(\frac{\prod_{i=1}^{k_1} (u_i^{2n+1} - 1) \prod_{j=1}^{k_2} (u_{k_1+j}^{2n+1} + 1)}{\prod_{i=1}^{k_1+k_2} u_i^{2n+1}} \right. \\ &\quad \left. + \frac{\prod_{i=1}^{k_1} (u_i^{2n+1} - 1) \prod_{j=1}^{k_2} (u_{k_1+j}^{2n+1} + 1)}{u_k^{2n+1} \prod_{i=1}^{k_1+k_2} u_i^{2n+1}} \right) c_{b,n}. \end{aligned}$$

Let

$$g_b(x) = K_b u \sum_{n \geq 0} \left(\frac{\prod_{i=1}^{k_1} (u_i^{2n+1} - 1) \prod_{j=1}^{k_2} (u_{k_1+j}^{2n+1} + 1)}{\prod_{i=1}^{k_1+k_2} u_i^{2n+1}} \right) c_{b,n} \frac{(u^2 x)^{2n}}{(2n)!}.$$

Then we have

$$(3.29) \quad c_m(x) = g_b(x) + \frac{1}{u_k} g_b(x/u_k).$$

In fact, by the argument in Case 1, we see that $g_b(x)$ is a linear combination of the functions $c_b(tx)$, where $t|u^2$ and the coefficient of $c_b(tx)$ equals $K_b t/u$.

Similarly, by (3.22) we find

$$\begin{aligned} d_{m,n} &= K_b u(u^2)^{2n-1} \frac{\prod_{i=1}^{k_1} (u_i^{2n} - 1) \prod_{j=1}^{k_2} (u_{k_1+j}^{2n} + 1)}{\prod_{i=1}^{k_1+k_2} u_i^{2n}} \times \frac{u_k^{2n} - 1}{u_k^{2n}} d_{b,n} \\ &= K_b u(u^2)^{2n-1} \left(\frac{\prod_{i=1}^{k_1} (u_i^{2n} - 1) \prod_{j=1}^{k_2} (u_{k_1+j}^{2n} + 1)}{\prod_{i=1}^{k_1+k_2} u_i^{2n}} \right. \\ &\quad \left. - \frac{\prod_{i=1}^{k_1} (u_i^{2n} - 1) \prod_{j=1}^{k_2} (u_{k_1+j}^{2n} + 1)}{u_k^{2n} \prod_{i=1}^{k_1+k_2} u_i^{2n}} \right) d_{b,n}. \end{aligned}$$

Let

$$h_b(x) = K_b u \sum_{n \geq 1} \left(\frac{\prod_{i=1}^{k_1} (u_i^{2n} - 1) \prod_{j=1}^{k_2} (u_{k_1+j}^{2n} + 1)}{\prod_{i=1}^{k_1+k_2} u_i^{2n}} \right) d_{b,n} \frac{(u^2 x)^{2n-1}}{(2n-1)!}.$$

We get

$$(3.30) \quad d_m(x) = h_b(x) - \frac{1}{u_k} h_b(x/u_k).$$

Again, from the reasoning in Case 1 it follows that $h_b(x)$ is a linear combination of the functions $d_b(tx)$, where $t|u^2$ and the coefficient of $d_b(tx)$ equals $K_b t/u$. Combining the equations (3.29) and (3.30) yields the following relation:

$$s_m(x) = g_b(x) + h_b(x) + \frac{1}{u_k} (g_b(x/u_k) - h_b(x/u_k)).$$

Thus we obtain that $s_m(x)$ can also be expressed as a linear combination of the functions $w(b, t)(\pm c_b(tx) \pm d_b(tx))$, where $t|u^2$ and $w(b, t) = K_b t/u$.

Finally, as mentioned before we shall show that the justification for the above two subcases can be applied to the case for $q > 1$. Let us give an example for $q = 2$. Suppose that u_{k-1} and u_k are the last two primes such that $u_{k-1} \equiv 3 \pmod{4}$,

$u_k \equiv 3 \pmod{4}$, and $\left(\frac{-b}{u_{k-1}}\right) = -1$, $\left(\frac{-b}{u_k}\right) = -1$. Using the functions $g_b(x)$ and $h_b(x)$ as given before, we can deduce the following formula for $s_m(x)$:

$$\begin{aligned} s_m(x) = & g_b(x) + h_b(x) + \frac{1}{u_{k-1}}(g_b(x/u_{k-1}) - h_b(x/u_{k-1})) \\ & + \frac{1}{u_k}(g_b(x/u_k) - h_b(x/u_k)) \\ & + \frac{1}{u_{k-1}u_k}(g_b(x/u_{k-1}u_k) + h_b(x/u_{k-1}u_k)). \end{aligned}$$

Clearly, now $s_m(x)$ can also be written as a linear combination of the functions $w(b, t)(\pm c_b(tx) \pm d_b(tx))$, where $t|u^2$ and $w(b, t) = K_b t/u$. For the other conditions on u_{k-1} and u_k , the computation can be done in the same spirit.

In summary, $s_m(x)$ can be expressed as a linear combination of the functions $w(b, t)(\pm c_b(tx) \pm d_b(tx))$, where $t|u^2$ and $w(b, t) = K_b t/u$. Since b is square-free, $c_b(tx)$ and $d_b(tx)$ can be written as a linear combination of the functions $\sec(btx) \cos((b-p)tx)$ and $\sec(btx) \sin(ptx)$, respectively, by using the formulas in the previous section. This completes the proof. $\square$

Here we give three examples corresponding to the above three cases. For Case 1, suppose that $m = 3(5 \times 13)^2 = 3(65)^2 = 3 \times 4225 = 12675$. Then we have

$$s_{12675}(x) = 65s_3(4225x) - 5s_3(325x) + 13s_3(845x) - s_3(65x),$$

where

$$s_3(x) = \sec(3x)(\sin 2x + \cos x).$$

For the first subcase of Case 2, suppose that $m = 6(5 \times 3)^2 = 3(15)^2 = 6 \times 225 = 1350$. Then we get

$$s_{1350}(x) = 15s_6(225x) - 3s_6(45x),$$

where

$$s_6(x) = \sec(6x)(\cos 5x + \sin x) + \sec(6x)(\cos x + \sin 5x).$$

For the second subcase of Case 2, assume that $m = 225 = (5 \times 3)^2$. We find

$$\begin{aligned} 2s_{225}(x) = & 15(c_1(225x) + d_1(225x)) - 3(c_1(45x) + d_1(45x)) \\ & + 5(c_1(75x) - d_1(75x)) - (c_1(15x) - d_1(15x)), \end{aligned}$$

where

$$c_1(x) = \sec x, \quad d_1(x) = \tan x.$$

4. A COMBINATORIAL INTERPRETATION FOR $s_{m,n}$

In this section, we aim to give a combinatorial interpretation of $s_{m,n}$ when m is square-free based on the generating function formula for $s_m(x)$. Let us recall the known combinatorial interpretations of $s_{m,n}$ when $m = 1, 2, 3, 4$.

For $m = 1$, $(s_{1,n})_{n \geq 0}$ is called the sequence of Euler numbers. Let E_n be the n -th Euler number, that is, the number of up-down permutations on $[n] = \{1, 2, \dots, n\}$, which are also called snakes of type A_{n-1} by Arnol'd [2]. The following generating function is due to André [1]:

$$\sum_{n \geq 0} E_n \frac{x^n}{n!} = \sec x + \tan x.$$

Note that Springer also gave an explanation of the Euler numbers in terms of the irreducible root system A_{n-1} and derived the generating function of André in this context.

For $m = 2$, the sequence $(s_{2,n})_{n \geq 0}$ turns out to be the sequence of Springer numbers of the irreducible root system B_n [20]. Purtill [13] has found an interpretation of this sequence. Let P_n be the n -th entry of this sequence, whereas Purtill used the notation $E_n^\pm$. He has shown that P_n equals the number of André signed permutations on $[n]$. On the other hand, it has been shown by Arnol'd [2] that S_n also counts the number of snakes of type B_n . Hoffman [7] has derived the generating function of the number of snakes of type B_n by giving a direct combinatorial proof.

For $m = 3$, the sequence $(s_{3,n})_{n \geq 0}$ has been studied by Ehrenborg and Readdy [4]. Let F_n denote the n -th Ehrenborg and Readdy number, which was denoted by $|ER_n|$; see Hoffman [7]. It has been shown that F_n equals the number Λ -alternating augmented 3-signed permutations on $[n]$. Meanwhile, Hoffman [7] presented another combinatorial interpretation of the sequence in the case $m = 3$ in terms of ER_n -snakes in the spirit of the snakes of type A_{n-1} and of type B_n .

For $m = 4$, the sequence $(s_{4,n})_{n \geq 0}$ has also been studied by Ehrenborg and Readdy [5, p. 719]. In fact, they introduced the concept of nonaugmented André R -signed permutations on a finite set N with $|N| = n$, where $R = (r_i)_{i \in N}$ is a vector indexed by N . They have shown that the exponential generating function for the number of nonaugmented André R -signed permutations with the n -tuple $R = (r, r, \dots, r)$ is $\sec(rx) + \tan(rx)$. Therefore, for the n -tuple $R = (4, 4, \dots, 4)$, it follows that the number of nonaugmented André R -signed permutations is counted by $s_{4,n}$.

We proceed to give a combinatorial interpretation for $s_{m,n}$ when m is square-free based on the generating function $s_m(x)$. More precisely, in this case we shall show that $s_m(x)$ can be expressed by the generating function $\Lambda_{m,p}(x)$ for the numbers $\Lambda_{m,p,n}$ defined by Ehrenborg and Readdy [4]. Consequently, when m is square-free, the numbers $s_{m,n}$ can be written as a linear combination of the numbers $\Lambda_{m,p,n}$ with coefficients being 1 or -1 . With the aid of the combinatorial interpretation of the generating function due to Ehrenborg and Readdy, we see that $s_{m,n}$ can be explained in terms of maximal chains in the m -cubical lattice.

When m is not square-free, write $m = bu^2$. The numbers $s_{m,n}$ or $2s_{m,n}$ can be written as a linear combination of the numbers $\Lambda_{bt,pt,n}$ with integer coefficients, where $t|u^2$ and $0 \leq p \leq b$. However, these integer coefficients are obtained from the generating function of $s_m(x)$. It would be interesting to find a combinatorial explanation of these coefficients when m is not square-free.

We now need some definitions. Ehrenborg and Readdy [4] defined a class of posets called the Sheffer posets, which can be viewed as a generalization of the binomial posets introduced by Stanley [21]. As an important example, they studied the r -cubical lattice, which is a set of ordered r -tuples $(A_1, A_2, \dots, A_r)$ of subsets from an infinite set I together with the reverse inclusion order and with a minimum element $\hat{0}$ adjoined. Note that the r -cubical lattice has been studied by Metropolis, Rota, Strehl and White [10]. Ehrenborg and Readdy further generalized the concept of R -labelings to linear edge-labelings. By considering the set of maximal chains in the interval $[\hat{0}, \hat{1}]$ on the Hasse diagram of the r -cubical lattice, they derived a formula for the number of Λ -alternating augmented r -signed permutations.

To describe the definition of Λ -alternating augmented r -signed permutations, Ehrenborg and Readdy constructed a linear edge-labeling on the Hasse diagram of the r -cubical lattice. To be more specific, for an edge corresponding to the cover relation $A < B$ with $A \neq \hat{0}$, let (i, j) denote its label where i equals the unique index such that $A_i \neq B_i$ and j is the singleton element in $A_i - B_i$. Let G be the label of the edge corresponding to $\hat{0} < A$, which is called the special element by Ehrenborg and Readdy. Then an augmented r -signed permutation is a list $(G, (i_1, j_1), (i_2, j_2), \dots, (i_n, j_n))$, where $i_1, i_2, \dots, i_n \in [r]$, and $(j_1, j_2, \dots, j_n)$ forms a permutation on $[n]$. In other words, r -signed permutations are permutations on $[n]$ in which each element is assigned one of the r signs.

To define the descent set of r -signed permutations, let Λ be the set of such labels of the edges on the Hasse diagram of the r -cubical lattice. As is easily seen,

$$\Lambda = ([r] \times [n]) \cup \{G\}.$$

Let p be an integer such that $0 \leq p \leq r$. For fixed positive integers r , n and p , we can define a linear order on Λ which satisfies the conditions

$$(4.31) \quad (i, j) <_{\Lambda} G \Rightarrow i \leq r - p$$

and

$$(4.32) \quad (i, j) >_{\Lambda} G \Rightarrow i > r - p,$$

where (i, j) is the label of the edge corresponding to the cover relation $A < B$ such that A covers $\hat{0}$, and G is the special element. For the remaining labels, we may arrange them in the lexicographic order. The descent set of an augmented r -signed permutation $(g_0, g_1, \dots, g_n)$ is defined as the set $\{k: g_{k-1} >_{\Lambda} g_k\}$, where $g_0 = G$ and $g_k = (i_k, j_k)$ for $1 \leq k \leq n$. Therefore, for an Λ -alternating augmented r -signed permutation, that is, permutation having descent set $\{2, 4, 6, \dots\}$, it is necessary to have the condition $(i_1, j_1) >_{\Lambda} G$, or $i_1 > r - p$. In other words, the labels above G in this ordering are those whose first coordinate belongs to the set $\{r - p + 1, r - p + 2, \dots, r\}$.

Recall that $\Lambda_{r,p,n}$ denotes the number of Λ -alternating augmented r -signed permutations. Since there is a one-to-one correspondence between the set of maximal chains in the r -cubical lattice and the set of augmented r -signed permutations, $\Lambda_{r,p,n}$ also equals the number of maximal chains with descent set $\{2, 4, 6, \dots\}$ in the r -cubical lattice for which the first nonspecial edge has label (i_1, j_1) with $r - p + 1 \leq i_1 \leq r$. Here the first nonspecial edge is defined as the edge corresponding to the cover relation $A < B$ such that A covers $\hat{0}$. Based on this observation, Ehrenborg and Readdy derived the following generating function for $\Lambda_{r,p,n}$:

$$(4.33) \quad \Lambda_{r,p}(x) = \sum_{n \geq 0} \Lambda_{r,p,n} \frac{x^n}{n!} = \frac{\cos((r-p)x) + \sin(px)}{\cos(rx)}.$$

In view of the above formula (4.33) and the generating functions for the generalized Euler and class numbers $s_{m,n}$ in the previous sections, for any square-free m we can give a combinatorial interpretation of $s_{m,n}$ in terms of the augmented m -signed permutations, or equivalently, in terms of the maximal chains in the Hasse diagram of the m -cubical lattice.

When m is square-free, as is shown in Section 2, $s_{m,n}$ is a linear combination of the numbers $\Lambda_{m,p,n}$ with coefficients being 1 or -1 . For example, when $m = 7$,

since

$$s_7(x) = \frac{\cos 3x + \sin 4x}{\cos 7x} + \frac{\cos x + \sin 6x}{\cos 7x} - \frac{\cos 5x + \sin 2x}{\cos 7x},$$

we obtain that

$$s_{7,n} = \Lambda_{7,4,n} + \Lambda_{7,6,n} - \Lambda_{7,2,n}.$$

When m is not square-free, assume that $m = bu^2$ and $u > 1$. By Theorem (3.1), $K_b^{-1}s_{m,n}$ can be written as a linear combination of the numbers $\Lambda_{bt,pt,n}$ with integer coefficients. Recall that $K_1^{-1} = 2$ and $K_b^{-1} = 1$ for $b \neq 1$. For instance, we have

$$2s_9(x) = 3(\sec(9x) + \tan(9x)) + (\sec(3x) - \tan(3x)),$$

which implies

$$2s_{9,n} = 3\Lambda_{9,9,n} + (-1)^n \Lambda_{3,3,n}.$$

ACKNOWLEDGMENTS

We wish to thank the referee for valuable suggestions. This work was supported by the 973 Project, the PCSIRT Project of the Ministry of Education, and the National Science Foundation of China.

REFERENCES

- [1] D. André, Sur les permutations alternées, *J. Math. Pures Appl.* 7 (1881), 167–184.
- [2] V.I. Arnol'd, The calculus of snakes and the combinatorics of Bernoulli, Euler, and Springer numbers of Coxeter groups, *Uspekhi Mat. Nauk.* 47 (1992), 3–45 (Russian); *Russian Math. Surveys* 47 (1992), 1–51. MR1171862 (93h:20042)
- [3] P.T. Bateman and R.A. Horn, A heuristic asymptotic formula concerning the distribution of prime numbers, *Math. Comp.* 16 (1962), 363–367. MR0148632 (26:6139)
- [4] R. Ehrenborg and M.A. Readdy, Sheffer posets and r -signed permutations, *Ann. Sci. Math. Québec* 19 (1995), 173–196. MR1365825 (96m:06005)
- [5] R. Ehrenborg and M.A. Readdy, The r -cubical lattice and a generalization of the cd -index, *Europ. J. Combin.* 17 (1996), 709–725. MR1415312 (97k:06014)
- [6] G.H. Hardy, Ramanujan, Twelve lectures on subjects suggested by his life and work, New York, Chelsea Publishing Company, 1959. MR0106147 (21:4881)
- [7] M.E. Hoffman, Derivative polynomials, Euler polynomials, and associated integer sequences, *Electron. J. Combin.* 6 (1999), R21. MR1685701 (2000c:11027)
- [8] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, Springer-Verlag, 1990. MR1070716 (92e:11001)
- [9] E. Landau, *Elementary Number Theory*, New York, Chelsea Publishing Company, 1958. MR0092794 (19:1159d)
- [10] N. Metropolis, G.-C. Rota, V. Strehl and N. White, Partitions into chains of a class of partially ordered sets, *Proc. Amer. Math. Soc.* 71 (1978), 193–196. MR0551483 (58:27667)
- [11] M.B. Nathanson, *Elementary Methods in Number Theory*, Springer-Verlag, 2000. MR1732941 (2001j:11001)
- [12] G. Pall, The distribution of integers represented by binary quadratic forms, *Bull. Amer. Math. Soc.* 49 (1943), 447–449. MR0008084 (4:240g)
- [13] M. Purtill, André permutations, lexicographic shellability and the cd -index of a convex polytope, *Trans. Amer. Math. Soc.* 338 (1993), 77–104. MR1094560 (93j:52017)
- [14] D. Shanks, On the conjecture of Hardy and Littlewood concerning the number of primes of the form $n^2 + a$, *Math. Comp.* 14 (1960), 321–332. MR0120203 (22:10960)
- [15] D. Shanks and J.W. Wrench, The calculation of certain Dirichlet series, *Math. Comp.* 17 (1963), 136–154. MR0159796 (28:3012)
- [16] D. Shanks, Polylogarithms, Dirichlet series, and certain constants, *Math. Comp.* 18 (1964), 322–324. MR0175275 (30:5460)
- [17] D. Shanks, Generalized Euler and class numbers, *Math. Comp.* 21 (1967), 689–694; Corrigenda, *ibid.*, 22 (1968), 699. MR0223295 (36:6343)

- [18] D. Shanks, Calculation and application of Epstein zeta functions, Math. Comp. 29 (1975), 271–287. MR0409357 (53:13114a)
- [19] N.J.A. Sloane, The On-Line Encyclopedia of Integer Sequences, <http://www.research.att.com/~njas/sequences>.
- [20] T.A. Springer, Remarks on a combinatorial problem, Nieuw Arch. Wisk. 19 (1971), 30–36. MR0338199 (49:2965)
- [21] R.P. Stanley, Binomial posets, Möbius inversion, and permutation enumeration, J. Combin. Theory Ser. A 20 (1976), 336–356. MR0409206 (53:12968)

CENTER FOR COMBINATORICS, LPMC-TJKLC, NANKAI UNIVERSITY, TIANJIN 300071, PEOPLE'S REPUBLIC OF CHINA

E-mail address: `chen@nankai.edu.cn`

CENTER FOR COMBINATORICS, LPMC-TJKLC, NANKAI UNIVERSITY, TIANJIN 300071, PEOPLE'S REPUBLIC OF CHINA

E-mail address: `fjy@cfc.nankai.edu.cn`

CENTER FOR COMBINATORICS, LPMC-TJKLC, NANKAI UNIVERSITY, TIANJIN 300071, PEOPLE'S REPUBLIC OF CHINA

E-mail address: `jyt@cfc.nankai.edu.cn`