

ANALYSIS OF SCHWARZ METHODS FOR A HYBRIDIZABLE DISCONTINUOUS GALERKIN DISCRETIZATION: THE MANY-SUBDOMAIN CASE

MARTIN J. GANDER AND SOHEIL HAJIAN

ABSTRACT. Schwarz methods are attractive parallel solution techniques for solving large-scale linear systems obtained from discretizations of partial differential equations (PDEs). Due to the iterative nature of Schwarz methods, convergence rates are an important criterion to quantify their performance. Optimized Schwarz methods (OSM) form a class of Schwarz methods that are designed to achieve faster convergence rates by employing optimized transmission conditions between subdomains. It has been shown recently that for a two-subdomain case, OSM is a natural solver for hybridizable discontinuous Galerkin (HDG) discretizations of elliptic PDEs. In this paper, we generalize the preceding result to the many-subdomain case and obtain sharp convergence rates with respect to the mesh size and polynomial degree, the subdomain diameter, and the zeroth-order term of the underlying PDE, which allows us for the first time to give precise convergence estimates for OSM used to solve parabolic problems by implicit time stepping. We illustrate our theoretical results with numerical experiments.

1. INTRODUCTION

For the numerical treatment of a parabolic equation, e.g.,

$$(1.1) \quad \begin{aligned} \frac{\partial u}{\partial t} - \nabla \cdot (a(x)\nabla u) &= f(x, t) && \text{in } \Omega \times (0, T], \\ u(x, t) &= 0 && \text{on } \partial\Omega \times (0, T], \\ u(x, 0) &= g(x) && \text{on } \Omega, \end{aligned}$$

one often first discretizes the spatial dimension using a finite difference (FD), finite element (FE) or discontinuous Galerkin (DG) method. This approach, called *method of lines*, results in a *semi-discrete* system where the unknown $u(x, t)$ is approximated by a finite dimensional vector $\mathbf{u}_h(t)$ and the differential operator $-\nabla \cdot (a(x)\nabla)$ by a stiffness matrix which we denote by A_h . More precisely we then have

$$(1.2) \quad \frac{\partial \mathbf{u}_h(t)}{\partial t} + A_h \mathbf{u}_h(t) = \mathbf{f}(t),$$

and $\mathbf{u}_h(t = 0) = \mathbf{g}_h$. We then discretize in time using for example a backward Euler method with time step τ , i.e.,

$$(1.3) \quad \left(\frac{1}{\tau} M_h + A_h \right) \mathbf{u}_n = \frac{1}{\tau} M_h \mathbf{u}_{n-1} + \mathbf{f}(t_n),$$

Received by the editor March 15, 2016, and, in revised form, February 6, 2017.

2010 *Mathematics Subject Classification*. Primary 65N22, 65F10, 65F08, 65N55, 65H10.

Key words and phrases. Additive Schwarz, optimized Schwarz, discontinuous Galerkin methods, scalability, parabolic problems.

where M_h is called the mass matrix and \mathbf{u}_n is an approximation of $\mathbf{u}_h(t_n)$. Therefore, at each time-step, a linear system has to be solved.

One approach for solving (1.3) efficiently is to use a domain decomposition method where we decompose the original spatial domain Ω into overlapping or non-overlapping subdomains and then solve smaller linear systems in parallel. In this paper we choose the spatial discretization to be a DG method, more precisely a hybridizable interior penalty (IPH) method.

It has been shown that optimized Schwarz methods are attractive and natural solvers for hybridizable DG discretizations; see [11], [16]. This is due to the fact that hybridizable DG methods impose continuity across elements and subdomains using a Robin transmission condition; see [10]. Robin transmission conditions and a suitable choice of the Robin parameter are the core of OSM to achieve fast convergence [9]. Special care is needed when OSM is used as a solver for classical FEM when cross-points are present; see, e.g., [12–14, 20]. Those are points which are shared by more than two subdomains. This is not the case when we apply OSM to a hybridizable DG method, e.g., IPH, since subdomains only communicate if they have a non-zero measure interface with each other.

We generalize here our previous results for a two-subdomain configuration in [11] to the case of many subdomains, perform an analysis with respect to the polynomial degree of the IPH, and study for the first time the influence of the time-step τ on the performance and scalability of the OSM. However, we do not make an attempt to optimize the solver with respect to the jumps in the $a(x)$ coefficient and therefore we work, without loss of generality in this context, with

$$(\eta - \Delta)u = f \quad \text{in } \Omega,$$

where $\eta = \tau^{-1}$ is a constant. In Section 2 we recall the definition of IPH in a *hybridizable* formulation and introduce the domain decomposition settings. In Section 3 we introduce an OSM for IPH and analyze its convergence properties. The main contributions of the paper are Theorem 3.4, Corollary 3.5 and the refined analysis in Section 3.3. We validate our theoretical findings by performing numerical experiments in Section 4.

2. THE IPH METHOD

IPH was first introduced in [7] as a stabilized discontinuous finite element method and later was studied as a member of the class of hybridizable DG methods in [6]. It has been shown that it is equivalent to a method called Ultra Weak Variational Formulation (UWVF) for the Helmholtz equation; see [15]. IPH also fits into the framework developed in [2] for a unified analysis of DG methods.

In this section we recall the IPH method and its properties. We can define many DG methods by two equivalent formulations, namely the *primal* and *flux* formulation; see for instance [2]. However, there is also a third equivalent formulation for a class of hybridizable DG methods introduced in [6]. For the sake of simplicity we use only the hybridized formulation for IPH and refer the reader to [16] and [18] for its primal and flux formulations.

2.1. Notation. We now define the necessary operators and function spaces needed to analyze DG methods. We follow the notation in [2]. Let $\mathcal{T}_h = \{K\}$ be a shape-regular and quasi-uniform triangulation of the domain Ω . We denote the diameter of an element of the triangulation by $h_K := \max_{x,y \in K} |x - y|$ and define

$h := \max_{K \in \mathcal{T}_h} h_K$. If e is an edge of an element, we denote the length of that edge by h_e . The quasi-uniformity of the mesh implies $h \approx h_K \approx h_e$. Let us denote the set of interior edges shared by two elements in \mathcal{T}_h by \mathcal{E}^0 , i.e., $\mathcal{E}^0 := \{e = \partial K_1 \cap \partial K_2, \forall K_1, K_2 \in \mathcal{T}_h\}$. Similarly we define the set of boundary edges by \mathcal{E}^∂ and all edges by $\mathcal{E} := \mathcal{E}^\partial \cup \mathcal{E}^0$.

We seek a DG approximation which belongs to the finite dimensional space

$$(2.1) \quad V_h := \{v \in L^2(\Omega) : v|_K \in \mathbb{P}^k(K) \forall K \in \mathcal{T}_h\},$$

where $\mathbb{P}^k(K)$ is the space of polynomials of degree less than k in the simplex $K \in \mathcal{T}_h$. Note that a function in V_h is not necessarily continuous. More precisely V_h is a finite dimensional subspace of a broken Sobolev space $H^l(\mathcal{T}_h) := \prod_{K \in \mathcal{T}_h} H^l(K)$, where $H^l(K)$ is the usual Sobolev space in $K \in \mathcal{T}_h$ and l is a positive integer. Since $H^l(\mathcal{T}_h)$ contains discontinuous functions, its trace space along \mathcal{E}^0 can be double-valued. We define the trace space of functions in $H^l(\mathcal{T}_h)$ by $T(\mathcal{E}) := \prod_{K \in \mathcal{T}_h} L^2(\partial K)$. Observe that $q \in T(\mathcal{E})$ can be double-valued on \mathcal{E}^0 but it is single-valued on \mathcal{E}^∂ .

We now define two trace operators: let $q \in T(\mathcal{E})$ and $q_i := q|_{\partial K_i}$. Then on $e = \partial K_1 \cap \partial K_2$ we define the average and jump operators

$$\{q\} := \frac{1}{2}(q_1 + q_2), \quad [q] := q_1 \mathbf{n}_1 + q_2 \mathbf{n}_2,$$

where \mathbf{n}_i is the unit outward normal from K_i on $e \in \mathcal{E}^0$. Note that the jump and average definition is independent of the element enumeration. Similarly for a vector-valued function $\sigma \in [T(\mathcal{E})]^2$ we define on interior edges

$$\{\sigma\} := \frac{1}{2}(\sigma_1 + \sigma_2), \quad (\text{jump}) \quad [\sigma] := \sigma_1 \cdot \mathbf{n}_1 + \sigma_2 \cdot \mathbf{n}_2.$$

On the boundary, we set the average and jump operators to $\{\sigma\} := \sigma$ and $[q] = q \mathbf{n}$ and we do not need to define $\{q\}$ and $[\sigma]$ on $e \in \mathcal{E}^\partial$ since they do not appear in the discrete formulation.

Since $H^l(\mathcal{T}_h)$ contains discontinuous functions, we need to define some piecewise gradient operators. For all $u, v \in H^l(\mathcal{T}_h)$ we define

$$\int_{\mathcal{T}_h} \nabla u \cdot \nabla v := \sum_{K \in \mathcal{T}_h} \int_K \nabla u \cdot \nabla v.$$

For $a, b \in T(\mathcal{E})$ and single-valued on \mathcal{E}^0 we define the edge integrals by

$$\int_{\mathcal{E}} a b := \sum_{e \in \mathcal{E}} \int_e a b.$$

2.2. Domain decomposition setting. In order to define IPH in a hybridizable form we first decompose the domain into N_s non-overlapping subdomains $\{\Omega_i\}_{i=1}^{N_s}$. We denote the interface between subdomains by Γ and assume the interface is a subset of internal edges, \mathcal{E}^0 . More precisely, we denote the interface between two subdomains by $\Gamma_{ij} := \partial\Omega_i \cap \partial\Omega_j$ for $i \neq j$ and the global interface by $\Gamma := \bigcup_{i \neq j} \Gamma_{ij} \subset \mathcal{E}^0$. In other words the domain decomposition does not go through any element of the triangulation. For convenience we denote the interface belonging to subdomain Ω_i by $\Gamma_i := \bigcup_{j \in N(i)} \Gamma_{ij}$ where $N(i)$ is a set containing neighbors of the subdomain Ω_i ; for an example see Figure 1.




FIGURE 1. An unstructured mesh with the interface Γ (blue-dashed) and cross-points.

This domain decomposition induces a set of non-overlapping triangulations $\{\mathcal{T}_i\}_{i=1}^{N_s}$. Moreover, we can define local DG spaces on each subdomain and represent each function in V_h as a direct sum,

$$V_h = V_{h,1} \oplus V_{h,2} \oplus \cdots \oplus V_{h,N_s},$$

where $V_{h,i}$ for $i = 1, \dots, N_s$ is a local space defined as

$$V_{h,i} := \{v \in L^2(\Omega_i) : v|_{K \in \mathcal{T}_i} \in \mathbb{P}^k(K)\}.$$

We also need a finite dimensional space on the interface which we denote by Λ_h ,

$$\Lambda_h := \{\varphi \in L^2(\Gamma) : \varphi|_{e \in \Gamma} \in \mathbb{P}^k(e)\}.$$

For the analysis of our Schwarz methods we also need to define local spaces on Γ_i for all $i = 1, \dots, N_s$,

$$\Lambda_i := \{\varphi \in L^2(\Gamma_i) : \varphi|_{e \in \Gamma_i} \in \mathbb{P}^k(e)\},$$

and its global counterpart $\prod_{i=1}^{N_s} \Lambda_i$. Note that Λ_h is single-valued across Γ while $\prod_{i=1}^{N_s} \Lambda_i$ is double-valued. We denote the maximum diameter of the subdomains by H and the diameter of the mono-domain Ω by H_Ω . We assume $0 < h \leq H < H_\Omega$. For convenience we define a function for the set of neighboring subdomains of Ω_i and denote it by $N(i)$.

2.3. Hybridizable formulation. We now present IPH in a hybridizable form. A DG method is hybridizable if one can eliminate the degrees of freedom inside each element and obtain a linear system in terms of a *single-valued* function along edges. Not all DG methods can be written in a hybridized form, for instance the classical IP method is not hybridizable. A hybridization procedure for DG methods has been developed and studied in [6] where IPH is also included.

In a DG context the continuity of the exact solution is imposed weakly through a Nitsche penalization technique. Penalization is regulated through a parameter, $\mu \in \mathbb{T}(\mathcal{E})$ and scaled like $\mu = \alpha k^2/h$ for $\alpha > 0$, independent of h and k and sufficiently large. This choice of μ guarantees coercivity of the DG bilinear form and optimal approximation. Let $(u, \lambda), (v, \varphi) \in V_h \times \Lambda_h$ and $u_i, v_i \in V_{h,i}$ be the restriction of u and v to Ω_i . Then the IPH bilinear form reads

$$(2.2) \quad a((u, \lambda), (v, \varphi)) := a_\Gamma(\lambda, \varphi) + \sum_{i=1}^{N_s} \left(a_i(u_i, v_i) + a_{i\Gamma}(v_i, \lambda) + a_{i\Gamma}(u_i, \varphi) \right),$$

where

$$(2.3) \quad a_\Gamma(\lambda, \varphi) := \mu \sum_{i=1}^{N_s} \int_{\Gamma_i} \lambda \varphi, \quad a_{i\Gamma}(v_i, \varphi) := \int_{\Gamma_i} \left(\frac{\partial v_i}{\partial \mathbf{n}_i} - \mu v_i \right) \varphi,$$

and the local solvers $a_i(\cdot, \cdot)$ are defined as

$$(2.4) \quad \begin{aligned} a_i(u_i, v_i) := & \int_{\mathcal{T}_i} \eta u_i v_i + \nabla u_i \cdot \nabla v_i - \int_{\mathcal{E}_i^0} \{ \nabla u_i \} \cdot [v_i] + \{ \nabla v_i \} \cdot [u_i] \\ & + \int_{\mathcal{E}_i^0} \frac{\mu}{2} [u_i] \cdot [v_i] - \frac{1}{2\mu} [\nabla u_i] [\nabla v_i] + \int_{\partial\Omega_i} \mu u_i v_i - \frac{\partial u_i}{\partial \mathbf{n}_i} v_i - \frac{\partial v_i}{\partial \mathbf{n}_i} u_i. \end{aligned}$$

This is an IPH discretization of the model problem in Ω_i , and $\partial\Omega_i$ is treated as a Dirichlet boundary. Observe that $a_i(\cdot, \cdot)$ and $a_\Gamma(\cdot, \cdot)$ are symmetric and therefore $a(\cdot, \cdot)$ is symmetric too.

The global bilinear form $a(\cdot, \cdot)$ is coercive at the discrete level. In order to show coercivity we first introduce a semi-norm on each subdomain for all $(v_i, \varphi) \in V_{h,i} \times \Lambda_h$,

$$(2.5) \quad \|(v_i, \varphi)\|_i^2 := \eta \|v_i\|_{\mathcal{T}_i}^2 + \|\nabla v_i\|_{\mathcal{T}_i}^2 + \mu \|[v_i]\|_{\mathcal{E}_i \setminus \Gamma_i}^2 + \mu \|v_i - \varphi\|_{\Gamma_i}^2$$

for $i = 1, \dots, N_s$. Note that if a subdomain is *floating*, that is, it does not touch the Dirichlet boundary condition, and $\eta = 0$, then $\|(v_i, \varphi)\|_i = 0$ implies v_i and φ are constants and not necessarily zero. The energy norm over the whole domain is defined by

$$(2.6) \quad \|(v, \varphi)\|^2 := \sum_{i=1}^{N_s} \|(v_i, \varphi)\|_i^2.$$

In order to verify that this is actually a norm, we just need to check its kernel: if $\|(v, \varphi)\| = 0$, then v and φ are both constants and $v|_\Gamma = \varphi$. Moreover, there are subdomains that touch the Dirichlet boundary condition and therefore $v|_{\partial\Omega} = 0$. Hence $(v, \varphi) = 0$.

The proof of coercivity for IPH is done subdomain by subdomain: we first collect the contribution of each subdomain

$$(2.7) \quad \begin{aligned} a((v, \varphi), (v, \varphi)) &= a_\Gamma(\varphi, \varphi) + \sum_{i=1}^{N_s} (a_i(v_i, v_i) + 2a_{i\Gamma}(v_i, \varphi)) \\ &= \sum_{i=1}^{N_s} (a_i(v_i, v_i) + 2a_{i\Gamma}(v_i, \varphi) + \mu \|\varphi\|_{\Gamma_i}^2). \end{aligned}$$

Each right-hand side can be bounded from below by semi-norms $\|(v_i, \varphi)\|_i$ for $i = 1, \dots, N_s$; for details see [11], [18]. Then we obtain

$$a((v, \varphi), (v, \varphi)) \geq c \sum_{i=1}^{N_s} \|(v_i, \varphi)\|_i^2 = c \|(v, \varphi)\|^2 \quad \forall (v, \varphi) \in V_h \times \Lambda_h,$$

where $0 < c < 1$ and c does not depend on h and k .

An IPH approximation of the exact solution is obtained by solving the following problem: find $(u_h, \lambda_h) \in V_h \times \Lambda_h$ such that

$$(2.8) \quad a((u_h, \lambda_h), (v, \varphi)) = \int_{\Omega} f v \quad \forall (v, \varphi) \in V_h \times \Lambda_h,$$

which has a unique solution since $a(\cdot, \cdot)$ is coercive on $V_h \times \Lambda_h$. We can also show that IPH has optimal approximation properties, i.e., if the weak solution u is regular enough, then

$$\|u_h - u\|_0 \leq ch^{k+1}|u|_{k+1,\Omega};$$

see details in [2], [18].

We now describe how subdomains in the discrete problem (2.8) communicate. If we test (2.8) with $\varphi = 0$ and $v = 0$ in all subdomains except Ω_i we obtain

$$(2.9) \quad a_i(u_i, v_i) + a_{i\Gamma}(v_i, \lambda_h) = \int_{\Omega_i} f v_i \quad \forall v_i \in V_{h,i}, \text{ for } i = 1, \dots, N_s,$$

where $u_i := u_h|_{\Omega_i}$. This shows that u_i is determined if λ_h is known. More precisely λ_h is used as a Dirichlet boundary data on $\partial\Omega_i$ in a weak sense using a Nitsche penalization technique. Now if we test (2.8) with $v = 0$ and $\varphi \neq 0$, we obtain an equation for λ_h :

$$(2.10) \quad a_{\Gamma}(\lambda_h, \varphi) + \sum_{i=1}^{N_s} a_{i\Gamma}(u_i, \varphi) = 0 \quad \forall \varphi \in \Lambda_h.$$

If we further let φ be non-zero only on Γ_{ij} , a segment shared by Ω_i and Ω_j , then (2.10) reads

$$(2.11) \quad \lambda_h = \frac{1}{2\mu} \left(\mu u_i - \frac{\partial u_i}{\partial \mathbf{n}_i} \right) + \frac{1}{2\mu} \left(\mu u_j - \frac{\partial u_j}{\partial \mathbf{n}_j} \right) \quad \text{on } \Gamma_{ij}.$$

In the language of HDG methods, equation (2.11) is called the continuity condition. The continuity condition (2.11) is the core of the optimized Schwarz method that we will describe in Section 3. We have shown in [11] for the case of two subdomains how to exploit (2.11) to design a fast solver which we extend to many subdomains in this paper.

2.4. Schur complement and matrix formulation. The discrete problem (2.8) can be written in an equivalent matrix form. We first choose nodal basis functions for $\mathbb{P}^k(K)$ and denote the space of degrees of freedoms (DOFs) of V_h by V and similarly for subspaces, denoted by $\{V_i\}$. Then the discrete problem (2.8) is equivalent to

$$(2.12) \quad \underbrace{\begin{bmatrix} A_I & A_{I\Gamma} \\ A_{I\Gamma}^{\top} & A_{\Gamma} \end{bmatrix}}_{A:=} \begin{pmatrix} \mathbf{u} \\ \boldsymbol{\lambda} \end{pmatrix} = \begin{pmatrix} \mathbf{f} \\ 0 \end{pmatrix},$$

where \mathbf{u} and $\boldsymbol{\lambda}$ are DOFs corresponding to u_h and λ_h , respectively. Here A_I corresponds to the bilinear form $\sum_{i=1}^{N_s} a_i(\cdot, \cdot)$, $A_{I\Gamma}$ corresponds to $\sum_{i=1}^{N_s} a_{i\Gamma}(\cdot, \cdot)$ and A_{Γ} corresponds to $a_{\Gamma}(\cdot, \cdot)$.

Since the bilinear form (2.2) is symmetric and positive definite we can conclude that A is s.p.d. Hence its diagonal blocks, A_I and A_{Γ} , are also s.p.d. If we eliminate the interface unknown, $\boldsymbol{\lambda}$, we arrive at a linear system in terms of *primal* variables

\mathbf{u} only. This coincides with the *primal* formulation of IPH; see [11], [18] for details. On the other hand we can eliminate \mathbf{u} and obtain a Schur complement formulation

$$(2.13) \quad S_\Gamma \boldsymbol{\lambda} = \mathbf{g},$$

where

$$(2.14) \quad S_\Gamma := A_\Gamma - A_{I\Gamma}^\top A_I^{-1} A_{I\Gamma}, \quad \mathbf{g} := -A_{I\Gamma}^\top A_I^{-1} \mathbf{f}.$$

The Schur complement matrix has smaller dimension compared to (2.12) and is also s.p.d. Therefore one approach in solving (2.13) is to use the conjugate gradient (CG) method. However, the convergence of CG is affected by the condition number of S_Γ , which is similar to the condition number of classical FEM Schur complement systems.

Proposition 1. *Let S_Γ be the Schur complement of the IPH discretization. Then for all $\varphi \in \Lambda_h$ we have*

$$(2.15) \quad c \frac{H}{H_\Omega^2} \|\varphi\|_\Gamma^2 \leq \boldsymbol{\varphi}^\top S_\Gamma \boldsymbol{\varphi} \leq C \alpha \frac{k^2}{h} \|\varphi\|_\Gamma^2,$$

and therefore the condition number $\kappa(S_\Gamma)$ is bounded by

$$(2.16) \quad \kappa(S_\Gamma) \leq C \alpha \frac{H_\Omega^2 k^2}{Hh} \kappa(M_\Gamma),$$

where M_Γ is the mass matrix along the interface. Moreover, all constants are independent of α , k , h , and H .

Proof. See [17, Appendix 3]. □

3. OPTIMIZED SCHWARZ METHOD FOR IPH

In this section we define and analyze an optimized Schwarz method (OSM) for IPH discretizations. Since an IPH discretization is s.p.d. we can use an additive Schwarz preconditioner in conjunction with CG. However, it was first observed in [10] that the convergence mechanism of the additive Schwarz method for IPH is different from classical FEM. For a FEM discretization, the overlap between subdomains makes the additive Schwarz method converge, while for IPH convergence is due to a Robin transmission condition in a non-overlapping setting, and the Robin parameter is exactly the penalty parameter of IPH, $\mu = \alpha k^2/h$.

Robin transmission conditions are the core of OSM to obtain faster convergence compared to the additive Schwarz method. It was shown in [9] that OSM's best performance is achieved if the Robin parameter is scaled like $1/\sqrt{h}$. This, however, poses a contradiction with the IPH discretization penalty parameter since the scaling of μ cannot be weakened, otherwise coercivity and optimal approximation properties are lost. In [11], the authors modified and analyzed an OSM while not changing the scaling of μ . They showed that for the two-subdomain case the OSM's contraction factor is $\rho \leq 1 - O(\sqrt{h})$. This is a superior convergence factor compared to additive Schwarz with $\rho \leq 1 - O(h)$. If OSM is used as preconditioner for a Krylov subspace method, then a contraction factor of $\rho \leq 1 - O(h^{1/4})$ is observed.

We will now define a many-subdomain OSM for IPH with this property and then analyze the solver and optimize the performance with respect to the mesh parameter h and the polynomial degree k .




FIGURE 2. A many-subdomain configuration with unknown duplication along interfaces.

3.1. Definition of OSM. We now construct an OSM for the IPH discretization. Observe that from (2.9), we can conclude that $u_i \in V_{h,i}$ is determined provided λ_h is known. Recall also from (2.11) that two subdomains, say Ω_i and Ω_j for $i \neq j$, are communicating using

$$\lambda_h = \frac{1}{2\mu} \left(\mu u_i - \frac{\partial u_i}{\partial \mathbf{n}_i} \right) + \frac{1}{2\mu} \left(\mu u_j - \frac{\partial u_j}{\partial \mathbf{n}_j} \right) \quad \text{on } \Gamma_{ij}.$$

Let us now assume that λ_h is double-valued across the interface Γ . Then we can assign an interface unknown to each subdomain which we call λ_i for $i = 1, \dots, N_s$, as illustrated in Figure 2. Therefore on each interface between two subdomains, say Γ_{ij} , we should introduce two conditions. We do so by splitting the continuity condition in the following fashion:

$$(3.1) \quad \begin{aligned} \gamma \lambda_i + (1 - \gamma) \lambda_j &= \frac{1}{2\mu} \left(\mu u_i - \frac{\partial u_i}{\partial \mathbf{n}_i} \right) + \frac{1}{2\mu} \left(\mu u_j - \frac{\partial u_j}{\partial \mathbf{n}_j} \right), \\ (1 - \gamma) \lambda_i + \gamma \lambda_j &= \frac{1}{2\mu} \left(\mu u_i - \frac{\partial u_i}{\partial \mathbf{n}_i} \right) + \frac{1}{2\mu} \left(\mu u_j - \frac{\partial u_j}{\partial \mathbf{n}_j} \right), \end{aligned}$$

where $\gamma \in \mathbb{R}^+$ is a “suitable” parameter that we will use to optimize the iterative method. Observe that if we subtract the two conditions in (3.1), we arrive at $(1 - 2\gamma)(\lambda_i - \lambda_j) = 0$. If $\gamma \neq \frac{1}{2}$, then we have $\lambda_i = \lambda_j = \lambda_h$ on all Γ_{ij} , i.e., we recover the single-valued λ_h and therefore the solution to the *augmented* system coincides with the original IPH approximation.

The conditions in (3.1) can be written in an equivalent variational form by multiplying them with appropriate test functions with support on $\partial\Omega_i \setminus \partial\Omega$. The advantage is that we can then use the original blocks of the IPH linear system. For a subdomain, e.g., Ω_i , we obtain

$$(3.2) \quad \gamma a_{\Gamma}^{(i)}(\lambda_i, \varphi_i) + a_{i\Gamma}(u_i, \varphi_i) + \sum_{j \in N(i)} 2\mu(1 - \gamma) \int_{\Gamma_{ij}} \lambda_j \varphi_i + \int_{\Gamma_{ij}} \left(\frac{\partial u_j}{\partial \mathbf{n}_j} - \mu u_j \right) \varphi_i = 0.$$

In order to clarify the definition of the new linear system we provide an example in the case of two subdomains.

Example 3.1 (Two-subdomain case). Suppose we have two subdomains and we call the interface between them Γ . Then an IPH discretization with this configuration looks like

$$(3.3) \quad \begin{bmatrix} A_1 & & A_{1\Gamma} \\ & A_2 & A_{2\Gamma} \\ A_{1\Gamma}^\top & A_{2\Gamma}^\top & A_\Gamma \end{bmatrix} \begin{pmatrix} \mathbf{u}_1 \\ \mathbf{u}_2 \\ \boldsymbol{\lambda} \end{pmatrix} = \begin{pmatrix} \mathbf{f}_1 \\ \mathbf{f}_2 \\ 0 \end{pmatrix}.$$

Observe that continuity between the two subdomains is imposed through the last row of (3.3). We now suppose $\boldsymbol{\lambda}$ is double-valued across the interface, $\boldsymbol{\lambda}_1, \boldsymbol{\lambda}_2$. Then we introduce two conditions for the two interface unknowns, $\boldsymbol{\lambda}_1$ and $\boldsymbol{\lambda}_2$,

$$\begin{aligned} \gamma A_\Gamma \boldsymbol{\lambda}_1 + (1 - \gamma) A_\Gamma \boldsymbol{\lambda}_2 + A_{1\Gamma}^\top \mathbf{u}_1 + A_{2\Gamma}^\top \mathbf{u}_2 &= 0, \\ (1 - \gamma) A_\Gamma \boldsymbol{\lambda}_1 + \gamma A_\Gamma \boldsymbol{\lambda}_2 + A_{1\Gamma}^\top \mathbf{u}_1 + A_{2\Gamma}^\top \mathbf{u}_2 &= 0, \end{aligned}$$

where $\gamma \neq \frac{1}{2}$. If we regroup the unknowns according to the subdomain enumeration, then the “augmented” linear system looks like:

$$(3.4) \quad \left[\begin{array}{cc|cc} A_1 & A_{1\Gamma} & A_{2\Gamma}^\top & (1 - \gamma) A_\Gamma \\ \hline A_{1\Gamma}^\top & \gamma A_\Gamma & A_2 & A_{2\Gamma} \\ \hline A_{1\Gamma}^\top & (1 - \gamma) A_\Gamma & A_{2\Gamma}^\top & \gamma A_\Gamma \end{array} \right] \begin{pmatrix} \mathbf{v}_1 \\ \boldsymbol{\lambda}_1 \\ \mathbf{v}_2 \\ \boldsymbol{\lambda}_2 \end{pmatrix} = \begin{pmatrix} \mathbf{f}_1 \\ 0 \\ \mathbf{f}_2 \\ 0 \end{pmatrix}.$$

Note that, provided $\gamma \neq \frac{1}{2}$, the linear systems (3.3) and (3.4) are equivalent in the sense that $\mathbf{v}_1 = \mathbf{u}_1$ and $\mathbf{u}_2 = \mathbf{v}_2$ and $\boldsymbol{\lambda}_1 = \boldsymbol{\lambda}_2 = \boldsymbol{\lambda}$.

The authors showed in [11] that for the convergence analysis of a block Jacobi method applied to (3.4) we need to obtain sharp bounds on the eigenvalues of $A_\Gamma^{-1} B_i$ where

$$(3.5) \quad B_i := A_{i\Gamma}^\top A_i^{-1} A_{i\Gamma}, \quad i = 1, 2.$$

Such bounds were obtained in [11, Lemma 3.7]. We will further improve the eigenvalue bounds and also obtain sharp estimates with respect to the time-step τ , $\eta = \tau^{-1}$.

We are now in the position to define the OSM for an IPH discretization. Formally we first construct the augmented system with double-valued interface unknowns along the interfaces, i.e., (3.4). Then we rearrange the unknowns subdomain by subdomain, i.e., collect $\{(\mathbf{u}_i, \boldsymbol{\lambda}_i)\}_{i=1}^{N_s}$ and finally we perform a block Jacobi method on the augmented linear system with a suitable optimization parameter γ .

Algorithm 3.2. Let $\{(u_i^{(0)}, \lambda_i^{(0)})\}_{i=1}^{N_s}$ be a set of initial guesses for all subdomains. Then for $n = 1, 2, \dots$ find $\{(u_i^{(n)}, \lambda_i^{(n)})\}_{i=1}^{N_s}$ such that

$$(3.6) \quad a_i(u_i^{(n)}, v_i) + a_{i\Gamma}(v_i, \lambda_i^{(n)}) = \int_{\Omega_i} f v_i \quad \forall v_i \in V_{h,i},$$

and the continuity condition on Γ_{ij} reads

$$(3.7) \quad \gamma \lambda_i^{(n)} - \frac{1}{2\mu} \left(\mu u_i - \frac{\partial u_i}{\partial \mathbf{n}_i} \right)^{(n)} = -(1 - \gamma) \lambda_j^{(n-1)} + \frac{1}{2\mu} \left(\mu u_j - \frac{\partial u_j}{\partial \mathbf{n}_j} \right)^{(n-1)}.$$

Since the solution of the augmented system coincides with the original IPH linear system, we can conclude that Algorithm 3.2 has the same fixed point as the solution of the IPH discretization.

We call $u_i^{(n)}$ satisfying (3.6) with $f = 0$ a *discrete harmonic extension* of λ_i in Ω_i . This definition helps us in analyzing OSM.

Definition 3.3 (Discrete harmonic extension). For all $\varphi_i \in \Lambda_i$, we denote by $\mathcal{H}_i(\varphi_i) \in V_{h,i}$ the discrete harmonic extension into Ω_i ,

$$(3.8) \quad \mathcal{H}_i(\varphi) \equiv -A_i^{-1}A_{i\Gamma}\varphi_i,$$

where A_i and $A_{i\Gamma}$ correspond to the bilinear forms $a_i(\cdot, \cdot)$ and $a_{i\Gamma}(\cdot, \cdot)$. The corresponding φ_i is called generator. In other words $u_i := \mathcal{H}_i(\varphi_i)$ is an approximation obtained from the IPH discretization in Ω_i using φ_i as Dirichlet data, i.e., $A_i u_i + A_{i\Gamma} \varphi_i = 0$.

There are some questions to be addressed concerning Algorithm 3.2, e.g.,

- (1) Is Algorithm 3.2 well-posed?
- (2) Does Algorithm 3.2 converge? If yes, then can we obtain a contraction factor?
- (3) How do we use the optimization parameter γ to improve the contraction factor?
- (4) How do different choices of η affect the algorithm and its scalability?

We will answer these questions now in Section 3.2.

3.2. Analysis of OSM. The main goal of this section is to analyze Algorithm 3.2 and answer the questions regarding its well-posedness and convergence. Our analysis is inspired by a similar result for FEM in [21], [22], [23], and we refer the reader to the original work of Lions in [19] for an analysis at the continuous level. Our analysis is, however, substantially different since DG methods impose continuity across elements weakly. We will first prove the following.

Theorem 3.4 (Convergence estimate). *Let the optimization parameter satisfy $\frac{1}{2} < \gamma \leq 1$. Then Algorithm 3.2 is well-posed and converges. More precisely the following contraction estimate holds:*

$$\|\mathcal{R}(\varphi^{(n)})\|^2 \leq \rho \|\mathcal{R}(\varphi^{(n-1)})\|^2,$$

where $\|\mathcal{R}(\varphi)\|^2 := \sum_{i=1}^{N_s} \|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2$ and $\mathcal{R}_i(\varphi_i) := \gamma\varphi_i - \frac{1}{2\mu} \left(\mu - \frac{\partial}{\partial n_i} \right) \mathcal{H}_i(\varphi_i)$. Here the contraction factor, i.e., ρ , is

$$(3.9) \quad \rho = 1 - \min \frac{(2\gamma - 1)}{\mu(2\gamma - 1)^2 C(H, \eta) + 1},$$

where $\mu = \alpha k^2/h$ is the penalization parameter and

$$C(H, \eta) := \begin{cases} H & \text{in the case of no floating subdomains,} \\ \frac{1}{H\eta} & \text{in the case of floating subdomains.} \end{cases}$$

The choice $\gamma = 1$ is a special case. It is shown in [11], [16] that in this case Algorithm 3.2 is equivalent to a non-overlapping additive Schwarz method¹ applied to the primal formulation of IPH. The theory for s.p.d. preconditioners, i.e., the abstract Schwarz framework, shows that the condition number of the one-level additive Schwarz method for IPH is bounded by $k^2 h^{-1} H^{-1}$. This is equivalent to a contraction factor $\rho \leq 1 - O(\frac{hH}{k^2})$. More precisely, suppose A is the original system matrix in primal form and A_{add} is the corresponding additive Schwarz preconditioner (see for instance [24, Section 1.5]), then the block Jacobi method converges with the aforementioned contraction factor in the A -norm. It is easy to see that

¹Non-overlapping additive Schwarz method for DG methods means non-overlapping both at the algebraic level as well as continuous level in contrast to FEM.

our analysis also reveals the same contraction factor (in the $\|\mathcal{R}(\cdot)\|$ norm) in this special case: let $\gamma = 1$ in (3.9) and recall that $\mu = \alpha k^2/h$. Then, we have

$$(3.10) \quad \rho \leq 1 - O\left(\frac{hH}{k^2}\right),$$

as h and H go to zero or k goes to infinity.

Our second objective of this section is to minimize the contraction factor through a suitable choice of the optimization parameter γ . This is stated in the following.

Corollary 3.5 (Optimized contraction factor). *Let $\eta = \tau^{-1}$ where τ is the time-step which is chosen to be $O(1)$ or $O(H)$ or $O(H^2)$. Then the optimized contraction factor for Algorithm 3.2 for the different choices of τ is*

$$(3.11) \quad \rho_{opt} \leq \begin{cases} 1 - O\left(\frac{\sqrt{hH}}{k}\right) & \text{for } \tau = O(1), & \text{if } \gamma_{opt} = \frac{1}{2}\left(1 + \frac{\sqrt{hH}}{k}\right), \\ 1 - O\left(\frac{\sqrt{h}}{k}\right) & \text{for } \tau = O(H), & \text{if } \gamma_{opt} = \frac{1}{2}\left(1 + \frac{\sqrt{h}}{k}\right), \\ 1 - O\left(\sqrt{\frac{h}{H}}\frac{1}{k}\right) & \text{for } \tau = O(H^2), & \text{if } \gamma_{opt} = \frac{1}{2}\left(1 + \sqrt{\frac{h}{H}}\frac{1}{k}\right). \end{cases}$$

Observe that the h -dependency and k -dependency is weakened by a square root compared to (3.10). Moreover, if the time-step is chosen to scale like a forward Euler time-step, i.e., $O(H^2)$, then Algorithm 3.2 is scalable.

Proof of Theorem 3.4. We first show that Algorithm 3.2 is well-posed, i.e., we can actually iterate. By linearity we assume that $f = 0$. We proceed by eliminating $u_i^{(n)}$ for all subdomains and simplify Algorithm 3.2 to: for all subdomains, find $\lambda_i^{(n)}$ such that

$$(3.12) \quad \gamma \lambda_i^{(n)} - \frac{1}{2\mu} \left(\mu - \frac{\partial}{\partial \mathbf{n}_i} \right) \mathcal{H}_i(\lambda_i^{(n)}) = -(1-\gamma) \lambda_j^{(n-1)} + \frac{1}{2\mu} \left(\mu - \frac{\partial}{\partial \mathbf{n}_j} \right) \mathcal{H}_j(\lambda_j^{(n-1)}),$$

on Γ_{ij} for all $j \in N(i)$, where $N(i)$ is the set of neighboring subdomains of Ω_i . Let us denote the linear operator on the left-hand side by $\mathcal{R}_i : \Lambda_i \rightarrow \Lambda_i$, that is,

$$(3.13) \quad \mathcal{R}_i(\varphi_i) := \gamma \varphi_i - \frac{1}{2\mu} \left(\mu - \frac{\partial}{\partial \mathbf{n}_i} \right) \mathcal{H}_i(\varphi_i).$$

If we show that $\mathcal{R}_i(\cdot)$ is an invertible operator, then Algorithm 3.2 is well-posed. We show $\mathcal{R}_i(\cdot)$ is invertible by showing that it is injective.

Lemma 3.6. *If $\gamma > \frac{1}{2}$, then the operator $\mathcal{R}_i(\cdot)$ is injective for all $i = 1, \dots, N_s$. More precisely we have the estimate*

$$(3.14) \quad \|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i} \geq \left(\gamma - \frac{1}{2} + c(h, H, k) \right) \|\varphi_i\|_{\Gamma_i} \quad \forall \varphi_i \in \Lambda_i,$$

where

$$c(h, H, k) := \begin{cases} c \frac{h}{H} \frac{1}{k^2} & \text{for non-floating subdomains,} \\ 0 & \text{for floating subdomains.} \end{cases}$$

Proof. We multiply $\mathcal{R}_i(\varphi_i)$ by φ_i and integrate over Γ_i to get

$$\int_{\Gamma_i} \mathcal{R}_i(\varphi_i) \varphi_i = \gamma \|\varphi_i\|_{\Gamma_i}^2 + \frac{1}{2\mu} a_{i\Gamma}(u_i, \varphi_i),$$

where $u_i := \mathcal{H}_i(\varphi_i)$. Recall that if u_i is the harmonic extension of φ_i , then $a_i(u_i, u_i) + a_{i\Gamma}(u_i, \varphi_i) = 0$. Therefore we have $\int_{\Gamma_i} \mathcal{R}_i(\varphi_i) \varphi_i = \gamma \|\varphi_i\|_{\Gamma_i}^2 - \frac{1}{2\mu} a_i(u_i, u_i)$.

We can show that $a(u_i, u_i) \leq (1 - c(h, H, k))\mu\|\varphi_i\|_{\Gamma_i}^2$ (see Appendix A), and obtain

$$\int_{\Gamma_i} \mathcal{R}_i(\varphi_i) \varphi_i \geq \left(\gamma - \frac{1}{2} + c(h, H, k)\right)\|\varphi_i\|_{\Gamma_i}^2.$$

If $\gamma > \frac{1}{2}$, then the right-hand side is positive. Now we apply the Cauchy-Schwarz inequality to the left-hand side and obtain $\|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i} \geq \left(\gamma - \frac{1}{2} + c(h, H, k)\right)\|\varphi_i\|_{\Gamma_i}$ which completes the proof. \square

Note that Lemma 3.6 provides a lower bound for the norm-equivalence between $\|\mathcal{R}_i(\cdot)\|_{\Gamma_i}$ and the L^2 -norm, i.e., $\|\cdot\|_{\Gamma_i}$. The upper bound in the norm-equivalence can also be obtained, as we show in the following proposition.

Proposition 2 (Norm-equivalence). *The two norms $\|\mathcal{R}_i(\cdot)\|_{\Gamma_i}$ and $\|\cdot\|_{\Gamma_i}$ are equivalent, i.e.,*

$$C\|\varphi_i\|_{\Gamma_i} \geq \|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i} \geq \left(\gamma - \frac{1}{2} + c(h, H, k)\right)\|\varphi_i\|_{\Gamma_i} \quad \forall \varphi_i \in \Lambda_i,$$

where $C > 0$ is independent of h, H, α , and η . Here $c(h, H, k)$ is the constant defined in Lemma 3.6.

Proof. The lower bound estimate is from Lemma 3.6. For the upper bound we use the estimate from Lemma 3.10 (which will appear in Section 3.3). More precisely we have

$$\varphi_i^\top B_i \varphi_i \geq c\mu\|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2,$$

where $B_i := A_{i\Gamma}^\top A_i^{-1} A_{i\Gamma}$ (see Example 3.1). We then use the estimate for the eigenvalues of B_i , i.e., [11, Lemma 3.7] to obtain

$$\mu\|\varphi_i\|_{\Gamma_i}^2 \geq \varphi_i^\top B_i \varphi_i \geq c\mu\|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2.$$

This completes the proof. \square

Since $\mathcal{R}_i(\cdot)$ is linear and injective we conclude that it induces a local norm on Λ_i . We can also define a global norm on the space of $\prod_{i=1}^{N_s} \Lambda_i$ by

$$(3.15) \quad \|\mathcal{R}(\varphi)\|^2 := \sum_{i=1}^{N_s} \|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2 \quad \forall \varphi \in \prod_{i=1}^{N_s} \Lambda_i,$$

where $\varphi := (\varphi_1, \varphi_2, \dots, \varphi_{N_s})$. This turns out to be the right norm for the convergence analysis of Algorithm 3.2.

We can now show that Algorithm 3.2 converges with a concrete contraction factor estimate. The right-hand side of the iteration equation (3.12) can be simplified to

$$(3.16) \quad \mathcal{R}_i(\varphi_i^{(n)}) = (2\gamma - 1)\varphi_j^{(n-1)} - \mathcal{R}_j(\varphi_j^{(n-1)})$$

on Γ_{ij} for all $j \in N(i)$. Note that $2\gamma - 1$ is strictly-positive with our condition $\gamma > \frac{1}{2}$. For a given subdomain, say Ω_i , we take the L^2 -norm on both sides. To simplify the presentation, we suppress the iteration index for the moment, but terms on the left-hand side are evaluated at iteration (n) while on the right-hand side they are

evaluated at iteration index $(n - 1)$:

$$\begin{aligned}
\|\mathcal{R}_i(\varphi_i)\|_{\Gamma_{ij}}^2 &= \|\mathcal{R}_j(\varphi_j) - (2\gamma - 1)\varphi_j\|_{\Gamma_{ij}}^2 \\
&= \|\mathcal{R}_j(\varphi_j)\|_{\Gamma_{ij}}^2 + \|(2\gamma - 1)\varphi_j\|_{\Gamma_{ij}}^2 - 2(2\gamma - 1) \int_{\Gamma_{ij}} \mathcal{R}_j(\varphi_j) \varphi_j \\
&= \|\mathcal{R}_j(\varphi_j)\|_{\Gamma_{ij}}^2 + [(2\gamma - 1)^2 - 2(2\gamma - 1)\gamma] \|\varphi_j\|_{\Gamma_{ij}}^2 \\
&\quad + \frac{1}{\mu}(2\gamma - 1) \int_{\Gamma_{ij}} \left(\mu u_j - \frac{\partial u_j}{\partial \mathbf{n}_j}\right) \varphi_j \\
&= \|\mathcal{R}_j(\varphi_j)\|_{\Gamma_{ij}}^2 - (2\gamma - 1) \left[\|\varphi_j\|_{\Gamma_{ij}}^2 - \frac{1}{\mu} \int_{\Gamma_{ij}} \left(\mu u_j - \frac{\partial u_j}{\partial \mathbf{n}_j}\right) \varphi_j \right].
\end{aligned}$$

Then we sum over all interfaces of Ω_i and all subdomains to obtain

$$\begin{aligned}
\|\mathcal{R}(\varphi)\|^2 &= \sum_{i=1}^{N_s} \sum_{j \in N(i)} \|\mathcal{R}_j(\varphi_j)\|_{\Gamma_{ij}}^2 \\
&\quad - (2\gamma - 1)\mu^{-1} \left[\mu \|\varphi_j\|_{\Gamma_{ij}}^2 - \int_{\Gamma_{ij}} \left(\mu u_j - \frac{\partial u_j}{\partial \mathbf{n}_j}\right) \varphi_j \right] \\
&= \|\mathcal{R}(\varphi)\|^2 \\
(3.17) \quad &\quad - (2\gamma - 1)\mu^{-1} \sum_{m=1}^{N_s} \left[\mu \|\varphi_m\|_{\Gamma_m}^2 + a_m \Gamma(u_m, \varphi_m) \right] \\
&= \|\mathcal{R}(\varphi)\|^2 \\
&\quad - (2\gamma - 1)\mu^{-1} \sum_{m=1}^{N_s} \left[\mu \|\varphi_m\|_{\Gamma_m}^2 - a_m(u_m, u_m) \right] \\
&\leq \|\mathcal{R}(\varphi)\|^2 - c(2\gamma - 1)\mu^{-1} \sum_{m=1}^{N_s} \|(u_m, \varphi_m)\|_m^2,
\end{aligned}$$

where for the left-hand side we used

$$\sum_{i=1}^{N_s} \sum_{j \in N(i)} \|\mathcal{R}_i(\varphi_i)\|_{\Gamma_{ij}}^2 = \sum_{i=1}^{N_s} \|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2 =: \|\mathcal{R}(\varphi)\|^2$$

and for the right-hand side we used the coercivity inequality

$$\mu \|\varphi_m\|_{\Gamma_m}^2 - a_m(u_m, u_m) \geq c \|(u_m, \varphi_m)\|_m^2;$$

see Appendix A for details. Note that $\|(u_m, \varphi_m)\|_m$ is *subdomain-wise* positive definite if $\eta > 0$. More precisely we can show that if $\eta > 0$ then for all subdomains, even *floating* ones, we have the estimate

$$(3.18) \quad \|\mathcal{R}_m(\varphi_m)\|_{\Gamma_m}^2 \leq \left((2\gamma - 1)^2 C(H, \eta) + \mu^{-1} \right) \|(u_m, \varphi_m)\|_m^2,$$

where

$$(3.19) \quad C(H, \eta) := \begin{cases} H & \text{for non-floating subdomain,} \\ \frac{1}{H\eta} & \text{for floating subdomain.} \end{cases}$$

Note that (3.18) makes sense only if $\eta > 0$ since $\|(\cdot, \cdot)\|_m$ is only a semi-norm for floating subdomains if $\eta = 0$ while $\|\mathcal{R}_m(\cdot)\|_{\Gamma_m}$ is a norm (see Appendix A), in particular (A.9) and (A.7). We have ignored the $\eta \|u_i\|_{\Omega_i}$ term in (A.7) for simplicity of the exposition; the $\eta \|u_i\|_{\Omega_i}$ term in (A.7) will be exploited in Section 3.3.

We then insert the norm estimate (3.18) into the last inequality of (3.17) and reintroduce the iteration index to obtain

$$(3.20) \quad \|\mathcal{R}(\varphi^{(n)})\|^2 \leq \left(1 - \min \left\{ \frac{2\gamma - 1}{\mu(2\gamma - 1)^2 H + 1}, \frac{2\gamma - 1}{\mu(2\gamma - 1)^2 (H\eta)^{-1} + 1} \right\}\right) \|\mathcal{R}(\varphi^{(n-1)})\|^2,$$

which shows convergence and proves Theorem 3.4. □

Proof of Corollary 3.5. We need to choose a suitable $\gamma > \frac{1}{2}$ to achieve the best possible contraction factor. In order to weaken dependencies on the mesh parameter, subdomain diameter, and polynomial degree, we make for the optimization parameter the ansatz

$$(3.21) \quad \gamma = \frac{1}{2} \left(1 + \frac{h^\xi H^\zeta}{k^\psi}\right),$$

with $\xi, \zeta, \psi \in \mathbb{R}$ to be chosen. We would like to minimize the contraction factor, i.e.,

$$(3.22) \quad \rho_{\text{opt}} \leq 1 - \max_{\xi, \zeta, \psi} \min \left\{ \frac{h^\xi H^\zeta}{k^{2-\psi} h^{2\xi-1} H^{2\zeta+1} + k^\psi}, \frac{h^\xi H^\zeta}{k^{2-\psi} h^{2\xi-1} H^{2\zeta-1} \eta^{-1} + k^\psi} \right\}.$$

Remark 3.7 (On the choice of γ). It has been shown in [16] and [17, Section 3.2] that the transmission condition between two subdomains in Algorithm 3.2 is equivalent at the continuous level to

$$\left((2\gamma - 1)\mu u_1 + \frac{\partial u_1}{\partial \mathbf{n}_1} \right)^{(n)} = \left((2\gamma - 1)\mu u_2 + \frac{\partial u_2}{\partial \mathbf{n}_1} \right)^{(n-1)}.$$

It has been shown (at the continuous level [9]) that the optimal choice of the Robin parameter is $(2\gamma - 1)\mu = O(h^{-1/2})$. This translates to choosing $\gamma = \frac{1}{2}(1 + \sqrt{h})$. We will show that this is also the optimal scaling at the discrete level. In [9], it has been shown that the optimal scaling of the Robin parameter is $O((hL)^{-1/2})$ where L is the length of the interface and it can be viewed as a measure of the diameter of a subdomain, i.e., H . This motivates our choice of optimization parameter, i.e., γ .

When dealing with parabolic problems, $\eta = \tau^{-1}$ and τ is the time-step. Therefore it is reasonable to optimize γ for different choices of the time-step.

- $\tau = O(1)$: we start with the dependence on the polynomial degree. Observe that the weakest dependence is achieved if we let $\psi = 1$. This leads to $\rho \leq 1 - O(\frac{1}{k})$, which compares very favorably to (3.10). Now we consider the case where H is fixed and we refine the mesh, $h \rightarrow 0$. Then $\xi = \frac{1}{2}$ is the optimal choice which yields $\rho \leq 1 - O(\frac{\sqrt{h}}{k})$. This leads to a simplified bound for ρ_{opt} , namely

$$\rho_{\text{opt}} \leq 1 - \max_{\zeta} \min \left\{ \frac{H^\zeta}{H^{2\zeta+1} + 1}, \frac{H^\zeta}{H^{2\zeta-1} + 1} \right\} O\left(\frac{\sqrt{h}}{k}\right).$$

The optimal value for ζ is therefore $\frac{1}{2}$. We thus obtain the optimal parameter and corresponding contraction factor

$$(3.23) \quad \gamma_{\text{opt}} := \frac{1}{2} \left(1 + \frac{\sqrt{hH}}{k}\right), \quad \rho_{\text{opt}} \leq 1 - O\left(\frac{\sqrt{hH}}{k}\right), \quad \text{if } \tau = O(1).$$

- $\tau = O(H)$: The best parameters with respect to k and h follow the same argument as before. For optimization with respect to H we now have

$$\rho_{\text{opt}} \leq 1 - \max_{\zeta} \min \left\{ \frac{H^\zeta}{H^{2\zeta+1} + 1}, \frac{H^\zeta}{H^{2\zeta} + 1} \right\} O\left(\frac{\sqrt{h}}{k}\right).$$

In this case we can eliminate the H -dependence by choosing $\zeta = 0$. Hence we have

$$(3.24) \quad \gamma_{\text{opt}} := \frac{1}{2} \left(1 + \frac{\sqrt{h}}{k} \right), \quad \rho_{\text{opt}} \leq 1 - O\left(\frac{\sqrt{h}}{k}\right), \quad \text{if } \tau = O(H).$$

- $\tau = O(H^2)$: This case is comparable to using a forward Euler method where τ is required to be proportional to h^2 . This is a typical constraint when dealing with parabolic problems and accurate trajectories in time are needed, but one could still take larger time steps in our setting than with forward Euler due to a larger constant. We proceed as before by choosing the same parameters with respect to k and h . For the H -dependence we have

$$\rho_{\text{opt}} \leq 1 - \max_{\zeta} \min \left\{ \frac{H^\zeta}{H^{2\zeta+1} + 1}, \frac{H^\zeta}{H^{2\zeta} + 1} \right\} O\left(\frac{\sqrt{h}}{k}\right).$$

The optimal parameter hence is $\zeta = -\frac{1}{2}$ which yields

$$(3.25) \quad \gamma_{\text{opt}} := \frac{1}{2} \left(1 + \sqrt{\frac{h}{H} \frac{1}{k}} \right), \quad \rho_{\text{opt}} \leq 1 - O\left(\sqrt{\frac{h}{H} \frac{1}{k}}\right), \quad \text{if } \tau = O(H^2).$$

Note that this choice of γ_{opt} is still feasible since $h \leq H$ and therefore $\gamma_{\text{opt}} \leq 1$. This shows that the method is weakly scalable if we choose a small enough time-step, without the need of a coarse solver. A similar result for the additive Schwarz method and FEM exists; see [4, Theorem 4].

This completes the proof of Corollary 3.5. □

3.3. A refined contraction factor with respect to the time-step. In this section we would like to investigate the effect of the time-step, $\tau = \eta^{-1}$, on the contraction factor while the number of subdomains is *fixed*, e.g., in the case of two subdomains. This has so far not been addressed, neither in [21] nor in the authors' paper [11] which deals with two subdomains only.

Suppose for the moment that we have two subdomains. Then as mentioned in Example 3.1 and proved in [11] the convergence of the OSM is governed by the eigenvalues of $A_\Gamma^{-1} B_i$ where $B_i := A_{i\Gamma}^\top A_i^{-1} A_{i\Gamma}$. We would like to obtain eigenvalue estimates that depend on η . This is stated in the following lemma which improves the estimate in [11, Lemma 3.7].

Lemma 3.8. *Let $B_i := A_{i\Gamma}^\top A_i^{-1} A_{i\Gamma}$ for $i = 1, 2$, where A_i and $A_{i\Gamma}$ correspond to the bilinear forms defined in (2.4) and (2.3), respectively. Then for $\eta \geq 0$ we have the estimate*

$$\varphi^\top B_i \varphi \leq \left(\frac{1}{1 + \frac{C\eta h^2}{1 + C\eta h^2}} \right) \left(1 - c \frac{h}{H\alpha} \right) \mu \|\varphi\|_\Gamma^2,$$

where c and C are positive constants which are independent of h, H, α , and η .

Proof. Recall the definition of A_i from (2.4), and let us decompose A_i into the mass matrix M_i and the stiffness matrix K_i ,

$$A_i := \eta M_i + K_i,$$

where $\mathbf{v}_i^\top M_i \mathbf{u}_i := \int_{\Omega_i} u_i v_i$ and K_i is defined as $\mathbf{v}_i^\top K_i \mathbf{u}_i := a_i(u_i, v_i) - \eta \mathbf{v}_i^\top M_i \mathbf{u}_i$. Now consider

$$\hat{A} := \begin{bmatrix} K_i + \eta M_i & A_{i\Gamma} \\ A_{i\Gamma}^\top & \frac{1}{2} A_\Gamma \end{bmatrix},$$

which is coercive, i.e., for all $\mathbf{w} := (\mathbf{u}_i, \boldsymbol{\varphi})$ we have (see [11, equation 3.6]),

$$(3.26) \quad \mathbf{w}^\top \hat{A} \mathbf{w} \geq c \|(u_i, \varphi)\|_i^2 \geq \eta \mathbf{u}_i^\top M_i \mathbf{u}_i + \frac{c}{H} \|\varphi\|_\Gamma^2,$$

where the last inequality is Lemma A.1. On the other hand we can easily verify that for $u_i := \mathcal{H}_i(\varphi)$ we have

$$(3.27) \quad \boldsymbol{\varphi}^\top B_i \boldsymbol{\varphi} = \mathbf{u}_i^\top (K_i + \eta M_i) \mathbf{u}_i \leq (Ch^{-2} + \eta) \mathbf{u}_i^\top M_i \mathbf{u}_i,$$

where we have used the fact that $\sigma(M_i^{-1} K_i) \in [c_1, c_2 h^{-2}]$, which is usual for elliptic operators; see for instance [5, Theorem 3.4]. For $\mathbf{w} := (\mathcal{H}_i(\varphi), \varphi)$, observing that

$$\frac{1}{2} \boldsymbol{\varphi}^\top A_\Gamma \boldsymbol{\varphi} - \boldsymbol{\varphi}^\top B_i \boldsymbol{\varphi} = \mathbf{w}^\top \hat{A} \mathbf{w},$$

and using (3.26) we have

$$\frac{1}{2} \boldsymbol{\varphi}^\top A_\Gamma \boldsymbol{\varphi} - \boldsymbol{\varphi}^\top B_i \boldsymbol{\varphi} \geq \frac{\eta}{Ch^{-2} + \eta} \boldsymbol{\varphi}^\top B_i \boldsymbol{\varphi} + \frac{c}{H} \|\varphi\|_\Gamma^2.$$

Recalling that $\frac{1}{2} \boldsymbol{\varphi}^\top A_\Gamma \boldsymbol{\varphi} = \mu \|\varphi\|_\Gamma^2$ we can conclude

$$(3.28) \quad \left(\frac{1}{1 + \frac{C\eta h^2}{1 + C\eta h^2}} \right) \left(1 - c \frac{h}{H\alpha} \right) \mu \|\varphi\|_\Gamma^2 \geq \boldsymbol{\varphi}^\top B_i \boldsymbol{\varphi}.$$

This completes the proof. □

We can use Lemma 3.8 to obtain a sharper contraction factor for the two-subdomain case with respect to η . In the following corollary, we study the effect of η on the contraction factor. We consider only the case when $\gamma = 1$ for clarity of the presentation. However, it is possible to use a combination of γ and the time-step $\tau = \eta^{-1}$ to optimize the contraction factor. Observe that in the following corollary, if $\eta = O(h^{-2})$, then the contraction factor is independent of the mesh size.

Theorem 3.9. *Consider the two-subdomain case and let $\gamma = 1$. Then the error of the interface variable satisfies the contraction estimate*

$$\|\mathbf{e}_i^{(n)}\|_{E_i^\top E_i} \leq \rho(h, H, \eta) \|\mathbf{e}_i^{(n-1)}\|_{E_i^\top E_i} \quad \text{for } i = 1, 2,$$

where $E_i := A_\Gamma^{1/2} (1 - A_\Gamma^{-1} B_i)$ and

$$\rho(h, H, \eta) := \begin{cases} 1 - c \frac{h}{H\alpha} & \text{for } \eta = O(1) \text{ and } O(h^{-1}), \\ 1 - C & \text{for } \eta = O(h^{-2}). \end{cases}$$

Proof. The proof relies on the proof given in [11, Section 4.1]. In the case of the two-subdomain case with $\gamma = 1$ we have from [11, Section 4.1] that

$$\|\mathbf{e}_i^{(n)}\|_{E_i^\top E_i} \leq \rho(h, H, \eta) \|\mathbf{e}_i^{(n-1)}\|_{E_i^\top E_i} \quad \text{for } i = 1, 2,$$

where

$$\rho(h, H, \eta) := \left[\left(\frac{1}{1 + \frac{C\eta h^2}{1 + C\eta h^2}} \right) \left(1 - c \frac{h}{H\alpha} \right) \right]^2,$$

which is the square of the upper bound constant in (3.28) divided by μ . Choosing $\eta = O(1)$, $O(h^{-1})$ and $O(h^{-2})$ completes the proof. In particular, observe that for $\eta = O(h^{-2})$ we have

$$\left(\frac{1}{1 + \frac{C\eta h^2}{1 + C\eta h^2}} \right) \left(1 - c \frac{h}{H\alpha} \right) \leq \left(\frac{1 - C_2 h}{1 + C_3} \right) \leq \left(\frac{1}{1 + C_3} \right).$$

This shows that with a time-step of the size of a forward Euler method, the algorithm converges in a *fixed number* of iterations since $\frac{1}{1 + C_3} < 1$ uniformly in h . \square

Let us now extend the above result to the case of many non-floating subdomains. In order to do so, we first need the following lemma that relates the $\mathcal{R}_i(\cdot)$ operator to B_i .

Lemma 3.10. *Let $B_i := A_{i\Gamma}^\top A_i^{-1} A_{i\Gamma}$ and $\mathcal{R}_i(\varphi_i) := \gamma \varphi_i - \frac{1}{2\mu} \left(\mu - \frac{\partial}{\partial \mathbf{n}_i} \right) \mathcal{H}_i(\varphi_i)$ for Ω_i which is a non-floating subdomain. Then the following estimate holds:*

$$(3.29) \quad c\mu \|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2 \leq \varphi_i^\top B_i \varphi_i,$$

where c is independent of h, α and η . Moreover, let $u_i = \mathcal{H}_i(\varphi_i)$; then we have

$$(3.30) \quad \left(\frac{\eta}{\eta + C h^{-2}} \right) \cdot c\mu \|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2 \leq \eta \|u_i\|_{\Omega_i}^2.$$

Proof. We take the L^2 -norm of $\mathcal{R}_i(\varphi_i)$ and use the triangle and Young’s inequality to obtain

$$\|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2 \leq 2\gamma^2 \|\varphi_i\|_{\Gamma_i}^2 + \frac{1}{2\mu^2} \|z_i\|_{\Gamma_i}^2,$$

where $z_i := (\mu - \partial_{\mathbf{n}_i})u_i \in \Lambda_i$. We know from [11, Proposition 2.4] that $z_i = M_{\Gamma_i}^{-1} B_i \varphi_i$. Then we have

$$\begin{aligned} \|z_i\|_{\Gamma_i}^2 &= \varphi_i^\top B_i M_{\Gamma_i}^{-1} M_{\Gamma_i} M_{\Gamma_i}^{-1} B_i \varphi = \varphi_i^\top B_i M_{\Gamma_i}^{-1} B_i \varphi_i \\ &= \varphi_i^\top B_i^{1/2} (B_i^{1/2} M_{\Gamma_i}^{-1} B_i^{1/2}) B_i^{1/2} \varphi_i, \end{aligned}$$

since B_i is s.p.d. A simple calculation shows that $\sigma(B_i^{1/2} M_{\Gamma_i}^{-1} B_i^{1/2}) = \sigma(M_{\Gamma_i}^{-1} B_i)$. Recall that $A_{\Gamma_i} = 2\mu M_{\Gamma_i}$. Then for z_i we have from the eigenvalues of $A_{\Gamma_i}^{-1} B_i$ (see [11, equation 3.1])

$$\|z_i\|_{\Gamma_i}^2 \leq 2\mu \cdot \sigma_{\max}(A_{\Gamma_i}^{-1} B_i) \cdot \varphi_i^\top B_i \varphi_i \leq 2\mu \cdot \varphi_i^\top B_i \varphi_i.$$

This yields

$$\|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2 \leq 2\gamma^2 \|\varphi_i\|_{\Gamma_i}^2 + \mu^{-1} \varphi_i^\top B_i \varphi_i \leq 2\|\varphi_i\|_{\Gamma_i}^2 + \mu^{-1} \varphi_i^\top B_i \varphi_i,$$

since $\gamma \leq 1$. The last step is to use $\|\varphi_i\|_{\Gamma_i}^2 \leq c_B^{-1} \mu^{-1} \varphi_i^\top B_i \varphi_i$, i.e., the lower bound for the eigenvalues of $A_{\Gamma_i}^{-1} B_i$ (see [11, equation 3.1]), where c_B is independent of η . Hence we proved (3.29). Using (3.27), we obtain for $u_i = \mathcal{H}_i(\varphi_i)$,

$$\left(\frac{\eta}{\eta + C h^{-2}} \right) \cdot c\mu \|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2 \leq \eta \|u_i\|_{\Omega_i}^2.$$

This completes the proof. \square

TABLE 1. Convergence of OSM for 110 subdomains (h -dependence).

Mesh size	h_0	$h_0/2$	$h_0/4$	$h_0/8$
# iterations	1057	1297	1951	2734

Lemma 3.10 enables us to prove the following theorem. Note that similar to the two-subdomain case, one can obtain a contraction factor independent of h by choosing $\eta = O(h^{-2})$.

Theorem 3.11. *Suppose the number of subdomains is fixed and they consist of only non-floating subdomains. Moreover, let $\gamma = 1$ and $\eta \geq 0$, then OSM converges, and we have the refined contraction estimate*

$$\|\mathcal{R}(\varphi^{(n)})\|^2 \leq \left(1 - C_1 \frac{\eta}{\eta + C_2 h^{-2}} - c \frac{h}{H\alpha}\right) \|\mathcal{R}(\varphi^{(n-1)})\|^2,$$

where $\|\mathcal{R}(\varphi)\|^2 := \sum_{i=1}^{N_s} \|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2$.

Proof. We consider the case when $\gamma = 1$. Inserting (3.30) into (A.7) and then into (3.17) gives

$$\|\mathcal{R}(\varphi^{(n)})\|^2 \leq \left(1 - C_1 \frac{\eta}{\eta + C_2 h^{-2}} - c \frac{h}{H\alpha}\right) \|\mathcal{R}(\varphi^{(n-1)})\|^2.$$

Note that for $\eta = O(h^{-2})$ the above estimate provides a contraction factor independent of the mesh parameter. This completes the proof. \square

4. NUMERICAL EXPERIMENTS

We now illustrate our theoretical results by performing some numerical experiments for the model problem

$$(4.1) \quad \begin{aligned} (\eta - \Delta)u &= f && \text{in } \Omega, \\ u &= 0 && \text{on } \partial\Omega, \end{aligned}$$

where Ω is either the unit square, i.e., $\Omega = (0,1)^2$, or the domain presented in Figure 1. The interface is such that it does not cut through any element, therefore $\Gamma \subset \mathcal{E}$. We use \mathbb{P}^k elements and $\alpha = c(k+1)(k+2)$, where $c > 0$ is a constant independent of h and k . We choose also a randomized initial guess for Algorithm 3.2.

4.1. Dependence on the mesh size. In [11, Section 6.3], we have already investigated numerically the convergence behavior of OSM for IPH for a many-subdomain configuration, and we show in Table 1 that indeed for a unit square domain decomposed into 110 subdomains (see Figure 3 left) the number of iterations grows like $h^{-1/2}$ when we refine the mesh, provided that $\gamma = \frac{1}{2}(1 + \sqrt{h})$, as our new theoretical analysis predicts.

4.2. Dependence on the polynomial degree. We next illustrate how the contraction factor of Algorithm 3.2 depends on the polynomial degree. First, we choose a two-subdomain configuration with a non-straight interface (see Figure 3 right) for $\Omega = (0,1)^2$. Then we choose $\gamma = \frac{1}{2}(1 + \frac{1}{k})$, $\eta = 1$ and run Algorithm 3.2. We expect from our analysis to obtain $\rho \leq 1 - O(\frac{1}{k})$, which is indeed observed in Figure 4.




FIGURE 3. An unstructured mesh with the interface Γ (blue-dashed).




FIGURE 4. Convergence of the OSM with respect to polynomial degree.

TABLE 2. Convergence of Algorithm 3.2 with $\gamma = 1$ and different choices of η .

	h_0	$h_0/2$	$h_0/4$	$h_0/8$
case $\eta = O(1)$	103	214	405	820
case $\eta = O(h^{-1})$	41	60	83	115
case $\eta = O(h^{-2})$	16	16	15	14

Then we choose Ω to be the domain in Figure 1 with seven subdomains (including floating ones). We observe in Figure 4 that the number of iterations grows like $O(k^{-1})$, which is expected from our analysis.

4.3. Effect of the time-step on convergence. In Section 3.3 we showed how the convergence of the two-subdomain algorithm is affected by the choice of η . In Table 2 we see the number of iterations required to reach a given accuracy for different choices of η . The domain decomposition setting is same as Figure 3 (right).

Observe that for $\eta = O(1)$, the number of iterations grows like $O(h^{-1})$ while for $\eta = O(h^{-1})$ we observe $O(h^{-1/2})$ for the growth of the number of iterations. If

TABLE 3. Comparison of contraction factors between the theoretical estimates of Section 3.3 and the numerical experiments.

	experiments	theoretical	
case $\eta = O(1)$	$1 - h$	$1 - h$	(sharp)
case $\eta = O(h^{-1})$	$1 - \sqrt{h}$	$1 - h$	(not sharp)
case $\eta = O(h^{-2})$	$1 - c$	$1 - c$	(sharp)

TABLE 4. Convergence of OSM for four subdomains with $\eta = O(h^{-2})$.

Mesh size	h_0	$h_0/2$	$h_0/4$	$h_0/8$
# iterations	144	157	168	164

TABLE 5. Convergence of OSM while the ratio H/h is constant with $\eta = O(H^{-2})$.

Mesh size	h_0	$h_0/2$	$h_0/4$	$h_0/8$
# iterations	105	95	99	104

we choose $\eta = O(h^{-2})$, we obtain an optimal solver since the number of iterations does not depend on the mesh parameter.

In Table 3 we compare the theoretical estimate in (3.28) with the numerical experiments of Table 2. Note that the estimates of Section 3.3 can capture the optimality of the solver when $\eta = O(h^{-2})$. However, it is not sharp when $\eta = O(h^{-1})$.

We perform the same experiment with four subdomains on $\Omega = (0, 1)^2$ and we choose $\eta = O(h^{-2})$. We see in Table 4 that the number of iterations remains constant as we refine the mesh.

Finally we perform numerical experiments on the weak scaling of the algorithm. According to Corollary 3.5, when $\tau = O(H^2)$ and the ratio H/h is constant, i.e., we refine the mesh and the subdomain at the same time, one obtains a contraction factor independent of the mesh size. This can be achieved also using ASM applied to FEM. In Table 5, we illustrate the convergence of the OSM on a sequence of fine and coarse meshes such that the ratio H/h remains constant.

5. CONCLUSION

We designed and analyzed an optimized Schwarz method (OSM) for the solution of elliptic problems discretized by hybridizable interior penalty (IPH) discontinuous Galerkin methods. Our results are a generalization of the two-subdomain analysis in [11] to the case of many subdomains, and we also study theoretically for the first time the influence of the polynomial degree of IPH discretizations, and the effect of the time-step on the convergence of OSM when solving parabolic problems. We derived the optimized parameter and corresponding contraction factor for various asymptotic regimes of the mesh and subdomain size and the time-step, and obtained scalability without a coarse space and also mesh independent solvers in certain specific regimes. We validated our theoretical results by numerical experiments. The optimized contraction factor shows a clear advantage of OSM compared to the additive Schwarz method applied to the primal formulation, e.g., see the one-level

ASM version of [8] or [1]. The next step is to design and analyze a coarse correction for these OSM solvers applied to IPH in the regimes where Algorithm 3.2 is not scalable.

APPENDIX A. PROOF OF SOME ESTIMATES

We now prove several technical estimates we used in the analysis of the OSM for IPH. For all subdomains when $\eta \geq 0$ we have the inequalities

$$(A.1) \quad \mu \|\varphi_i\|_{\Gamma_i}^2 - a_i(v_i, v_i) \geq c \|(v_i, \varphi_i)\|_i^2 \quad \forall \varphi_i \in \Lambda_i, v_i = \mathcal{H}_i(\varphi_i),$$

$$(A.2) \quad (1 - c(h, H, k))\mu \|\varphi_i\|_{\Gamma_i}^2 \geq a_i(v_i, v_i) \quad \forall \varphi_i \in \Lambda_i, v_i = \mathcal{H}_i(\varphi_i),$$

where

$$c(h, H, k) := \begin{cases} \frac{h}{H} \frac{1}{k^2} & \text{for non-floating subdomains,} \\ 0 & \text{for floating subdomains.} \end{cases}$$

We also have for all subdomains when $\eta > 0$ the estimate

$$(A.3) \quad \|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2 \leq \left((2\gamma - 1)^2 C(H, \eta) + \mu^{-1} \right) \|(u_i, \varphi_i)\|_i^2,$$

where

$$C(H, \eta) := \begin{cases} H & \text{for non-floating subdomain,} \\ \frac{1}{H\eta} & \text{for floating subdomain.} \end{cases}$$

We first recall an inequality related to the coercivity of the IPH method, that is,

$$(A.4) \quad a_i(v_i, v_i) + 2a_{i\Gamma}(v_i, \varphi_i) + \mu \|\varphi_i\|_{\Gamma_i}^2 \geq c \|(v_i, \varphi_i)\|_i^2 \quad \forall \varphi_i \in \Lambda_i, v_i \in V_{h,i}.$$

For a proof see [18] and [11]. The proof of (A.1) is obtained by choosing $v_i := \mathcal{H}_i(\varphi_i)$ in (A.4) and recalling the definition of the harmonic extension which leads to $a_i(v_i, v_i) + a_{i\Gamma}(v_i, \varphi_i) = 0$. Substituting this into (A.4) proves (A.1).

In order to prove (A.2) we decompose the proof into two parts: floating subdomains and non-floating subdomains. Recall that $\|(\cdot, \cdot)\|_i$ is a semi-norm for floating subdomains if $\eta = 0$, i.e., the kernel consists of constant functions. This concludes the proof for floating subdomains with $c(h, H, k) = 0$. For non-floating subdomains we recall a trace inequality for totally discontinuous functions; see [11, Lemma 3.6] and [3].

Lemma A.1. *Let $\varphi_i \in \Lambda_i$ and $u_i \in V_{h,i}$. Let H_i be the diameter of a non-floating subdomain. Then we have*

$$(A.5) \quad \frac{c}{H_i} \|\varphi_i\|_{\Gamma_i}^2 \leq \|\nabla u_i\|_{\Omega_i}^2 + \mu \|\llbracket u_i \rrbracket\|_{\mathcal{E}_i \setminus \Gamma_i}^2 + \mu \|u_i - \varphi_i\|_{\Gamma_i}^2.$$

We then substitute (A.5) into (A.1) and recalling the definition of $\|(u_i, \varphi_i)\|_i$ proves (A.2) for non-floating subdomains with $c(h, H, k) = \frac{h}{H} \frac{1}{k^2}$.

We now prove (A.3). Recall that the L^2 -norm of the $\mathcal{R}_i(\cdot)$ is a norm while $\|(\cdot, \cdot)\|_i$ is only a semi-norm for floating subdomains if $\eta = 0$. Therefore (A.3) makes sense for $\eta > 0$. Recall the definition of the $\mathcal{R}_i(\cdot)$ operator,

$$\mathcal{R}_i(\varphi_i) := \gamma \varphi_i - \frac{1}{2\mu} \left(\mu - \frac{\partial}{\partial \mathbf{n}_i} \right) u_i = \left(\gamma - \frac{1}{2} \right) \varphi_i + \frac{1}{2} (\varphi_i - u_i) + \frac{1}{2\mu} \frac{\partial u_i}{\partial \mathbf{n}_i},$$

where $u_i = \mathcal{H}_i(\varphi_i)$. We then take the L^2 -norm over Γ_i and apply the triangle inequality,

$$\begin{aligned} \|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2 &\leq 4\left(\gamma - \frac{1}{2}\right)^2 \|\varphi_i\|_{\Gamma_i}^2 + 4\left(\frac{1}{2}\right)^2 \|\varphi_i - u_i\|_{\Gamma_i}^2 + 4\left(\frac{1}{2\mu}\right)^2 \left\|\frac{\partial u_i}{\partial \mathbf{n}_i}\right\|_{\Gamma_i}^2, \\ (A.6) \quad &\leq 4\left(\gamma - \frac{1}{2}\right)^2 \|\varphi_i\|_{\Gamma_i}^2 + \|\varphi_i - u_i\|_{\Gamma_i}^2 + c\mu^{-1} \|\nabla u_i\|_{\Omega_i}^2, \\ &\leq 4\left(\gamma - \frac{1}{2}\right)^2 \|\varphi_i\|_{\Gamma_i}^2 + c\mu^{-1} (\mu \|\varphi_i - u_i\|_{\Gamma_i}^2 + \|\nabla u_i\|_{\Omega_i}^2). \end{aligned}$$

For non-floating subdomains we use Lemma A.1 for the first term on the right-hand side and obtain

$$(A.7) \quad \left((2\gamma - 1)^2 H_i + c\mu^{-1}\right) \eta \|u_i\|_{\Omega_i}^2 + \|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2 \leq \left((2\gamma - 1)^2 H_i + c\mu^{-1}\right) \|(u_i, \varphi_i)\|_i^2.$$

For floating subdomains we use a trace inequality by Feng and Karakashian [8, Lemma 3.1],

$$(A.8) \quad \|u_i\|_{\Gamma_i}^2 \leq c \left[H_i^{-1} \|u_i\|_{\Omega_i}^2 + H_i (\|\nabla u_i\|_{\Omega_i}^2 + h^{-1} \|[u_i]\|_{\mathcal{E}_i \setminus \Gamma_i}^2) \right].$$

We then invoke $\|\varphi_i\|_{\Gamma_i}^2 \leq 2\|u_i\|_{\Gamma_i}^2 + 2\|u_i - \varphi_i\|_{\Gamma_i}^2$, use (A.8) and recall the definition of $\|(u_i, \varphi_i)\|$ to obtain

$$\|\varphi_i\|_{\Gamma_i}^2 \leq \frac{C}{H_i \eta} \|(u_i, \varphi_i)\|_i^2.$$

Substituting this estimate back into (A.6) yields

$$(A.9) \quad \|\mathcal{R}_i(\varphi_i)\|_{\Gamma_i}^2 \leq \left((2\gamma - 1)^2 \cdot C \cdot (H_i \eta)^{-1} + c\mu^{-1}\right) \|(u_i, \varphi_i)\|_i^2.$$

REFERENCES

- [1] P. F. Antonietti and B. Ayuso, *Schwarz domain decomposition preconditioners for discontinuous Galerkin approximations of elliptic problems: non-overlapping case*, M2AN Math. Model. Numer. Anal. **41** (2007), no. 1, 21–54, DOI 10.1051/m2an:2007006. MR2323689
- [2] D. N. Arnold, F. Brezzi, B. Cockburn, and L. D. Marini, *Unified analysis of discontinuous Galerkin methods for elliptic problems*, SIAM J. Numer. Anal. **39** (2001/02), no. 5, 1749–1779, DOI 10.1137/S0036142901384162. MR1885715
- [3] S. C. Brenner, *Poincaré-Friedrichs inequalities for piecewise H^1 functions*, SIAM J. Numer. Anal. **41** (2003), no. 1, 306–324, DOI 10.1137/S0036142902401311. MR1974504
- [4] X.-C. Cai, *Additive Schwarz algorithms for parabolic convection-diffusion equations*, Numer. Math. **60** (1991), no. 1, 41–61, DOI 10.1007/BF01385713. MR1131498
- [5] P. Castillo, *Performance of discontinuous Galerkin methods for elliptic PDEs*, SIAM J. Sci. Comput. **24** (2002), no. 2, 524–547, DOI 10.1137/S1064827501388339. MR1951054
- [6] B. Cockburn, J. Gopalakrishnan, and R. Lazarov, *Unified hybridization of discontinuous Galerkin, mixed, and continuous Galerkin methods for second order elliptic problems*, SIAM J. Numer. Anal. **47** (2009), no. 2, 1319–1365, DOI 10.1137/070706616. MR2485455
- [7] R. E. Ewing, J. Wang, and Y. Yang, *A stabilized discontinuous finite element method for elliptic problems*, Numer. Linear Algebra Appl. **10** (2003), no. 1-2, 83–104, DOI 10.1002/nla.313. Dedicated to the 60th birthday of Raytcho Lazarov. MR1964287
- [8] X. Feng and O. A. Karakashian, *Two-level additive Schwarz methods for a discontinuous Galerkin approximation of second order elliptic problems*, SIAM J. Numer. Anal. **39** (2001), no. 4, 1343–1365, DOI 10.1137/S0036142900378480. MR1870847
- [9] M. J. Gander, *Optimized Schwarz methods*, SIAM J. Numer. Anal. **44** (2006), no. 2, 699–731, DOI 10.1137/S0036142903425409. MR2218966
- [10] M. J. Gander and S. Hajian, *Block Jacobi for discontinuous Galerkin discretizations: no ordinary Schwarz methods*, Domain Decomposition Methods in Science and Engineering XXI, Lect. Notes Comput. Sci. Eng., Springer, 2013.

- [11] M. J. Gander and S. Hajian, *Analysis of Schwarz methods for a hybridizable discontinuous Galerkin discretization*, SIAM J. Numer. Anal. **53** (2015), no. 1, 573–597, DOI 10.1137/140961857. MR3313831
- [12] M. J. Gander and F. Kwok, *Best Robin parameters for optimized Schwarz methods at cross points*, SIAM J. Sci. Comput. **34** (2012), no. 4, A1849–A1879, DOI 10.1137/110837218. MR2970388
- [13] M. J. Gander and F. Kwok, *On the applicability of Lionsâ energy estimates in the analysis of discrete optimized Schwarz methods with cross points*, Domain Decomposition Methods in Science and Engineering XX, Springer, 2013, pp. 475–483.
- [14] M. J. Gander and K. Santugini, *Cross-points in domain decomposition methods with a finite element discretization*, revised (2015).
- [15] C. J. Gittelsohn, R. Hiptmair, and I. Perugia, *Plane wave discontinuous Galerkin methods: analysis of the h-version*, M2AN Math. Model. Numer. Anal. **43** (2009), no. 2, 297–331, DOI 10.1051/m2an/2009002. MR2512498
- [16] S. Hajian, *An optimized Schwarz algorithm for discontinuous Galerkin methods*, Domain Decomposition Methods in Science and Engineering XXII, Springer, 2014.
- [17] S. Hajian, *Analysis of Schwarz methods for discontinuous Galerkin discretizations*, Ph.D. thesis, 06/04 2015, ID: unige:75225.
- [18] C. Lehrenfeld, *Hybrid discontinuous Galerkin methods for incompressible flow problems*, Master’s thesis, RWTH Aachen, 2010.
- [19] P.-L. Lions, *On the Schwarz alternating method. III. A variant for nonoverlapping subdomains*, Third International Symposium on Domain Decomposition Methods for Partial Differential Equations (Houston, TX, 1989), SIAM, Philadelphia, PA, 1990, pp. 202–223. MR1064345
- [20] S. Loisel, *Condition number estimates for the nonoverlapping optimized Schwarz method and the 2-Lagrange multiplier method for general domains and cross points*, SIAM J. Numer. Anal. **51** (2013), no. 6, 3062–3083, DOI 10.1137/100803316. MR3129755
- [21] L. Qin, Z. Shi, and X. Xu, *On the convergence rate of a parallel nonoverlapping domain decomposition method*, Sci. China Ser. A **51** (2008), no. 8, 1461–1478, DOI 10.1007/s11425-008-0103-2. MR2426076
- [22] L. Qin and X. Xu, *On a parallel Robin-type nonoverlapping domain decomposition method*, SIAM J. Numer. Anal. **44** (2006), no. 6, 2539–2558, DOI 10.1137/05063790X. MR2272605
- [23] L. Qin and X. Xu, *Optimized Schwarz methods with Robin transmission conditions for parabolic problems*, SIAM J. Sci. Comput. **31** (2008), no. 1, 608–623, DOI 10.1137/070682149. MR2460791
- [24] A. Toselli and O. Widlund, *Domain Decomposition Methods—Algorithms and Theory*, Springer Series in Computational Mathematics, vol. 34, Springer-Verlag, Berlin, 2005. MR2104179

SECTION DE MATHÉMATIQUES, UNIVERSITÉ DE GENÈVE, GENEVA, SWITZERLAND

E-mail address: martin.gander@unige.ch

INSTITUT FÜR MATHEMATIK, HUMBOLDT-UNIVERSITÄT ZU BERLIN, BERLIN, GERMANY

E-mail address: soheil.hajian@hu-berlin.de