THE DIFFERENCE BETWEEN CONSECUTIVE PRIME NUMBERS. IV

R. A. RANKIN

1. **Introduction.** Let p_n denote the n th prime, and let Θ be the lower bound of all positive numbers σ_0 such that no Dirichlet L -function $L(s, \chi)$ has a zero at $s = \sigma + it$, where $\sigma > \sigma_0$. Write

$$l = \liminf_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\log p_n}.$$

The purpose of this paper is to combine the methods used in two earlier papers¹ in order to prove the following theorem.

THEOREM.

$$(1) \quad l \leq c(1 + 4\Theta)/5,$$

where $c < 42/43$.

The precise definition of c is given by (18) and (8) below.

The theorem is an improvement on the result $l \leq (1 + 4\Theta)/5$ obtained in II. It was shown in III that

$$(2) \quad l \leq d = 0.966 \dots < 57/59.$$

The numbers c and d are connected by the relation

$$1 - d = (3 + 2^{3/2})(1 - c)/4 > 1 - c,$$

so that (1) is an improvement on (2) only if Θ is not too close to unity, in fact, if $\Theta < 0.986 \dots$. In particular, if the "grand Riemann hypothesis" is true, that is, if $\Theta = 1/2$, we have (using the value of c given by (18))

$$l \leq 3c/5 < 109/186 = 0.58602 \dots$$

2. **Notation.** Since (2) is sharper than (1) for $\Theta = 1$, we shall assume that $\Theta < 1$, and write

$$(3) \quad \sigma = (1 + 4\Theta)/5.$$

Let N be a large positive integer, and define

$$(4) \quad X = N^{1-\sigma}(\log N)^{-s(1+\gamma)}$$

Received by the editors January 4, 1949.

¹ See [2] and [3]; I refer to these as II and III respectively. Numbers in brackets refer to the references cited at the end of the paper.

as done in equation (6.7) of II; here γ is a positive constant which may be taken to be 2.

Let $\tilde{\omega}(\nu)$ be the number of prime pairs p, p' which satisfy

$$N < p' \leq p \leq 2N, \quad p - p' = \nu,$$

where ν is an integer. We suppose that h, k, m, n and r are integers such that, for some positive A_1, A_2 independent of N ,

$$(5) \quad A_1 \log N < 2k/3 < h < k < A_2 \log N,$$

$$(6) \quad 0 \leq m < n < A_2 \log N,$$

$$r = k - h.$$

The letters a, c, α, β denote positive numbers independent of N , and we write

$$(7) \quad D = \prod_{p>2} \left(1 + \frac{1}{p(p-2)} \right),$$

and take

$$(8) \quad \lambda = 7.3566 > e^{-2\gamma} \Delta_0 / 2,$$

where γ is Euler's constant and Δ_0 is Buchstab's number 46.67347 (see relation (4) of III).

For $\mu = 1, 2, \dots, k$ we define a_μ as follows:

$$(9) \quad a_\mu = \begin{cases} \alpha & (0 < \mu \leq r, k - r < \mu \leq k), \\ \beta & (r < \mu \leq k - r), \end{cases}$$

and write

$$(10) \quad \Psi(\theta) = \sum_{\mu=1}^k a_\mu e^{4\pi i \mu \theta}.$$

Then

$$(11) \quad |\Psi(\theta)|^2 = \xi(0) + 2 \sum_{\nu=1}^{k-1} \xi(\nu) \cos 4\pi \nu \theta,$$

where

$$\xi(\nu) = \sum_{i+j=\nu, 1 \leq i \leq k} a_i a_j.$$

3. Lemmas.

LEMMA 1. *We have*

$$\xi(\nu) = \begin{cases} 2(r - \nu)\alpha^2 + 2\nu\alpha\beta + (k - 2r - \nu)\beta^2 & (0 \leq \nu < r), \\ 2r\alpha\beta + (k - 2r - \nu)\beta^2 & (r \leq \nu < k - 2r), \\ (2r - k + \nu)\alpha^2 + 2(k - r - \nu)\alpha\beta & (k - 2r \leq \nu < k - r), \\ (k - \nu)\alpha^2 & (k - r \leq \nu < k), \end{cases}$$

and

$$\sum_{\nu=1}^{k-1} \xi(\nu) = \frac{1}{2} \{2r\alpha + (k - 2r)\beta\}^2 + O(\log N).$$

This follows easily from (9) and (11), and we omit the proof. It is in this lemma that the restriction $k - r = h > 2k/3$ made in (5) is used.

LEMMA 2.

$$\sum_{1 \leq q \leq X} \frac{\mu^2(q)}{\phi^2(q)} \sum_{f=0, (f, q)=1}^{q-1} e^{4\pi i f/q} = \frac{2}{D} \prod_{p|r, p>2} \left(\frac{p-1}{p-2} \right) + O(\nu X^{-1} (\log \log X)^2).$$

Here $\mu(q)$ is the Möbius function, $\phi(q)$ is Euler's function, and D is defined by (7).

It follows from the well known relation

$$\sum_{f=0, (f, q)=1}^{q-1} e^{4\pi i f/q} = \sum_{d|(2r, q)} d\mu \left(\frac{q}{d} \right)$$

that

$$\begin{aligned} \sum_{1 \leq q \leq X} \frac{\mu^2(q)}{\phi^2(q)} \sum_{f=0, (f, q)=1}^{q-1} e^{4\pi i f/q} &= \sum_{q \leq X} \frac{\mu(q)}{\phi^2(q)} \prod_{p|(2r, q)} (1 - p) \\ &= \sum_{q=1}^{\infty} \frac{\mu(q)}{\phi^2(q)} \prod_{p|(2r, q)} (1 - p) \\ &\quad - \sum_{q > X} \frac{\mu(q)}{\phi^2(q)} \prod_{p|(2r, q)} (1 - p). \end{aligned}$$

By a straightforward reduction the first term on the right may be shown to equal

$$\frac{2}{D} \prod_{p|r, p>2} \left(\frac{p-1}{p-2} \right).$$

See, for example, §§3.21–3.23 of [1]. Since $1/\phi(n) = O(n^{-1} \log \log n)$, the second term on the right is

$$O\left\{X^{-1}(\log \log X)^2 \prod_{p|2^s} (p-1)\right\} = O\{\nu X^{-1}(\log \log X)^2\}.$$

LEMMA 3. *If $s \geq 0$ and is independent of N , then*

$$\sum_{\nu=m}^{n-1} \nu^s \prod_{p|\nu, p>2} \left(\frac{p-1}{p-2}\right) = D \sum_{\nu=m}^{n-1} \nu^s + O\{(\log N)^s (\log \log N)^2\}.$$

The proof is similar to that of the lemma in III. We suppose that d is any product $p_{i_1} p_{i_2} \cdots p_{i_j}$ of distinct odd primes, and let d' be the associated product

$$d' = (p_{i_1} - 2)(p_{i_2} - 2) \cdots (p_{i_j} - 2).$$

We permit d to take the value unity and then put $d' = 1$. Then, clearly,

$$\begin{aligned} \sum_{d \leq n} \frac{1}{dd'} &= \sum_{d \geq 1} \frac{1}{dd'} + O\left\{\frac{1}{n} (\log \log 3n)^2\right\} \\ &= D + O\left\{\frac{1}{n} (\log \log 3n)^2\right\}, \end{aligned}$$

since $1/d' = O\{d^{-1}(\log \log d)^2\}$ for large² d ; and

$$\begin{aligned} \prod_{p|\nu, p>2} \left(\frac{p-1}{p-2}\right) &= \prod_{p|\nu, p>2} \left(1 + \frac{1}{p-2}\right) \\ &= \sum_{d|\nu} \frac{1}{d'}. \end{aligned}$$

For a given d' , ν is a multiple of d , say $\nu = \mu d$, and, since $m \leq \nu < n$, μ takes all integer values in the interval $m/d \leq \mu < n/d$, which we denote by $I(d)$. Clearly

$$\sum_{\mu \in I(d)} (\mu d)^s = \frac{n^{s+1} - m^{s+1}}{(s+1)d} + O(n^s),$$

and

$$\sum_{\nu=m}^{n-1} \nu^s = \frac{n^{s+1} - m^{s+1}}{s+1} + O(n^s).$$

² It is possible to dispense with the factor $(\log \log 3n)^2$.

Hence

$$\begin{aligned}
 \sum_{r=m}^{n-1} \nu^s \prod_{p|\nu, p>2} \left(\frac{p-1}{p-2} \right) &= \sum_{d<n} \frac{1}{d'} \sum_{\mu \in I(d)} (\mu d)^s \\
 &= \frac{n^{s+1} - m^{s+1}}{s+1} \sum_{d<n} \frac{1}{dd'} + O\left(n^s \sum_{d<n} \frac{1}{d'}\right) \\
 &= \frac{n^{s+1} - m^{s+1}}{s+1} \left\{ D + O\left(\frac{1}{n} (\log \log 3n)^2\right) \right\} \\
 &\quad + O\{n^s (\log \log 3n)^2 \log 2n\} \\
 &= D \sum_{r=m}^{n-1} \nu^s + O\{n^s (\log 2n)^2\},
 \end{aligned}$$

and the lemma follows from (6).

LEMMA 4.

$$\begin{aligned}
 \sum_{1 \leq q \leq X} \frac{\mu^2(q)}{\phi^2(q)} \sum_{f=0, (f,q)=1}^{q-1} \sum_{r=1}^{k-1} \xi(\nu) e^{4\pi i r f / q} \\
 = 2 \sum_{r=1}^{k-1} \xi(\nu) + O\{\log N(\log \log N)^2\} \\
 = \{2r\alpha + (k-2r)\beta\}^2 + O\{\log N(\log \log N)^2\}.
 \end{aligned}$$

By Lemma 2, the left-hand member is equal to

$$\frac{2}{D} \sum_{r=1}^{k-1} \xi(\nu) \prod_{p|\nu, p>2} \left(\frac{p-1}{p-2} \right) + O\left\{kX^{-1}(\log \log X)^2 \sum_{r=1}^{k-1} \xi(\nu)\right\}.$$

It follows from Lemmas 1 and 3 (with $s=0, 1$) that this equals

$$2 \sum_{r=1}^{k-1} \xi(\nu) + O\{\log N(\log \log N)^2\} + O\{X^{-1}(\log N)^2(\log \log X)^2\},$$

and the lemma follows from (4) and the second part of Lemma 1.

4. **Proof of the theorem.** By considering the expression

$$I(N, k) = \int_0^1 |S(\theta)\Psi(\theta)|^2 d\theta,$$

where

$$S(\theta) = \sum_{N < p \leq 2N} e^{2\pi i \theta p} \log p \quad (p \text{ prime}),$$

and $\Psi(\theta)$ is defined by (10), it is possible to show that, for large N ,

$$\begin{aligned}
 I(N, k) &= \xi(0)N \log N \{1 + o(1)\} \\
 &\quad + 2\{1 + o(1)\} \log^2 N \sum_{\nu=1}^{k-1} \xi(\nu) \bar{\omega}(2\nu) \\
 &> N\{1 + o(1)\} \sum_{1 \leq q \leq x} \frac{\mu^2(q)}{\phi^2(q)} \sum_{f=0, f, q=1}^{q-1} \left| \Psi\left(\frac{f}{q}\right) \right|^2 \\
 (12) \quad &\quad + o(N \log^2 N) \\
 &= \xi(0)N\{1 + o(1)\} \sum_{1 \leq q \leq x} \frac{\mu^2(q)}{\phi(q)} \\
 &\quad + 2N\{1 + o(1)\} \Re \sum_{1 \leq q \leq x} \frac{\mu^2(q)}{\phi^2(q)} \\
 &\quad \cdot \sum_{f=0, f, q=1}^{q-1} \sum_{\nu=1}^{k-1} \xi(\nu) e^{4\pi i f \nu / q} + o(N \log^2 N).
 \end{aligned}$$

The proof of this is similar in every respect to the proof of (4.1) and (6.8) of II, except that $\Psi(\xi)$ is defined by (10) and is not $\sum_{\nu=1}^{k-1} e^{2\pi i \nu \theta}$ as in II. The restriction $k < \log N$ made in II is unessential and can be replaced by $k < A_2 \log N$.

It follows from (3), (4), (12), Lemma 4, and Lemma 4 of³ II that

$$\begin{aligned}
 &2\{1 + o(1)\} \log^2 N \sum_{\nu=1}^{k-1} \xi(\nu) \bar{\omega}(2\nu) \\
 &> 4N\{1 + o(1)\} \sum_{\nu=1}^{k-1} \xi(\nu) - \sigma N \log N \{1 + o(1)\} \xi(0) \\
 (13) \quad &\quad + o(N \log^2 N) \\
 &= 2N\{2r\alpha + (k - 2r)\beta\}^2 \{1 + o(1)\} \\
 &\quad - \sigma N \log N \{2r\alpha^2 + (k - 2r)\beta^2\} \{1 + o(1)\} \\
 &\quad + o(N \log^2 N).
 \end{aligned}$$

Now, just as in III, it can be shown by Buchstab's form of Brun's method⁴ that, for sufficiently large N ,

³ There is a misprint in the enunciation of the lemma which should read: "If X is large $\sum_{1 \leq q \leq x} \mu^2(q)/\phi(q) \sim \log X$."

⁴ We must consider the interval $(N, 2N)$ instead of the interval $(x/\log x, x)$. Our only reason for considering an interval of the form $(N, 2N)$ is that it enables us to quote results directly from II. As far as Brun's method is concerned, the inequality stated remains true if we replace the left-hand side $\bar{\omega}(2\nu)$ by the number of prime pairs p, p' which satisfy $AN < p' \leq p \leq (A+1)N$, $p - p' = 2\nu > 0$ where $A \geq 0$.

$$\bar{\omega}(2\nu) < \frac{4\lambda N}{D(\log N)^2} \prod_{p|\nu, p>2} \left(\frac{p-1}{p-2}\right),$$

where λ is defined by (8). It follows from Lemmas 1 and 3 that

$$(14) \quad \sum_{\nu=k-r}^{k-1} \xi(\nu) \bar{\omega}(2r) < \frac{4\lambda N}{(\log N)^2} \sum_{\nu=k-r}^{k-1} \xi(\nu) + O\left\{\frac{N}{\log N} (\log \log N)^2\right\} \\ = \frac{2\lambda\alpha^2 r^2 N}{(\log N)^2} + O\left\{\frac{N}{\log N} (\log \log N)^2\right\}$$

since $r = O(\log N)$.

Accordingly, if we make the assumption that $\bar{\omega}(2\nu) = 0$ for all $\nu < h = k - r$ we have, from (13) and (14), that

$$(15) \quad \left\{ \frac{2r\alpha + (k - 2r)\beta}{\log N} \right\}^2 \{1 + o(1)\} \\ - \frac{1}{2} \sigma \left\{ \frac{2r\alpha^2 + (k - 2r)\beta^2}{\log N} \right\} \{1 + o(1)\} \\ < 2\lambda\alpha^2 \left(\frac{r}{\log N} \right)^2 \{1 + o(1)\}.$$

We now take $\alpha = 1$, $\beta = 2^{1/2}$, and put

$$(16) \quad k = \left[\frac{1}{2} \sigma \left(1 + \frac{2^{1/2}a}{\lambda} \right) \log N \right], \\ h = \left[\frac{1}{2} \sigma \left(1 - \frac{a}{\lambda} \right) \log N \right],$$

where a is a fixed positive number satisfying

$$(17) \quad a < 3 - 2^{3/2}.$$

It is easily verified that, with this choice of a , $h > 2k/3$.

Then, for large N ,

$$r = \frac{a\sigma}{2\lambda} (2^{1/2} + 1) \log N + O(1),$$

$$2r\alpha + (k - 2r)\beta = (2 - 2^{1/2})(k + 2^{1/2}h) = \frac{1}{2^{1/2}} \sigma \log N + O(1),$$

$$2r\alpha^2 + (k - 2r)\beta^2 = 2h = \sigma \left(1 - \frac{a}{\lambda} \right) \log N + O(1).$$

It follows from (15) that

$$\frac{1}{2}\sigma^2 - \frac{1}{2}\sigma^2\left(1 - \frac{a}{\lambda}\right) \leq 2\lambda \left\{ \frac{\sigma a(2^{1/2} + 1)}{2\lambda} \right\}^2,$$

that is, that

$$a \geq 3 - 2^{3/2},$$

which contradicts (17).

We conclude, accordingly, that $\tilde{\omega}(2\nu) > 0$ for some integer ν satisfying

$$\nu < \frac{1}{2}\sigma\left(1 - \frac{a}{\lambda}\right)\log N.$$

It follows from this inequality, and from (17), that

$$l \leq \sigma c$$

where

$$(18) \quad c = 1 - (3 - 2^{3/2})/\lambda = 0.97667 \dots < \frac{42}{43},$$

and this proves the theorem.

Note added 24 April 1949. Professor Erdős has informed me that considerable improvements have recently been made in Brun's method by A. Selberg and others; this will presumably mean that Buchstab's constant Λ_0 , and consequently my constants c and d (see equations (2) and (18)), can be replaced by smaller values.

REFERENCES

1. G. H. Hardy and J. E. Littlewood, *Partitio Numerorum* III: *On the expression of a number as the sum of primes*, Acta Math. vol. 44 (1923) pp. 1-70.
2. R. A. Rankin, *The difference between consecutive prime numbers*, II, Proc. Cambridge Philos. Soc. vol. 36 (1940) pp. 255-266.
3. ———, *The difference between consecutive prime numbers*, III, J. London Math. Soc. vol. 22 (1947) pp. 226-230.

CLARE COLLEGE, CAMBRIDGE