POLYNOMIAL ITERATIONS TO ROOTS OF ALGEBRAIC EQUATIONS

ALSTON S. HOUSEHOLDER

If ξ is a root of the equation

$$(1) \quad f(x) = 0,$$

and if $\phi(x)$ satisfies

$$(2) \quad \phi(\xi) = \xi, \quad \phi^{(s)}(\xi) = 0 \quad (s = 1, 2, \dots, r-1),$$

then $\phi(x)$ is said to define an iteration of order r to the root ξ . In fact, for $r > 1$, when x_0 is in a sufficiently small neighborhood of ξ the sequence

$$(3) \quad x_{i+1} = \phi(x_i)$$

converges to ξ with

$$(4) \quad x_{i+1} = \xi + O(x_i - \xi)^r.$$

For analytic f , iterations of all orders exist and can be constructed in many ways. Domb [2]¹ has shown further that for polynomial f it is always possible to make ϕ a polynomial. The purpose of this note is to describe a simple algorithm:

Let $f(x)$ be a polynomial with no multiple factors; let $p(x)$ and $q(x)$ be any polynomials satisfying

$$(5) \quad pf' + qf \equiv 1;$$

let

$$(6) \quad \begin{aligned} \phi_1 &= x - p_1 f, & \phi_r &= \phi_{r-1} + p_r (-f)^{r/r!}, \\ p_1 &= p, & p_r &= p p'_{r-1} - (r-1) q p_{r-1}. \end{aligned}$$

Then $\phi_r(x)$ is clearly a polynomial, and it defines an iteration of order $r+1$ to any root ξ of the equation (1).

While a direct induction is possible, it is simpler to proceed otherwise. Schröder [3] (see also [1]) has shown that

$$(7) \quad \begin{aligned} \phi_r &= x + \sum_1^r (-F)^r \delta^{r-1} F / \nu!, \\ \delta^0 F &= 1/F', & \delta^{r+1} F &= (\delta^r F)' / F', \end{aligned}$$

Presented to the Society, November 25, 1950; received by the editors October 23, 1950.

¹ Numbers in brackets refer to the bibliography at the end of the paper.

defines an iteration of order $r+1$ to the roots of

$$F = 0.$$

We ask, then, whether a function $g(x)$ can be found such that for

$$(8) \quad F = gf$$

the functions ϕ_r defined by (7) are polynomials. For $r=1$ we require that

$$(9) \quad F/F' = pf$$

for some polynomial p . To achieve this it is sufficient to choose p and f to satisfy (5) while

$$(10) \quad g'/g = q/p.$$

Suppose, now, it has been verified that

$$F^r \delta^{r-1} F = p_r f^r,$$

or, by (8), that

$$\delta^{r-1} F = p_r g^{-r}.$$

Then

$$\delta^r F = (p'_r g^{-r} - r p_r g' g^{-r-1})/F',$$

and after a few manipulations

$$F^{r+1} \delta^r F = f^{r+1} (p p'_r - r q p_r).$$

Hence

$$p_{r+1} = p p'_r - r q p_r$$

and the induction is complete.

BIBLIOGRAPHY

1. D. R. Blaskett and H. Schwerdtfeger, *A formula for the solution of an arbitrary analytic equation*, Quarterly of Applied Mathematics vol. 3 (1945) pp. 266-268.
2. C. Domb, *On iterative solutions of algebraic equations*, Proc. Cambridge Philos. Soc. vol. 45 (1949) pp. 237-240.
3. Ernst Schröder, *Über unendliche viele Algorithmen zur Auflösung der Gleichungen*, Math. Ann. vol. 2 (1870) pp. 317-365.

OAK RIDGE NATIONAL LABORATORY