ON THE DISTRIBUTION OF THE ROOTS OF A POLYNOMIAL WITH INTEGRAL COEFFICIENTS

ROBERT BREUSCH

The following question has been raised by D. H. Lehmer¹ in connection with prime number problems: "If ϵ is a positive quantity, to find a polynomial of the form $f(z) \equiv z^r + a_1 z^{r-1} + \cdots + a_r$, where the a 's are integers, such that the absolute value of the product of those roots of f which lie outside the unit circle, lies between 1 and $1 + \epsilon$." Lehmer calls this absolute value $\Omega(f)$. Since, for $f(z) = z^2 - z - 1$, $\Omega(f) = 1.6 \cdots$, we may assume $\epsilon < 1$, and therefore $a_r = \pm 1$. Then $\Omega(f)$ may be written in the form $\prod_{n=1}^r |z_n|^{[|z_n|]}$, where the z_n are the roots of $f(z)$. It is shown in this paper that no nonreciprocal polynomial $f(z)$ exists with $\Omega(f) < 1.179$. Lehmer himself gives an example of a reciprocal polynomial of degree 10 for which $\Omega(f)$ is less than this, namely $1.176 \cdots$; no nonreciprocal polynomial seems to be known for which $\Omega(f) < 1.32$. Thus the present result is still far from being a complete answer to the question.

Let $f(z) \equiv z^r + a_1 z^{r-1} + \cdots \pm 1$ be a nonreciprocal polynomial with integral coefficients, irreducible in the rational field. The proof makes use of the resultant $R(f, g)$ of $f(z)$ and the polynomial $g(z) \equiv \pm z^r f(1/z) = z^r + \cdots$. All the roots of $g(z)$ are of the form $z_n^* \equiv \bar{z}_n^{-1}$ where the $\bar{z}_n$ are the conjugates of the roots z_n of $f(z)$. Since $f(z)$ and $g(z)$ are irreducible and different, have integral coefficients and highest coefficients 1, $|R(f, g)| \equiv \prod_{n=1}^r \prod_{m=1}^r |z_n - z_m^*|$ must be not less than 1.

$$|R(f, g)| \equiv \prod_{n=1}^r \prod_{m>n} |z_n - z_m^*| \cdot \prod_{m=1}^r |z_m - z_n^*| \equiv P_1 \cdot P_2.$$

If $\Omega(f) = k$, it will be shown in Lemma 2 that $P_1 < k^{2r} \cdot r^r$, and in Lemma 3 that $P_2 \leq (4(k-1)/r)^r$.

LEMMA 1. $P_1 \equiv \prod_{n=1}^r \prod_{m>n} |z_n - z_m^*| \cdot |z_m - z_n^*|$ has the greatest value compatible with the two conditions (I) $\prod_{n=1}^r |z_n| = 1$, (II) $\prod_{n=1}^r |z_n|^{[|z_n|]} = k$ ($1 < k < 2$) for $|z_1| = k$, $|z_2| = \cdots = |z_{r-1}| = 1$, $|z_r| = 1/k$.

PROOF. For $n = 1, 2, \cdots, r$, call b_n values of the z_n for which P_1 takes its greatest value when restricted by conditions I and II.

Presented to the Society, April 28, 1951; received by the editors January 2, 1951.

¹ D. H. Lehmer, *Factorization of certain cyclotomic functions*, Ann. of Math. vol. 34 (1933) p. 476.

Assume that at least two of the b 's are absolutely greater than 1, for example, $|b_1| > 1$, $|b_2| > 1$. P_1 may be rewritten in the form

$$P_1 \equiv \left| (z_1 - \bar{z}_2^{-1})(\bar{z}_2 - z_1^{-1}) \cdot \prod_{n=3}^r (z_1 - z_n^*)(\bar{z}_2 - z_n^{-1})(\bar{z}_2 - z_n^{-1})(z_n - \bar{z}_2^{-1}) \right. \\ \left. \cdot \prod_{n=3}^r \prod_{m>n} (z_n - z_m^*)(z_m - z_n^*) \right|.$$

The expression within the absolute bars is now an analytic function of z_1 and $\bar{z}_2$ (but not of the remaining z 's). Calling this expression $F(z_1, \bar{z}_2, z_3, \dots, z_r)$, and introducing $A \equiv b_1 \bar{b}_2$ (with $1 < |A| \leq k$), our assumption implies that $F(z, A/z, b_3, \dots, b_r)$ which is analytic in $1 \leq |z| \leq |A|$ takes its largest value in this region for $z = b_1$. But this is impossible, for b_1 is by assumption an interior point of this region, because $1 < |b_1| < |A|$.

Interchanging the z_n with the z_n^* does not change P_1 ; thus not two of the $|b_n^*|$ can be greater than 1, that is, not two of the $|b_n|$ can be less than 1.

Therefore (by proper arrangement of the subscripts) P_1 takes its greatest value for $z_1 = ke^{i\theta_1}$, $z_1^* = (1/k)e^{i\theta_1}$, $z_r = (1/k)e^{i\theta_r}$, $z_r^* = ke^{i\theta_r}$, $z_n = e^{i\theta_n}$, $z_n^* = z_n$ ($1 < n < r$).

LEMMA 2. $P_1 < k^{2r} \cdot r^r$.

PROOF. Using the values found in Lemma 1, we get, for $1 < n < r$,

$$\begin{aligned} |z_1 - z_n^*| \cdot |z_n - z_1^*| &= |ke^{i\theta_1} - e^{i\theta_n}| \cdot \frac{1}{k} \cdot |ke^{i\theta_n} - e^{i\theta_1}| \\ &= \frac{1}{k} |z_1 - z_n|^2 \end{aligned}$$

(since $|ke^{i\theta_n} - e^{i\theta_1}| = |e^{i\theta_n} - ke^{i\theta_1}|$). Similarly, for $1 < n < r$,

$$\begin{aligned} |z_r - z_n^*| \cdot |z_n - z_r^*| &= k \cdot |z_r - z_n|^2, \\ |z_1 - z_r^*| \cdot |z_r - z_1^*| &= |e^{i\theta_1} - e^{i\theta_r}|^2 < |z_1 - z_r|^2. \end{aligned}$$

Therefore

$$\begin{aligned} P_1 &< |z_1 - z_r|^2 \cdot \prod_{n=2}^{r-1} |z_1 - z_n|^2 \cdot |z_r - z_n|^2 \cdot \prod_{n=2}^{r-1} \prod_{n < m < r} |z_n - z_m|^2 \\ &= \prod_{n=1}^r \prod_{m>n} |z_n - z_m|^2. \end{aligned}$$

The function

$$G(z) \equiv \frac{1}{z^{2r}} \left(z - \frac{z_1 z_r}{z} \right)^2 \cdot \prod_{n=2}^{r-1} (z - z_n)^2 \cdot \left(\frac{z_1 z_r}{z} - z_n \right)^2 \\ \cdot \prod_{n=2}^{r-1} \prod_{n < m < r} |z_n - z_m|^2$$

is analytic for $1 \leq |z| \leq \infty$, and has therefore its greatest absolute value in this region for $|z| = 1$. But for $|z| = 1$, $|G(z)|$ is the absolute value of the discriminant of the polynomial whose r roots $z, z_1 z_r/z, z_2, \dots, z_{r-1}$, all lie on the unit circle. This is known² to be not greater than r^r . Therefore $|G(z)| < r^r$ for $|z| > 1$, and $P_1 < |z_1^{2r}| \cdot |G(z_1)| < k^{2r} \cdot r^r$.

LEMMA 3. *If the z_n satisfy conditions I and II of Lemma 1, then $P_2 \equiv \prod_{n=1}^r |z_n - z_n^*| \leq (4(k-1)/r)^r$.*

PROOF. If $|z_n| = 1 + \epsilon_n \geq 1$ for $n = 1, 2, \dots, s$, and $|z_n^*| = |z_n|^{-1} = 1 + \epsilon_n > 1$ for $n = s+1, \dots, r$, then $|z_n - z_n^*| = \left| |z_n| - |z_n|^{-1} \right| = 1 + \epsilon_n - 1/(1 + \epsilon_n) \leq 2\epsilon_n$ for $n = 1, 2, \dots, r$. Also, since $\prod_{n=1}^s (1 + \epsilon_n) = k$, and $\prod_{n=s+1}^r (1 + \epsilon_n) = k, 1 + \sum_{n=1}^s \epsilon_n \leq k, 1 + \sum_{n=s+1}^r \epsilon_n \leq k$, and $\sum_{n=1}^r \epsilon_n \leq 2(k-1)$. Therefore $P_2 \leq \prod_{n=1}^r (2\epsilon_n) \leq (4(k-1)/r)^r$.

THEOREM. *If $f(z)$ is a nonreciprocal irreducible polynomial with integral coefficient and highest coefficient 1, then $\Omega(f) = k > 1.179$.*

PROOF. If r is the degree of $f(z)$, the statement is obviously correct for $|a_r| > 1$, since $k \geq |a_r|$. Thus we may assume $|a_r| = 1$. Then by Lemmas 2 and 3, $|R(f, g)| \equiv P_1 P_2 < k^{2r} r^r (4(k-1)/r)^r = (4k^2(k-1))^r$. This must be not less than 1; therefore $4k^2(k-1) \geq 1, k > 1.179$.

AMHERST COLLEGE

² I. Schur, *Ueber die Verteilung der Wurzeln bei gewissen algebraischen Gleichungen mit ganzzahligen Koeffizienten*, Math. Zeit. vol. 1 (1918) p. 385.