PATTERN INTEGRATION WITH IMPROPER RIEMANN INTEGRALS

R. E. CARR

1. Introduction. In a previous paper [1]¹ we have recently introduced and discussed a type of integration, which we have called pattern integration, and have established the following theorem:

THEOREM 1.1. *Let $f(x)$ be (proper) Riemann integrable, $0 \leq x \leq 1$. Let $(k-1)/n \leq \xi_k^{(n)} \leq k/n$ ($k=1, 2, 3, \dots, n$). Let the pattern P be characterized by the dyadic number*

$$t = 0.\alpha_1\alpha_2\alpha_3 \dots \alpha_n \dots (2),$$

such that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \alpha_k = \alpha.$$

Then the pattern integral

$$(P) \int_0^1 f(x) dx \equiv \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \alpha_k f(\xi_k^{(n)}) = \alpha (R) \int_0^1 f(x) dx.$$

It seems desirable to extend this result, if possible, to improper Riemann integrals. In this paper we shall confine our remarks to improper integrals of the form

$$\oint_0^1 f(x) dx \equiv \lim_{\epsilon \rightarrow 0+} (R) \int_{\epsilon}^1 f(x) dx.$$

(R) in front of any integral will indicate a proper Riemann integral.

2. The improper integral defined as the limit of a sum. We first consider the case where $\alpha_k = 1$ ($k=1, 2, 3, \dots$). A fundamental difficulty not encountered in the previous paper is apparent from the definition of $\oint_0^1 f(x) dx$. Bromwich and Hardy have considered this problem [2]. As they point out, the ordinary definition of this improper integral is by means of a double (repeated) limit; and in order to replace this limit by the single limit,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n f(\xi_k^{(n)}),$$

Received by the editors December 11, 1950.

¹ Numbers in brackets refer to the bibliography at the end of the paper.

some restrictions are certainly necessary. In the first place, if $0 < \xi_1^{(n)} \leq 1/n$, we can clearly choose $\xi_1^{(n)}$ so that $|(1/n)f(\xi_1^{(n)})|$ is greater than any assigned number. It is then suggested that the most obvious choice is $\xi_k^{(n)} = k/n$, and with this restriction they establish:

THEOREM 2.1. *Let $f(x)$ be positive and tend steadily to ∞ as x tends to zero. Let $\oint_0^1 f(x)dx$ be convergent. Then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n f\left(\frac{k}{n}\right) = \oint_0^1 f(x)dx.$$

The proof rests on the fact that for all n ,

$$(2.1) \quad \frac{1}{n} \sum_{k=2}^n f\left(\frac{k}{n}\right) \leq (R) \int_{1/n}^1 f(x)dx \leq \frac{1}{n} \sum_{k=1}^{n-1} f\left(\frac{k}{n}\right),$$

while the difference between the two sums in (2.1) is

$$(2.2) \quad \frac{1}{n} \left\{ f\left(\frac{1}{n}\right) - f(1) \right\}.$$

That the limit of (2.2), as $n \rightarrow \infty$, is zero, follows from the well known result that if $f(x)$ steadily increases as x decreases and $\oint_0^1 f(x)dx$ is convergent, then $\lim_{x \rightarrow 0+} xf(x) = 0$. From this the conclusion is immediate.

It appears that the following extension of the above theorem is new, although the theorem is quite elementary.

THEOREM 2.2. *Let $f(x)$ be positive and tend steadily to ∞ as x tends to zero. Let $\oint_0^1 f(x)dx$ be convergent. Let $(1/Mn) \leq \xi_1^{(n)} \leq (1/n)$ ($M \geq 1$ and fixed), and $(k-1)/n \leq \xi_k^{(n)} \leq (k/n)$ ($k = 2, 3, \dots, n$). Then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n f(\xi_k^{(n)}) = \oint_0^1 f(x)dx.$$

PROOF. Define

$$A_n \equiv \frac{1}{n} \sum_{k=1}^n f\left(\frac{k}{n}\right)$$

and

$$\begin{aligned} B_n &\equiv \frac{1}{n} \left\{ f\left(\frac{1}{Mn}\right) + \sum_{k=2}^n f\left(\frac{k-1}{n}\right) \right\} \\ &\equiv \frac{1}{n} \left\{ f\left(\frac{1}{Mn}\right) + \sum_{k=1}^n f\left(\frac{k}{n}\right) - f(1) \right\}. \end{aligned}$$

Then

$$A_n \leq \frac{1}{n} \sum_{k=1}^n f(\xi_k^{(n)}) \leq B_n.$$

But, clearly, $\lim_{n \rightarrow \infty} A_n = \lim_{n \rightarrow \infty} B_n = \oint_0^1 f(x) dx$, and our conclusion follows.

A sharper criterion, in order that the relation

$$(2.3) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n f\left(\frac{k}{n}\right) = \oint_0^1 f(x) dx$$

should hold whenever the improper integral on the right is convergent, has been given by Wintner [3] in the following result:

THEOREM 2.3. *If $f(x)$ is of bounded variation on every interval $\epsilon \leq x \leq 1$, where $\epsilon > 0$, and behaves, as $x \rightarrow 0$, so as to satisfy the restriction*

$$\int_{\epsilon}^1 |df(x)| = o(\epsilon^{-1}),$$

then the convergence of $\oint_0^1 f(x) dx$ implies that

$$(2.4) \quad \lim_{\epsilon \rightarrow 0^+} \epsilon \sum_{k\epsilon \leq 1} f(k\epsilon)$$

exists and equals $\oint_0^1 f(x) dx$.

The result (2.3) would follow from (2.4) by letting $\epsilon = 1/n$.

The following extension of Wintner's result appears to be new:

THEOREM 2.4. *Let $f(x)$ be of bounded variation on every interval $\epsilon \leq x \leq 1$, where $\epsilon > 0$, and let $f(x)$ behave, as $x \rightarrow 0^+$, so as to satisfy the restriction*

$$(2.5) \quad \int_{\epsilon}^1 |df(x)| = o(\epsilon^{-1}).$$

Let $\xi_1 = 1/Mn$ ($M \geq 1$ and fixed), and $(k-1)/n \leq \xi_k^{(n)} \leq (k/n)$ ($k = 2, 3, \dots, n$). Let $\oint_0^1 f(x) dx$ exist.

Then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n f(\xi_k^{(n)}) = \oint_0^1 f(x) dx.$$

PROOF. From the hypothesis it follows that

$$(2.6) \quad \lim_{x \rightarrow 0} xf(x) = 0.$$

For,

$$\epsilon |f(1) - f(\epsilon)| = \epsilon \left| \int_{\epsilon}^1 df(x) \right| \leq \epsilon \int_{\epsilon}^1 |df(x)| \rightarrow 0.$$

From Wintner's result, we know

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n f\left(\frac{k}{n}\right) = \oint_0^1 f(x) dx.$$

Consider

$$\left\{ \frac{1}{n} \sum_{k=1}^n f(\xi_k^{(n)}) - \frac{1}{n} \sum_{k=1}^n f\left(\frac{k}{n}\right) \right\}.$$

For all n ,

$$\begin{aligned} \left| \sum_{k=1}^n f(\xi_k^{(n)}) - \sum_{k=1}^n f\left(\frac{k}{n}\right) \right| &\leq \sum_{k=1}^n \left| f(\xi_k^{(n)}) - f\left(\frac{k}{n}\right) \right| \\ &< \int_{(1/n)}^1 |df(x)| + \left| f\left(\frac{1}{Mn}\right) \right| + \left| f\left(\frac{1}{n}\right) \right|. \end{aligned}$$

Hence, using (2.5) and (2.6),

$$\lim_{n \rightarrow \infty} \left| \frac{1}{n} \sum_{k=1}^n f(\xi_k^{(n)}) - \frac{1}{n} \sum_{k=1}^n f\left(\frac{k}{n}\right) \right| = 0,$$

and our conclusion follows.

It should be noted that a number of theorems are available in the "converse" direction, that is, in which the convergence of the improper integral $\oint_0^1 f(x) dx$ follows from the knowledge of the existence of $\lim_{\epsilon \rightarrow 0+} \epsilon \sum_{n\epsilon \leq 1} f(n\epsilon)$. In this paper, however, we are not concerned with such results.

3. A pattern integral theorem. To hope for a complete extension of Theorem 1.1 to improper integrals of the form $\oint_0^1 f(x) dx$ is too much. Counter examples are not difficult to construct. For example, see §5 of the paper by Bromwich and Hardy [2].

The following theorem partially extends the earlier results.

THEOREM 3.1. *Let $f(x)$ be positive and tend steadily to ∞ as x tends to zero. Let $\oint_0^1 f(x) dx$ be convergent. Let $(1/Mn) \leq \xi_1^{(n)} \leq (1/n)$ ($M \geq 1$ and fixed), and $(k-1)/n \leq \xi_k^{(n)} \leq k/n$ ($k = 2, 3, \dots, n$). Let the pattern*

P be characterized by the dyadic number

$$t = 0.\alpha_1\alpha_2\alpha_3 \cdots \alpha_n \cdots (2),$$

such that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \alpha_k = \alpha.$$

Then

$$(P) \oint_0^1 f(x) dx \equiv \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \alpha_k f(\xi_k^{(n)}) = \alpha \oint_0^1 f(x) dx.$$

PROOF. Define $f_N(x) = f(1/MN)$, $0 \leq x < 1/N$, $f_N(x) = f(x)$, $1/N \leq x \leq 1$. For all values of $N > 0$, $(R) \int_0^1 f_N(x) dx$ exists and

$$(3.1) \quad (P) \int_0^1 f_N(x) dx = \alpha(R) \int_0^1 f_N(x) dx.$$

Consider the expression

$$(3.2) \quad \frac{1}{n} \sum_{k=1}^n \alpha_k f(\xi_k^{(n)}).$$

Let the subinterval in which $1/N$ occurs be the $\lambda(n)$ th. (Then $\lim_{n \rightarrow \infty} \lambda(n)/n = 1/N$.) Hence (3.2) can be written as $I_n + II_n - III_n + IV_n$, where

$$I_n = \frac{1}{n} \alpha_{\lambda(n)} f(\xi_{\lambda(n)}^{(n)}), \quad II_n = \frac{1}{n} \sum_{k=1}^n \alpha_k f_N(\xi_k^{(n)}),$$

$$III_n = \frac{1}{n} \sum_{k=1}^{\lambda(n)} \alpha_k f_N(\xi_k^{(n)}), \quad IV_n = \frac{1}{n} \sum_{k=\lambda(n)}^{\lambda(n)-1} \alpha_k f(\xi_k^{(n)}).$$

Let us consider the behavior of I_n , II_n , III_n , and IV_n , as $n \rightarrow \infty$. Clearly $f(\xi_{\lambda(n)}^{(n)}) \leq f(1/Mn)$, and since $\lim_{x \rightarrow 0+} xf(x) = 0$, $\lim_{n \rightarrow \infty} I_n = 0$. From (3.1), we see that

$$\lim_{n \rightarrow \infty} II_n = \alpha(R) \int_0^1 f_N(x) dx = \alpha \left\{ \frac{1}{N} f\left(\frac{1}{MN}\right) + (R) \int_{1/N}^1 f(x) dx \right\}.$$

Now

$$III_n = f\left(\frac{1}{MN}\right) \frac{\lambda(n) - 1}{n} \frac{1}{\lambda(n) - 1} \sum_{k=1}^{\lambda(n)-1} \alpha_k + \frac{1}{n} \alpha_{\lambda(n)} f_N(\xi_{\lambda(n)}^{(n)}).$$

Hence,

$$\lim_{n \rightarrow \infty} III_n = \alpha \frac{1}{N} f\left(\frac{1}{MN}\right).$$

We have

$$IV_n \leq \frac{1}{n} \sum_{k=1}^{\lambda(n)-1} f(\xi_k^{(n)}).$$

Hence

$$0 \leq IV_n \leq \oint_0^{1/N} f(x) dx + \frac{1}{n} f\left(\frac{1}{Mn}\right)$$

for all n . We then see that, as $n \rightarrow \infty$, the expression (3.2) at worst oscillates between

$$(3.3) \quad \alpha(R) \int_{1/N}^1 f(x) dx$$

and

$$(3.4) \quad \alpha(R) \int_{1/N}^1 f(x) dx + \oint_0^{1/N} f(x) dx.$$

But this is true for all $N > 0$, so letting $N \rightarrow \infty$, we see from (3.3) and (3.4) that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \alpha_k f(\xi_k^{(n)}) = \alpha(R) \oint_0^1 f(x) dx,$$

which is the desired result.

4. Miscellaneous results. The following is an extension of Theorem 3.1.

THEOREM 4.1. *Let $\oint_0^1 f(x) dx$ be convergent. Let there be a $g(x)$ with the following properties:*

1. *In the interval $0 < x \leq r$, where $0 < r \leq 1$, $g(x)$ is positive and tends steadily to ∞ as x tends to zero.*

2. *$\oint_0^r g(x) dx$ is convergent.*

Let $|f(x)| \leq g(x)$, $0 < x \leq r$. Let $1/Mn \leq \xi_1^{(n)} \leq 1/n$ ($M \geq 1$ and fixed), and $(k-1)/n \leq \xi_k^{(n)} \leq k/n$ ($k = 2, 3, \dots, n$). Let the pattern P be characterized by the dyadic number

$$t = 0.\alpha_1\alpha_2\alpha_3 \cdots \alpha_n \cdots (2),$$

such that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \alpha_k = \alpha.$$

Then,

$$(P) \oint_0^1 f(x) dx \equiv \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \alpha_k f(\xi_k^{(n)}) = \alpha \oint_0^1 f(x) dx.$$

Since the proof is similar to that of Theorem 3.1, it will be omitted. It is of interest to note that if one chooses

$$f(x) = \frac{1}{x} \cos \frac{1}{x},$$

then $\oint_0^1 f(x) dx$ converges, but

$$(4.1) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n f(\xi_k^{(n)})$$

fails to exist.

Define

$$A_n = \frac{1}{n} f\left(\frac{1}{Mn}\right) + \frac{1}{n} \sum_{k=2}^n f\left(\frac{k-1}{n}\right) \quad (M \geq 1 \text{ and fixed}),$$

$$B_n = \frac{1}{n} \sum_{k=1}^n f\left(\frac{k}{n}\right).$$

Then $A_n - B_n = (1/n) \{f(1/Mn) - f(1)\}$. If (4.1) exists,

$$\lim_{n \rightarrow \infty} M \frac{1}{Mn} f\left(\frac{1}{Mn}\right) = 0,$$

which is certainly not the case when $f(x) = (1/x) \cos (1/x)$.

BIBLIOGRAPHY

1. R. E. Carr and J. D. Hill, *Pattern integration*, Proceedings of the American Mathematical Society vol. 2 (1951) pp. 242-245.
2. T. J. I'a Bromwich and G. H. Hardy, *The definition of an infinite integral as the limit of a finite or infinite series*, The Quarterly Journal of Pure and Applied Mathematics vol. 39 (1907-1908) pp. 222-240.
3. A. Wintner, *The sum formula of Euler-Maclaurin and the inversions of Fourier and Möbius*, Amer. J. Math. vol. 69 (1947) pp. 685-708.

MICHIGAN STATE COLLEGE