MOMENTS WHICH ARE INTEGERS

R. M. REDHEFFER

Let $\alpha(t)$ be of bounded variation and have infinitely many points of increase on $(0, a)$. If $\int_0^a t^n d\alpha(t)$ is an integer for $n=0, 1, 2, \dots$, then $a \geq 4$.

To prove this let I_n be the integral in the theorem, so that

$$(1) \quad f(z) = \sum I_n z^n = \int_0^a \frac{d\alpha(t)}{1 - zt}.$$

The right side of (1) shows that $f(z)$ is regular except for $1/a \leq z < \infty$ and this region has image-radius greater than 1 if $a < 4$. The substitution $z=1/s$ turns (1) into a Hilbert transform (as was pointed out to the author by G. M. Wing) and the complex inversion formula shows then that $f(z)$ can be rational only when the integral reduces to a finite sum. The result now follows from the Pólya-Carlson theorem. Letting $t=4 \sin^2 \theta$ in

$$(2) \quad \pi I_n = \int_0^1 t^{n+1/2} (4-t)^{-1/2} dt$$

shows that *the condition $a \geq 4$ cannot be replaced by $a > 4$.*

If $q(n)$ is a polynomial in n , the operator (zd/dz) applied to $(1-z)^{-1}$ shows that $\sum q(n)z^n$ has, as sole singularity, a pole at $z=1$. Hence *the condition $I_n = \text{integer}$ can be replaced by $I_n = p_n/q(n)$ where p_n are integers and $q(n)$ is a polynomial with integral coefficients.* Here we have the beginnings of a way to show that certain definite integrals (satisfying suitable recursion relations) are irrational. The discussion can be extended to $\int_0^a f_n(t) d\alpha(t)$, incidentally, where f_n has a known generating function $g(z, t) = \sum f_n(t) z^n$.

In case the limits of integration are (a, b) , the conclusion of the theorem is $b-a \geq 4$, as was pointed out by R. Steinberg. For integral a an example can be constructed after the manner of (2).

Open questions are: Is $b-a \geq 4$ the optimum result for nonintegral a ? How far can proofs of irrationality be carried by these methods? Is there a counter-example like (2) belonging to L^2 (or L^∞) on $(0, 4)$? Is the counter-example essentially unique? Must it always be simply related to the mapping function (as (2) is)?

UNIVERSITY OF CALIFORNIA, LOS ANGELES

Received by the editors May 11, 1951.