MEAN CONVERGENCE OF ORTHOGONAL SERIES

JEROME NEWMAN AND WALTER RUDIN

I. **Introduction.** Marcel Riesz's theorem [9, p. 153]¹ about the mean convergence of Fourier series in L^p spaces has been extended by Harry Pollard to series of classical orthogonal polynomials [2; 3; 4; 5]. In particular, it was shown that the Legendre expansion of a function $f \in L^p$ on $(-1, 1)$ converges in the mean of order p to f if $4/3 < p < 4$, but that this is false for any $p > 4$ (or $1 \leq p < 4/3$). The question of the status of the limiting values $p=4$ and $p=4/3$ was raised but not settled by Pollard, and is repeated in the reviews of [3] and [5].

Entirely similar results were established for Jacobi series (for the limiting values of p , see the statements of Theorems 2 and 3 below), and for series of Bessel functions (Theorem 4), where the problem was also left open for $p=4/3$ and $p=4$ [8, pp. 797 and 802].

The purpose of the present paper is to show by a very simple argument that in all the above cases the property of mean convergence fails to hold at the limiting values of p .

II. **A necessary condition.** Suppose the sequence $\{\phi_n(x)\}$ is orthonormal with respect to a measure function $\mu(x)$ on (a, b) , that is,

$$\int_a^b \phi_n(x) \phi_m(x) d\mu(x) = \delta_{mn}.$$

Let L_μ^p denote the space of functions f for which

$$\|f\|_{p, \mu} = \left\{ \int_a^b |f(x)|^p d\mu(x) \right\}^{1/p} < \infty \quad (p \geq 1),$$

and put

$$(1) \quad a_n(f) = \int_a^b f(x) \phi_n(x) d\mu(x).$$

Then if the series

$$(2) \quad \sum_{n=0}^{\infty} a_n(f) \phi_n(x)$$

Presented to the Society by Jerome Newman, November 25, 1950, under the title *Remark on some papers by H. Pollard*; received by the editors April 16, 1951 and, in revised form, July 18, 1951.

¹ Numbers in brackets refer to the bibliography at the end of the paper.

converges in L_μ^p for every $f \in L_\mu^p$, we must have

$$(3) \quad a_n(f) \|\phi_n\|_{p,\mu} = O(1)$$

for every $f \in L_\mu^p$. Hence the Banach-Steinhaus theorem [1, p. 80, Theorem 5] applied to the functionals (3) shows:

THEOREM 1. *If the series (2) converges in L_μ^p for every $f \in L_\mu^p$, then*

$$(4) \quad \|\phi_n\|_{p,\mu} \|\phi_n\|_{q,\mu} = O(1) \quad \left(\frac{1}{p} + \frac{1}{q} = 1 \right).$$

III. Jacobi series and Bessel series. Let $\alpha \geq -1/2$, $\beta \geq -1/2$, be fixed. The polynomials $P_n^{(\alpha,\beta)}(x)$ (we use Szegő's notation [6, p. 57]) are orthogonal on $(-1, 1)$ with respect to the weight function

$$w(x) = (1-x)^\alpha (1+x)^\beta.$$

If the sequence $\{c_n\}$ is determined by the conditions

$$\int_{-1}^1 [c_n P_n^{(\alpha,\beta)}(x)]^2 w(x) dx = 1 \quad (n = 0, 1, 2, \dots; c_n > 0),$$

we put

$$\Phi_n(x) = c_n P_n^{(\alpha,\beta)}(x),$$

$$\Psi_n(x) = c_n P_n^{(\alpha,\beta)}(x) [w(x)]^{1/2}.$$

It is known that c_n is of order $n^{1/2}$ [5, formula (4.3.3)].

We suppose that $\alpha \geq \beta$ (the results are completely analogous if $\alpha < \beta$), and put

$$r = \frac{4(\alpha+1)}{2\alpha+3}, \quad s = \frac{4(\alpha+1)}{2\alpha+1}.$$

THEOREM 2. *If $p=r$, or if $p=s$, there exists a function $f \in L_\mu^p$ on $(-1, 1)$ ($d\mu(x) = w(x)dx$), such that the series $\sum a_n(f) \Phi_n(x)$ does not converge in L_μ^p .*

Here

$$a_n(f) = \int_{-1}^1 f(x) \Phi_n(x) w(x) dx.$$

THEOREM 3. *If $p=4/3$, or if $p=4$, there exists a function $f \in L^p$ on $(-1, 1)$ such that the series $\sum a_n(f) \Psi_n(x)$ does not converge in L^p .*

Here

$$a_n(f) = \int_{-1}^1 f(x) \Psi_n(x) dx.$$

Next, let ξ be real, and $\nu \geq -1/2$. Let $0 < \lambda_1 < \lambda_2 < \dots$ be the positive real zeros of the function $J_\nu(x) \cos \xi + x J'_\nu(x) \sin \xi$. The functions $J_\nu(\lambda_n x)$ ($n=1, 2, 3, \dots$) are orthogonal on $(0, 1)$ with respect to the weight function x . If the sequence $\{b_n\}$ is determined by the conditions

$$\int_0^1 [b_n J_\nu(\lambda_n x)]^2 x dx = 1 \quad (n = 1, 2, 3, \dots; b_n > 0),$$

we put

$$B_n(x) = b_n J_\nu(\lambda_n x).$$

Let us note that the numbers λ_n are interlaced with the positive zeros of $J_\nu(x)$ [7, p. 480]. Since the latter are of order n [7, p. 506], so is λ_n . It follows that b_n is of order $n^{1/2}$ [7, p. 577].

THEOREM 4. *If $p=4/3$, or if $p=4$, there exists a function $f \in L_\mu^p$ on $(0, 1)$ ($d\mu(x) = x dx$) such that the series $\sum a_n(f) B_n(x)$ does not converge in L_μ^p .*

PROOFS. For Theorems 2 and 3 we use the formula [5, (8.21.18)]

$$(5) \quad P_n^{(\alpha, \beta)}(\cos \theta) = n^{-1/2} k(\theta) \{ \cos(N\theta + \gamma) + (n \sin \theta)^{-1} O(1) \},$$

where $N = n + (\alpha + \beta + 1)/2$, $\gamma = -(2\alpha + 1)\pi/4$, and $k(\theta) = \pi^{-1/2} \cdot (\sin(\theta/2))^{-\alpha-1/2} (\cos(\theta/2))^{-\beta-1/2}$.

Putting $x = \cos \theta$, and applying (5) on the interval $[n^{-1}, \pi - n^{-1}]$ on which the error term is uniform, a simple calculation shows that

$$(6) \quad \left\{ \int_{-1}^1 |\Phi_n(x)|^s w(x) dx \right\}^{1/s} > A_1 (\log n)^{1/s},$$

$$(7) \quad \left\{ \int_{-1}^1 |\Phi_n(x)|^r w(x) dx \right\}^{1/r} > A_2,$$

$$(8) \quad \left\{ \int_{-1}^1 \Psi_n^4(x) dx \right\}^{1/4} > A_3 (\log n)^{1/4},$$

$$(9) \quad \left\{ \int_{-1}^1 |\Psi_n(x)|^{4/3} dx \right\}^{3/4} > A_4.$$

Here $A_1, A_2, \dots$ are positive constants. Theorems 2 and 3 fol-

low since (4) is violated in each case.

It should be noted that (6) was derived on the tacit assumption that $s < \infty$. If $s = \infty$, then $\alpha = -1/2$, $P_n^{(\alpha, \beta)}$ is a Tchebichef polynomial, and Theorem 2 reduces to a statement about trigonometric series which is well known [9, p. 155].

Finally, to prove Theorem 4, we put $y = \lambda_n x$ in the integrals involving $J_\nu(\lambda_n x)$, use the asymptotic formula [7, p. 199]

$$J_\nu(y) \sim \left(\frac{2}{\pi y}\right)^{1/2} \cos\left(y - \frac{\nu\pi}{2} - \frac{\pi}{4}\right) \quad (y \rightarrow \infty),$$

and find that

$$(10) \quad \left\{ \int_0^1 B_n^4(x) x dx \right\}^{1/4} > A_5 (\log n)^{1/4},$$

$$(11) \quad \left\{ \int_0^1 |B_n(x)|^{4/3} x dx \right\}^{3/4} > A_6.$$

Thus (4) is again violated.

BIBLIOGRAPHY

1. S. Banach, *Théorie des opérations linéaires*, Warsaw, 1932.
2. H. Pollard, *The mean convergence of series of orthogonal polynomials*, Proc. Nat. Acad. Sci. U.S.A. vol. 32 (1946) pp. 8-10.
3. ———, *The mean convergence of orthogonal series*. I, Trans. Amer. Math. Soc. vol. 62 (1947) pp. 387-403.
4. ———, *The mean convergence of orthogonal series*. II, Trans. Amer. Math. Soc. vol. 63 (1948) pp. 355-367.
5. ———, *The mean convergence of orthogonal series*. III, Duke Math. J. vol. 16 (1949) pp. 189-191.
6. G. Szegő, *Orthogonal polynomials*, Amer. Math. Soc. Colloquium Publications, vol. 23, 1939.
7. G. N. Watson, *A treatise on the theory of Bessel functions*, 2d ed., Cambridge, 1944.
8. G. M. Wing, *The mean convergence of orthogonal series*, Amer. J. Math. vol. 72 (1950) pp. 792-808.
9. A. Zygmund, *Trigonometrical series*, Warsaw, 1935.

COLUMBIA UNIVERSITY AND
MASSACHUSETTS INSTITUTE OF TECHNOLOGY