A NOTE ON PARTITIONS IN $GF[q, x]$

L. CARLITZ

1. Introduction. In this note we consider problems of the following kind. For given $A \in GF[q, x]$, $A \neq 0$, let $p(A)$ denote the number of solutions U_i of

$$(1.1) \quad A = U_1 + U_2 + \cdots \quad (\deg A = \deg U_1 > \deg U_2 > \cdots).$$

Also let $p'(A)$ denote the number of solutions of (1.1) subject to the additional restriction that all the polynomials are *primary*. We may also consider various modifications of these problems; for example the degree of some or all the U_i may be specified. In particular let $r(A)$ denote the number of solutions of (1.1) such that

$$(1.2) \quad \deg U_i = m - i + 1 \quad (i = 1, 2, \dots, m + 1),$$

where $m = \deg A$, and let $r'(A)$ denote the corresponding number with all U_i primary. Again returning to (1.1) we may restrict the number of parts; thus let $p_k(A)$ denote the number of solutions of

$$(1.3) \quad A = U_1 + \cdots + U_k \quad (\deg U_1 > \deg U_2 > \cdots),$$

and let $p'_k(A)$ denote the corresponding number when all the polynomials are primary. Other such problems are readily devised.

We find that the numbers just defined can be evaluated rather simply; thus these partition problems are considerably simpler than the corresponding problems involving integers (see for example [1, Chap. 19]). We prove the following results ($\deg A = m$):

$$(1.4) \quad p(A) = \prod_{i=0}^{m-1} (1 + (q - 1)q^i),$$

$$(1.5) \quad p'(A) = \prod_{i=0}^{m-1} (1 + q^i),$$

$$(1.6) \quad r(A) = (q - 1)^m q^{m(m-1)/2},$$

$$(1.7) \quad r'(A) = q^{m(m-1)/2},$$

$$(1.8) \quad p_k(A) = q^{(k-1)(k-2)/2} (q - 1)^{k-1} \begin{bmatrix} m \\ k - 1 \end{bmatrix} \quad (k \geq 1),$$

$$(1.9) \quad p'_k(A) = q^{(k-1)(k-2)/2} \begin{bmatrix} m \\ k - 1 \end{bmatrix} \quad (k \geq 1),$$

Presented to the Society, February 28, 1953; received by the editors October 27, 1952.

where in (1.8) and (1.9)

$$(1.10) \quad \begin{bmatrix} m \\ k \end{bmatrix} = \frac{(q^m - 1) \cdots (q - 1)}{(q - 1) \cdots (q^k - 1)} = \begin{bmatrix} m \\ m - k \end{bmatrix}, \quad \begin{bmatrix} m \\ 0 \end{bmatrix} = 1.$$

Note that in each case the number of partitions depends only on the degree of A . It will be recalled that, if q is replaced by an indeterminate, then

$$\begin{bmatrix} m \\ k \end{bmatrix}$$

is the generating function for partitions into at most k parts, each part being $\leq m - k$ [2, p. 5]. We also note that (1.9) is equivalent to

$$(1 + x)(1 + qx) \cdots (1 + q^{m-1}x) = \sum_{k=0}^m p'_{k+1}(m)x^k.$$

Similar generating functions can be constructed for the remaining partition functions.

In §5 we consider a special quadratic partition problem similar to that concerning $r(A)$ and $r'(A)$.

2. Proof of (1.4) and (1.5). Put

$$(2.1) \quad \begin{aligned} A &= a_0x^m + a_1x^{m-1} + \cdots + a_m, \\ U_1 &= b_0x^m + b_1x^{m-1} + \cdots + b_m \end{aligned} \quad (a_i, b_i \in GF(q)).$$

Then $b_0 = a_0$ but U_1 is otherwise arbitrary. It is then clear that $p(A)$ depends only on the degree of A ; we may accordingly put $p(A) = p(m)$. Now if $b_1 \neq a_1$, $\deg(A - U_1) = m - 1$; otherwise $\deg(A - U_1) < m - 1$. Thus for $q - 1$ choices of b_1 and arbitrary choice of $b_2, \dots, b_m$, $\deg(A - U_1) = m - 1$. In the next place when $b_1 = a_1$, then for $q - 1$ choices of b_2 and arbitrary choice of $b_3, \dots, b_m$, $\deg(A - U_1) = m - 2$. It is clear that in this way we get the recursion

$$(2.2) \quad \begin{aligned} p(m) &= (q - 1)q^{m-1} \cdot p(m - 1) + \cdots \\ &+ (p - 1)q \cdot p(1) + (p - 1) \cdot p(0). \end{aligned}$$

Clearly (2.2) implies

$$p(m) = (q - 1)q^{m-1} \cdot p(m - 1) + p(m - 1),$$

or what is the same thing

$$(2.3) \quad p(m) = (1 + (q - 1)q^{m-1})p(m - 1) \quad (m \geq 1).$$

Since $p(0) = 1$, (1.4) follows immediately.

Next let $a_0 = b_0 = 1$. We require that all U_i be primary. Thus either $a_1 - b_1 = 1$, and $b_2, \dots, b_m$ are arbitrary, or $a_1 = b_1$. In the latter case either $a_2 - b_2 = 1$, $b_3, \dots, b_m$ arbitrary, or $a_2 = b_2$, and so on. Thus corresponding to (2.2) we get

$$(2.4) \quad p'(m) = q^{m-1} \cdot p'(m-1) + \dots + q \cdot p'(1) + p'(0),$$

since again it is clear that $p'(A) = p'(m)$. It follows from (2.4) that

$$(2.5) \quad p'(m) = (1 + q^{m-1})p'(m-1) \quad (m \geq 1).$$

Since $p'(0) = 1$, (1.5) follows at once.

3. Proof of (1.6) and (1.7). We seek U_i satisfying both (1.1) and (1.2). Using the notation (2.1), again $a_0 = b_0$; however it is now necessary that $a_1 \neq b_1$. Thus there are $q-1$ choices of b_1 while $b_2, \dots, b_m$ are arbitrary. Consequently

$$(3.1) \quad r(A) = r(m) = (q-1)q^{m-1} \cdot r(m-1) \quad (m \geq 1).$$

Since $r(0) = 1$, we see that (3.1) implies

$$r(m) = (q-1)^m q^{m(m-1)/2},$$

which is the same as (1.6).

As for $r'(A)$ it is clear that in place of (3.1) we get

$$(3.2) \quad r'(A) = r'(m) = q^{m-1} \cdot r'(m-1) \quad (m \geq 1);$$

(1.7) is an immediate consequence of (3.2).

The definition of $r(A)$ and $r'(A)$ can be modified in various ways. For example, let $e_1, \dots, e_k$ be fixed integers such that $m = e_1 > e_2 > \dots > e_k \geq 0$. Then it is easily proved that the number of solutions of

$$(3.3) \quad A = U_1 + \dots + U_k \quad (\deg U_i = e_i)$$

is furnished by

$$(3.4) \quad (q-1)^k q^{e_1 + \dots + e_k},$$

while if A is primary, then the number of primary solutions of (3.3) is

$$(3.5) \quad q^{e_1 + \dots + e_k}.$$

For $e_i = m - i + 1$, (3.4) and (3.5) reduce to (1.6) and (1.7) respectively.

4. Proof of (1.8) and (1.9). It is convenient to consider

$$(4.1) \quad \begin{aligned} w_k(A) &= p_k(A) + \dots + p_1(A), \\ w'_k(A) &= p'_k(A) + \dots + p'_1(A), \end{aligned}$$

so that $w_k(A)$, $w'_k(A)$ enumerate the number of partitions of the respective kinds into at most k parts. Then as before we have

$$(4.2) \quad w_k(m) - w_k(m-1) = (q-1)q^{m-1} \cdot w_{k-1}(m-1) \quad (k \geq 2).$$

If we introduce the generating function

$$(4.3) \quad f_k(x) = \sum_{m=0}^{\infty} w_k(m)x^m,$$

we find that (4.2) implies

$$(4.4) \quad (1-x)f_k(x) = 1 + (q-1)xf_{k-1}(qx) \quad (k \geq 2).$$

Since $f_1(x) = (1-x)^{-1}$, (4.4) yields

$$(4.5) \quad f_k(x) = \sum_{i=0}^{k-1} \frac{(q-1)^i q^{i(i-1)/2} x^i}{(1-x)(1-qx) \cdots (1-q^i x)}.$$

Using the familiar formula

$$\frac{1}{(1-x)(1-qx) \cdots (1-q^{k-1}x)} = \sum_{m=0}^{\infty} \begin{bmatrix} m+k-1 \\ k-1 \end{bmatrix} x^m,$$

where

$$\begin{bmatrix} m \\ k \end{bmatrix}$$

is defined by (1.10), (4.5) becomes

$$f_k(x) = \sum_{i=0}^{k-1} (q-1)^i q^{i(i-1)/2} x^i \sum_{m=0}^{\infty} \begin{bmatrix} m+i \\ i \end{bmatrix} x^m.$$

Comparison with (4.3) gives

$$(4.6) \quad w_k(m) = \sum_{i=0}^{k-1} (q-1)^i q^{i(i-1)/2} \begin{bmatrix} m \\ i \end{bmatrix}.$$

Consequently by (4.1)

$$p_k(m) = w_k(m) - w_{k-1}(m) = (q-1)^{k-1} q^{(k-1)(k-2)/2} \begin{bmatrix} m \\ k-1 \end{bmatrix},$$

which proves (1.8).

The proof of (1.9) is quite similar. In place of (4.2) we now have

$$w'_k(m) - w'_k(m-1) = q^{m-1} \cdot w_{k-1}(m-1) \quad (k \geq 2).$$

Then if we put

$$f'_k(x) = \sum_{m=0}^{\infty} w'_k(m) x^m,$$

we find that

$$(1-x)f'_k(x) = 1 + xf_{k-1}(qx) \quad (k \geq 2),$$

whence

$$\begin{aligned} f'_k(x) &= \sum_{i=0}^{k-1} \frac{q^{i(i-1)/2} x^i}{(1-x)(1-qx) \cdots (1-q^i x)} \\ &= \sum_{i=0}^{k-1} q^{i(i-1)/2} x^i \sum_{m=0}^{\infty} \begin{bmatrix} m+i \\ i \end{bmatrix} x^m, \end{aligned}$$

so that

$$(4.7) \quad w'_k(m) = \sum_{i=0}^{k-1} q^{i(i-1)/2} \begin{bmatrix} m \\ i \end{bmatrix},$$

and therefore

$$p'_k(m) = w'_k(m) - w'_{k-1}(m) = q^{(k-1)(k-2)/2} \begin{bmatrix} m \\ k-1 \end{bmatrix}.$$

Note that by definition $w_{m+1}(m) = p(m)$, $w'_{m+1}(m) = p'(m)$, and it is easy to verify that (4.6) and (4.7) reduce to (1.4) and (1.5) for $k = m+1$. Also by definition $p_{m+1}(m) = r(m)$, $p'_{m+1}(m) = r'(m)$, and it is evident that (1.8) and (1.9) reduce to (1.6) and (1.7) for $k = m+1$.

5. A quadratic problem (q odd). We now briefly consider the following special quadratic partition problem. Let $\deg A = 2m$ and for simplicity take A primary. We seek the number of solutions in primary U_i of

$$(5.1) \quad A = U_1^2 + \cdots + U_m^2 + c \quad (\deg U_i = m - i + 1),$$

where c is some number $\in GF(q)$. Now if we put

$$A = x^{2m} + a_1 x^{2m-1} + \cdots + a_{2m}, \quad U_1 = x^m + b_1 x^{m-1} + \cdots + b_m x,$$

then $A - U_1^2$ will be primary of degree $2m-2$ provided $2b_1 = a_1$, $b_1^2 + 2b_2 + 1 = a_2$. Thus b_1, b_2 are uniquely determined while $b_3, \dots, b_m$ are arbitrary. If then we denote the number of solutions of (5.1) by $s'(m) = s(A)$, it follows that

$$(5.2) \quad s'(m) = q^{m-2} \cdot s'(m-1) \quad (m \geq 2).$$

Since $s'(1) = 1$, we see that (5.2) implies

$$(5.3) \quad s'(m) = q^{(m-1)(m-2)/2} \quad (m \geq 1).$$

Similarly if $s(m)$ denotes the number of solutions of (5.1) in which the U_i are no longer required to be primary we find that

$$s(m) = (q - 1)^{[m/2]} q^{(m-1)(m-2)/2},$$

where $[m/2]$ is the greatest integer $\leq m/2$.

In the same way we can consider special partitions of higher degree.

REFERENCES

1. G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, Oxford, 1938.
2. P. A. MacMahon, *Combinatory analysis*, vol. 2, Cambridge, 1916.

DUKE UNIVERSITY