A UNIFORM APPROXIMATION TO FOURIER COEFFICIENTS

M. L. JUNCOSA AND D. M. YOUNG

Let $f(x)$ be a real-valued function which is continuous in the interval $I: -1 \leq x \leq 1$ except at N points at which it may have a finite jump. The sequence of its ordinary complex Fourier coefficients is given by

$$a_n = \frac{1}{2} \int_{-1}^1 f(x) e^{-in\pi x} dx, \quad n = 0, \pm 1, \dots$$

It is the purpose of this note to show that the elements of the sequence of Riemann sum approximations to a_n ,

$$b_n(M) = \begin{cases} \frac{1}{2M} \sum_{j=-M+1}^M f\left(\frac{j}{M}\right) e^{-in\pi j/M}, & n = 0, \pm 1, \dots, \pm M, \\ 0, & |n| > M \end{cases}$$

tend to a_n uniformly as $M \rightarrow \infty$.

Let

$$s_m = \sum_{j=-m}^m a_j e^{ij\pi x}, \quad m = 0, 1, \dots,$$

and

$$\sigma_k(x) = \frac{1}{k+1} \sum_{m=0}^k s_m(x), \quad k = 0, 1, \dots$$

Then

$$\sigma_k(x) = \sum_{n=-k}^k A_{k,n} e^{in\pi x}$$

where

$$A_{k,n} = \begin{cases} (1 - |n|/k+1)a_n, & n = 0, \pm 1, \dots, \pm k, \\ 0, & |n| > k. \end{cases}$$

Let us cover the set of points of discontinuity of $f(x)$ in I by a set E which is the sum of a finite number of open intervals of lengths η_i , $i=1, 2, \dots, N$, each of which covers at most one point of discontinuity and such that $\sum_{i=1}^N \eta_i \equiv \eta < \epsilon/6F$, where $F = \text{LUB}|f(x)|$ in

Received by the editors June 16, 1952.

I. Then, on $I - IE$, by a generalization of Fejér's theorem on summability $(C, 1)$ of Fourier series (see, e.g., [1, p. 66]) $\sigma_k(x)$ converges uniformly to $f(x)$. Therefore, let M_1 be such that $|\sigma_k(x) - f(x)| < \epsilon/6$ uniformly on $I - IE$ for $k \geq M_1$. Henceforth, let $M \geq \max \{M_1, 3NF/\epsilon\}$. Since, in general, $|\sigma_k(x)| \leq F$ (see [1, p. 58]), on IE we have $|\sigma_k(x) - f(x)| \leq 2F$.

For $|n| \leq M$, using the above inequalities, we obtain

$$\begin{aligned} |b_n(M) - A_{M,n}| &= \frac{1}{2M} \left| \sum_{j=-M+1}^M \left[f\left(\frac{j}{M}\right) - \sigma_M\left(\frac{j}{M}\right) \right] e^{-in\pi j/M} \right| \\ &\leq \frac{1}{2M} \sum_{j=-M+1}^M \left| f\left(\frac{j}{M}\right) - \sigma_M\left(\frac{j}{M}\right) \right| \\ &\leq \frac{1}{2M} \left[\frac{2M\epsilon}{6} + \sum_{i=1}^N (M\eta_i + 1)2F \right] < \frac{2\epsilon}{3}. \end{aligned}$$

For $|n| > M$, $|b_n(M) - A_{M,n}| = 0$. For all n , we have

$$\begin{aligned} |A_{M,n} - a_n| &= \frac{1}{2} \left| \int_{-1}^1 [\sigma_M(x) - f(x)] e^{-in\pi x} dx \right| \\ &< \frac{1}{2} \left(\frac{\epsilon}{3} + 2\eta F \right) < \frac{\epsilon}{3}. \end{aligned}$$

Putting these results in the triangle inequality, we get $|b_n(M) - a_n| < \epsilon$ uniformly for all integral n .

REFERENCE

1. Dunham Jackson, *The theory of approximation*, Amer. Math. Soc. Colloquium Publications, vol. 11, 1930.

BALLISTIC RESEARCH LABORATORIES, ABERDEEN PROVING GROUND, AND
UNIVERSITY OF MARYLAND