THE ALGEBRA OF BOUNDED FUNCTIONS

KENNETH G. WOLFSON

1. **Introduction.** The Banach algebra $B(X)$ of all bounded complex-valued functions on a set X has been considered by several writers (cf., e.g. [10]). It is our purpose to give an ideal-theoretic characterization of this algebra. The main result (Theorem 3) states necessary and sufficient conditions that a B^* -algebra be equivalent to a $B(X)$ on an essentially unique set X .

Our method consists in establishing a lattice anti-isomorphism between the lattice of subsets of X and the lattice of all those ideals of $B(X)$ which are defined by annihilation.

These results are also applied to the study of certain p -rings.

2. By a B^* -algebra is meant a Banach algebra over the complex numbers such that, to each element x , there corresponds a unique element x^* (the adjoint of x), with the following properties: (i) $(x^*)^* = x$, (ii) $(xy)^* = y^*x^*$, (iii) if α, β are complex numbers, and $\bar{\alpha}, \bar{\beta}$ are their complex conjugates, then $(\alpha x + \beta y)^* = \bar{\alpha}x^* + \bar{\beta}y^*$, (iv) $\|x^*x\| = \|x\|^2$.

Let X be any set and denote by $B(X)$ the set of all bounded complex-valued functions defined on X . For $f(x) \in B(X)$ define $\|f(x)\| = \sup_{x \in X} |f(x)|$. Under the usual definitions of addition and multiplication and with the norm defined above, and $[f(x)]^* = \overline{f(x)}$, $B(X)$ is a commutative B^* -algebra with identity.

If G is any subset of an arbitrary commutative ring K , denote by $R(G)$ the set of elements k in K which satisfy $Gk = 0$. $R(G)$ is an ideal of K which we call an *annulet*. It is closed if K is a topological ring.

$E(X)$ will denote an arbitrary algebra of bounded functions on the set X , while the algebra of all functions which are zero at all but a finite set of points of X is denoted by $B_0(X)$.

If S is any subset of X , and G a subset of $E(X)$, let $A(S)$ be the set of all $f \in E(X)$ which satisfy $f(S) = 0$, and $N(G)$ the set of all $x \in X$ for which $G(x) = 0$. It is clear that $A(S)$ is a closed ideal of $E(X)$.

If S, T are any subsets of X , the logical sum and product are denoted by $S + T$ and $S \cap T$ respectively, while $X - S$ denotes the set of those elements in X which are not in S .

LEMMA 1. *If $E(X)$ is an algebra of bounded functions containing the*

Received by the editors June 10, 1953.

algebra $B_0(X)$, then

- (1) $R[A(S)] = A(X - S)$ for all $S \leq X$,
- (2) $N[A(S)] = S$ for all $S \leq X$,
- (3) $R(G) = A[X - N(G)]$ for every subset G of E .

PROOF. (1) $A(S)A(X - S) = 0$ implies that $A(X - S) \leq R[A(S)]$. If $S = X$, then $A(S) = 0$ so that $R[A(S)] = E$. If we accept the convention that all functions vanish on the empty set, we see that (1) holds for $S = X$. Assume $S < X$ and let $f \in R[A(S)]$ so that $A(S)f = 0$. Let $x \in X - S$. Then there exists $g \in E$ satisfying $g(S) = 0$ and $g(x) = 1$. Thus $g \in A(S)$ but since $gf = 0$ we must have $f(x) = 0$. That is, $f(X - S) = 0$ or $f \in A(X - S)$ proving (1).

(2) Since $A(S)[S] = 0$ we have $S \leq N[A(S)]$. To complete the proof we must show $N[A(S)] \leq S$. We shall assume $x \notin S$ and show $x \notin N[A(S)]$. Write $X = x + S + U$ and define f by

$$f(S + U) = 0 \quad \text{and} \quad f(x) = 1.$$

Such an $f \in E(X)$ by assumption. Then $f \in A(S)$. But $f(x) = 1$ implies $A(S)(x) \neq 0$ so that $x \notin N[A(S)]$ which shows (2).

(3) Let $f \in R(G)$ so that $Gf = 0$. Let $x \in [X - N(G)]$ so that $x \notin N(G)$. This means there exists a function $g \in G$ such that $g(x) \neq 0$. But since $g(x)f(x) = 0$ this implies $f(x) = 0$. Hence $f[X - N(G)] = 0$ and $f \in A[X - N(G)]$. The reverse implication follows by reversing the steps.

THEOREM 1. *Let $E(X)$ be an algebra of bounded functions which contains $B_0(X)$. Then the mapping $S \rightarrow A(S)$ is a duality (lattice anti-isomorphism) between the lattice of all subsets S of X and the lattice of all annulets of $E(X)$.*

PROOF. The mapping $S \rightarrow A(S)$ is one-one by (2) of Lemma 1. Clearly $S_1 \geq S_2$ implies $A(S_1) \leq A(S_2)$ and the mapping is a duality. Every annulet has the form $A(S)$ by (3) of Lemma 1, and every $A(S)$ is an annulet by (1) of Lemma 1.

THEOREM 2. *In $B(X)$ the sum of two annulets is an annulet.*

PROOF. Denote by (a) the ideal generated by the element a . An ideal of $B(X)$ is an annulet if and only if it is a principal ideal generated by an idempotent. For, if e is idempotent $(e) = R(1 - e)$, and if G is an annulet, $G = (e)$ for idempotent e by Theorem 2.3 of [3]. It is clear that if $G = A(S)$, the idempotent e is just the characteristic function of $X - S$. Now, let the given annulets be $A(S)$, $A(T)$ where $A(S) = (e)$, $A(T) = (f)$ for idempotents e and f which satisfy

$$\begin{aligned} e(S) &= 0, & f(T) &= 0, \\ e(X - S) &= 1, & f(X - T) &= 1. \end{aligned}$$

Since $A(S)$ and $A(T)$ are contained in $A(S \cap T)$ it follows that $A(S) + A(T) \leq A(S \cap T)$. It remains to show that $A(S \cap T) \leq A(S) + A(T)$. Let $A(S \cap T) = (g)$ where $g(S \cap T) = 0$, $g\{X - [S \cap T]\} = 1$. Write $X = [S \cap T] + [S - (S \cap T)] + [T - (S \cap T)] + [X - (S + T)]$, a decomposition of X into four mutually disjoint sets. Then $g = e + f - ef$ on these four parts separately and hence $g = e + f - ef$. Hence $g \leq A(S) + A(T)$ and $(g) \leq A(S) + A(T)$, which completes the proof.

THEOREM 3. *A commutative B^* -algebra K containing an identity is isomorphic (in a norm and $*$ preserving manner) to an algebra $B(X)$ of all bounded complex-valued functions on an essentially unique set X if and only if:*

- (1) *Every nonzero closed ideal of K contains a minimal ideal.*
- (2) *The sum of two annulets is an annulet.*

PROOF. The minimal ideals of K are of the form $e_i K$ for e_i a projection (self-adjoint idempotent) [2, Lemma 4.2]. Each $e_i K$ is a field and by the Gelfand-Mazur theorem is isomorphic and isometric to the complex field. Consider the mapping $x \rightarrow \{e_i x\}$ for $x \in K$. Since each e_i is of norm 1, if we define $*$ and norm in the obvious way we have a $*$ homomorphism of K into the algebra of all bounded sequences. Since $\prod_i R(e_i K)$ is an intersection of maximal and hence [1, p. 8] closed ideals, it follows from (1) and a theorem of McCoy [5, p. 873], that K is isomorphic to a special subdirect sum of the $e_i K$. That is, K is isomorphic to an algebra $E(X)$ of bounded functions defined on the set X of minimal ideals $e_i K$, where if $k \in K$, by $k(e_i K)$ is meant the complex number λ defined by $e_i k = \lambda e_i$. Moreover, to each $x \in X$ and complex number λ , there exists $f \in E(X)$ such that $f(x) = \lambda$, and $f(X - x) = 0$. Since $E(X)$ is a ring we have $E(X) \geq B_0(X)$.

We wish to show that $E(X)$ contains all characteristic functions of subsets of X . For any subset $S \leq X$, consider the annulets $A(S)$, $A(X - S)$. By Theorem 1,

$$\begin{aligned} A(S) \cup A(X - S) &= A(0) = E(X), \\ A(S) \cap A(X - S) &= A(X) = 0, \end{aligned}$$

where $\cup$ indicates the lattice join of annulets. But, by (2),

$$A(S) \cup A(X - S) = A(S) + A(X - S).$$

Hence $E(X) = A(S) \oplus A(X - S)$ in the sense of direct sum of ideals. By a theorem of von Neumann [7, p. 7] it follows, since $E(X)$ contains an identity, that there exists an idempotent $e \leq E(X)$ such that

$$A(S) = (1 - e)E, \quad A(X - S) = eE.$$

Since $e \leq A(X - S)$ this implies $e(X - S) = 0$. $1 - e \in A(S)$ implies $(1 - e)(S) = 0$ or $e(S) = 1$. Hence $E(X)$ contains all characteristic functions and is therefore clearly dense in $B(X)$. Applying Theorem 6.4 of [2] the proof of sufficiency is complete.

By Theorem 1, if K were isomorphic to $B(X_1)$ and $B(X_2)$ there would exist a lattice isomorphism of the lattice of all subsets of X_1 upon the lattice of all subsets of X_2 . In this sense the set X is unique.

The algebra $B(X)$ is certainly a B^* -algebra with identity, and satisfies (2) by Theorem 2. Every nonzero ideal of $B(X)$ (not only the closed ones) contains a minimal ideal, as in Theorem 16 of [5] mentioned previously. This completes the proof.

REMARK. The set X may also be considered to be the set of all maximal annulets of $B(X)$ (those maximal ideals which are annihilators of minimal ideals). Since $B(X)$ may be represented as the algebra $C(M)$ of all continuous functions on the compact space M of its maximal ideals, it is seen that M is the Stone-Čech compactification [9, p. 463] of the discrete space X of maximal annulets. An alternative proof of Theorem 3 could start with the representation $C(M)$.

3. The preceding results with slight modification apply to rings of functions from a discrete set X to a finite field.

In [6] it was shown that a ring of prime characteristic p , all of whose elements x satisfy $x^p = x$ (called a p -ring), is isomorphic to a subdirect sum of fields $GF(p)$. The ring $P(X)$ of all functions from a set X to $GF(p)$ (the complete direct sum of the fields) is generated by its idempotents, since an arbitrary function in $P(X)$ assumes only a finite number of distinct values and each is an integral multiple of the identity of the field $GF(p)$. Hence we have:

A p -ring K containing an identity element is isomorphic to a $P(X)$ if and only if:

- (1) *Every annulet of K contains a minimal ideal.*
- (2) *The sum of two annulets is an annulet.*

Specializing this to Boolean rings ($p = 2$) we get:

A Boolean ring K containing an identity is isomorphic to the ring of all subsets of a set S if and only if it satisfies conditions (1) and (2) above.

This result is similar to one obtained by Stone [8, p. 98, Theorem 62].

The complete direct sum of the fields $GF(p^n)$ is generated by its idempotents when considered as an algebra over $GF(p^n)$ (although not as a ring). Hence we have:

Let K be an algebra (containing an identity element) over the Galois field $GF(p^n)$, all of whose elements satisfy the equation $x^{p^n} = x$. Then K is isomorphic to the algebra $W(X)$ of all functions from a set X to the field $GF(p^n)$ if and only if K satisfies the conditions (1) and (2) (cf. [4, p. 379, Theorem 3]).

REFERENCES

1. I. Gelfand, *Normierte Ringe*, Mat. Sbornik N.S. vol. 9 (1941) pp. 3–24.
2. Irving Kaplansky, *Normed algebras*, Duke Math. J. vol. 16 (1949) pp. 399–418.
3. ———, *Projections in Banach algebras*, Ann. of Math. vol. 53 (1951) pp. 235–249.
4. N. H. McCoy, *Subrings of direct sums*, Amer. J. Math. vol. 60 (1938) pp. 374–382.
5. ———, *Subdirect sums of rings*, Bull. Amer. Math. Soc. vol. 53 (1947) pp. 856–877.
6. N. H. McCoy and D. Montgomery, *A representation of generalized Boolean rings*, Duke Math. J. vol. 3 (1937) pp. 455–459.
7. J. von Neumann, *Continuous geometry*, vol. 2, Princeton Lectures, 1936–37.
8. M. H. Stone, *The theory of representations for Boolean algebras*, Trans. Amer. Math. Soc. vol. 40 (1936) pp. 37–111.
9. ———, *Applications of the theory of Boolean rings to general topology*, Trans. Amer. Math. Soc. vol. 41 (1937) pp. 375–481.
10. Bertram Yood, *Banach algebras of bounded functions*, Duke Math. J. vol. 16 (1949) pp. 151–163.

RUTGERS UNIVERSITY