ON PARACOMPACT SPACES¹

CHUNG-TAO YANG

1. Stone² has proved that a space is fully normal T_1 if and only if it is paracompact T_2 . If throughout his proof T_1 is deleted and T_2 is replaced by T_4 (normality), we obtain that *a space is fully normal if and only if it is paracompact T_4* . In this note we prove that T_2 can also be replaced by any one of the following:

T'_2 . Every two points with disjoint closures have disjoint neighborhoods.

T'_3 . For each point x and neighborhood U of $\bar{x}$ there is a neighborhood V of x whose closure is contained in U .

LT_4 . Every point has a neighborhood whose closure is normal.

Moreover, we shall also study a natural decomposition of a space with certain properties and prove that a space is paracompact T_4 if and only if it has a retract which is paracompact T_2 and meets every non-null closed set.

The following terminology will be used.

PROPERTY *. If x, y, z are any three points such that $\bar{x} \cap \bar{y} \neq \emptyset$ and $\bar{x} \cap \bar{z} \neq \emptyset$, then $\bar{y} \cap \bar{z} \neq \emptyset$.

PROPERTY **. Every point has a compact closure.

We remark that

- (1) $T_2 \rightarrow T'_2$ and $T_3 \rightarrow T'_3$.
- (2) $T_4 \rightarrow LT_4 \rightarrow T'_3 \rightarrow T'_2 \rightarrow \text{property } *$.

2. THEOREM 1. *In a paracompact space, T'_2 , T'_3 , LT_4 and T_4 are equivalent.*

PROOF. It is immediate that T_4 implies T'_2 , T'_3 , and LT_4 . To prove that T'_2 plus paracompactness implies T'_3 and that T'_3 plus paracompactness implies T_4 we can simply follow Dieudonné's proof³ with T_2 and T_3 replaced by T'_2 and T'_3 respectively. For any paracompact LT_4 space there is a locally finite open covering, say $\{U_\alpha\}$, such that each $\overline{U_\alpha}$ is normal. Since $\{\overline{U_\alpha}\}$ is still locally finite, the un-

Received by the editors October 23, 1952 and, in revised form, August 11, 1953.

¹ One part of this work was done under Contract N7-434 Task Order III, Naval Department, Office of Naval Research.

² Cf. A. H. Stone, *Paracompactness and product spaces*, Bull. Amer. Math. Soc. vol. 54 (1948) pp. 977-982.

³ Cf. J. Dieudonné, *Une généralisation des espaces compacts*, J. Math. Pures Appl. (9) vol. 23 (1944) pp. 65-76.

proved part of Theorem 1 follows from the following more general result.

LEMMA.⁴ *A space which is covered by a locally finite system of normal closed sets is normal.*

PROOF. Let E, F be disjoint closed sets in a topological space X covered by a locally finite system of closed normal sets A_α . For each α , there exist, by the normality of A_α , open sets U_α, V_α in X such that

$$E \cap A_\alpha \subset U_\alpha, \quad F \cap A_\alpha \subset V_\alpha, \text{ and } U_\alpha \cap V_\alpha \cap A_\alpha = \emptyset.$$

Now we define

$$\begin{aligned} P(x) &= X - \bigcup \{A_\alpha : A_\alpha \ni x\}, & x \in X; \\ Q(x) &= \bigcap \{U_\alpha : A_\alpha \ni x\}, & x \in E; \\ R(x) &= \bigcap \{V_\alpha : A_\alpha \ni x\}, & x \in F. \end{aligned}$$

It follows by the local finiteness of the system $\{A_\alpha\}$ that $P(x)$, $Q(x)$, and $R(x)$ are open. Let

$$U = \bigcup \{P(x) \cap Q(x) : x \in E\}, \quad V = \bigcup \{P(x) \cap R(x) : x \in F\}.$$

Then U, V are disjoint neighborhoods of E, F .

REMARK. We can show by examples that in a pointwise paracompact space, T'_2 , T'_3 , LT_4 , and T_4 are not equivalent to one another.

COROLLARY.⁵ *The product of a paracompact T_4 space and a compact T_4 space is paracompact T_4 .*

To prove this corollary we have only to observe that the product of a paracompact space and a compact space is paracompact⁶ and that the product of two T'_2 spaces is T'_2 .

3. Given any space X with property * we can define an equivalence relation such that two points x, y of X are equivalent if and only if $\bar{x} \cap \bar{y} \neq \emptyset$. This relation yields a decomposition D of X , that is, a system of sets, pairwise disjoint, whose union is X and such that two points of X are contained in a same member of D if and only if they are equivalent. Denote by ϕ the *projection* of X onto D , i.e., the function of X into D such that $f(x) = p$ if $x \in p$. There is a topology on D such that a subset G of D is open if and only if $\phi^{-1}(G)$ is open. D with this topology is the *natural quotient space* of X . If X is T_4 and then

⁴ This is suggested by the referee as a known but unpublished result.

⁵ Cf. H. Miyazaki, *A note on paracompact spaces*, Tôhoku Math. J. (2) vol. 4 (1952) pp. 88–92.

⁶ Cf. J. Dieudonné, loc. cit.

has property *, the natural quotient space of X agrees with one constructed by Čech.⁷ Throughout the rest of this note, D and ϕ always denote the natural quotient space and the projection for space X provided X has property *.

LEMMA. *Let X be a space with property * and property **.*

(3) *Whenever $p \in D$, $\phi^{-1}(p)$ contains a smallest non-null closed set F_p . $F_p = \bar{x}$ for $x \in F_p$.*

(4) *Every open set which meets F_p contains $\phi^{-1}(p)$. Hence for any $x \in X$, if $\bar{x}$ is covered by a system of open sets, $\bar{x}$ is contained in one of them.*

(5) *If E, F are disjoint closed sets in X , then $\phi(E) \cap \phi(F) = \emptyset$.*

PROOF. Fix $p \in D$ and let $x_0 \in \phi^{-1}(p)$. Clearly $\{\bar{x} \cap \bar{x}_0 : x \in \phi^{-1}(p)\}$ is a system of closed sets in $\bar{x}_0$ and it has, by property *, the finite intersection property. It follows by property ** that

$$F_p = \cap \{x : x \in \phi^{-1}(p)\} = \cap \{\bar{x} \cap \bar{x}_0 : x \in \phi^{-1}(p)\}$$

is a non-null closed set contained in $\phi^{-1}(p)$. For any non-null closed set F contained in $\phi^{-1}(p)$, we have $F \supset \bar{x} \supset F_p$ whenever $x \in F$. Hence F_p is the smallest. If $x \in F_p$, then $\bar{x} \subset F_p \subset \bar{x}$. Hence (3) is proved.

By construction $F_p \subset \bar{x}$ for each $x \in \phi^{-1}(p)$. Hence, if U is open and meets F_p , U meets $\{x\}$, i.e., U contains each $x \in \phi^{-1}(p)$, proving that $\phi^{-1}(p) \subset U$. If $x \in X$ and $\bar{x}$ is covered by a system $\mathfrak{U}$ of open sets, then some open set of $\mathfrak{U}$ meets F_p with $p = \phi(x)$, and hence contains $\bar{x}$. This proves (4).

If E and F are disjoint closed sets in X , then for $x \in E$ and $y \in F$, $\bar{x} \cap \bar{y} \subset E \cap F = \emptyset$ and so $\phi(x) \neq \phi(y)$. Hence $\phi(E) \cap \phi(F) = \emptyset$, proving (5).

LEMMA. *Let X be a T'_3 space with property **.*

(6) *ϕ is closed.*

(7) *Given any open covering $\{U_\alpha\}$ of X , $\{D - \phi(X - U_\alpha)\}$ is an open covering of D .*

(8) *D is T_{23} (i.e., T_2 and T_3).*

(9) *D is T_{24} (i.e., T_2 and T_4) or LT_{24} (i.e., T_2 and LT_4) according as X is T_4 or LT_4 .*

PROOF. Let F be a closed subset of X . Given any point p of $D - \phi(F)$ there is, by (3), a smallest non-null closed set F_p contained in $\phi^{-1}(p)$. Take a point x_p of F_p ; there is, by T'_3 , a neighborhood V_p of x_p whose closure is contained in $X - F$. Therefore, by (4) and (5), $\phi^{-1}(p) \subset V_p \subset \bar{V}_p \subset X - \phi^{-1}\phi(F) = \phi^{-1}(D - \phi(F))$. Hence $\phi^{-1}(D - \phi(F))$

⁷ Cf. E. Čech, *On bicomact spaces*, Ann. of Math. (2) vol. 38 (1937) pp. 823-844.

$= \bigcup \{ V_p : p \in D - \phi(F) \}$ is open and so $\phi(F)$ is closed. This proves (6).

Given any open covering $\{U_\alpha\}$ of X , $\{D - \phi(X - U_\alpha)\}$ is a system of open sets in D by (6). For each $p \in D$ there is, by (4), some α such that $\phi^{-1}(p) \subset U_\alpha$ and so $p \in D - \phi(X - U_\alpha)$. Hence (7) is proved.

By (6), D is a T_1 space. Therefore D is T_{23} if we can show that for a point p of D and a neighborhood G of p there is a neighborhood of p whose closure is contained in G . Take a point x of F_p and let U be a neighborhood of x with $\bar{U} \subset \phi^{-1}(G)$. Then $D - \phi(X - U)$ is a neighborhood of p by (4) and (6) and its closure is contained in G by (5).

Suppose that X is T_4 . Given any two disjoint closed subsets P, Q of D , $\phi^{-1}(P)$ and $\phi^{-1}(Q)$ are disjoint closed subsets of X and they have disjoint neighborhoods U and V . Applying (6), we can easily see that $D - \phi(X - U)$ and $D - \phi(X - V)$ are disjoint neighborhoods of P and Q , proving that D is T_4 and hence T_{24} .

Suppose now that X is LT_4 . Let $p \in D$ and $x \in F_p$; there is, by hypothesis, a neighborhood U of x whose closure is normal. By (4) and (6), $G = D - \phi(X - U)$ is a neighborhood of p . Then there is, by T_3 , a neighborhood V of p such that $\bar{V} \subset G$. Since $\phi^{-1}(\bar{V}) \subset \bar{U}$ is normal, it follows by the preceding result that $\bar{V}$ is normal, proving that D is LT_4 and hence LT_{24} .

LEMMA. Let X be a pointwise paracompact (paracompact) space.

(10) X has property **.

(11) If X is T'_3 , then D is pointwise paracompact (paracompact).

PROOF. Fix a point x_0 of X . Given any system $\mathcal{U}$ of open sets whose union contains $\bar{x}_0$, $\mathcal{U} \cup \{X - \bar{x}_0\}$ is an open covering of X and it admits a point-finite refinement $\mathcal{V}$. Let $\mathcal{V}' = \{V : V \in \mathcal{V}, V \cap \bar{x}_0 \neq \emptyset\} = \{V : V \in \mathcal{V}, V \ni x_0\}$ and for each $V \in \mathcal{V}'$ we take a $U_V \in \mathcal{U}$ such that $V \subset U_V$. Then $\mathcal{U}' = \{U_V : V \in \mathcal{V}'\}$ is a finite subsystem of $\mathcal{U}$ which covers $\bar{x}_0$, proving the compactness of $\bar{x}_0$. Hence X has property **.

Given any open covering $\{G_\alpha\}$ of D , $\{\phi^{-1}(G_\alpha)\}$ is an open covering of X and it admits, by hypothesis, a point-finite (locally finite) refinement $\{U_\beta\}$. By (7), $\{D - \phi(X - U_\beta)\}$ is an open covering of D which is obviously point-finite (locally finite) and refines $\{G_\alpha\}$. Hence D is pointwise paracompact (paracompact).

THEOREM 2. A space X is pointwise paracompact (paracompact) T'_3 if and only if it has a pointwise paracompact (paracompact) T_{23} retract A which meets every non-null closed subset of X . Moreover, such a retract A is LT_{24} or T_{24} if and only if X is LT_4 or T_4 . Finally, A is unique up to a homeomorphism and the related retraction is uniquely determined.

PROOF. Suppose that X is pointwise paracompact (paracompact) T_3' . By (2), and (10), X has property $*$ and property $**$; it follows by (3) that for each $p \in D$ there is a smallest non-null closed set F_p contained in $\phi^{-1}(p)$. We take, for each $p \in D$, a point x_p of F_p and denote by A the set of these points x_p . Clearly A meets every non-null closed subset of X and ϕ defines a 1-1 mapping ψ of A onto D . For any closed subset F of X , we have $\psi(F \cap A) = \phi(F)$ which is closed by (6). Hence ψ is a homeomorphism. From this result, we obtain a retraction $f = \psi^{-1}\phi$ of X onto A . Moreover, it follows by (8) and (9) that A is T_{23} and that A is T_{24} or LT_4 according as X is T_4 or LT_4 .

Conversely suppose that X has a pointwise paracompact (paracompact) T_{23} retract A which meets every non-null closed subset of X . Let f be a retraction of X into A . For any $x \in X$, $\bar{x} \cap A$ is non-null and T_2 ; therefore it contains exactly one point. From this result, it is easily seen that for any two points x, y of X , $\bar{x} \cap \bar{y} \neq \emptyset$ if and only if $\bar{x} \cap A = \bar{y} \cap A$. Hence property $*$ as well as property $**$ holds for X , and for each $p \in D$, $\phi^{-1}(p) \cap A$ contains exactly one point contained in F_p . From the latter result, we have, for $x \in X$, $f(x) = f(\bar{x}) = \bar{x} \cap A$. Let $x \in X$ and let U be a neighborhood of $\bar{x}$. Then there is a neighborhood G of $\bar{x} \cap A$ in A such that $\bar{x} \cap A \subset G \subset \bar{G} \cap A \subset U \cap A$. Hence $f^{-1}(G)$ is a neighborhood of x whose closure is contained in U . This proves that X is T_3' .

If A is T_4 , then for any two disjoint closed subsets E, F of X , $E \cap A$ and $F \cap A$ have disjoint neighborhoods G and H in A and so $f^{-1}(G)$ and $f^{-1}(H)$ are disjoint neighborhoods of E and F . Hence X is also T_4 . If A is LT_4 , then for any $x \in X$, $f(x)$ has a neighborhood G in A whose closure in A is normal. It follows by the preceding result that $f^{-1}(G)$ is a neighborhood of x whose closure is normal. Hence X is also LT_4 .

Given any open covering $\{U_\alpha\}$ of X , $\{U_\alpha \cap A\}$ is an open covering of A and it admits, by hypothesis, a point-finite (locally finite) refinement $\{G_\beta\}$. We can easily see that $\{f^{-1}(G_\beta)\}$ is a point-finite (locally finite) refinement of $\{U_\alpha\}$. Hence X is pointwise paracompact (paracompact).

According to the above argument, A is homeomorphic to D and hence is unique up to a homeomorphism. Moreover, for any $x \in X$, $\bar{x} \cap A = f(x)$. Hence the retraction f is uniquely determined.

The author is much indebted to the referee for improving several results as well as for simplifying some proofs.