ON INTERVALS CONTAINING AN AFFINELY EQUIVALENT SET OF n INTEGERS MOD k

L. J. MORDELL

1. Let $n > 2$ be a given positive integer and $(x) = (x_1, x_2, \dots, x_n)$ a given set of n integers. Let k be a given positive integer. Then L. Redei [1], improving an old result of Thue [2], has recently proved the

THEOREM. *If k is a prime, there exist integers a, b , $(a, k) = 1$, such that a set of integers (y) , defined by*

$$y_r \equiv ax_r + b \pmod{k} \quad (r = 1, 2, \dots, n),$$

lies in an interval of width L where

$$nL^{n-1} \leq 2^{n-1}k^{n-2},$$

i.e.

$$|y_r - y_s| \leq L \quad (r, s = 1, 2, \dots, n).$$

The estimate is trivial if it can give a value of $L \geq k$, i.e. if $nk^{n-1} \leq 2^{n-1}k^{n-2}$ or $k \leq 2^{n-1}/n$. There is also a trivial result $L = 0$ if we had allowed $a \equiv 0 \pmod{k}$.

I notice that Redei's proof can be presented in a rather simpler form and also extended to the case where k is not a prime. We have now

THEOREM I. *Let δ be the greatest divisor of k such that*

$$x_1 \equiv x_2 \equiv \dots \equiv x_n \pmod{\delta},$$

and so $\delta = (x_1 - x_n, x_2 - x_n, \dots, x_{n-1} - x_n, k)$. Then integers $a \not\equiv 0 \pmod{k}$, b exist such that an integer set (y) , defined by

$$y_r \equiv ax_r + b \pmod{k} \quad (r = 1, 2, \dots, n),$$

lies in an interval of width $L = L(\delta, k)$, where

$$nL^{n-1} \leq 2^{n-1}\delta k^{n-2}.$$

There is no loss of generality in theorems of this kind, if we suppose hereafter that $\delta = 1$. For clearly

$$L(\delta, k) = \delta L(1, k/\delta),$$

since $y_r - y_s \equiv a(x_r - x_s) \pmod{k}$, and so we can put $y_r = \delta y'_r + c$ where

Presented to the Society, September 5, 1953; received by the editors April 5, 1953.

y_r' is an integer and c is a constant.

Results of this kind can also be found by an application of Dirichlet's classical distribution result and this leads to

THEOREM II. *Let λ be the least positive integer which satisfies the inequality*

$$(\lambda + 1)^n - \lambda^n \geq k^{n-2}.$$

Then we can take $L = 2\lambda$ or $L = 2\lambda + 1$ according as the inequality or equality sign holds. Under certain conditions, as stated in Theorem III, the result $L = 2\lambda + 1$ is best possible.¹

Further the result $L = 2\lambda$ can be replaced by $L = 2\lambda - 1$ if now λ is the least positive integer which satisfies the inequality

$$(\lambda + 1)^n - (\lambda - 1)^n \geq 2k^{n-2}.$$

Write $(k^{n-2}/n)^{1/(n-1)} = I + f$, where I is an integer and $0 \leq f < 1$. When $0 \leq f < 1/2$, the result of Theorem I is included in the first part above. For on putting $\lambda + 1/2 = \mu$, the first inequality is satisfied by $\lambda + 1/2 \geq I + f$, i.e. $\lambda = I$ and so $L = 2I$. Theorem I gives $L = [2I + 2f] = 2I$.

When $1/2 \leq f < 1$, the result of Theorem I is given in the second part above. For the second equality is satisfied by $\lambda \geq I + f$, i.e. $\lambda = I + 1$ and so $L = 2I + 1$. Theorem I, however, gives $L = [2I + 2f] = 2I + 1$.

2. For Theorem I, put

$$y_r = ax_r + b + k\xi_r \quad (r = 1, 2, \dots, n),$$

where ξ_r is an integer. Then we have to satisfy

$$|ax_r - x_s + k(\xi_r - \xi_s)| \leq L \quad (r, s = 1, 2, \dots, n).$$

Write

$$\xi_r = \eta_r + \xi_n, \quad x_r = x_r' + x_n,$$

where now, and in the remainder of this chapter, r and s take the values $1, 2, \dots, n-1$. Then

$$|ax_r' + k\eta_r| \leq L, \quad |a(x_r' - x_s') + k(\eta_r - \eta_s)| \leq L.$$

Define an $(n-1)$ -dimensional region R by

¹ Redei [1, p. 81] states that this is easily proved when k is a prime and $n=3$. The referee notes that this is obvious even when k is not a prime on taking $x_1=1$, $x_2=3\lambda+2$, $x_3=0$. I should like to thank him for this remark and also for some other valuable comments.

$$|X_r| \leq L, \quad |X_r - X_s| \leq L.$$

Then R contains a point (x) with coordinates

$$x_r = ax'_r + k\eta_r \quad (r = 1, 2, \dots, n-1),$$

where a and the η are integers. We define a $(2n-1)$ -dimensional region S by combining R and

$$|X_r - ax'_r - k\eta_r| < 1, \quad |a| < k,$$

where the η and a are the additional n variables.

We use Minkowski's convex region theorem to find L such that S contains an integer set $[(X), a, (\eta)] \neq [(0), 0, (0)]$. Redei states that the content of R is obviously nL^{n-1} , but a proof is contained in the lemma following. Then clearly on integrating first for η and a , the content V of S is given by

$$V = nL^{n-1}(2/k)^{n-1}2k,$$

and so $V \geq 2^{2n-1}$ if

$$nL^{n-1} \geq 2^{n-1}k^{n-2}.$$

This is really Redei's result; and on noting the equation defining R , it suffices to take

$$L = (2^{n-1}k^{n-2}/n)^{1/(n-1)},$$

where the square bracket denotes the integer part. Then S contains an integer set and for this, obviously,

$$X_r - ax'_r - k\eta_r = 0,$$

i.e.

$$X_r \equiv ax'_r \pmod{k}.$$

It remains only to show that the trivial result $a=0$ has been excluded. If $a=0$, $X_r \equiv 0 \pmod{k}$. By Minkowski's theorem, all the X_r are not zero, since if they were, all the η_r would be zero. Hence the trivial solution is excluded if $L < k$, i.e. $k > 2^{n-1}/n$.

3. Theorem II follows from a simple application of a general principle. Let a bounded n -dimensional convex region $R(\lambda)$, symmetrical about the origin, be defined by $f(X) \leq \lambda$, where $f(X)$ is a distance function and so $f(X' - X'') \leq f(X') + f(X'')$, and λ is a positive parameter. We find an estimate for L such that the region $R(L)$ will contain an integer set $(X) \neq 0$, satisfying r given linear homogeneous congruences $\pmod{k}$, say,

$$L_1(X) \equiv 0, L_2(X) \equiv 0, \dots, L_r(X) \equiv 0.$$

Let $N(\lambda)$ be the number of lattice points in $R(\lambda)$. Then if (X) runs through these $N(\lambda)$ points, the L 's assume $N(\lambda)$ sets of residues (mod k). If $N(\lambda) > k^r$, two different lattice points (X') , (X'') give the same set of residues for the L 's. Hence $(X) = (X' - X'')$ satisfies the homogeneous congruences and clearly lies in the region $R(2\lambda)$.

The $R(2\lambda)$ may sometimes be improved to $R(2\lambda - 1)$ as follows. The region $R(\lambda - 1)$ contains $N(\lambda - 1)$ lattice points. We adjoin to these, say, $N_1(\lambda)$ lattice points lying in $R(\lambda)$ and not in $R(\lambda - 1)$, taken in such a way that $(X' - X'')$ does not lie on the boundary of $R(2\lambda)$. Then if $N(\lambda - 1) + N_1(\lambda) > k^r$, the point $(X' - X'')$ will lie in $R(2\lambda - 1)$. In fact, we take $N_1(\lambda) = 1$ in the proof of Theorem II.

4. We require the

LEMMA. Let $R(\lambda)$, where λ is a positive integer, denote the region

$$|x_r| \leq \lambda, \quad |x_r - x_s| \leq \lambda \quad (r, s = 1, 2, \dots, n-1).$$

Then $R(\lambda)$ contains exactly

$$N(\lambda) = (\lambda + 1)^n - \lambda^n$$

lattice points, i.e. points with integer coordinates.

My original proof was rather lengthy so I replace it by a very simple one found subsequently by Dr. Cassels and kindly put at my disposal. Write $x_0 = 0$ and define $y_0, y_1, \dots, y_{n-1}$ by $y_0 = \max(x_r)$, $y_r + x_r = y_0$ ($r = 1, 2, \dots, n-1$). If y_0 arises from an x_s , then $y_s = 0$. Also $0 \leq y_r \leq \lambda$. Then the y 's can take all integer values satisfying

$$\max y_r \leq \lambda, \quad \min y_r = 0 \quad (r = 0, 1, 2, \dots, n-1).$$

The first inequality gives $(\lambda + 1)^n$ possibilities for the y 's, and the second excludes $y_r = 1, 2, \dots, \lambda$, i.e. λ^n possibilities. Hence the result.

COROLLARY. It follows on inscribing $(n-1)$ -dimensional cubes of sides $1/\lambda$ in R that the content V of $R(1)$ is given by

$$V = \lim_{\lambda \rightarrow \infty} N(\lambda)/\lambda^{n-1} = n.$$

5. We now find λ so that the region $R(\lambda)$ contains an integer set (X) congruent mod k to at least one of the sets (ax') , where (x') is any given set of $n-1$ integers with $(x'_1, x'_2, \dots, x'_{n-1}, k) = 1$, and a takes the values $1, 2, \dots, k-1$, i.e.

$$X_r \equiv ax'_r \pmod{k} \quad (r = 1, 2, \dots, n-1).$$

Write $Y_r = X_r - ax'_r$. When (X) runs through the $N(\lambda)$ lattice

points in $R(\lambda)$ and a runs through the values $0, 1, \dots, k-1$, then the (Y) runs through $kN(\lambda)$ residue sets mod k . Hence if $kN(\lambda) > k^{n-1}$, there must be at least two different sets (X', a') , (X'', a'') giving the same residue sets for (Y) . Then a solution of the congruences $(Y) \equiv 0$ is given by $(X) = (X' - X'')$, $a = a' - a''$. This set (X) will be in $R(2\lambda)$. If this occurs with the trivial solution $a=0$, then $(X) \equiv 0 \pmod{k}$, and so this will be excluded if $k > 2\lambda$. Hence if λ is a positive integer such that

$$(\lambda + 1)^n - \lambda^n > k^{n-2},$$

(X) will be in the region $R(2\lambda)$.

If, however, there exists a positive integer λ such that

$$(\lambda + 1)^n - \lambda^n = k^{n-2},$$

we show that there will be a suitable point (X) in $R(2\lambda+1)$. For, take a lattice point $(\bar{X})$ lying in $R(\lambda+1)$ but not in $R(\lambda)$. This one combined with those in $R(\lambda)$ gives more than k^{n-2} sets of residues. But now the solution $(X) = (X' - X'')$ will lie in $R(2\lambda+1)$.

We prove now the

THEOREM III. *The result $R(2\lambda+1)$ is best possible if the values assumed by $(X) - a(x)$ form a complete set of residues mod k when (X) runs through the lattice points of $R(\lambda)$ and a takes the values $0, 1, 2, \dots, k-1$.*

It suffices to prove that if (X) is any lattice point in $R(2\lambda)$, then

$$(X) = (Z) + (Z'),$$

where (Z) and (Z') are lattice points in $R(\lambda)$. For then the condition in the theorem implies that the relation

$$(Z) - a(x) \equiv -(Z')$$

holds only when $a=0$, $(Z) = -(Z')$. Then $(X)=0$, and so the only solution of $(X) \equiv a(x)$, where (X) is in $R(2\lambda)$, is the trivial one.

Let $\alpha, \alpha', \dots; \beta, \beta', \dots; \gamma, \gamma', \dots$ typify indices $1, 2, \dots, n-1$ for which

$$\begin{aligned} X_\alpha &= 2Y_\alpha \equiv 0 \pmod{2}, & X_\beta &\leq 0, \\ X_\beta &= 2Y_\beta - 1 \equiv 1 \pmod{2}, & X_\gamma &\geq 0, \\ X_\gamma &= 2Y_\gamma + 1 \equiv 1 \pmod{2}, \end{aligned}$$

When the same argument applies to both the β and γ sets, we use the letter δ and put

$$X_\delta = 2Y_\delta + \epsilon_\delta, \quad \epsilon_\delta = \epsilon_{\delta'} = \dots = \pm 1.$$

We take (Z) , (Z') to be lattice points which we can write as

$$(Z) = (Y_\alpha, \dots, Y_\beta - 1, \dots, Y_\gamma \dots),$$

$$(Z') = (Y_\alpha, \dots, Y_\beta, \dots, Y_\gamma + 1, \dots),$$

where the notation indicates the various types of coordinates. These points will be in $R(\lambda)$, i.e. $|Z_r| \leq \lambda$, $|Z_r - Z_s| \leq \lambda$, etc., if the following five sets of conditions are satisfied.

$$(1) \quad |Y_\alpha| \leq \lambda, \quad |Y_\delta| \leq \lambda - 1.$$

These hold, as follows from

$$|2Y_\alpha| \leq 2\lambda, \quad |2Y_\delta + \epsilon_\delta| \leq 2\lambda.$$

For $2|Y_\delta| + 1 \leq 2\lambda$, $|Y_\delta| \leq \lambda - 1/2$, and since Y_δ is an integer, $|Y_\delta| \leq \lambda - 1$.

$$(2) \quad |Y_\alpha - Y_{\alpha'}| \leq \lambda, \quad |Y_\delta - Y_{\delta'}| \leq \lambda.$$

This holds since

$$|2Y_\alpha - 2Y_{\alpha'}| \leq 2\lambda, \quad |2Y_\delta + \epsilon_\delta - (2Y_{\delta'} + \epsilon_{\delta'})| \leq 2\lambda, \quad \epsilon_\delta = \epsilon_{\delta'}.$$

$$(3) \quad |Y_\alpha - Y_\beta + 1| \leq \lambda, \quad |Y_\alpha - Y_\beta| \leq \lambda.$$

Now $|2Y_\alpha - 2Y_\beta + 1| \leq 2\lambda$ which gives

$$|2Y_\alpha - 2Y_\beta + 2| \leq 2\lambda + 1, \quad |2Y_\alpha - 2Y_\beta| \leq 2\lambda + 1, \\ |Y_\alpha - Y_\beta + 1| \leq \lambda + 1/2, \quad |Y_\alpha - Y_\beta| \leq \lambda + 1/2, \text{ etc.}$$

$$(4) \quad |Y_\beta - 1 - Y_\gamma| \leq \lambda.$$

This results from

$$|2Y_\beta - 1 - 2Y_\gamma - 1| \leq 2\lambda, \\ (5) \quad |Y_\alpha - Y_\gamma| \leq \lambda, \quad |Y_\alpha - Y_\gamma - 1| \leq \lambda.$$

This follows from

$$|2Y_\alpha - 2Y_\gamma - 1| \leq 2\lambda,$$

which gives

$$|2Y_\alpha - 2Y_\gamma| \leq 2\lambda + 1, \quad |2Y_\alpha - 2Y_\gamma - 2| \leq 2\lambda + 1, \text{ etc.}$$

This finishes the proof.

BIBLIOGRAPHY

1. L. Redei, *Über eine Verschärfung eines Zahlentheoretischen Satzes von Thue*, Acta Mathematica Academiae Scientiarum Hungaricae vol. 2 (1951).

2. A. Thue, *Et par antydninger til en taltheoretisk metode*, Christiania Videnskabs Selskabs Forh., 1902, no. 7.

ST. JOHNS COLLEGE, CAMBRIDGE UNIVERSITY