A NOTE ON DENDRITES AND TREES

L. E. WARD, JR.

By a *tree* we mean a continuum (=compact connected Hausdorff space) in which every pair of distinct points is separated by a third point. It is well known [4] that a dendrite is simply a metric tree. In this note we give a new characterization of dendrites and trees in terms of partially ordered spaces.

Recall from [3] that a POTS (=partially ordered topological space) is a topological space X endowed with a partial order, $\leq$, which is semicontinuous in the sense that $L(x) = \{a: a \leq x\}$ and $M(x) = \{a: x \leq a\}$ are closed sets for each $x \in X$. A partial order is *order dense* if $x < y$ implies there is a z such that $x < z < y$. A *chain* of a partially ordered set is a subset which is linear with respect to the partial order. An element in a partially ordered set is *minimal* (*maximal*) if it has no proper predecessor (successor).

Suppose that K is a locally connected continuum and $e \in K$. Define a relation, $\leq$, in K by $x \leq y$ if, and only if, $x = e$, or $x = y$, or x separates e and y in K . From [3] we have

LEMMA 1. *The relation $\leq$ is a semicontinuous partial order.*

We shall require two more results from [3]. Lemma 2 is originally due to Wallace [2].

LEMMA 2. *A compact POTS contains a maximal element.*

LEMMA 3. *A compact order dense POTS is connected if its set of minimal elements is connected.*

LEMMA 4. *A tree is locally connected.*

PROOF. Let X be a tree; by [1, p. 140] X is regular in the sense that if $x \in U \subset X$, U open, then there is a neighborhood V of x , with $\bar{V} \subset U$ and such that the frontier of V is finite. By [4, p. 19] X is locally connected. Since the above arguments are valid in our situation, the proof is complete. $\square$

We may now prove our chief result.

THEOREM 1. *Let X be a compact Hausdorff space. A necessary and sufficient condition that X be a tree is that X admit a partial order, $\leq$, satisfying*

(i) $\leq$ *is semicontinuous,*

Presented to the Society, June 19, 1954; received by the editors April 23, 1954.

- (ii) $\leq$ is order dense,
- (iii) for $x \in X$, $y \in X$, it follows that $L(x) \cap L(y)$ is a non-null chain,
- (iv) $M(x) - x$ is an open set, for each $x \in X$.

PROOF. Let X be a tree; choose $e \in X$ and let X have the semi-continuous partial order of Lemma 1. To establish (ii), let $x < y$ and suppose z separates x and y , i.e.,

$$X - z = C \cup D, \quad C \mid D, \quad x \in C, \quad y \in D.$$

If $x = e$, then, by definition, $x < z < y$. Otherwise, we have

$$X - x = A \cup B, \quad A \mid B, \quad e \in A, \quad y \in B.$$

It is easy to see that $x \in C$ and $y \in B$ imply $A \subset C$. Therefore $e \in C$, so that $z < y$. Likewise, $z \in B$, so that $x < z$. To prove (iii), we first note that by definition of the partial order, $e \in L(x) \cap L(y)$ for any x and y in X . It remains to show that $L(x)$ is a chain. If $e < p_1, p_2 < x$, we have

$$\begin{aligned} X - p_i &= A_i \cup B_i, & A_i \mid B_i & \quad (i = 1, 2), \\ x &\in B_1 \cap B_2, & e &\in A_1 \cap A_2. \end{aligned}$$

It is clear that $p_1 \in \overline{A_2}$ implies $p_1 \leq p_2$ and $p_1 \in \overline{B_2}$ implies $p_2 \leq p_1$. To establish (iv), we note simply that $M(x) - x = X - x$ if $x = e$, and if $x \neq e$, then

$$M(x) - x = \bigcup \{ B_\alpha : X - x = A_\alpha \cup B_\alpha, A_\alpha \mid B_\alpha, e \in A_\alpha \}.$$

In either case, $M(x) - x$ is open. Thus the condition is necessary.

Suppose now that X admits a partial order satisfying (i)–(iv). By Lemma 3, X is connected and hence is a continuum. If $x < y$, then by (ii) there is $z \in X$ such that $x < z < y$. By (i) and (iv), $M(z)$ is closed and $M(z) - z$ is open, so that z separates x and y . If x and y are not comparable, then by (iii) and Lemma 2, there exists $z = \max(L(x) \cap L(y))$. Choose t such that $z < t < x$. Then $x \in M(t) - t$ and $y \in X - M(t)$, whence t separates x and y . This completes the proof.

By Theorem 1 and Lemma 4, we have at once:

COROLLARY. *Let X be a compactum. A necessary and sufficient condition that X be a dendrite is that X admit a partial order satisfying (i)–(iv) of Theorem 1.*

BIBLIOGRAPHY

1. R. L. Moore, *Foundations of point set theory*, New York, 1932.

2. A. D. Wallace, *A fixed point theorem*, Bull. Amer. Math. Soc. vol. 51 (1945) pp. 413-416.
3. L. E. Ward, Jr., *Partially ordered topological spaces*, Proc. Amer. Math. Soc. vol. 5 (1954) pp. 144-161.
4. G. T. Whyburn, *Analytic topology*, New York, 1942.

THE UNIVERSITY OF NEVADA

CORRECTION TO "UNIQUENESS OF THE PROJECTIVE PLANE WITH 57 POINTS"

MARSHALL HALL, JR.

Professor Gunter Pickert has pointed out to me that there is an error in my paper [Proc. Amer. Math. Soc. vol. 4 (1953) pp. 912-916]. The equation on page 915 which reads $12 + 3s + 2t = 24$ should read $12 + 3s + 2t + u = 24$. This invalidates my conclusion $u = 0$ from which I deduce $U = 0$, which is necessary for the rest of the paper. I give here a new proof that $U = 0$.

We must show that in a 57 point plane a line containing points ACC is not possible, or that $U = 0$. The proof is by showing that if there is a line ACC we reach a contradiction. We may take ACC as $A_1C_4C_6$ by numbering points appropriately. We now reletter by the following rule:

$$\begin{array}{cccccccccc} A_1 & A_2 & A_3 & A_4 & B_1 & B_2 & B_3 & C_4 & C_6 \\ A_1 & B_1 & B_5 & A_0 & B_4 & B_2 & A_2 & B_3 & B_6 \end{array}$$

and obtain the following configuration:

$$\begin{array}{cccc} A_0 & B_1 & B_2 & B_3 \\ A_0 & B_4 & B_5 & B_6 \\ A_0 & A_1 & A_2 & \\ A_1 & B_1 & B_4 & \\ A_1 & B_2 & B_5 & \\ A_1 & B_3 & B_6 & \\ A_2 & B_1 & B_5 & \\ A_2 & B_2 & B_6 & \\ A_2 & B_3 & B_4 & \end{array}$$