STRICT SEPARATION OF CONVEX SETS¹

V. L. KLEE, JR.

A *hyperplane* in a (real) topological linear space E is a set of the form $f^{-1}c$, where c is a real number and f is a not-identically-zero continuous linear functional on E . The hyperplane determines two *closed half-spaces*— $f^{-1}] - \infty, c]$ and $f^{-1}[c, \infty[$, and two *open half-spaces*— $f^{-1}] - \infty, c[$ and $f^{-1}]c, \infty[$. Subsets A and B of E are said to be *separated* (*strictly separated*) by the hyperplane provided A lies in one of the closed (*open*) half-spaces and B in the other. An account of the principal known results on separation and strict separation of convex sets may be found in [1] and [3]. Our purpose here is to set forth some apparently new results on strict separation, including the following: *If A and B are disjoint closed convex subsets of Euclidean n -space and neither contains a ray in its boundary, then A and B can be strictly separated by a hyperplane.* A related problem was considered several years ago by the author [3, p. 459], but the present investigation was prompted by a question of Mr. Isaac Namioka.

The proof of our principal result (4) is based on several lemmas, of which the first is given below. A discussion of net-convergence (used below) is contained in [2]. The origin of the linear space under consideration will always be denoted by ϕ .

(1) Suppose C is a locally compact closed convex subset of a topological linear space, $\phi \in C$, U is a neighborhood of ϕ , and S is a net in $C \setminus U$. Then there is a subnet $(x, >)$ of S and a corresponding net $(t, >)$ of positive numbers such that t_n converges to a number $t_0 < \infty$ and $t_n x_n$ converges to a point y_0 of $C \setminus \{\phi\}$.

PROOF. Since C is closed and locally compact, U contains a neighborhood V of ϕ such that the intersection B of C with the boundary of V is compact. For each member k of the domain of S we have $a_k S_k \in B$ for some $a_k \in]0, 1[$. The desired conclusion then follows from the fact that $[0, 1]$ and B are compact, and $\phi \notin B$.

A subset of a linear space is *linearly bounded* provided its intersection with each line is bounded. For convex sets, this is equivalent to saying that the set contains no ray.

Presented to the Society, June 18, 1955; received by the editors February 21, 1955.

¹ Sponsored by the Office of Ordnance Research, U. S. Army, under Contract DA-04-200-ORD-292.

(2) With C as in (1), C is compact if and only if C is linearly bounded.

PROOF. The "only if" part is trivial, and we wish to show that if C is not compact, then C contains a ray. Now if C is not compact, there exist S and U as in (1) such that S has no convergent subnet. And if $t_0 \neq 0$, x_n converges to $t_0^{-1}y_0$, so it follows that $t_0 = 0$. Let $y_n = t_n x_n$ for each n ; then $[0, t_n^{-1}]y_n \subset C$, whence it follows that $[0, \infty[y_0 \subset C$, completing the proof of (2).

(3) Suppose A and B are disjoint closed convex subsets of a topological linear space, and there exists a hyperplane H separating A from B such that $A \cap H$ is nonempty and linearly bounded. Then if either A or B is locally compact, there is an open $V \ni \phi$ such that $A + V$ misses B .

PROOF. Let the continuous linear functional f and the real number a be such that $H = f^{-1}a$ and $A \subset f^{-1}[-\infty, a]$. Consider the case in which A is locally compact. We wish first to show that the set $C = A \cap f^{-1}[a-1, a]$ is compact, and see by (1) that it suffices to show that C is linearly bounded. Suppose C contains a ray T with initial point x , and let $y \in A \cap H$. From the fact that A is closed and convex it follows readily that A contains the ray $T' = y + (T - x)$. And since $T \subset f^{-1}[a-1, a]$, it is clear that f is constant on T , whence $T' \subset A \cap H$, a contradiction showing that C is compact. Now since C is compact and disjoint from the closed set B , there exists open $U \ni \phi$ such that $C + U$ misses B . Let $V = U \cap f^{-1}[-1/2, 1/2]$. Then V is open, $\phi \in V$, $f^{-1}[-\infty, a-1] + V$ misses B , and $C + V$ misses B , so V is the desired open set.

Consider now the case in which B is locally compact. We may assume $\phi \in B$. Suppose the desired V does not exist, and let $\mathfrak{W}$ be the set of all open $W \ni \phi$. Then there exists $U \in \mathfrak{W}$ such that for each $W \in \mathfrak{W}$ there are points $b_w \in B \setminus U$ and $a_w \in A$ with $b_w \in a_w + W$. Now let us apply (1), with the roles of C and S taken by B and $(b_w | W \in \mathfrak{W}, \subset)$ respectively, to obtain nets $(x, >)$ and $(t, >)$ as described in (1), and let $(u, >)$ be the corresponding subnet of $(a_w | W \in \mathfrak{W}, \subset)$. If $t_0 > 0$, then x_n converges to $t_0^{-1}y_0$, whence u_n converges to $t_0^{-1}y_0$ and $A \cap B \neq \Lambda$, contradicting one of our hypotheses. Thus $t_0 = 0$, and as in (2), $[0, \infty[y_0 \subset B$. Let $p \in A \cap H$ and for each n let $z_n = t_n(u_n - p)$. Then z_n converges to y_0 and always $p + [0, t_n^{-1}]z_n \subset A$, whence $p + [0, \infty[y_0 \subset A$. Since the parallel rays $[0, \infty[y_0$ and $p + [0, \infty[y_0$ are separated by the hyperplane $H = f^{-1}a$, f must be constant on each ray and it follows that $p + [0, \infty[y_0 \subset A \cap H$, a contradiction completing the proof of (3).

It is now easy to prove our principal result, which is

(4) For a locally compact closed convex subset A of a locally convex topological linear space E , the following two assertions are equivalent: (i) Whenever H is a hyperplane in E which supports A , then $A \cap H$ is linearly bounded. (ii) Whenever B is a closed convex subset of $E \setminus A$ which can be separated from A , then B can be separated from A by a hyperplane which misses A .

PROOF. That (i) implies (ii) follows at once from (3), the local convexity of E , and the basic separation theorem for convex bodies. That (ii) implies (i) is obvious when E is the plane, and general proof can be based on this case.

From (4) and the known special separation properties of finite-dimensional spaces, we have

(5) For a closed convex subset A of E^n , the following two assertions are equivalent: (i) A contains no ray in its boundary. (ii) Whenever B is a closed convex subset of E^n which misses A , then B can be separated from A by a hyperplane which misses A .

A partial analogue of (4) can be obtained by applying (3) for the case in which B is locally compact. Other corollaries are obtained by observing that if A and B can be separated by two hyperplanes, one missing A and the other missing B , then A and B can be strictly separated. (Together with (5), this yields the result stated in the introduction.) Another consequence of (5) is

(6) Suppose A is a closed convex subset of E^n and that A contains no ray in its boundary. Then every linear manifold in E^n which misses A is contained in a hyperplane which misses A .

In conclusion, we remark that there are examples showing that the assumptions of local compactness cannot be avoided.

REFERENCES

1. N. Bourbaki, *Espaces vectoriels topologiques*, Actualités Scientifiques et Industrielles, No. 489, Paris, 1953.
2. J. L. Kelley, *Convergence in topology*, Duke Math. J. vol. 17 (1950) pp. 277-283.
3. V. L. Klee, Jr., *Convex sets in linear spaces*, Duke Math. J. vol. 18 (1951) pp. 443-466.

UNIVERSITY OF WASHINGTON