THE EXPANSION OF CERTAIN PRODUCTS

L. CARLITZ

1. **Introduction.** The familiar identities

$$(1.1) \quad \prod_{n=0}^{\infty} (1 + x^n t) = \sum_{m=0}^{\infty} x^{m(m-1)/2} t^m (x)_m \quad (|x| < 1),$$

$$(1.2) \quad \prod_{n=0}^{\infty} (1 - x^n t)^{-1} = \sum_{m=0}^{\infty} t^m / (x)_m \quad (|x| < 1),$$

where

$$(1.3) \quad (x)_m = (1 - x)(1 - x^2) \cdots (1 - x^m), \quad (x_0) = 1,$$

suggest the problem of expanding the products

$$(1.4) \quad \prod_{m,n=0}^{\infty} (1 + x^m y^n t), \quad \prod_{m,n=0}^{\infty} (1 - x^m y^n t)^{-1} \quad (|x| < 1, |y| < 1).$$

Indeed Bellman [1] has recently proposed the expansion of the second of (1.4) as a research problem. We put

$$(1.5) \quad \prod_{m,n=0}^{\infty} (1 + x^m y^n t) = \sum_{m=0}^{\infty} t^m G_m(x, y) / (x)_m (y)_m,$$

$$(1.6) \quad \prod_{m,n=0}^{\infty} (1 - x^m y^n t)^{-1} = \sum_{m=0}^{\infty} t^m H_m(x, y) / (x)_m (y)_m.$$

In this note we derive some properties of $G_m(x, y)$ and $H_m(x, y)$ as well as some related functions.

2. To begin with we set up some recursion formulas satisfied by G_m and H_m . Logarithmic differentiation of (1.5) yields

$$\sum_0^{\infty} \frac{x^m y^n t}{1 + x^m y^n t} = \frac{\sum_0^{\infty} m t^n G_m / (x)_m (y)_m}{\sum_0^{\infty} t^n G_m / (x)_m (y)_m}.$$

Since the left member

$$= \sum_{m,n=0}^{\infty} \sum_{r=1}^{\infty} (-1)^{r-1} (x^m y^n t)^r = \sum_{r=1}^{\infty} (-1)^{r-1} \frac{t^r}{(1 - x^r)(1 - y^r)},$$

Received by the editors August 3, 1955.

we get

$$(2.1) \quad mG_m = \sum_{r=1}^m (-1)^{r-1} \frac{(x)_m (y)_m}{(x)_{m-r} (y)_{m-r}} \frac{G_{m-r}}{(1-x^r)(1-y^r)}.$$

In exactly the same way logarithmic differentiation of (1.6) leads to

$$(2.2) \quad mH_m = \sum_{r=1}^m \frac{(x)_m (y)_m}{(x)_{m-r} (y)_{m-r}} \frac{H_{m-r}}{(1-x^r)(1-y^r)}.$$

It follows at once from (2.1) and (2.2) that G_m and H_m are polynomials in x and y . We may easily compute the first few values using the recursion formulas:

$$\begin{aligned} G_0 &= 1, & G_1 &= 1, & G_2 &= x + y, \\ G_3 &= xy + x^3 + x^2y + xy^2 + y^3 + x^2y^2, \\ H_0 &= 1, & H_1 &= 1, & H_2 &= 1 + xy, \\ H_3 &= 1 + xy + xy^2 + x^2y + x^2y^2 + x^3y^3. \end{aligned}$$

Next if $f(x)$ denotes the left member of (1.5), it is evident that

$$(2.3) \quad f(x) = f(xt) \prod_{n=0}^{\infty} (1 + y^n t).$$

It follows from (2.3), (1.1), and (1.5) that

$$\sum_0^{\infty} \frac{t^m G_m}{(x)_m (y)_m} = \sum_0^{\infty} \frac{x^m t^m G_m}{(x)_m (y)_m} \sum_0^{\infty} \frac{y^{n(n-1)/2} t^n}{(y)_n}.$$

Equating coefficients we get

$$G_m = \sum_{r=0}^m x^{m-r} y^{r(r-1)/2} \frac{(x)_m}{(x)_{m-r}} \frac{(y)_m}{(y)_r (y)_{m-r}} G_{m-r},$$

which yields

$$(2.4) \quad G_m = \sum_{r=1}^m x^{m-r} y^{r(r-1)/2} \frac{(x)_{m-1}}{(x)_{m-r}} \left[\begin{matrix} m \\ r \end{matrix} \right]_y G_{m-r},$$

where

$$(2.5) \quad \left[\begin{matrix} m \\ r \end{matrix} \right]_y = \frac{(y)_m}{(y)_r (y)_{m-r}}.$$

The coefficients (2.5) occur in the identity

$$(2.6) \quad \prod_{r=0}^{m-1} (1 + y^r t) = \sum_{r=0}^m y^{r(r-1)/2} \left[\begin{matrix} m \\ r \end{matrix} \right]_y t^r.$$

Interchanging x and y in (2.4) we have also

$$(2.7) \quad G_m = \sum_{r=1}^m x^{r(r-1)/2} y^{m-r} \begin{bmatrix} m \\ r \end{bmatrix}_x \frac{(y)_{m-1}}{(y)_{m-r}} G_{m-r}.$$

To get similar recursions for H_m , let $g(t)$ denote the left member of (1.6); then

$$g(t) = g(xt) \prod_0^{\infty} (1 - y^n t)^{-1}.$$

Using (1.2) we get

$$(2.8) \quad H_m = \sum_{r=1}^m x^{m-r} \frac{(x)_{m-1}}{(x)_{m-r}} \begin{bmatrix} m \\ r \end{bmatrix}_y H_{m-r}$$

and on interchanging x and y

$$(2.9) \quad H_m = \sum_{r=1}^m y^{m-r} \begin{bmatrix} m \\ r \end{bmatrix}_y \frac{(y)_{m-1}}{(y)_{m-1}} H_{m-r}.$$

Since by (2.6),

$$\begin{bmatrix} m \\ r \end{bmatrix}_y$$

is a polynomial in y with integral coefficients, it follows from (2.4) and (2.8) that G_m and H_m are polynomials in x and y with integral coefficients; this last fact is not evident from (2.1) and (2.2).

3. It is clear from (1.5) and (1.6) that

$$(3.1) \quad \begin{aligned} G_m(x, y) &= G_m(y, x), \\ H_m(x, y) &= H_m(y, x). \end{aligned}$$

We now prove that

$$(3.2) \quad G_m(x, y) = (xy)^{m(m-1)/2} G_m\left(\frac{1}{x}, \frac{1}{y}\right),$$

$$(3.3) \quad H_m(x, y) = (xy)^{m(m-1)/2} H_m\left(\frac{1}{x}, \frac{1}{y}\right)$$

and

$$(3.4) \quad H_m(x, y) = y^{m(m-1)/2} G_m\left(x, \frac{1}{y}\right).$$

We prove (3.2) by induction. In the right member of (2.1) replace

x, y by $1/x, 1/y$; we get using the inductive hypothesis

$$\sum_{r=1}^m (-1)^{r-1} \frac{(x)_m}{(x)_{m-r}} \frac{(y)_m}{(y)_{m-r}} \frac{G_{m-r}(x, y)}{(1-x)(1-y^r)} (xy)^{-k},$$

where

$$\begin{aligned} k &= m(m+1)/2 - (m-r)(m-r+1)/2 - (m-r)(m-r-1)/2 - r \\ &= m(m+1)/2 - (m-r) - r = m(m-1)/2. \end{aligned}$$

This evidently completes the induction. In the same way we use (2.2) to prove (3.3). To prove (3.4) we replace y by $1/y$ in the right member of (2.1). Then by the inductive hypothesis we get

$$\sum_{r=1}^m \frac{(x)_m (y)_m}{(x)_{m-r} (y)_{m-r}} \frac{H_{m-r}(x, y)}{(1-x^r)(1-y^r)} y^{-m(m-1)/2} = m y^{-m(m-1)/2} H_m(x, y),$$

by (2.2). This completes the proof of (3.4).

If we put

$$(3.5) \quad \prod_{m, n=0}^{\infty} \frac{1 + x^m y^n t}{1 - x^m y^n t} = \sum_{m=0}^{\infty} K_m(x, y) t^m / (x)_m (y)_m,$$

then by (1.5), (1.6) and (2.5)

$$(3.6) \quad K_m = \sum_{r=0}^m \begin{bmatrix} m \\ r \end{bmatrix}_x \begin{bmatrix} m \\ r \end{bmatrix}_y G_r H_{m-r}.$$

Thus K_m is a polynomial in x, y with integral coefficients. Moreover K_m has the following symmetry properties

$$(3.7) \quad K_m(x, y) = K_m(y, x),$$

$$(3.8) \quad K_m(x, y) = y^{m(m-1)/2} K_m\left(x, \frac{1}{y}\right);$$

the latter is an immediate consequence of (3.4) and (3.6). Incidentally (3.5) implies the quadratic relation

$$(3.9) \quad \sum_{r=0}^m (-1)^r \begin{bmatrix} m \\ r \end{bmatrix}_x \begin{bmatrix} m \\ r \end{bmatrix}_y K_r K_{m-r} = 0 \quad (m > 1).$$

Logarithmic differentiation of the left member of (3.5) gives

$$\begin{aligned} \sum_{m, n=0}^{\infty} \left(\frac{x^m y^n}{1 + x^m y^n t} + \frac{x^m y^n}{1 - x^m y^n t} \right) &= 2 \sum_{m, n=0}^{\infty} \sum_{r=0}^{\infty} (x^m y^n)^{2r+1} t^{2r} \\ &= 2 \sum_{r=0}^{\infty} \frac{t^{2r}}{(1 - x^{2r+1})(1 - y^{2r+1})}. \end{aligned}$$

We accordingly get the recursion

$$(3.10) \quad mK_m = 2 \sum_{0 \leq 2r < m} \frac{(x)_m (y)_m}{(x)_{m-2r-1} (y)_{m-2r-1}} \frac{K_{m-2r-1}}{(1-x^{2r+1})(1-y^{2r+1})}.$$

It evidently follows from (3.10) that for m even, K_m is divisible by $(1+x^{m/2})(1+y^{m/2})$; indeed if $m=2^e k$, then K_m is divisible by

$$\frac{1-x^m}{1-x^k} \frac{1-y^m}{1-y^k}.$$

By means of (3.10) we may compute the first few values of K_m :

$$\begin{aligned} K_0 &= 1, & K_1 &= 2, & K_2 &= 2(1+x)(1+y), \\ K_3 &= 2(1+x)(1+y)(1+2xy+x^2+y^2+x^2y^2), \\ K_4 &= (1+x)(1+x^2)(1+y)(1+y^2)\{(1-x)(1-x^2)(1-y)(1-y^2) \\ &\quad + 1+2xy+x^2+y^2+x^2y^2\}. \end{aligned}$$

Expanding K_4 we find that some of the coefficients are negative. Since the coefficients of

$$\begin{bmatrix} m \\ r \end{bmatrix}_x$$

are non-negative it therefore follows from (3.6) that some of the coefficients of G_4 and H_4 are negative.

4. In (1.6) take $y=0$. Comparison with (1.2) gives

$$(4.1) \quad H_m(x, 0) = 1;$$

similarly from (1.1) we get

$$(4.2) \quad G_m(x, 0) = x^{m(m-1)/2}.$$

In view of (3.2) and (3.3) it follows that G_m and H_m are of degree $m(m-1)/2$ in each of x and y . As for K_m , (3.6) implies

$$K_m(x, 0) = \sum_{r=0}^m \begin{bmatrix} m \\ r \end{bmatrix}_x x^{m(m-1)/2} = \prod_0^{m-1} (1+x^r)$$

by (2.6), and therefore $K_m(x, y)$ is also of degree $m(m-1)/2$ in each of x and y .

In the next place if we take $y=1$ in (2.1) we get

$$G_m(x, 1) = \frac{1-x^m}{1-x} G_{m-1}(x, 1),$$

which yields

$$(4.3) \quad G_m(x, 1) = (x)_m / (1 - x)^m.$$

If we put $y = \zeta$, a primitive r th root of unity, where $r \mid m$, (2.1) implies

$$mG_m(x, \zeta) = (-1)^{r-1} C_m \frac{(x)_m}{(1 - x^r)(x)_{m-r}} G_{m-r}(x, \zeta),$$

where

$$C_m = \left\{ \frac{(y)_m}{(1 - y^r)(y)_{m-r}} \right\}_{y=\zeta} = m.$$

It follows that

$$(4.4) \quad G_m(x, \zeta) = (-1)^{m-m/r} \frac{(x)_m}{(1 - x^r)^{m/r}},$$

which reduces to (4.3) for $r = 1$. The corresponding formula for H_m is

$$(4.5) \quad H_m(x, \zeta) = (x)_m / (1 - x^r)^{m/r}.$$

For K_m it follows from (3.10) that

$$(4.6) \quad K_m(x, \zeta) = 0 \quad (m \geq 2, r \text{ even}),$$

while

$$(4.7) \quad K_m(x, \zeta) = 2^{m/r} \frac{(x)_m}{(1 - x^r)^{m/r}} \quad (r \text{ odd}).$$

We also note that when m is odd, (3.10) implies

$$K_m(x, -1) = \frac{1 - x^m}{1 - x} K_{m-1}(x, -1) = 0$$

by (4.6). Thus $K_m(x, y)$ is divisible by $(1+x)(1+y)$ for all $m \geq 2$.

The following consequence of (4.3) and (4.5) may be noted:

$$(4.8) \quad G_m(1, 1) = m! = H_m(1, 1).$$

Since, as we have seen, $G_m(x, y)$ is of degree $m(m-1)/2$ in each of x and y , the number of terms is $O(m^4)$. Therefore it follows from (4.8) that at least one coefficient of G_m is unbounded; the same is also true of H_m and K_m . Indeed by (4.7) we have

$$(4.9) \quad K_m(1, 1) = 2^m \cdot m!.$$

If ζ is a primitive r th root of unity and η a primitive s th root of unity, $r \mid m$, $s \mid m$, then (4.4) implies the following curious property.

$$(4.10) \quad G_m(\eta, \zeta) = \begin{cases} 0 & (r \neq s), \\ (-1)^{m-m/r} r^{m/r} (m/r)! & (r = s). \end{cases}$$

Like results hold for H_m and K_m .

5. If $x=y$, (1.5) and (1.6) become

$$(5.1) \quad \prod_0^\infty (1 + x^m t)^{m+1} = \sum_0^\infty t^m G_m(x)/(x)_m^2,$$

$$(5.2) \quad \prod_0^\infty (1 - x^m t)^{-m-1} = \sum_0^\infty t^m H_m(x)/(x)_m^2,$$

where $G_m(x) = G_m(x, x)$, $H_m(x) = H_m(x, x)$. Thus (2.1) and (2.2) reduce to

$$(5.3) \quad mG_m(x) = \sum_{r=1}^m (-1)^{r-1} \left\{ \frac{(x)_m}{(1-x^r)(x)_{m-r}} \right\}^2 G_{m-r}(x),$$

$$(5.4) \quad mH_m(x) = \sum_{r=1}^m \left\{ \frac{(x)_m}{(1-x^r)(x)_{m-r}} \right\}^2 H_{m-r}(x),$$

and similarly for (2.4) and (2.8). Also (3.2) and (3.3) imply

$$(5.5) \quad G_m(x) = x^{m(m-1)} G_m\left(\frac{1}{x}\right), \quad H_m(x) = x^{m(m-1)} H_m\left(\frac{1}{x}\right).$$

Since $H_m(0)=1$, it follows that $H_m(x)$ is of degree $m(m-1)$. On the other hand using (2.4) we can show that $G_m(x)$ is divisible by exactly x^{m-1} which implies that $G_m(x)$ is of degree $(m-1)^2$ for $m \geq 1$. We also note that (4.8) gives

$$(5.6) \quad G_m(1) = m! = H_m,$$

so that at least one coefficient of $G_m(x)$ and of $H_m(x)$ is unbounded.

In conclusion we remark that the left member of (5.2) may be compared with the generating function for unrestricted plane partitions [2; 3, p. 243]:

$$(5.7) \quad \prod_1^\infty (1 - x^m)^{-m} = \sum_0^\infty P(m) x^m.$$

REFERENCES

1. R. Bellman, Bull. Amer. Math. Soc. Research problem 61-1-5.
2. T. W. Chaundy, *The unrestricted plane partition*, Quart. J. Math. Oxford Ser. vol. 3 (1932) pp. 76-80.
3. P. A. MacMahon, *Combinatory analysis*, vol. 2, Cambridge, 1916.

DUKE UNIVERSITY